

Production of Prostaglandin E₂ and I₂ Is Coupled with Cyclooxygenase-2 in Human Follicular Dendritic Cells

Whajung Cho¹, Jini Kim¹, Kyu-Bong Cho² and Jongseon Choe^{1*}

¹Department of Microbiology and Immunology, School of Medicine, Kangwon National University, Chuncheon 200-701,

²Department of Clinical Laboratory Science, Shinheung College, Uijeongbu 480-701, Korea

Background: Prostaglandins (PGs) play pathogenic and protective roles in inflammatory diseases. The novel concept of PGs as immune modulators is being documented by several investigators. By establishing an *in vitro* experimental model containing human follicular dendritic cell-like cells, HK cells, we reported that HK cells produce prostaglandin E₂ (PGE₂) and prostaglandin I₂ (PGI₂) and that these PGs regulate biological functions of T and B cells. **Methods:** To investigate the respective contribution of cyclooxygenase-1 (COX-1) and COX-2 to PGE₂ and PGI₂ production in HK cells, we performed siRNA technology to knock down COX enzymes and examined the effect on PG production. **Results:** Both PGE₂ and PGI₂ productions were almost completely inhibited by the depletion of COX-2. In contrast, COX-1 knockdown did not significantly affect PG production induced by lipopolysaccharide (LPS). **Conclusion:** The current results suggest that mPGES-1 and PGIS are coupled with COX-2 but not with COX-1 in human follicular dendritic cell (FDC) and may help understand the potential effects of selective COX inhibitors on the humoral immunity.

[Immune Network 2011;11(6):364-367]

INTRODUCTION

Prostanoids are small lipid molecules, playing important roles in various physiological and pathological processes, such as kidney function, vasodilation, platelet aggregation, and cancer progress (1). Prostanoids are derived from membrane glycerophospholipids in response to inflammatory stimuli via three sequential enzymatic reactions. The first reaction is mediated

by phospholipase A₂, which liberates arachidonic acids from membrane phospholipids (2). The arachidonic acids are subject to the second reactions and converted to prostaglandin (PG) G₂ and then PGH₂ by cyclooxygenases (COXs), COX-1 or COX-2. The last steps of prostanoid production depend on tissue- and cell-specific terminal synthases. PGH₂ are metabolized to PGE₂, PGI₂, PGD₂, PGF_{2α}, and thromboxane A₂ (TXA₂) by respective actions of PGE₂ synthase (PGES), PGI₂ synthase (PGIS), PGD₂ synthase (PGDS), PGF₂ synthase (PGFS), and thromboxane synthase (3). In particular, PGE₂ is catalyzed by three different PGESs, cytosolic PGES (cPGES) and two membrane-bound PGESs, mPGES-1 and mPGES-2 (4). The expression of cPGES and mPGES-2 are constitutive, while mPGES-1 is induced during inflammation.

In addition to the well-known roles of prostanoids as inflammatory and vascular mediators (5,6), they are recognized as important immune modulators (7-9). For instance, PGE₂ exhibits immunostimulatory as well as immunosuppressive activities depending on its concentrations. PGE₂ inhibits activation, proliferation, and differentiation of T cells at micromolar concentrations, whereas it potentiates Th1 and Th17 differentiation and proliferation of T cells at nanomolar concentrations (10).

Using an *in vitro* model of germinal center reactions that contains human follicular dendritic cell (FDC)-like cells HK (11), we have demonstrated that both FDC and HK cells express PGI₂ synthase (12), PGI₂ production is not controlled by the induction of PGIS but by COX-2 (13), HK cells secrete PGI₂ and PGE₂ but not TXA₂ (14), PGs produced by HK cells

Received on October 5, 2011. Revised on October 21, 2011. Accepted on October 25, 2011.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Corresponding Author. Tel: 82-33-250-8862; Fax: 82-33-244-3286; E-mail: jchoe@kangwon.ac.kr

Keywords: Human, Stromal cells, Lipid mediator

inhibit proliferation and apoptosis T cells (14), T cells control PG production from HK cells via IL-4-Janus kinase 1 (JAK1)-Signal transducer and activator of transcription 6 (STAT6)-COX-2 pathway (15), and PGI₂ and its analogues enhance CD86 expression on the surface of activated B cells (16). These results support the emerging concept of PGs as critical immune modulators.

In this study, we investigated the relative contribution of COX-1 and COX-2 to PGI₂ and PGE₂ synthesis in HK cells. Several studies demonstrated the coupling between COXs and terminal prostanoid synthases. However, most studies were performed using murine cells (2). The current results suggest that mPGES-1 and PGIS are coupled with COX-2 but not with COX-1 in human FDC and imply that chronic administration of selective COX-2 inhibitors might disturb the normal humoral immune responses taking place in the culminating site of germinal centers.

MATERIALS AND METHODS

Culture of HK cells

HK cells are primary cells obtained from human tonsils and used until they display degenerate features in culture. They are prepared as described by Kim et al. (17) and maintained in RPMI-1640 (Irvine Scientific, Santa Ana, CA) containing 10% fetal calf serum (Hyclone, Logan, UT), 2 mM L-glutamine (Invitrogen, Carlsbad, CA), 100 U/ml penicillin G (Sigma-Aldrich, St. Louis, MO), and 100 µg/ml streptomycin (Invitrogen). Lipopolysaccharide (LPS) was purchased from Sigma-Aldrich.

Immunoblotting

The whole cell lysates of HK cells were subject to immunoblotting as previously described (18). The protein concentrations of the each fraction were assayed with a bicinchoninic acid (BCA) assay. Used antibodies were against COX-1, COX-2 (Cayman Chemical, Ann Arbor, MI), β-actin (Sigma-Aldrich), and horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Jackson ImmunoResearch, West Grove, PA). The membranes were incubated with SuperSignal West Pico Chemiluminescent Substrate (Pierce, Rockford, IL) and exposed to X-ray films.

siRNA transfection

The siRNA duplexes used (Ambion Inc, Austin, TX) were con-

structed with the following target sequences. Control (Neg-siRNA#2, sequence not disclosed by Ambion); COX-1, sense (5'-GCUCUUUAAGGAUGGGAAATT-3'), antisense (5'-UUCCCAUCCUAAAAGAGCCG-3'); COX-2, sense (5'-CCA CC CAUGUCAAAACCGATT-3'), antisense (5'-UCG GUUUUGA CAUGGGUGGGA-3'). HK cells were cultured to 50~60% confluence in 100 mm plates. For each plate, 40 nM of each siRNA and 24 µl Lipofectamine™ (Invitrogen) were separately diluted in 400 µl serum-free medium without antibiotics, mixed together, and incubated at RT for 45 min. The plates were then washed with serum-free medium, added with 5 ml serum-free medium, and then with the diluted solutions. The plates were incubated at 37°C for 8 h, followed by the addition of a growth medium containing 10% serum. After 48 h of additional incubation, cells were used for experiments. The degree of gene-silencing was assayed by immunoblotting.

Enzyme immunoassay to measure prostaglandins

HK cells were cultured with LPS for 48 h to harvest the supernatants. The amounts of PGE₂ and 6-keto-PGF_{1α}, stable metabolite of PGI₂, were measured using enzyme immunoassay (EIA) kits as described previously (14). PG concentration was normalized to total cellular protein and expressed as ng/mg protein.

Statistical analysis

Statistical analysis and graphic presentation were carried out with GraphPad Prism 4.0. The statistical significance of differences was determined by Student's *t*-test; *p*<0.05 was considered significant.

RESULTS

To investigate the relative contribution of COX-1 and COX-2 to the production of PGE₂ and PGI₂, we carried out siRNA technology to knock down COX-1 and COX-2 proteins in HK cells. HK cells were transfected with siRNA duplexes specific to COX-1 and COX-2 and cultured for 48 h, followed by further cultures in the presence or absence of LPS. The silencing of target proteins was demonstrated by immunoblotting. As shown in Fig. 1, COX-2 protein levels were up-regulated by LPS stimulation in control siRNA-transfected HK cells, whereas COX-1 levels were unaffected by LPS treatment. Transfection with COX-1-specific siRNA resulted in significant reduction of COX-1 protein levels regardless of LPS stimulation. Interestingly, LPS-induced COX-2 levels in COX-1 siRNA-trans-




Figure 1. COX-1 and COX-2 proteins were markedly depleted in HK cells by siRNA technology. Control, COX-1, or COX-2 siRNA-transfected HK cells (1×10^5 cells) were cultured for 48 h, followed by further incubation in the presence or absence of LPS ($1 \mu\text{g/ml}$) for 8 h. The depletion degree of target proteins was evaluated by immunoblotting. β -actin was used to demonstrate equal loading of lysates. The results were reproducible in three independent experiments.




Figure 2. Depletion of COX-2 but not COX-1 prevents LPS-induced production of PGE₂ and PGI₂. Control, COX-1, or COX-2 siRNA-transfected HK cells (1×10^5 cells) were cultured for 48 h, followed by further incubation in the presence or absence of LPS ($1 \mu\text{g/ml}$) for another 48 h. The amounts of PGE₂ (A) and 6-keto-PGF_{1 α} (B) in the culture supernatants were measured by EIA and normalized by protein concentration in each well. Data are presented as means \pm standard error of the mean (SEM) of duplicates. Representative results of three reproducible experiments. Asterisks indicate significant differences (* $p < 0.05$, ** $p < 0.01$). NS, non-significant.

ected cells were markedly higher compared to control cells. Transfection with COX-2-specific siRNA almost completely prevented induction of COX-2 proteins that was triggered by LPS. COX-2 silencing did not significantly affect COX-1 expression levels. These results indicate that COX-1 and COX-2 proteins were successfully knocked down in HK cells by siRNA duplexes. We next measured the concentrations of PGE₂ and 6-keto-PGF_{1 α} in the culture supernatants after incubation of siRNA-transfected cells in the presence or absence of LPS for 48 h. 6-keto-PGF_{1 α} is the hydrolysis product of unstable PGI₂. PGE₂ and 6-keto-PGF_{1 α} concentrations in control cells were increased 4- and 5-folds, respectively, by LPS stimulation (Fig. 2). Similar levels of enhancement were obtained in COX-1 siRNA-transfected cells. In contrast, COX-2 silencing almost completely prevented the PG production to background levels. We observed that knock down of COX-1 resulted in a significant increase of PGE₂ but not PGI₂ without LPS stimulation (Fig. 2A). Whether this result reflects a preferential coupling of COX-2 with PGES over PGIS is currently unclear. Based upon these results, we conclude that PGE₂ and PGI₂ production is coupled with COX-2 but not with COX-1 in HK cells.

DISCUSSION

In this study, we demonstrate that LPS stimulates FDC-like cells to produce PGE₂ and PGI₂, which depends on the presence of COX-2. Our results imply that FDC is a physiologic source of PG during inflammatory immune responses to bacterial infection. PGs secreted from FDC may exert regulatory roles in the various processes taking place in germinal centers of the secondary lymphoid tissues (7-9).

PGE₂ and PGI₂ are biosynthesized by mPGES-1 and PGIS. In line with the current data, we previously provided evidence that COX-2 and mPGES-1 are functionally associated. COX-2 and mPGES-1 mRNA and proteins were increased by LPS treatment, which was inhibited by IL-4 (14,15). mPGES-1 induction by LPS stimulation occurred following COX-2 induction in HK cells (14). This coordinated kinetics of COX-2 and mPGES-1 induction may contribute to the efficient production of PGE₂. In contrast, PGIS is constitutive in HK cells (13,14). Therefore, the PGI₂ production in HK cells may depend on the substrate supply for downstream PGIS from upstream COX-2, suggesting the coupling of COX-2 with PGIS. This interpretation is compatible with the general belief that inducible COX-2 is required for delayed PG synthesis whereas

constitutive COX-1 is necessary for immediate PG production after inflammatory stimulation (2). Further investigation is necessary to determine whether the functional association of COX-2 with mPGES-1 and PGIS results from physical coupling of these enzymes. Confocal microscopy may be a useful method to study their possible colocalization and subcellular localization. However, the coupling between COX-2 and PGIS is not absolute and appears to vary depending on cell types. Different from HK cells, Bolego et al, reported that PGI₂ production is coupled with COX-1 but not with COX-2 in human endothelial cells (19). PGIS is constitutively expressed in human endothelial cells (20). Furthermore, stimulus types and the degree of substrate abundance seem to affect the functional coupling between COX and PGIS or mPGES-1.

Considering the accumulating reports on the side effects of selective COX inhibitors (21), the current data may help understand the potential effects of selective COX inhibitors on the humoral immunity.

ACKNOWLEDGEMENTS

This work was supported by the Grants of the Korean Ministry of Education, Science and Technology (The Regional Core Research Program/Medical & Bio-Materials Research Center and the Basic Research Promotion Fund No. 2009-0068105).

CONFLICTS OF INTEREST

The authors have no financial conflict of interest.

REFERENCES

1. Fosslien E: Cardiovascular complications of non-steroidal anti-inflammatory drugs. *Ann Clin Lab Sci* 35;347-385, 2005.
2. Ueno N, Takegoshi Y, Kamei D, Kudo I, Murakami M: Coupling between cyclooxygenases and terminal prostanoid synthases. *Biochem Biophys Res Commun* 338;70-76, 2005.
3. Nakashima K, Ueno N, Kamei D, Tanioka T, Nakatani Y, Murakami M, Kudo I: Coupling between cyclooxygenases and prostaglandin F(2alpha) synthase. Detection of an inducible, glutathione-activated, membrane-bound prostaglandin F(2alpha)-synthetase activity. *Biochim Biophys Acta* 1633; 96-105, 2003.
4. Chandrasekharan S, Foley NA, Jania L, Clark P, Audoly LP, Koller BH: Coupling of COX-1 to mPGES1 for prostaglandin E2 biosynthesis in the murine mammary gland. *J Lipid Res* 46;2636-2648, 2005.
5. Cheng Y, Wang M, Yu Y, Lawson J, Funk CD, Fitzgerald GA: Cyclooxygenases, microsomal prostaglandin E synthase-1, and cardiovascular function. *J Clin Invest* 116;1391-1399, 2006.
6. Vane J, Corin RE: Prostacyclin: a vascular mediator. *Eur J Vasc Endovasc Surg* 26;571-578, 2003.
7. Harizi H, Gualde N: The impact of eicosanoids on the cross-talk between innate and adaptive immunity: the key roles of dendritic cells. *Tissue Antigens* 65;507-514, 2005.
8. Harris SG, Padilla J, Koumas L, Ray D, Phipps RP: Prostaglandins as modulators of immunity. *Trends Immunol* 23; 144-150, 2002.
9. Rocca B, FitzGerald GA: Cyclooxygenases and prostaglandins: shaping up the immune response. *Int Immunopharmacol* 2;603-630, 2002.
10. Sreeramkumar V, Fresno M, Cuesta N: Prostaglandin E(2) and T cells: friends or foes? *Immunol Cell Biol* 2011. [Epub ahead of print]
11. Lee IY, Lee JH, Park WS, Nam EC, Shin YO, Choe JS: 3C8, a new monoclonal antibody directed against a follicular dendritic cell line, HK. *Immune Netw* 1;26-31, 2001.
12. Lee IY, Ko EM, Kim SH, Jeoung DI, Choe J: Human follicular dendritic cells express prostacyclin synthase: a novel mechanism to control T cell numbers in the germinal center. *J Immunol* 175;1658-1664, 2005.
13. Lee IY, Bae YD, Jeoung DI, Kang D, Park CH, Kim SH, Choe J: Prostacyclin production is not controlled by prostacyclin synthase but by cyclooxygenase-2 in a human follicular dendritic cell line, HK. *Mol Immunol* 44;3168-3172, 2007.
14. Lee IY, Cho W, Kim J, Park CS, Choe J: Human follicular dendritic cells interact with T cells via expression and regulation of cyclooxygenases and prostaglandin E and I synthases. *J Immunol* 180;1390-1397, 2008.
15. Cho W, Kim Y, Jeoung DI, Kim YM, Choe J: IL-4 and IL-13 suppress prostaglandins production in human follicular dendritic cells by repressing COX-2 and mPGES-1 expression through JAK1 and STAT6. *Mol Immunol* 48;966-972, 2011.
16. Kim J, Park CS, Park CH, Jeoung DI, Kim YM, Choe J: Beraprost enhances the APC function of B cells by upregulating CD86 expression levels. *J Immunol* 186;3866-3873, 2011.
17. Kim HS, Zhang X, Choi YS: Activation and proliferation of follicular dendritic cell-like cells by activated T lymphocytes. *J Immunol* 153;2951-2961, 1994.
18. Lee IY, Ha KS, Choe J: 3C8 antigen is a novel protein expressed by human follicular dendritic cells. *Biochem Biophys Res Commun* 303;624-630, 2003.
19. Bolego C, Buccellati C, Prada A, Gaion RM, Folco G, Sala A: Critical role of COX-1 in prostacyclin production by human endothelial cells under modification of hydroperoxide tone. *FASEB J* 23;605-612, 2009.
20. Spisni E, Bartolini G, Orlandi M, Belletti B, Santi S, Tomasi V: Prostacyclin (PGI₂) synthase is a constitutively expressed enzyme in human endothelial cells. *Exp Cell Res* 219;507-513, 1995.
21. Fosbol EL, Kober L, Torp-Pedersen C, Gislason GH: Cardiovascular safety of non-steroidal anti-inflammatory drugs among healthy individuals. *Expert Opin Drug Saf* 9;893-903, 2010.