

Prevalence of Cardiovascular Disorders Among Iranian Elite Athletes

Shahin Salehi,¹ Farhad Moradi Shahpar,^{2,3,*} Gholamreza Norouzi,³ Farshad Ghazalian,⁴ Mehrshad Poursaid Esfehiani,⁵ and Amir Hosein Abedi Yekta⁵

¹Imam Hossein Medical and Educational Center, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

²Physical Education Department, University of Isfahan, Isfahan, IR Iran

³Iran Sports Medicine Federation, Tehran, IR Iran

⁴Department of Physical Education, Science and Research Branch, Islamic Azad University, Tehran, IR Iran

⁵Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

*Corresponding author: Farhad Moradi Shahpar, Physical Education Department, University of Isfahan, Isfahan, IR Iran. E-mail: drfmoradi@yahoo.com

Received 2015 December 27; Revised 2016 February 07; Accepted 2016 March 06.

Abstract

Background: Athletes' health is an important issue and for promoting it, pre-participation examination (PPE) is widely performed by responsible bodies around the world.

Objectives: This study was to determine prevalence of cardiovascular disorders among athletes participating in the Asian games and answering the question whether the electrocardiogram (ECG) is a necessary part of pre-participation examination (PPE) for prevention of sudden cardiac death.

Materials and Methods: All athletes participated at Asian games came to sports medicine federation for a PPE including a comprehensive questionnaire, physical examination and ECG. In this retrospective study all profiles of 338 athletes have been studied as well as their electrocardiograms. Multiple logistic regressions as well as Firth's bias reduction were used with R statistical software and SPSS. For predicting the changes in ECG, receiver operating characteristic (ROC) curve has done.

Results: Among 388 athletes, 80 (20.6%) were female and 308 (79.4%) male with mean age of 23.2 + 8 years. Nine athletes (2.3%) were smokers, 28 of them (7.2%) experienced chest pain and discomfort, 45 of them (13.3%) had palpitations and 28 (7.2%) had history of anemia. Study of their electrocardiograms showed that long Q-T interval was not seen for anyone, but evidence of left ventricular hypertrophy was seen in 12 (3.1%), inverted T wave in 6 (1.5%), and right bundle branch block in 45 (13.3%).

Conclusions: PPE provides very important information of athletes' health. This study has shown that there was not any significant relation between current examination and electrocardiogram changes but regarding the ECG changes we recommend it as a routine part of PPE.

Keywords: Sudden Cardiac Death, Prevention, Electrocardiogram

1. Background

Athletes' health is an important issue and for promoting it, pre-participation examination (PPE) is widely performed by responsible bodies around the world. While the sudden cardiac death (SCD) incidence in sports is rare (1:50,000 to 1:200,000 annually) (1, 2), it must be considered as an important cause of death among athletes, so early diagnosis of any fatal disorder could prevent sudden death. Sudden cardiac death has been defined by world health organization (WHO) as a fatal condition that happens within 24h after onset of symptoms (3). At the 26th Bethesda conference "athletes' SCD" was defined as: "non-traumatic, nonviolent, unexpected death due to cardiac causes within one hour of the onset of symptoms. Sports-related SCD are defined as those with symptoms occurring within one hour of sports participation" (4).

Most of the deaths reported in sports are related to cardiovascular conditions (5-11) and hypertrophic cardiomy-

opathy is the most common cause among them. The other fatal cardiovascular conditions which result in SCD include major Vascular anomalies, valvular disorders, coronary ischemic heart disease, Wolf Parkinson white syndrome, arrhythmogenic cardiomyopathy, long Q-T interval, aortic aneurysm, cardiac conduction disorders, myocarditis, and the other congenital heart diseases (12). That is obviously clear that most of SCD causes could be diagnosed by electrocardiogram (ECG), and it should be considered that under the age of 35, genetic and congenital disorders are more common and for people above 35 years old, coronary artery disease is the most common causes of SCD (13). The risk of SCD increases by rising of age and exercise intensity (11, 12). The other non-cardiac causes of sudden death among athletes are head, chest and abdominal traumas (9-11, 14). It has estimated that sudden death cases are 2 times more common among athletes comparing to non-athlete people (3). Therefore it could be concluded that

early diagnosis of cardiac disorders has an important preventive role in sudden death. On the other hand many of the above mentioned causes are hereditary, so in case of any SCD the other family members should be examined for screening (15). It is a critical issue to find a reliable and specific approach to diagnose people exposed to SCD (4). Many countries have their own attitude to PPE for detecting possible causes of sudden death such as Spain (16), USA (17-24), Sweden (25), Denmark (26, 27), Norway (28), Italy (26, 29, 30), Switzerland (31), Germany (32), France (33), Netherlands (34), China (35), Japan (36, 37), and UAE (38). These examinations are also routinely performed for all athletes participating at major sports events like Asian games and Olympic games by Iran's Sports medicine federation. The examination consists of a comprehensive questionnaire, full physical examination (by cardiologist as well as orthopedic surgeon, sports medicine specialist, optometrist and internist), a 12 lead 2-dimensional electrocardiogram (ECG), routine blood and urine laboratory examinations (such as blood cell count, hemoglobin, serum iron, fasting blood sugar, lipid profile, liver function tests, blood urea nitrogen, creatinine, and urine analysis). In recent years an echocardiography has been added to these examinations. Supplementary exams and para-clinicals such as radiography and magnetic resonance imaging (MRI) are also done upon request of physicians.

2. Objectives

Our aim was to investigate the prevalence of cardiovascular abnormalities in history, examination and electrocardiography of Iranian elite athletes participating in Asian games.

3. Materials and Methods

In this retrospective study, all profiles of athletes participating at Asian games who referred to Iran sports medicine federation during 2012 to 2013 were reviewed. Most of the athletes were at the peak of exercise and physical fitness. All athletes were examined one by one by expert general practitioners and specialists, the setting of examinations was based on station-type which means multiple examiners conducted discrete portions of PPE in sequence. We had two stations for heart examination. In the first station, medical history was taken and also cardiovascular system was examined entirely by a cardiologist. In the second station ECG and blood pressure were recorded by an expert. The 6-channel ECG device (brand name: CONTEC®, Model Number: ECG600G) as well as Richter® mercury sphygmomanometer were used. Then all data were

reviewed and evaluated by a sports medicine specialist. All profiles with missing data were omitted. All data related to cardiovascular system including past history, familial history, and physical examination as well as ECG interpretation were extracted and analyzed again. Sokolov-Lyon criteria was used for the detection of cardiac hypertrophy. Electrocardiographic criteria for right atrial hypertrophy was a peaked P wave (P pulmonale) with amplitude bigger than 2.5 mm in the inferior leads (II, III and aVF) or bigger than 1.5 mm in V₁ and V₂. Diagnostic criteria for right atrial hypertrophy was bifid P wave with more than 40 milliseconds between the two peaks or total P wave duration more than 110 milliseconds in lead II or biphasic P wave with terminal negative portion bigger than 40 milliseconds duration or biphasic P wave with terminal negative portion more than 1mm deep in V₁. For long QT syndrome we used from QTc) Corrected QT). QTc is prolonged if it was more than 440 milliseconds in men or more than 460 milliseconds in women. Inverted T was important for us in inferior leads (II, III, aVF), lateral leads (I, aVL, V5-6) and anterior leads (V2-6). Standard for left axis deviation was QRS axis less than -30 degrees and for right axis deviation, QRS axis greater than +90 degrees.

To study single variable impact, IBM SPSS Statistics® software version 21 and multiple logistics regression of R® statistical software were used at the same time. In order to determine any relation between clinical signs and symptoms and ECG changes, Chi-squared (X^2) test as well as Fisher exact test were performed. First bias reductions also performed because despite the likelihood function converges, some of the parameters are not convergent. In order to predict ECG changes by taking history and physical examination, receiver operating characteristic (ROC) curve was performed. Level of significance was set as 0.05.

This study was approved by ethical subcommittee of Iran Sports medicine federation education and research committee.

4. Results

In total, 388 elite athletes including 80 females (20.6%) and 308 males (79.4%) participated in this study. The youngest one was 16 and oldest one 36. Comparing 21 various sports disciplines, dragon boat athletes were 48 persons and team of diving and tennis consisted of 4 athletes. Baseline characteristics are shown in Tables 1 and 2.

Two athletes (0.5%) suffered from asthma, and in 8 persons (2%) who didn't have any other cardiovascular symptom, cardiac murmur was reported. Based on physical examination 11 athletes (2.8%) had heart murmur. Investigation of their ECG showed that there was not any evidence

Table 1. [Part 1] Demographic and Descriptive Characteristics

Characteristics	No. (%)
Demographic data	
Male	308 (79.4)
Female	80 (20.6)
Sport category	
Power ^a	200 (51.5)
Endurance ^b	7 (1.8)
Team Sports ^c	151 (46.6)
Sport Discipline	
Dragon Boat	48 (12.4)
Basketball	46 (11.9)
Kabaddi	34 (8.8)
Wushu	34 (8.8)
Volleyball	22 (5.7)
Jujitsu	21 (4.4)
Wrestling	19 (4.9)
Taekwondo	18 (4.6)
Archery	16 (4.1)
Karate	15 (3.9)
water polo	13 (3.4)
shooting	12 (3.1)
Others	90 (23.19)
Total	388 (100)
History findings	
Smoking	9 (2.3)
Dizziness	43 (11.2)
Faint	8 (2.09)
Chest wall pain	21 (5.6)
Anemia ^d	28 (7.4)
Death of Female Family Member under 65	11 (2.8)
Death of Male Family Member under 55	23 (6)
Palpation	49 (13.1)
Extra sound (described by athlete)	8 (2.1)

^aWeightlifting, Wrestling, Wushu, Karate, Taekwondo, Boxing, Judo, Jujitsu.

^bDragon Boat, Cycling, Distance Running, others.

^cVolleyball, Basketball, Kabaddi, Water polo.

^dAnemia is a medical condition in which the red blood cell count or hemoglobin is less than normal. For men, anemia is typically defined as hemoglobin level of less than 13.5 g/100 mL and in women as hemoglobin of less than 12.0 g/100 mL.

of right ventricular hypertrophy in athletes while 12 sportsmen (3%) had signs of left ventricular hypertrophy. First degree heart block was detected among 45 (11.6%), and there

Table 2. [Part 2] Demographic and Descriptive Characteristics

Characteristics	No. (%)
Physical Exam Finding	
Murmur	11 (3.1)
ECG Findings Axis	
NL	380 (97.9)
Right Deviation	4 (1)
Left Deviation	4 (1)
Left Ventricular hypertrophy	12 (3.1)
Atrial hypertrophy	3 (8)
Heart block	
block level 1	35 (11.6)
Branch block	
RBBB	38 (9.7)
Sinus Rhythm	
Bradycardia	124 (32)
Tachycardia	5 (1.3)
Arrhythmia	
PVC	2 (0.5)
WPW	1 (0.3)
T change	
T invert	6 (1.5)
Jpoint elevation	17 (4.4)

was nobody with second degree heart block. Right bundle branch block (RBBB) was shown in 38 (9.8%) but nobody had left bundle branch block (LBBB). Only 2 (0.5%) of them had arrhythmia which was benign premature ventricular contraction (PVC) according to their ECG. Six sportsmen (1.5%) had inverted T wave in v_1 and v_2 leads and nobody had long corrected Q-T interval.

Regarding to the distribution of symptoms among various sports disciplines, water polo players had the highest rate of chest wall pain compliant (30.8% of athletes). The highest rate of palpitation was reported among weightlifters with (40%) and after that were shooting (25%) and water polo (23.1%). Regarding the physical examination, highest resting heart rate belonged to soccer players (64.2% of them had heart rate above 75/min) and lowest heart rate was for water polo players (23.3% had heart rate below 50/min) (Table 3).

There was not any significant relation between ECG changes and variables including sports discipline, history of hypertension, history of lightheadedness, vertigo, unconsciousness, chest pain, history of death below age of

Table 3. Quantitative Parameters

Quantitative Parameters	Mean (SD)	Range
Heart rate	66 (8.4)	48 - 91
Respiratory rate	12 (1.07)	10 - 15
Blood pressure max	11.1 (1.1)	9 - 15.6
Blood pressure min	7.1 (0.8)	5 - 10

65 in female and 55 in male family members due to cardiovascular causes, metabolic disorders, palpitation, and exhaustion. There was a significant relation between gender and ECG changes ($P < 0.0001$) that means 58% of male athletes had ECG variations (which most of them were considered normal changes) but these changes are lesser in female sportspersons (13.8%) which have been shown in Tables 4 and 5. The mean age of athletes with and without ECG changes didn't have any significant correlation ($P = 0.201$) which were 23 and 23.5 years old respectively.

In accordance to Table 6 and Figure 1, it is shown that clinical signs and symptoms could not be used as predictors for ECG changes ($P = 0.863$).




Figure 1. It is shown That Clinical Signs and Symptoms Could Not be Used as Predictors for ECG Changes ($P = 0.863$)

5. Discussion

As above mentioned, two purposes were followed in this study; to determine prevalence of cardiovascular disorders among athletes participating at Asian games and answering the question whether an electrocardiogram (ECG) is a necessary part of pre-participation examination (PPE) for prevention of sudden cardiac death. In other words, the current study was to find any correlation between ECG changes and cardiovascular signs and symptoms. In Iran there are diverse standards for pre-participation examination. For example for school level athletes PPE includes of history taking and physical examination but for professional and championship level athletes the PPE consists of ECG and some blood and urine tests in addition to history taking and physical examination. Sometimes in accordance to level and number of athletes, echocardiography is also performed for them. Among various reasons for performing examinations, predicting and prevention of sudden cardiac death is very important due to its incidence. While the incidence of SCD is low and reported only about 90 annually in United States (39) but regarding the importance of athletes' health, PPE is performed in accordance to 26th Bethesda conference (40, 41) which is emphasized on family history and examination and there is not any clear recommendation for performing ECG (42). Based on American heart association (AHA) guidelines the most effective and cautious method of screening for detecting cardiovascular abnormalities, is history taking and physical examination (43). In Italy taking ECG is required for all athletes prior to participation at sports events (42). In Veneto region of Italy, performing this guideline resulted in 89% decrease of sudden death among athletes (29). European society of cardiology also recommended ECG in European countries for cardiovascular disorder screening (44).

International Olympic committee (IOC) also recommended ECG for athletes participating in Olympic games. For screening of SCD, 42 Diversity of methods and strategies around the world is due to different common causes of sudden cardiac death in various countries. In USA the most common cause of SCD is hypertrophic cardiomyopathy, but in Italy arrhythmogenic right ventricular cardiomyopathy is the most common cause of it (9, 29, 39, 45). Wheeler et al showed that performing ECG is both cost effective and diagnostic for SCD (42). In our current study and regarding to the results we concluded that there is not any correlation between positive signs and symptoms and ECG changes. On the other hand the cost of taking ECG is lesser than 5 US\$, therefore ECG could be considered as a mandatory part of PPE. Although there is not any consensus about the PPE (24) and more studies are recommended

Table 4. Electrocardiogram Changes According to Other Variables^a

	ECG Change		Total	P Value
	No	Yes		
Gender				< 0.001 ^b
Male	128 (41.7)	179 (58.3)	307	
Female	69 (86.3)	11 (13.8)	80	
Sport				0.707 ^c
power speed	82 (45.8)	97 (54.2)	179	
Endurance	4 (57.1)	3 (42.9)	7	
Team	70 (49.6)	71 (50.4)	141	
HTN history				> 0.99 ^c
Yes	1 (100.0)	0	1	
No	194 (50.7)	189 (49.3)	383	
Dizziness history				0.339 ^b
Yes	19 (44.2)	24 (55.8)	43	
No	176 (51.9)	163 (48.1)	339	
Light headedness				0.608 ^b
Yes	7 (58.3)	5 (41.7)	12	
No	187 (50.8)	181 (49.2)	368	
Faint history				> 0.99 ^c
Yes	4 (50.0)	4 (50.0)	8	
No	191 (51.2)	182 (48.8)	373	
Chest pain				0.453 ^b
Yes	9 (42.9)	12 (57.1)	21	
No	181 (51.3)	172 (48.7)	353	
Dead 55 male^d				0.446 ^b
Yes	10 (43.5)	13 (56.5)	23	
No	185 (51.7)	173 (48.3)	358	
Dead 65 female^d				0.323 ^b
Yes	4 (36.4)	7 (63.6)	11	
No	190 (51.5)	179 (48.5)	369	
Metabolic history^e				0.98 ^b
Yes	27 (51.9)	25 (48.1)	52	
No	164 (51.7)	153 (48.3)	317	
Tired history				0.892 ^b
Yes	18 (50.0%)	18 (50.0)	36	
No	172 (51.2%)	164 (48.8)	336	
Palpitation				0.753 ^b
Yes	24 (49.0%)	25 (51.0)	49	
No	166 (51.4%)	157 (48.6)	323	
Extra sound history				0.007 ^c
Yes	8 (100.0)	0	8	
No	182 (50.0)	182 (50.0)	364	
Age				0.201 ^f
Mean (SD)	22.96 (4.44)	23.54 (4.32)		
Median (Range)	23 (16 - 36)	23 (16 - 36)		

^a Values are expressed as No. (%).^b Mann-Whitney U test.^c Kruskal-Wallis Test.^d Family history of early coronary heart disease (heart attack, stroke percutaneous coronary catheter interventional procedure, CABG, treated angina or SCD) in a first-degree family in a female parent or sibling before age 65 years or a male parent or sibling before age 55 years.^e The metabolic syndrome refers to the accompaniment of several cardiovascular risk factors, including: atherogenic dyslipidemia, obesity, insulin resistance and hypertension.^f Spearman's rho.

Table 5. Variables' Regression

	Regression Coefficient	S. E.	Chi Square	P Value	OR
Age	0.006	0.029	0.039	0.844	1.006
Gender					
Male	1.975	0.369	37.301	< 0.0001	7.213
Female					
Sport					
Power-Speed	0.330	0.249	1.805	0.179	1.391
Endurance	-0.323	0.792	0.191	0.662	0.724
Team					
HTN history					
Yes	-1.712	2.542	0.885	0.347	0.181
No					
Dizziness history					
Yes	-0.182	0.392	0.23	0.632	0.833
No					
Light headedness					
Yes	-0.041	0.837	0.003	0.957	0.959
No					
Faint history					
Yes	-0.519	1.091	0.277	0.599	0.595
No					
Chest pain					
Yes	0.069	0.541	0.018	0.894	1.072
No					
Dead 55 male					
Yes	0.207	0.511	0.179	0.672	1.23
No					
Dead 65 female					
Yes	0.559	0.702	0.735	0.319	1.75
No					
Metabolic history					
Yes	0.397	0.354	0.013	0.909	1.04
No					
Tired					
Yes	0.316	0.435	0.556	0.456	1.372
No					
Palpitation					
Yes	0.118	0.364	0.11	0.740	1.126
No					
Extra sound history					
Yes	-2.307	1.661	3.874	0.049	0.099
No					

Table 6. Area Under the ROC Curve (AUC)

Parameters	Results
Area under the ROC curve (AUC)	0.505
Standard Error	0.0289
95% Confidence interval	0.450 to 0.560
Z statistic	0.173
Significance level P (Area = 0.5)	0.8627

in accordance to application of different methods, their costs and efficacies (46). Performing echocardiography is still expensive (47), controversial and needs more investigations.

While 2.3% of athletes were smokers and their average age was 4 years more than mean age of all athletes, regarding the increase of cardiovascular disorders' risk, the educational programs could be conducted for confronting this problem.

Acknowledgments

Authors of this study appreciate Iran sports medicine federation and Iran National Olympic committee for their support in conducting the current investigation.

Footnotes

Authors' Contribution: Study concept and design: Shahin Salehi; administrative, technical, and material support: Farhad Moradi Shahpar; study supervision and technical, and material support: Gholamreza Norouzi; consultant: other authors.

Funding/Support: All expenses for this study were covered by Iran sports medicine federation as part of a mandatory pre-participation examination.

References

1. Maron BJ, Haas TS, Doerer JJ, Thompson PD, Hodges JS. Comparison of U.S. and Italian experiences with sudden cardiac deaths in young competitive athletes and implications for preparticipation screening strategies. *Am J Cardiol.* 2009;**104**(2):276–80. doi: [10.1016/j.amjcard.2009.03.037](#). [PubMed: [19576360](#)].
2. Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults?. *J Am Coll Cardiol.* 2003;**42**(11):1959–63. [PubMed: [14662259](#)].
3. Ferreira M, Santos-Silva PR, de Abreu LC, Valenti VE, Crispim V, Imaizumi C, et al. Sudden cardiac death athletes: a systematic review. *Sports Med Arthrosc Rehabil Ther Technol.* 2010;**2**:19. doi: [10.1186/1758-2555-2-19](#). [PubMed: [20682064](#)].
4. 26th Bethesda Conference. Recommendations for determining eligibility for competition in athletes with cardiovascular abnormalities. *Medicine and science in sports and exercise.*
5. Burke AP, Farb A, Virmani R, Goodin J, Smialek JE. Sports-related and non-sports-related sudden cardiac death in young adults. *Am Heart J.* 1991;**121**(2 Pt 1):568–75. [PubMed: [1825009](#)].
6. Maron BJ, Carney KP, Lever HM, Lewis JF, Barac I, Casey SA, et al. Relationship of race to sudden cardiac death in competitive athletes with hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2003;**41**(6):974–80. [PubMed: [12651044](#)].
7. Maron BJ, Mitchell JH. Revised eligibility recommendations for competitive athletes with cardiovascular abnormalities. *J Am Coll Cardiol.* 1994;**24**(4):848–50. [PubMed: [7930213](#)].
8. Maron BJ, Pelliccia A. The heart of trained athletes: cardiac remodeling and the risks of sports, including sudden death. *Circulation.* 2006;**114**(15):1633–44. doi: [10.1161/CIRCULATIONAHA.106.613562](#). [PubMed: [17030703](#)].
9. Maron BJ, Shirani J, Poliac LC, Mathenge R, Roberts WC, Mueller FO. Sudden death in young competitive athletes. Clinical, demographic, and pathological profiles. *JAMA.* 1996;**276**(3):199–204. [PubMed: [8667563](#)].
10. Mueller FO, Cantu RC. Catastrophic injuries and fatalities in high school and college sports, fall 1982-spring 1988. *Med Sci Sports Exerc.* 1990;**22**(6):737–41. [PubMed: [2287249](#)].
11. Van Camp SP, Bloor CM, Mueller FO, Cantu RC, Olson HG. Nontraumatic sports death in high school and college athletes. *Med Sci Sports Exerc.* 1995;**27**(5):641–7. [PubMed: [7674867](#)].
12. Thompson PD, Franklin BA, Balady GJ, Blair SN, Corrado D, Estes N3, et al. Exercise and acute cardiovascular events placing the risks into perspective: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism and the Council on Clinical Cardiology. *Circulation.* 2007;**115**(17):2358–68. doi: [10.1161/CIRCULATIONAHA.107.181485](#). [PubMed: [17468391](#)].
13. Inklaar H, Panhuyzen-Goedkoop NM. Periodical cardiovascular screening is mandatory for elite athletes. *Neth Heart J.* 2007;**15**(6):221–3. [PubMed: [17612687](#)].
14. Borjesson M, Pelliccia A. Incidence and aetiology of sudden cardiac death in young athletes: an international perspective. *Br J Sports Med.* 2009;**43**(9):644–8. doi: [10.1136/bjism.2008.054718](#). [PubMed: [19734497](#)].
15. van der Werf C, van Langen IM, Wilde AA. The relationship between pre-participation screening of young competitive athletes and family screening. *J Am Coll Cardiol.* 2009;**53**(24):2309. doi: [10.1016/j.jacc.2008.12.080](#). [PubMed: [19520259](#)] author reply 2309–10.
16. Hevia AC, Fernandez MM, Palacio JM, Martin EH, Castro MG, Reguero JJ. ECG as a part of the preparticipation screening programme: an old and still present international dilemma. *Br J Sports Med.* 2011;**45**(10):776–9. doi: [10.1136/bjism.2009.063958](#). [PubMed: [19858111](#)].
17. Chaitman BR. An electrocardiogram should not be included in routine preparticipation screening of young athletes. *Circulation.* 2007;**116**(22):2610–4. doi: [10.1161/CIRCULATIONAHA.107.71465](#). [PubMed: [18040040](#)] discussion 2615.

18. Link MS, Estes NA. Sudden cardiac death in the athlete: bridging the gaps between evidence, policy, and practice. *Circulation*. 2012;**125**(20):2511-6. doi: [10.1161/CIRCULATIONAHA.111.023861](https://doi.org/10.1161/CIRCULATIONAHA.111.023861). [PubMed: [22615422](https://pubmed.ncbi.nlm.nih.gov/22615422/)].
19. Maron BJ. Cardiovascular risks to young persons on the athletic field. *Ann Intern Med*. 1998;**129**(5):379-86. [PubMed: [9735066](https://pubmed.ncbi.nlm.nih.gov/9735066/)].
20. Myerburg RJ, Vetter VL. Electrocardiograms should be included in preparticipation screening of athletes. *Circulation*. 2007;**116**(22):2616-26. doi: [10.1161/CIRCULATIONAHA.107.733519](https://doi.org/10.1161/CIRCULATIONAHA.107.733519). [PubMed: [18040041](https://pubmed.ncbi.nlm.nih.gov/18040041/)].
21. Estes N3, Link MS. Preparticipation athletic screening including an electrocardiogram: an unproven strategy for prevention of sudden cardiac death in the athlete. *Prog Cardiovasc Dis*. 2012;**54**(5):451-4. doi: [10.1016/j.pcad.2012.01.008](https://doi.org/10.1016/j.pcad.2012.01.008). [PubMed: [22386297](https://pubmed.ncbi.nlm.nih.gov/22386297/)].
22. Maron BJ, Bodison SA, Wesley YE, Tucker E, Green KJ. Results of screening a large group of intercollegiate competitive athletes for cardiovascular disease. *J Am Coll Cardiol*. 1987;**10**(6):1214-21. [PubMed: [2960727](https://pubmed.ncbi.nlm.nih.gov/2960727/)].
23. Baggish AL, Hutter AMJr, Wang F, Yared K, Weiner RB, Kupperman E, et al. Cardiovascular screening in college athletes with and without electrocardiography: A cross-sectional study. *Ann Intern Med*. 2010;**152**(5):269-75. doi: [10.7326/0003-4819-152-5-201003020-00004](https://doi.org/10.7326/0003-4819-152-5-201003020-00004). [PubMed: [20194232](https://pubmed.ncbi.nlm.nih.gov/20194232/)].
24. Gupta S, Baman T, Day SM. Cardiovascular health, part 1: preparticipation cardiovascular screening. *Sports Health*. 2009;**1**(6):500-7. doi: [10.1177/1941738109350405](https://doi.org/10.1177/1941738109350405). [PubMed: [23015913](https://pubmed.ncbi.nlm.nih.gov/23015913/)].
25. Wisten A, Messner T. Symptoms preceding sudden cardiac death in the young are common but often misinterpreted. *Scand Cardiovasc J*. 2005;**39**(3):143-9. doi: [10.1080/14017430510009168](https://doi.org/10.1080/14017430510009168). [PubMed: [16146977](https://pubmed.ncbi.nlm.nih.gov/16146977/)].
26. Corrado D, Basso C, Schiavon M, Pelliccia A, Thiene G. Preparticipation screening of young competitive athletes for prevention of sudden cardiac death. *J Am Coll Cardiol*. 2008;**52**(24):1981-9. doi: [10.1016/j.jacc.2008.06.053](https://doi.org/10.1016/j.jacc.2008.06.053). [PubMed: [19055989](https://pubmed.ncbi.nlm.nih.gov/19055989/)].
27. Holst AG, Winkel BG, Theilade J, Kristensen IB, Thomsen JL, Ottesen GL, et al. Incidence and etiology of sports-related sudden cardiac death in Denmark-implications for preparticipation screening. *Heart Rhythm*. 2010;**7**(10):1365-71. doi: [10.1016/j.hrthm.2010.05.021](https://doi.org/10.1016/j.hrthm.2010.05.021). [PubMed: [20580680](https://pubmed.ncbi.nlm.nih.gov/20580680/)].
28. Patel A, Lantos JD. Can we prevent sudden cardiac death in young athletes: the debate about preparticipation sports screening. *Acta Paediatr*. 2011;**100**(10):1297-301. doi: [10.1111/j.1651-2227.2011.02337.x](https://doi.org/10.1111/j.1651-2227.2011.02337.x). [PubMed: [21535127](https://pubmed.ncbi.nlm.nih.gov/21535127/)].
29. Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden cardiovascular death in young competitive athletes after implementation of a preparticipation screening program. *JAMA*. 2006;**296**(13):1593-601. doi: [10.1001/jama.296.13.1593](https://doi.org/10.1001/jama.296.13.1593). [PubMed: [17018804](https://pubmed.ncbi.nlm.nih.gov/17018804/)].
30. Corrado D, Basso C, Schiavon M, Thiene G. Screening for hypertrophic cardiomyopathy in young athletes. *N Engl J Med*. 1998;**339**(6):364-9. doi: [10.1056/NEJM199808063390602](https://doi.org/10.1056/NEJM199808063390602). [PubMed: [9691102](https://pubmed.ncbi.nlm.nih.gov/9691102/)].
31. Bille K, Figueiras D, Schamasch P, Kappenberger L, Brenner JI, Meijboom FJ, et al. Sudden cardiac death in athletes: the Lausanne Recommendations. *Eur J Cardiovasc Prev Rehabil*. 2006;**13**(6):859-75. doi: [10.1097/01.hjr.0000238397.50341.4a](https://doi.org/10.1097/01.hjr.0000238397.50341.4a). [PubMed: [17143117](https://pubmed.ncbi.nlm.nih.gov/17143117/)].
32. Thunenkotter T, Schmied C, Dvorak J, Kindermann W. Benefits and limitations of cardiovascular pre-competition screening in international football. *Clin Res Cardiol*. 2010;**99**(1):29-35. doi: [10.1007/s00392-009-0072-7](https://doi.org/10.1007/s00392-009-0072-7). [PubMed: [19756813](https://pubmed.ncbi.nlm.nih.gov/19756813/)].
33. Marijon E, Tafflet M, Celermajer DS, Dumas F, Perier MC, Mustafic H, et al. Sports-related sudden death in the general population. *Circulation*. 2011;**124**(6):672-81. doi: [10.1161/CIRCULATIONAHA.110.008979](https://doi.org/10.1161/CIRCULATIONAHA.110.008979). [PubMed: [21788587](https://pubmed.ncbi.nlm.nih.gov/21788587/)].
34. Bessem B, Groot FP, Nieuwland W. The Lausanne recommendations: a Dutch experience. *Br J Sports Med*. 2009;**43**(9):708-15. doi: [10.1136/bjism.2008.056929](https://doi.org/10.1136/bjism.2008.056929). [PubMed: [19549617](https://pubmed.ncbi.nlm.nih.gov/19549617/)].
35. Ma JZ, Dai J, Sun B, Ji P, Yang D, Zhang JN. Cardiovascular preparticipation screening of young competitive athletes for prevention of sudden death in China. *J Sci Med Sport*. 2007;**10**(4):227-33. doi: [10.1016/j.jsams.2006.07.001](https://doi.org/10.1016/j.jsams.2006.07.001). [PubMed: [16914373](https://pubmed.ncbi.nlm.nih.gov/16914373/)].
36. Tanaka Y, Yoshinaga M, Anan R, Tanaka Y, Nomura Y, Oku S, et al. Usefulness and cost effectiveness of cardiovascular screening of young adolescents. *Med Sci Sports Exerc*. 2006;**38**(1):2-6. [PubMed: [16394946](https://pubmed.ncbi.nlm.nih.gov/16394946/)].
37. Haneda N, Mori C, Nishio T, Saito M, Kajino Y, Watanabe K, et al. Heart diseases discovered by mass screening in the schools of Shimane Prefecture over a period of 5 years. *Jpn Circ J*. 1986;**50**(12):1325-9. [PubMed: [3820545](https://pubmed.ncbi.nlm.nih.gov/3820545/)].
38. Alattar A, Ghani S, Mahdy N, Hussain H, Maffulli N. Pre-participation musculoskeletal and cardiac screening of male athletes in the United Arab Emirates. *Transl Med UniSa*. 2014;**9**:43-9. [PubMed: [24809035](https://pubmed.ncbi.nlm.nih.gov/24809035/)].
39. Maron BJ. Hypertrophic cardiomyopathy and other causes of sudden cardiac death in young competitive athletes, with considerations for preparticipation screening and criteria for disqualification. *Cardiol Clin*. 2007;**25**(3):399-414. doi: [10.1016/j.ccl.2007.07.006](https://doi.org/10.1016/j.ccl.2007.07.006). [PubMed: [17961794](https://pubmed.ncbi.nlm.nih.gov/17961794/)] vi.
40. Maron BJ, Douglas PS, Graham TP, Nishimura RA, Thompson PD. Task Force 1: preparticipation screening and diagnosis of cardiovascular disease in athletes. *J Am Coll Cardiol*. 2005;**45**(8):1322-6. doi: [10.1016/j.jacc.2005.02.007](https://doi.org/10.1016/j.jacc.2005.02.007). [PubMed: [15837281](https://pubmed.ncbi.nlm.nih.gov/15837281/)].
41. Maron BJ, Thompson PD, Ackerman MJ, Balady G, Berger S, Cohen D, et al. Recommendations and considerations related to preparticipation screening for cardiovascular abnormalities in competitive athletes: 2007 update: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. *Circulation*. 2007;**115**(12):1643-455. doi: [10.1161/CIRCULATIONAHA.107.181423](https://doi.org/10.1161/CIRCULATIONAHA.107.181423). [PubMed: [17353433](https://pubmed.ncbi.nlm.nih.gov/17353433/)].
42. Wheeler MT, Heidenreich PA, Froelicher VF, Hlatky MA, Ashley EA. Cost-effectiveness of preparticipation screening for prevention of sudden cardiac death in young athletes. *Ann Intern Med*. 2010;**152**(5):276-86. doi: [10.7326/0003-4819-152-5-201003020-00005](https://doi.org/10.7326/0003-4819-152-5-201003020-00005). [PubMed: [20194233](https://pubmed.ncbi.nlm.nih.gov/20194233/)].
43. Koester MC. A Review of Sudden Cardiac Death in Young Athletes and Strategies for Preparticipation Cardiovascular Screening. *J Athl Train*. 2001;**36**(2):197-204. [PubMed: [12937463](https://pubmed.ncbi.nlm.nih.gov/12937463/)].
44. Corrado D, Pelliccia A, Bjornstad HH, Vanhees L, Biffi A, Borjesson M, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J*. 2005;**26**(5):516-24. doi: [10.1093/eurheartj/ehi018](https://doi.org/10.1093/eurheartj/ehi018). [PubMed: [15689345](https://pubmed.ncbi.nlm.nih.gov/15689345/)].
45. Pelliccia A, Maron BJ. Preparticipation cardiovascular evaluation of the competitive athlete: perspectives from the 30-year Italian experience. *Am J Cardiol*. 1995;**75**(12):827-9. [PubMed: [7717289](https://pubmed.ncbi.nlm.nih.gov/7717289/)].
46. Borriore P, Quaranta F, Ciminelli E. Pre-participation screening for the prevention of sudden cardiac death in athletes. *World J Methodol*. 2013;**3**(1):1-6. doi: [10.5662/wjmv.v3.i1.1](https://doi.org/10.5662/wjmv.v3.i1.1). [PubMed: [25237617](https://pubmed.ncbi.nlm.nih.gov/25237617/)].
47. Anderson JB, Grenier M, Edwards NM, Madsen NL, Czosek RJ, Spar DS, et al. Usefulness of combined history, physical examination, electrocardiogram, and limited echocardiogram in screening adolescent athletes for risk for sudden cardiac death. *Am J Cardiol*. 2014;**114**(11):1763-7. doi: [10.1016/j.amjcard.2014.09.011](https://doi.org/10.1016/j.amjcard.2014.09.011). [PubMed: [25307198](https://pubmed.ncbi.nlm.nih.gov/25307198/)].