

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/25458>

Please be advised that this information was generated on 2017-07-15 and may be subject to change.

# Cloning of the Human Carnitine-Acylcarnitine Carrier cDNA and Identification of the Molecular Defect in a Patient

Marjan Huizing,<sup>1</sup> Vito Iacobazzi,<sup>2</sup> Lodewijk IJlst,<sup>3</sup> Paul Savelkoul,<sup>1</sup> Wim Ruitenbeek,<sup>1</sup> Lambert van den Heuvel,<sup>1</sup> Cesare Indiveri,<sup>2</sup> Jan Smeitink,<sup>1</sup> Frans Trijbels,<sup>1</sup> Ronald Wanders,<sup>3</sup> and Ferdinando Palmieri<sup>2</sup>

<sup>1</sup>University Hospital Nijmegen, Department of Pediatrics, Nijmegen, The Netherlands; <sup>2</sup>University of Bari, Department of Pharmaco-Biology, Bari, Italy; and <sup>3</sup>University of Amsterdam, Departments of Clinical Chemistry and Pediatrics, Amsterdam

## Summary

The carnitine-acylcarnitine carrier (CAC) catalyzes the translocation of long-chain fatty acids across the inner mitochondrial membrane. We cloned and sequenced the human CAC cDNA, which has an open reading frame of 903 nucleotides. Northern blot studies revealed different expression levels of CAC in various human tissues. Furthermore, mutation analysis was performed for a CAC-deficient infant. Direct sequencing of the patient's cDNA revealed a homozygous cytosine nucleotide insertion. This insertion provokes a frameshift and an extension of the open reading frame with 23 novel codons. This is the first report documenting a mutation, in the CAC cDNA, responsible for mitochondrial  $\beta$ -oxidation impairment.

## Introduction

The carnitine-acylcarnitine carrier (CAC) shuttles acylcarnitine esters, in exchange for free carnitine, across the inner mitochondrial membrane (Pande 1975; Ramsay and Tubbs 1975). This transport is an essential step in the process of long-chain fatty-acid oxidation (Stanley 1987; Coates and Tanaka 1992; Stanley et al. 1992). The oxidation of fatty acids in mitochondria plays an important role in energy production. During fasting, fatty acids are used for hepatic ketone-body synthesis. Furthermore, fatty acids are an important source of energy for heart muscle and also for skeletal muscle, during exercise, whereas ketone bodies are excellent substrates

for the brain (Stanley 1987; Coates and Tanaka 1992). The overall fatty-acid oxidation in mitochondria requires the concerted action of at least 17 different proteins, including 16 enzymes and the transporter CAC. Genetic defects have been identified in most of these proteins (Stanley 1987; Stanley et al. 1992). These defects generally present in early infancy, with acute, potentially life-threatening episodes of hypoketotic hypoglycemic coma induced by fasting. The clinical phenotypes are very similar and can be attributed to one of three major types of presentation, with predominantly hepatic, cardiac, or skeletal muscle involvement (Kelly and Strauss 1994; Pande and Murthy 1994; Pollit 1995). So far, six cases of CAC deficiency have been reported (Stanley et al. 1992; Pande et al. 1993; Brivet et al. 1994, 1996; Niezen-Koning et al. 1995; Ogier de Baulney et al. 1995). The main features in these severely affected patients with onset in the neonatal period are hypoketotic hypoglycemia, mild hyperammonemia, variable dicarboxylic aciduria, hepatomegaly with abnormal liver functions, various cardiac symptoms, and skeletal muscle weakness. In all cases, CAC activity in cultured skin fibroblasts is below detectable levels. However, so far, the CAC deficiency has not been characterized at the molecular level, in any patient.

Fundamental properties of eukaryotic CAC have been investigated extensively in intact mitochondria (Pande 1975; Ramsay and Tubbs 1975; Pande and Parvin 1980; Idell-Wenger 1981; Murthy and Pande 1984) and after purification and reconstitution into liposomes (Indiveri et al. 1990, 1991a, 1991b, 1992, 1994, 1995). The carrier is embedded in the inner mitochondrial membrane and has an apparent molecular mass of 32.5 kD in rat liver (Indiveri et al. 1990). It governs a one-to-one exchange, between long-chain acylcarnitine esters and nonesterified carnitine, across the inner mitochondrial membrane and also the unidirectional transport of carnitine across this membrane, although less efficiently (Pande and Parvin 1980; Indiveri et al. 1991a). Incorporated into liposomes, the purified carrier protein has substrate specificity and inhibitor sensitivities similar to

Received February 13, 1997; accepted for publication September 16, 1997; electronically published November 7, 1997.

Address for correspondence and reprints: Dr. Ferdinando Palmieri, University of Bari, Department of Pharmaco-Biology, Laboratory of Biochemistry and Molecular Biology, Via E. Orabona 4, 70125 Bari, Italy. E-mail: fpalm@farmbiol.uniba.it

© 1997 by The American Society of Human Genetics. All rights reserved.  
0002-9297/97/6106-0006\$02.00

those in intact mitochondria (Indiveri et al. 1990, 1991a, 1991b). CAC operates according to a ping-pong mechanism (Indiveri et al. 1994).

Very recently we described the cDNA and amino acid sequence of the rat CAC (Indiveri et al. 1997). These studies have shown that CAC belongs to a protein family that, so far, has been found to comprise 10 biochemically well-characterized mitochondrial carriers and also several other members of unknown function that are beginning to emerge with the advance of genomic DNA sequencing (Walker and Runswick 1993; Palmieri 1994; Crabeel et al. 1996; Palmieri et al. 1996). These proteins have evolved from a common ancestor, by two-tandem gene duplication, and have related structures and mechanisms (Walker and Runswick 1993; Palmieri 1994).

We report on the nucleotide sequence of the human CAC cDNA and the corresponding amino acid sequence, as well as the distribution of CAC mRNA in human tissues. For the first time, a mutation has been found in the CAC cDNA of a unique, now 9-year-old, CAC-deficient patient.

## Patient and Methods

### Case Report

This 9-year-old girl with a normal family history survived a severe neonatal condition consisting of hypoglycemia, cardiac arrest, hepatomegaly, and hepatic dysfunction. These features are often observed in patients with a fatty-acid oxidation disorder. Thereafter episodes of lethargy and hepatomegaly occurred only during mild viral infections. Acylglycines were nondiagnostic, and acylcarnitines showed increased medium- to long-chain hydroxy and unsaturated derivatives. A liver biopsy showed fatty infiltration. During the last few years, she has had a rather normal physical and neurophysiological development: she has done well with prolonged exercise, and in between the attacks she is an alert, cooperative, and pleasant young girl. Her symptoms are much milder than those in the other CAC-deficient patients who have been described. Cultured skin fibroblasts were sent to one of us (R.W.) to investigate a possible defect in mitochondrial fatty-acid oxidation.

### Fibroblast Study

Overall  $\beta$ -oxidation in fibroblasts was measured as described previously by Olpin et al. (1992). The CAC activity in fibroblasts was measured essentially according to the study by Pande et al. (1993).

### Cloning and Sequencing of the Human CAC

Oligonucleotides were designed on the basis of the cDNA sequence of the rat liver CAC (Indiveri et al. 1997), at the following nucleotide positions: 125–150


```

1  GGGCTCTGAGC  GGGCCGCCCG  GCAGGTCTGAG  AACTGACAGA  CCGAGTGA  GACCGACTGA  CC ATG  GCC  GAC  CAG
1  M  A  R  Q
75  CCA  AAA  CCC  ATC  AGC  CCG  CTC  AAG  AAC  CTG  CTG  GCC  GGC  GGC  TTT  GGC  GGC  GTG  TGC  CTG  GTG  TTC  GTC  GGT
5  P  K  P  I  S  P  L  K  N  L  L  A  G  G  F  G  G  V  C  L  V  F  V  G
147  CAC  CCF  CTG  GAC  AGC  GTC  AAG  GTC  CGA  CTG  CAG  ACA  CAG  CCA  CCG  AGT  TTG  CCF  GGA  CAA  CCF  CCC  ATG  TAC
29  H  P  L  D  T  V  K  V  R  L  Q  T  Q  P  P  S  L  P  G  Q  P  P  M  Y
219  TCT  GGG  ACC  TTT  GAC  TGT  TTC  CGG  AAG  ACT  CTT  TTT  AGA  GAG  GGC  ATC  ACG  GGG  CTA  TAT  CCG  GGA  ATG  GCT
53  S  G  T  E  D  C  F  R  K  T  L  F  R  R  G  I  T  G  L  Y  R  G  M  A
291  GCC  CCF  ATC  ATC  GGG  GTC  ACT  CCC  ATG  TTT  GGC  GTG  TGC  TTC  TTT  GGG  TTT  GGT  TTG  GGG  AAG  AAA  CTA  CAA
77  A  P  I  I  G  V  T  P  M  F  A  V  C  F  F  G  F  G  L  G  K  E  L  Q
363  CAG  AAA  CAC  CCA  GAA  GAT  GTG  CTC  AGC  TAT  CCC  CAG  CTT  TTT  GCA  GCT  GGG  ATG  TTA  TCT  GGC  GTA  TTC  ACC
101  Q  K  H  P  E  D  E  L  S  Y  P  Q  I  P  A  A  M  L  S  G  V  F  T
435  ACA  GGA  ATC  ATG  ACT  CCF  GGA  GAA  CCG  ATC  AAG  TGC  TTA  TTA  CAG  ATT  CAG  GCT  TCT  TCA  GGA  GAA  AGC  AAG
125  T  G  I  M  T  P  G  E  R  I  K  C  L  L  Q  I  Q  A  S  S  G  E  S  K
507  TAC  ACT  GGT  ACC  TTG  GAC  TGT  GCA  AAG  AAG  CTG  TAC  CAG  GAG  TTT  GGG  ATC  CGA  GGC  ATC  TAC  AAA  GGG  ACT
149  Y  T  G  T  L  D  C  A  K  K  L  Y  Q  R  F  G  I  R  G  L  Y  K  G  T
579  GTG  CTT  ACC  CTT  ATG  CGA  GAT  GTC  CCA  GCT  AGT  GGA  ATG  TAT  TTC  ATG  ACA  TAT  GAA  TGG  CTG  AAA  AAT  ATC
173  L  L  T  L  M  R  D  V  P  A  S  G  M  V  F  M  T  Y  R  W  L  K  N  I
651  TTC  ACF  CCG  GAG  CGA  AAG  AAG  GTC  AGT  GAG  CTC  AGT  GGC  CCF  CCG  ATC  TTG  GTG  GCF  GGG  GGC  ATT  GCA  GGG
197  F  T  P  E  G  K  E  V  S  E  L  S  A  P  R  L  L  V  A  G  G  L  A  G
723  ATC  TTC  AAC  TGG  GCT  GTG  GCA  ATC  CCC  CGA  GAT  GTG  CTC  AAG  TCT  CGA  TTC  CAG  ACT  GCA  CCF  CCF  GGG  AAA
221  I  F  N  W  A  V  A  I  P  P  D  V  L  K  S  R  F  Q  T  A  P  P  O  K
795  TAT  CCF  AAT  GGT  TTC  AGA  GAT  GTG  CTG  AGG  GAG  CTG  ATC  CCG  GAT  GAA  GGA  GTC  ACA  TCC  TTG  TAC  AAA  GGG
245  Y  P  N  G  F  R  D  V  L  R  R  L  I  E  D  E  G  V  T  S  L  Y  K  G
867  TTC  AAT  GCA  GTG  ATG  ATC  CGA  GGC  TTC  CGA  GGC  AAT  GGC  GCT  TTT  TTC  CTT  GGC  TTT  GAA  GTF  GGC  ATG  AAG
269  F  N  A  V  M  I  R  A  F  P  A  N  A  A  C  F  L  G  F  E  E  A  M  K
939  TTC  CTT  AAT  TGG  GGC  ACC  CCC  AAC  TTG  TGA  GGC  TGA  AGGC  TGC  TCA  AGTT  CACT  TCT  GGA  TGC  TGG  AAG  C  TGT  CG
293  E  L  N  W  A  T  P  N  L
1014  TTGAGGAGAA  GGAATAGTAA  GCAGAACTAA  GCAGTCTTGG  AGGGCAAGGG  GAGGGGAATG  GTGAGATCCG  AGGCC
1089  TGTGCAATGA  CTTGCTGAGA  CTGTTGCCCT  AATGACATCC  TGCACCGTGT  ATAACCTAGT  GTGTCATTT  GAAACTT
1166  GAATTCAT  TCTTATCAAT  TTAAGGGATC  TTAAGGGAT  TTGAATGAA  CAAGTAGCT  CCAAGCCAG  TAC  TACCTGT

```

**Figure 1** Nucleotide sequence of human CAC cDNA. The deduced amino acid sequence of the open reading frame of 903 bp (301 codons) is shown below the nucleotide sequence. Human CAC amino acids that differ from those of rat CAC (Indiveri et al. 1997) are shown in underlined, italicized print. The cDNA and deduced protein sequences are in GenBank (accession no. Y10319; <http://www.ncbi.nlm.nih.gov/Web/Genbank/index.htm>).

(sense 1F, 5'-CCTGGTGTGTTTGTGGGGCACCCCTTG-3'); 455–480 (antisense 1R, 5'-CTGAATCTGCAGTAAGCATTGATC-3'); 405–430 (sense 2F, 5'-GGGATGTTATCTGGTGTGTTCCACCA-3'); and 937–962 (antisense 2R, 5'-ACAAGTTGGGGGCAATCCAATTGAG-3'). These primers (Pharmacia Biotech) were used in PCRs, to amplify cDNA fragments encoding the human liver CAC. The template was first-strand cDNA reverse transcribed, with oligo(dT) by AMV reverse transcriptase (Boehringer Mannheim) for 1 h at 55°C, from human liver mRNA (Clontech). PCR reactions were carried out for 30 cycles (94°C for 30 s, 60°C for 1 min, and 72°C for 2 min). At the end, a single incubation at 72°C for 7 min was added. The extension to the 3' end was performed by priming from poly(A), with two nested forward primers (AGAAGAAGGAGTCA-CCTCCTTGTA and CAAAGGGTTCAATGCAGTCA-TGA, corresponding to nucleotides 831–854 and 855–877, respectively, of the rat cDNA sequence). To extend the sequence to the 5' end, a touchdown PCR (Don et al. 1991) was performed on an adaptor-ligated



**Figure 2** Expression of CAC mRNA in human tissues. Northern blots (2  $\mu$ g poly[A] RNA/tissue) hybridized with a human CAC probe (A) and an actin probe (B) are shown. The molecular weight is indicated in the figure: human CAC mRNA is ~1.8 kb. Results for the following tissues are shown: heart (H), brain (B), placental (P), lung (Lu), liver (L), skeletal muscle (SM), kidney (K), and pancreatic (Pa).

double-stranded human liver cDNA (Clontech). In this case we employed two nested forward adaptor primers and two reverse-specific primers (TTGTGTCTG-CAGTCGGACCTTGAC and CCAGAGGGTGACCG-ACGAACACCAGGC, corresponding to nucleotides 257–278 and 121–146, respectively, of the rat cDNA sequence). DNA-sequence analysis was performed according to established procedures (Indiveri et al. 1997).

#### Northern Blot Analysis

The sequence corresponding to nucleotides 421–967 of the human CAC cDNA was used as a probe in northern blots performed on mRNA derived from various human tissues (Clontech). Northern blot analysis was performed according to standard protocols, with high stringency. The filters were autoradiographed at  $-80^{\circ}\text{C}$ . For normalization of the hybridization signals a probe encoding part of human actin (Clontech) was employed.

#### Mutation Detection

Total RNA was extracted (Chomczynski and Sacchi 1987) from cultured skin fibroblasts and was stored as an ethanol precipitate at  $-80^{\circ}\text{C}$ . A 5- $\mu$ g sample of RNA was reverse transcribed to cDNA, in 1 h at  $42^{\circ}\text{C}$ , with 200 U of Superscript II reverse transcriptase (Life Technologies), by use of oligo(dT) and random hexamer primers. Five microliters of this first-strand cDNA was subjected to PCR amplification. By PCR, we generated three overlapping fragments that covered the entire coding region, using the following synthetic oligonucleotide primers (Perkin Elmer): for fragment 1, 5'-GCAGGT-CGAGAACTGACAGAC-3' (sense F1) and 5'-TTTCTC-CTGAAGAAGCCTGAA-3' (antisense R1), at positions

21–41 and 481–501, respectively, of the human CAC cDNA sequence (fig. 1); for fragment 2, 5'-CCTGGA-GAACGGATCAAGTG-3' (sense F2) and 5'-CAATTA-AGGAACTTCATGGCAA-3' (antisense R2), at positions 450–469 and 928–949, respectively; and, for fragment 3, 5'-GCAGTGATGATCCGAGCCTTC-3' (sense F3) and 5'-ACAGGTAGTATCTGGTCTGGAA-3' (antisense R3), at positions 873–893 and 1224–1245, respectively. For each fragment, 35 cycles of PCR were performed ( $92^{\circ}\text{C}$  for 60 s,  $60^{\circ}\text{C}$  for 60 s, and  $72^{\circ}\text{C}$  for 90 s). The cycles were preceded by an initial denaturation step at  $95^{\circ}\text{C}$  for 3 min and were followed by a final extension at  $72^{\circ}\text{C}$  for 10 min. Ten microliters of each PCR reaction was analyzed on 1.0% agarose gels with 0.5  $\mu$ g ethidium bromide/ml in  $1 \times$  Tris borate-EDTA. The nucleotide sequences of the PCR products were analyzed by direct sequencing using the *Taq* Dye Deoxy Terminator Cycle Sequencing Kit (Applied Biosystems), according to the manufacturer's recommendations.

## Results

### Cloning and Sequencing of Human CAC cDNA

The recently reported rat cDNA of CAC (Indiveri et al. 1997) was used for cloning and sequencing of the human CAC homologue. Four overlapping sequences were amplified, by PCR using human liver cDNA as a template, with synthetic oligonucleotide primers based on the rat cDNA sequence and, to extend the sequence to the 5' end, with two nested forward primers complementary to adaptors that were added to the 5' extremities of the human liver cDNA. The obtained cDNA sequence was 1,243 bp in length and had an open reading frame of 903 bp (fig. 1). Assignment of position 63 as the first nucleotide of the initiation codon was deduced from comparison with the CAC cDNA of rat liver (Indiveri et al. 1997), since no stop codon was found in the 5' UTR. The sequence, which extended into the 3' non-coding region, did not contain the polyadenylation signal. The protein encoded by the human CAC cDNA contains 301 amino acids, and its calculated molecular weight is 32.9 kD.

### Expression of CAC mRNA in Human Tissues

A hybridization probe consisting of nucleotides 421–967 of the human liver CAC cDNA was employed in northern blot experiments. Figure 2 shows the presence of one band for CAC mRNA, ~1.8 kb in length, in various human tissue types. The CAC mRNA was highly expressed in heart, skeletal muscle, and liver tissues. A much lower level of expression was found in brain, placental, pancreatic, and kidney tissues and especially in lung tissue. These differences in the level of expression of the CAC transcript in the various tissues

**Table 1****Overall  $\beta$ -Oxidation Rates and CAC Activities in Cultured Skin Fibroblasts of the Patient and of Controls**

	Patient	Controls ( <i>n</i> ) <sup>a</sup>
Overall $\beta$ -oxidation (nmol/h/mg):		
[9,10- <sup>3</sup> H]Myristic acid	4.1	6.0 $\pm$ 2.3 (61)
[9,10- <sup>3</sup> H]Palmitic acid	2.4	8.8 $\pm$ 4.1 (33)
CAC activity (pmol/min/mg):		
Produced <sup>14</sup> CO <sub>2</sub>	ND <sup>b</sup>	47 $\pm$ 16 (5)

<sup>a</sup> Data are the mean  $\pm$  SD. *n* = no. of controls.<sup>b</sup> ND = not detectable.

tested were not due to variations in the amounts of RNA loaded on the gel, since this was checked in a control experiment with an actin probe.

### Biochemical Studies in the Index Patient

Biochemical measurements of intact cultured skin fibroblasts showed a diminished oxidation rate of <sup>3</sup>H-labeled myristic and palmitic acid (66% and 27%, respectively, of the control mean), as shown in table 1. The oxidation rate of <sup>14</sup>C-labeled octanoate and butyrate was found to be normal (not shown). The activities of the various mitochondrial fatty acyl-CoA dehydrogenases, the enoyl-CoA hydratases, the 3-hydroxyacyl-CoA dehydrogenases, and the 3-ketoacyl-CoA thiolases, as well as of carnitine palmitoyltransferases I and II, were all normal (not shown). As shown in table 1, the activity of CAC was not detectable.

### Mutation Analysis

The CAC mRNA from cultured skin fibroblasts from the patient was reverse transcribed, and the cDNA was PCR amplified in three overlapping fragments. Sequencing of these fragments revealed an insertion of a cytosine in the cytosine-rich region of bp 955-959, as shown in figure 3. This insertion changes the sequence of the CAC protein from amino acid 300 (asparagine to glutamine) to the carboxy terminus and expands the length of the protein by 21 amino acids, to 322 amino acids (fig. 4A).

Hydrophilicity calculations were performed according to the Kyte-Doolittle method, for the peptide sequence of the wild-type and the patient's CAC. The values obtained were used for Chou-Fasman predictions of the secondary structure of the protein. The Chou-Fasman two-dimensional plot (fig. 4B) showed a dramatic conformational change, at the C-terminal region, between the wild-type and the patient's CAC protein.


## Discussion

### Primary Structure of Human CAC


The protein encoded by the human CAC gene is 301 amino acids long, and its calculated molecular weight is

32.9 kD. The amino acid sequences of human and rat CAC are highly conserved: there is ~90% identity between these species (fig. 1) (Indiveri et al. 1997). Human CAC differs from its rat counterpart in 29 amino acids, 15 of which are nonconserved. The identity between the human and rat CAC is less than that found for some other mitochondrial carriers. In fact, 97% identity was found between the human and rat 2-oxoglutarate carriers (Iacobazzi et al. 1992; Dolce et al. 1994b), and 95% identity was found between the human and rat citrate carriers (Kaplan et al. 1993; Iacobazzi et al. 1997) and phosphate carriers (Ferreira et al. 1989; Dolce et al. 1994a). Three repeated homologous domains, each ~100 amino acids in length, can be distinguished in the human CAC, a characteristic previously recognized in other mitochondrial transport proteins (Walker and Runswick 1993; Palmieri 1994). These domains are related to those found in the mitochondrial carrier-protein family (Walker and Runswick 1993; Palmieri 1994).

The cDNA and deduced protein sequences are in GenBank (accession no. Y10319; <http://www.ncbi.nlm.nih.gov/Web/Genbank/index.htm>). A Blast search for



**Figure 3** Sequences of CAC cDNA segments, with their accompanying amino acid codes, for a control (wild type) (A) and the patient (B). For the patient, a C insertion, resulting in a frameshift and an extension of the C terminus of the CAC protein (*underlined*), is shown (*arrow*).



**Figure 4** A, Primary C-terminal amino acid sequence of wild-type (*top*) and patient's (*bottom*) CAC protein. B, Predicted secondary structure, according to the Chou-Fasman algorithm, of wild-type (*top*) and patient's elongated (*bottom*) CAC.

homologous sequences revealed seven different expressed-sequence-tag (EST) clones (AA305590, R11780, F08483, Z28872, AA378439, N77642, and N87428) that originated from different human tissues. These EST clones have sequences that are homologous with different parts of the CAC protein and are presumed to encode a carrier protein of unknown function.

#### *Tissue Distribution of CAC mRNA*

The CAC defect was detected in the patient's fibroblasts. It is likely that CAC deficiency occurs in all tissues, since no evidence for the existence of various tissue-specific isoforms has been found. As shown in figure 2, high levels of CAC mRNA transcripts were found in

heart, skeletal muscle, and liver tissues, which is in fair agreement with the clinical involvement of these tissues in the patient. Much lower levels of expression were found in brain, placental, kidney, and pancreatic tissues and especially in lung tissue.

#### *CAC-Deficient Patient*

We have identified CAC deficiency in cultured skin fibroblasts from a child who survived a stormy neonatal period due to her fatty-acid oxidation disorder, which invariably has been fatal in other patients (Stanley et al. 1992; Pande et al. 1993; Brivet et al. 1994, 1996; Nizzen-Koning et al. 1995; Ogier de Baulney et al. 1995). Direct sequencing of the entire cDNA of the patient re-

vealed the presence of a homozygous insertion of a cytosine nucleotide in the only cytosine-containing region, bp 955–959, resulting in a frameshift. We repeated sequencing of the cDNA of this patient and of three controls, in three independent experiments, all of which revealed the homozygous insertion in the patient's cDNA. This excludes the possibility of a mistake made by the *Taq* DNA polymerase in the PCR reaction. The region containing the insertion consists of five cytosine nucleotides, which makes confirmation of the insertion by restriction-enzyme analysis impossible. Unfortunately, no material from the other family members was available for further investigations.

The CAC protein of the patient has an obviously changed C terminus: amino acids 300 and 301 are changed from asparagine and leucine, respectively, to glutamine and leucine, respectively, and the protein has been elongated by 21 amino acids. A Chou-Fasman prediction shows a dramatically changed secondary structure. The C terminus is changed from a turn (in wild-type CAC) to a helix structure, in the patient. The molecular basis of the transport process mediated by the CAC is still unrevealed. A proper folding and orientation of the CAC protein in the mitochondrial membrane is crucial for adequate functioning. The novel extension in the patient contains a hydrophobic domain, which may be embedded in the mitochondrial inner membrane instead of protruding into the intermembrane space. A diminished entrapping or binding capacity of the positive charge of the quaternary nitrogen of carnitine to the negative carboxylate of the CAC C-terminus can be hypothesized for the patient. The alterations of the patient's CAC also may lead to instability of the protein. An impairment of the patient's CAC in substrate binding or translocation also can be considered. The functional consequences of the molecular defect apparently are restricted, in view of the mild clinical phenotype of the patient. Surprisingly, no CAC activity was detectable in fibroblasts.

In the present study, we determined the sequence of the human cDNA of the mitochondrial CAC, showed differences in human tissue distribution, and defined the first molecular defect in a CAC-deficient patient. This study provides the techniques necessary to resolve the molecular basis of CAC deficiency. This may be of great importance for the reliable diagnosis of patients at risk and also may guide the way toward prenatal diagnosis of this severe type of inborn error in metabolism.

## Acknowledgments

We thank Jos Ruiten, Peter Veenhuizen, and Stef Mentzel for their expert technical and computer assistance. The financial support of Telethon-Italy (grant 985 to F.P.), the Ministero dell'Università e della Ricerca Scientifica e Tecnologica (to F.P.),

the Consiglio Nazionale delle Ricerche (to F.P.), and the Prinses Beatrix Fonds (grant 95-0501 to M.H.) is gratefully acknowledged.

## References

- Brivet M, Slama A, Millington DS, Roe CR, Demaugre F, Legendre A, Boutron A, et al (1996) Retrospective diagnosis of carnitine/acylcarnitine translocase deficiency by acylcarnitine analysis in the proband Guthrie card and enzymatic studies in the parents. *J Inher Metab Dis* 19:181–184
- Brivet M, Slama A, Ogier H, Boutron A, Demaugre F, Saudubray JM, Lemonnier A (1994) Diagnosis of carnitine acylcarnitine translocase deficiency by complementation analysis. *J Inher Metab Dis* 17:271–274
- Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156–159
- Coates PM, Tanaka K (1992) Molecular basis of mitochondrial fatty acid oxidation defects. *J Lipid Res* 33:1099–1110
- Crabeel M, Soetens O, De Rijcke M, Pratiwi R, Pankiewicz R (1996) The ARG11 gene of *Saccharomyces cerevisiae* encodes a mitochondrial integral membrane protein required for arginine biosynthesis. *J Biol Chem* 271:5141–5149
- Dolce V, Iacobazzi V, Palmieri F, Walker JE (1994a) The sequences of human and bovine genes of the phosphate carrier from mitochondria contain evidence of alternatively spliced forms. *J Biol Chem* 269:10451–10460
- Dolce V, Messina A, Cambria A, Palmieri F (1994b) Cloning and sequencing of the rat cDNA encoding the mitochondrial 2-oxoglutarate carrier protein. *DNA Seq* 5:103–109
- Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS (1991) "Touchdown" PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Res* 19:4008
- Ferreira GC, Pratt RD, Pedersen PL (1989) Energy-linked anion transport: cloning, sequencing, and characterization of a full length cDNA encoding the rat liver mitochondrial proton/phosphate symporter. *J Biol Chem* 264:15628–15633
- Iacobazzi V, Lauria G, Palmieri F (1997) Organisation and sequence of the human gene for the mitochondrial citrate transport protein. *DNA Seq* 7:127–139
- Iacobazzi V, Palmieri F, Runswick MJ, Walker JE (1992) Sequences of the human and bovine genes for the mitochondrial 2-oxoglutarate carrier. *DNA Seq* 3:79–88
- Idell-Wenger JA (1981) Carnitine:acylcarnitine translocase of rat heart mitochondria. *J Biol Chem* 256:5597–5603
- Indiveri C, Iacobazzi V, Giangregorio N, Palmieri F (1997) The mitochondrial carnitine carrier protein: cDNA cloning, primary structure and comparison with other mitochondrial transport proteins. *Biochem J* 321:713–719
- Indiveri C, Tonazzi A, Dierks T, Krämer R, Palmieri F (1992) The mitochondrial carnitine carrier: characterization of SH-groups relevant for its transport function. *Biochim Biophys Acta* 1140:53–58
- Indiveri C, Tonazzi A, Giangregorio N, Palmieri F (1995) Probing the active site of the reconstituted carnitine carrier from rat liver mitochondria with sulfhydryl reagents. *Eur J Biochem* 228:271–278
- Indiveri C, Tonazzi A, Palmieri F (1990) Identification and

- purification of the carnitine carrier from rat liver mitochondria. *Biochim Biophys Acta* 1020:81–86
- (1991a) Characterization of the unidirectional transport of carnitine catalyzed by the reconstituted carnitine carrier from rat liver mitochondria. *Biochim Biophys Acta* 1069:110–116
- (1994) The reconstituted carnitine carrier from rat liver mitochondria: evidence for a transport mechanism different from that of the other mitochondrial translocators. *Biochim Biophys Acta* 1189:65–73
- Indiveri C, Tonazzi A, Prezioso G, Palmieri F (1991b) Kinetic characterization of the reconstituted carnitine carrier from rat liver mitochondria. *Biochim Biophys Acta* 1065:231–238
- Kaplan RS, Mayor JA, Wood DO (1993) The mitochondrial tricarboxylate transport protein: cDNA cloning, primary structure and comparison with other mitochondrial transport proteins. *J Biol Chem* 268:13682–13690
- Kelly DP, Strauss AW (1994) Mechanisms of disease. *N Engl J Med* 330:913–919
- Murthy SR, Pande SV (1984) Mechanism of carnitine acylcarnitine translocase-catalyzed import of acylcarnitines into mitochondria. *J Biol Chem* 259:9082–9089
- Niezen-Koning KE, van Spronsen FJ, IJlst L, Wanders RJA, Brivet M, Duran M, Reijngoud DJ, et al (1995) A patient with lethal cardiomyopathy and a carnitine-acylcarnitine translocase deficiency. *J Inherit Metab Dis* 18:230–232
- Ogier de Baulney H, Slama A, Touati G, Turnbull DM, Pourfarzam M, Brivet M (1995) Neonatal hyperammonemia caused by a defect of carnitine-acylcarnitine translocase. *J Pediatr* 127:723–728
- Olpin SE, Manning NJ, Carpenter K, Middleton B, Pollitt RJ (1992) Differential diagnosis of hydroxydicarboxylic aciduria based on release of  $^3\text{H}_2\text{O}$  from [9,10- $^3\text{H}$ ]palmitic acids by intact cultured fibroblasts. *J Inherit Metab Dis* 15:883–890
- Palmieri F (1994) Mitochondrial carrier proteins. *FEBS Lett* 346:48–54
- Palmieri L, Palmieri F, Runswick MJ, Walker JE (1996) Identification by bacterial expression and functional reconstitution of the yeast genomic sequence encoding the mitochondrial dicarboxylate carrier protein. *FEBS Lett* 399:299–302
- Pande SV (1975) A mitochondrial carnitine acylcarnitine translocase system. *Proc Natl Acad Sci USA* 72:883–887
- Pande SV, Brivet M, Slama A, Demaugre F, Aufrant C, Saudubray JM (1993) Carnitine-acylcarnitine translocase deficiency with severe hypoglycemia and auriculo ventricular block. *J Clin Invest* 91:1247–1252
- Pande SV, Murthy MSR (1994) Carnitine-acylcarnitine translocase deficiency: implications in human pathology. *Biochim Biophys Acta* 1226:269–276
- Pande SV, Parvin R (1980) Carnitine-acylcarnitine translocase catalyzes an equilibrating unidirectional transport as well. *J Biol Chem* 255:2994–3001
- Pollitt RJ (1995) Disorders of mitochondrial long-chain fatty acid oxidation. *J Inherit Metab Dis* 18:473–490
- Ramsay RR, Tubbs PK (1975) The mechanism of fatty acid uptake by heart mitochondria: an acylcarnitine-carnitine exchange. *FEBS Lett* 54:21–25
- Stanley CA (1987) New genetic defects in mitochondrial fatty acid oxidation and carnitine deficiency. *Adv Pediatr* 34:59–88
- Stanley CA, Hale DE, Berry GT, DeLeeuw S, Boxer J, Bonfont JP (1992) A deficiency of carnitine-acylcarnitine translocase in the inner mitochondrial membrane. *N Engl J Med* 327:19–22
- Walker JE, Runswick MJ (1993) The mitochondrial transport protein superfamily. *J Bioenerg Biomembr* 25:435–445