

Clinical Trials Study

Clinicopathological and prognostic significance of aberrant Arpin expression in gastric cancer

Tao Li, Hong-Mei Zheng, Nai-Mei Deng, Ying-Jian Jiang, Jiang Wang, Dian-Liang Zhang

Tao Li, Hong-Mei Zheng, Ying-Jian Jiang, Jiang Wang, Dian-Liang Zhang, Center of Colon and Rectum, Qingdao Municipal Hospital, Qingdao University, Qingdao 266011, Shandong Province, China

Nai-Mei Deng, Infection Department, Qingdao Municipal Hospital, Qingdao 266011, Shandong Province, China

Author contributions: Zhang DL designed the research; Li T and Wang J performed the research; Zheng HM, Deng NM and Jiang YJ contributed new reagents and analytic tools; Li T analyzed the data and wrote the paper.

Institutional review board statement: The study was reviewed and approved by the Affiliated Hospital of Qingdao University Institutional Review Board.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors declare no conflict of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Dian-Liang Zhang, MD, PhD, Center of Colon and Rectum, Qingdao Municipal Hospital, Qingdao University, No. 1 Jiaozhou Road, Qingdao 266011, Shandong Province, China. phdzdl@yahoo.com
Telephone: +86-532-82852057
Fax: +86-532-82852058

Received: October 1, 2016

Peer-review started: October 5, 2016

First decision: October 20, 2016

Revised: November 20, 2016

Accepted: January 11, 2017

Article in press: January 11, 2017

Published online: February 28, 2017

Abstract**AIM**

To detect the expression of Arpin, and determine its correlation with clinicopathological characteristics and the prognosis of gastric cancer (GC) patients.

METHODS

A total of 176 GC patients were enrolled as study subjects and classified into groups according to different clinicopathological variables. GC mucosal tissues were obtained *via* surgery. Another 43 paraffin-embedded tissue blocks of normal gastric epithelium (> 5 cm away from the edge of the tumor) were included in the control group. Immunohistochemistry (IHC) for the Arpin and Arp3 proteins was performed on the formalin-fixed, paraffin-embedded GC tissues. Additionally, expression of the Arpin protein in 43 normal gastric tissues was also determined using IHC.

RESULTS

Expression of the Arpin protein in GC was lower than that in normal gastric mucosa (30.68% *vs* 60.47%, $P < 0.001$). A χ^2 test of the 176 GC samples used for IHC showed that decreased Arpin expression was associated with advanced TNM stage ($P < 0.01$) and the presence or absence of lymph node metastasis (80.92% *vs* 35.56%, $P < 0.001$). Additionally, a significant correlation was observed between the expression of Arpin and the presence of the Arp2/3 complex in GC tissues ($\chi^2 = 30.535$, $P < 0.001$). Moreover, a multivariate Cox

regression analysis revealed that Arpin expression [hazard ratio (HR) = 0.551, $P = 0.029$] and TNM stage (HR = 5.344, $P = 0.001$) were independent prognostic markers for overall survival of GC patients. Regarding the 3-year disease-free survival (DFS), the recurrence rate of GC patients with low Arpin expression levels (median DFS 19 mo) was higher than that in the high-Arpin-expression group (median DFS 34 mo, $P = 0.022$).

CONCLUSION

Low Arpin levels are associated with clinicopathological variables and a poor prognosis in GC patients. Arpin may be regarded as a potential prognostic indicator in GC.

Key words: Clinicopathological characteristics; Gastric cancer; Arpin; Arp2/3 complex; Prognosis

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Arpin, a newly found Arp2/3 complex inhibitor reported in Nature, in 2013, was shown to restrict the rate of actin polymerization and control cell migration. However, little is known about whether the expression of Arpin is altered in gastric cancer (GC) tissues, and the detailed mechanisms for invasion and metastasis of GC remain unknown. Our research shows that expression level of Arpin is low in GC, and decreased Arpin expression is associated with the characteristics of clinical pathology and poor prognosis of GC patients. It may be regarded as a potential prognosis indicator for clinical outcomes in GC.

Li T, Zheng HM, Deng NM, Jiang YJ, Wang J, Zhang DL. Clinicopathological and prognostic significance of aberrant Arpin expression in gastric cancer. *World J Gastroenterol* 2017; 23(8): 1450-1457 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i8/1450.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i8.1450>

INTRODUCTION

Gastric cancer (GC) remains the fourth most common malignancy and the third leading cause of cancer-related death worldwide, despite its steadily decreasing incidence and mortality since 1930s^[1,2]. Although the early detection rate of GC has increased, many patients still suffer from distant metastasis, resulting in a median survival of only 3-5 mo^[3-6]. At present, understanding of the multidisciplinary treatment of cancer is a concern of doctors and researchers. However, surgery is still the treatment of choice for most early solid tumors and even some advanced malignant tumors. With the progress made in molecular biology research, accumulating evidence has shown that the carcinogenesis of the gastric mucosa is

a multi-factor, multi-step, and multi-stage development process that involves a variety of related genes. Moreover, stages of GC development are affected by different genes. Therefore, it is necessary to identify novel markers that can accurately reflect the biological characteristics of GC tumors, provide new therapeutic strategies, and predict clinical outcomes.

Invasion and metastasis are the two main characteristics of malignant tumors. In general, invasion and metastasis accompany the movement of cancer cells from the primary focus in cancer tissues to other normal tissues or organs, and actin polymerization is necessary for this movement. Actin, a structural protein composed of actin filaments, exists in two forms, monomers and polymers. The actin polymerization process can be divided into two distinct steps: actin monomer polymerization in the nucleus, followed by the addition of actin monomers to the formed nuclei or fibers. Cell movement results from the mutual cooperation of polymerization and depolymerization of the cytoskeleton itself and adhesion and desorption between different parts of the cell and extracellular matrix. Moreover, the formation of lamellipodia is closely associated with actin polymerization. Lamellipodia play an important role in the process of exploration of the external environment by cancer cells and the formation of new adhesion contacts with the extracellular matrix, which allow motility and spreading. The process of lamellipodia formation is often driven by spatially and temporally regulated actin polymerization at the leading edge^[7]. The molecular mechanisms of cancer cell motility and migration are more complicated than that expected, the movement and migration of cancer cells is a result of multi-step process initiated by the formation of membrane protrusions in response to migratory and chemotactic stimuli. It is generally believed that lamellipodia in driving cancer cell migration plays a main role, and it is caused by adhesion force to pull the cell body forward in the basement membrane. However, many cytokines can form new actin fibers, and each factor can form a specific network. The actin-related protein 2/3 (Arp2/3) complex, the most investigated molecule by far, is the sole machine that generates branched actin networks, and is also considered to be a key regulator of cell motility^[8]. It has been reported to be involved in the development and migration of some cancers, such as pancreatic, gastric, colorectal, and breast cancer^[9-12]. By binding actin filaments to the side of an existing filament and initiating branch formation, the Arp2/3 complex accomplishes its role of nucleating actin filaments^[13]. Therefore, the Arp2/3 complex is thought to be involved in cancer cell invasion and metastasis and is controlled by the tumor-stromal interaction, but the specific guiding mechanism of cell invasion in tumors has not been extensively explored. Relevant research results show that the Arp2/3 complex and its activators, such as the WAVE complex, are deregulated

in diverse cancers. The Arp2/3 complex plays an important role in the invasion and metastasis of tumor cells. However, it is worth noting that the Arp2/3 complex is also vital for the maintenance of normal cell function.

In 2013, Dang *et al.*^[14] reported a new protein, Arpin, which contains 220 amino acid residues that are localized to the cell membrane. It is encoded by the *C15ORF38* gene, which is present in multicellular animals and unicellular amoebas. Interestingly, this protein contains a carboxyl structure region but no C-terminal end, helix, or single actin binding domain. However, it does inhibit the Arp2/3 complex, resulting in the inhibition of actin polymerization. Arpin contains the putative binding site of the Arp2/3 complex^[14], which has been reported to be closely related to the development and migration of cancer cells due to its key role in filopodia initiation^[15,16]. Arpin can guide the direction of cell movement, and it is also known as "the steering factor". However, this process is not dependent on the formation of lamellipodia and is not based on actin dynamics. Arpin acts in a dose-dependent manner and induces the cell to move. Arpin and its acidic motif can compete with nucleation promoting factors for Arp2/3 binding, thereby inhibiting the activation of the Arp2/3 complex^[14,17], resulting in the inhibition of actin filament polymerization. Related research has shown the significant relationship between decreased Arpin, the clinical features of breast cancer, and the poor prognosis of breast cancer patients with low Arpin expression relative to the 5-year relapse-free survival^[18]. However, little is known about the expression of Arpin in GC tissue and its relationship with the clinicopathological characteristics and prognosis of GC patients. In this study, we investigated the issues mentioned above, preliminarily determined the correlation between Arpin and GC as well as between Arpin and the Arp2/3 complex, and provided a theoretical and experimental basis for further research of the specific signaling transduction mechanism of Arpin and the Arp2/3 complex in advanced gastric carcinoma.

MATERIALS AND METHODS

Patients and specimens

The study cohort was composed of samples from 176 patients with gastric adenocarcinoma, including 110 men and 66 women (mean age 56.3 years) who had undergone a gastrectomy at Qingdao Municipal Hospital and the Affiliated Hospital of Qingdao University between March 2013 and August 2013. Following surgery, routine chemotherapy was administered to patients with advanced disease, but no radiation treatment was administered to any of the patients included in the present study. The eligibility criteria for this study included the following: (1) histologically proven adenocarcinoma; (2) no history of gastrectomy

or other malignancy; (3) no other gastric tumors such as gastric stromal tumors; (4) availability of complete clinicopathological and survival data; (5) patients had not received neoadjuvant chemotherapy; and (6) no distant metastasis. This study was approved by the Human Subjects Institutional Committee of the Affiliated Hospital of Qingdao University. All study participants or their legal guardian provided informed written consent prior to study enrollment.

Clinicopathological data

The clinicopathological data were recorded prospectively for the retrospective analysis. The clinicopathological data of the 176 GC patients included age, gender, size of the primary tumor, depth of invasion, lymph node metastasis, TNM stage, degree of differentiation, and location of the primary tumor. The TNM stages of the specimens subjected to immunohistochemistry (IHC) assays were as follows: 70 (39.78%) in TNM stage I, 50 (28.41%) in stage II, and 56 (31.81%) in stage III. Forty-five patients did not have lymph node metastasis, while 131 patients exhibited metastasis. Another 43 paraffin-embedded tissue blocks of normal gastric epithelium (> 5 cm away from the edge of the tumor) were included in the control group. The clinicopathological factors of GC patients are shown in Table 1.

IHC

IHC staining for Arpin was performed in the obtained specimens, which consisted of serial 4- μ m-thick sections of 10% formalin-fixed, paraffin-embedded tissue. After blocking endogenous peroxidase activity to reduce nonspecific binding by boiling in a pressure cooker for 3 min, the samples were placed in 3% hydrogen peroxide (H₂O₂) for 10 min. Then the samples washed with buffer and incubated with a goat polyclonal antibody specific to Arpin (sc-242049; Santa Cruz Biotechnology Inc., Bergheimer, Heidelberg, Germany, final dilution 1:25) at 4 °C for 24 h, followed by three washes in buffer. The secondary antibody consisted of biotinylated anti-goat immunoglobulin. Then, the slides were incubated with avidin-biotin-conjugated peroxidase. After washing and staining with 3,3-diaminobenzidine tetrahydrochloride and H₂O₂, a brown pigment was obtained after counterstaining with hematoxylin.

Because Arp2 exhibits the same distribution as Arp3 in tumorous tissue, we concluded that identical expression of Arp2 and Arp3 indicates the formation of Arp2/3 complex. Therefore, Arp3 IHC staining (Figure 1) represented the distribution of the Arp2/3 complex^[11]. The sections were incubated with an anti-Arp3 polyclonal antibody (diluted to 1:5000). Sections of GC tissue, in which Arp3 was confirmed to be expressed by immunoblot analysis, were used as the positive control for Arp3 immunostaining. For the negative control, normal rabbit serum was substituted

Table 1 Association analysis of immunohistochemistry staining for Arpin vs clinicopathological factors of gastric cancer, *n* (%)

Clinicopathological features	Number of patients (<i>n</i> = 176)	Arpin expression		χ^2	<i>P</i> value
		High (<i>n</i> = 54)	Low (<i>n</i> = 122)		
Age (yr)					
> 50	97 (55.11)	29 (29.90)	68 (70.10)	0.063	0.802
≤ 50	79 (44.89)	25 (31.65)	54 (68.35)		
Gender					
Male	110 (62.50)	31 (28.18)	79 (71.82)	0.865	0.353
Female	66 (37.50)	23 (34.85)	43 (65.15)		
Tumor size (cm)					
> 5	120 (68.18)	37 (30.83)	83 (69.17)	0.004	0.949
≤ 5	56 (31.82)	17 (30.36)	39 (69.64)		
Depth of invasion					
T1 + T2	44 (25.00)	24 (54.55)	20 (45.45)	15.79	0.000
T3 + T4	132 (75.00)	30 (22.73)	102 (77.27)		
Lymph node metastasis					
Yes	131 (74.43)	25 (19.08)	106 (80.92)	32.404	0.000
No	45 (25.57)	29 (64.44)	16 (35.56)		
TNM stage					
I	70 (39.77)	30 (42.86)	40 (57.14)	9.379	0.009
II	50 (28.41)	14 (28.00)	36 (72.00)		
III	56 (31.82)	10 (17.86)	16 (82.14)		
Histology					
Well + moderate	80 (45.45)	21 (26.25)	59 (73.75)	1.354	0.245
Poor	96 (54.55)	33 (34.38)	63 (65.62)		
Tumor site					
Upper	66 (37.50)	24 (36.36)	42 (63.64)	1.648	0.439
Middle	16 (9.09)	4 (25.00)	12 (75.00)		
Low	94 (53.41)	26 (27.66)	68 (72.34)		




Figure 1 Expression of Arp3 by tumor cells. Arp3 is expressed in the invasive cancer cells. Immunohistochemistry of Arp3; original magnification \times 400.

for the primary antibody. Smooth muscle cells of small blood vessels were used as endogenous-negative controls in each section. The secondary antibody consisted of biotinylated anti-goat immunoglobulin.

Immunohistochemical analysis and evaluation

Tumor cells in which the cytoplasm was stained dark brown under light microscopy were considered positive for Arpin IHC staining. Both the staining intensity and the percentage of stained cells were evaluated for the quantification of Arpin expression. Cells with no staining were scored as 0 points, 1 point represented weak staining intensity, 2 points represented moderate

staining intensity, and 3 points represented strong staining intensity. Additionally, we assessed the percentage of stained tumor cells: 0% corresponded to 0 points, less than 25% corresponded to 1 point, 25%-50% corresponded to 2 points, and more than 50% corresponded to 3 points. The final score for Arpin expression was equal to the sum of the two types of scores. A staining score ranging from 0 to 3 points represented low expression, and a score more than 3 points was considered a high expression level^[19].

If more than 10% of the tumor cells expressed Arp3, expression by the tumor cells was considered positive. Otherwise, Arp3 expression was considered negative.

Statistical analysis

SPSS 16.0 software (SPSS, Chicago Statistical IL, United States) was used for the statistical analysis. The differences in clinicopathological variables were analyzed by the χ^2 test. The McNemar test was applied to determine the correlation between expression of Arpin and the Arp2/3 complex. Survival curves for 3-year disease-free survival (DFS) were constructed using the Kaplan-Meier method and compared by a log-rank test. The significance of survival variables was evaluated using a multivariate Cox proportional hazard regression analysis, which further showed the independent effect of Arpin expression on DFS. A value of *P* < 0.05 represented a significant difference.




Figure 2 Representative photomicrographs of Arpin immunohistochemical staining. A: Indicates the high expression in normal gastric tissue; B: Indicates low expression in gastric carcinoma. Original magnification × 200.

Table 2 Immunohistochemistry staining for Arpin in gastric cancer and normal gastric tissues, <i>n</i> (%)					
Group	No. of patients (<i>n</i> = 219)	Arpin expression		χ^2	<i>P</i> value
		High (<i>n</i> = 80)	Low (<i>n</i> = 139)		
Normal	43	26 (60.47)	17 (39.53)	13.221	< 0.001
Cancer	176	54 (30.68)	122 (69.32)		

Table 3 Association analysis of expression of Arpin protein vs TNM stage of gastric cancer, <i>n</i> (%)				
TNM stage	No. of patients	Arpin expression		<i>P</i> value
		High (<i>n</i> = 54)	Low (<i>n</i> = 122)	
I	70 (39.77)	30 (42.86)	40 (57.14)	0.096 ¹
II	50 (28.41)	14 (28.00)	36 (72.00)	0.213 ²
III	56 (31.82)	10 (17.86)	46 (82.14)	0.003 ³

¹Stage I vs II; ²Stage II vs III; ³Stage I vs III.

Table 4 Correlation of Arpin and Arp2/3 complex expressions in gastric cancer patients				
Arp2/3 complex	Arpin		χ^2	<i>P</i> value
	High (<i>n</i> = 54)	Low (<i>n</i> = 122)		
Positive (<i>n</i> = 114)	27	87	30.535	0.000
Negative (<i>n</i> = 62)	27	35		

McNemar $\chi^2 = 30.535, P = 0.000$.

RESULTS

Expression of Arpin protein is decreased in GC tissues

The expression of Arpin in GC tissues was lower than that in normal gastric mucosa (30.68% vs 60.47%, *P* < 0.001, Figure 2 and Table 2).

Relationship between Arpin expression and clinicopathological parameters

Correlations between Arpin expression and clinicopathological characteristics are shown in Table 1. Analysis of the IHC results from 176 tumor samples showed that Arpin protein expression was lower in stage III (82.14%) than in stage I (57.14%) or II (72.00%). Low Arpin protein expression was significantly associated with advanced TNM stage (stage III vs I, *P* < 0.01, Table 3). The protein expression level of Arpin in tumor tissues was significantly correlated with the presence or absence of lymph node metastasis (80.92% vs 35.56%, *P* < 0.001). In addition, we found no significant correlation between the expression of Arpin protein and other clinical parameters, such as tumor location and size, patient age, gender, and histological type.

Correlation between the expression of Arpin and the Arp2/3 complex in GC tissues

The expression levels of Arpin and the Arp2/3 complex in gastric carcinoma were significantly correlated (McNemar $\chi^2 = 30.535, P < 0.001$, Table 4).

Low Arpin expression is associated with a poor prognosis in GC patients

The effects of Arpin expression and clinicopathological characteristics on DFS were evaluated by Kaplan-Meier analysis and log-rank tests. The results showed that GC patients in the low Arpin expression group had a higher recurrence rate (median DFS 19 mo) than those in the high Arpin expression group (median DFS 34 mo, *P* = 0.022, Figure 3). We found that the 3-year DFS was 31.00% in the low Arpin expression group and 46.00% in the high Arpin expression group. Univariate analyses of clinical variables considered as potential predictors of survival are shown in Table 5. Further analysis using a multivariate Cox proportional hazards model showed that Arpin expression, together with TNM stage, was strongly associated with DFS. Arpin expression [hazard ratio (HR) = 0.551, *P* = 0.029] and TNM stage (HR = 5.344, *P* = 0.001) were independent prognostic indicators of DFS in GC




Figure 3 Kaplan-Meier analysis of disease-free survival based on Arpin expression in all 176 patients.

patients (Table 5).

DISCUSSION

To our knowledge, the present study is the first to report the clinical significance of Arpin expression in GC patients, although reduced expression of Arpin and its influence on poor prognosis need to be confirmed in further follow-up studies with larger samples.

In this study, all participants with GC were treated surgically. The sections were obtained from postoperative gastric specimens. In addition, the Arpin and Arp3 IHC analysis was conducted using postoperative gastric slides. The results showed that the Arpin protein was significantly decreased in tumor samples relative to the normal tissues in the control group. Therefore, we speculate that due to the decreased expression of Arpin, tumor cells acquire the ability to invade and metastasize in GC, and the degree of malignancy of tumors is closely related to the clinical stage. We then assessed the correlation between Arpin expression and tumor staging. We found that low Arpin expression was significantly associated with advanced TNM stage, and we concluded that Arpin could play a significant role in tumor biology. In the later stage, lower Arpin expression levels indicate greater malignancy. Whether patients in TNM stage IV, with liver metastasis and peritoneal metastasis are correlated with the expression of Arpin deserves intensive study. Further research may accompany with the expression of Arpin with sensitivity of anti-cancer drug and the correlation of the expression of Arpin with cadherin. Because tumor invasion depth, lymph node metastasis, and distant metastasis are the main pathological basis of malignant tumor staging, we further analyzed the correlation between Arpin expression and local invasion of regional lymph nodes. The IHC results showed that low Arpin expression was significantly associated with the depth of invasion and local lymph node metastases. The results of this research may provide new approaches for determining

Table 5 Univariate and multivariate survival analysis of 3-year DFS in 176 patients with gastric cancer

Variable	Univariate analysis		Multivariate analysis	
	HR	P value	HR	P value
Age	1.14	0.488		
Gender	1.373	0.098		
tumor size	1.411	0.08		
Depth of invasion	1.818	0.017	1.117	0.732
Lymph node metastasis	2.498	0.035	1.413	0.225
tumor site	1.336	0.612		
TNM stage	4.985	< 0.001	5.344	0.001
Arpin expression	0.494	0.005	0.551	0.029

whether loss of Arpin function is associated with increased cell motility in the epithelial-mesenchymal transition during the progression of cancer. Moreover, the inhibitory effect of Arpin on cell migration could potentially be used to control metastasis. However, the tumor size had no significant effect on the test results. Due to the characteristics of the samples and other factors, our results may be different from previous experimental results.

The expression of the Arp2/3 complex was detected in gastric carcinoma tissue in our experiments. Some observations suggest that stromal cells that express Arp2/3 move and grow in advance of the cancer cells to prepare an environment that facilitates cancer cell invasion. Myofibroblasts, which are considered cancer-induced stromal cells, have been shown to affect the adhesion and movement of cancer cells^[19]. Additionally, Arpin inhibits the Arp2/3 complex at the lamellipodium tip. Our research results showed that the expression of Arpin and Arp2/3 complex expression in gastric carcinoma were significantly correlated. Determination of whether a negative correlation exists between these two proteins and the specific mechanism concerning how these proteins affect cell migration will require further study with larger samples. Our experiments may have also indirectly shown a decrease in Arpin expression in GC tissues.

More significantly, in order to investigate the effect of Arpin on the prognosis of GC patients, we compared the 3-year DFS in the Arpin low expression group and the high expression group. Consistently, our study demonstrated that Arpin expression and the TNM stage were independent prognostic indicators of DFS. We propose that Arpin likely plays a role in GC metastasis and the prognosis. However, the exact molecular events leading to cancer metastasis and a poor prognosis have not yet been well elucidated, and further research is required.

The present study had some limitations. We could not visualize the specific signal transduction and regulation mechanism of Arpin and the Arp2/3 complex in GC tissue due to technical difficulties. Arpin is a newly discovered protein, its impact on GC patients and the specific mechanism that it guides cell migration may

help us develop new therapeutic target. The study of mechanism about signal path is the direction of our future research. Thus, more work is required in the future.

In summary, the present study showed that Arpin was decreased in GC tissues. Additionally, this low expression pattern was found to be significantly correlated with aggressive clinicopathological features. In addition, the specific regulatory mechanisms of Rac, WAVE, Arpin and the Arp2/3 complex in the development of GC are still unclear. In recent years, targeted therapeutics for key molecular drivers of cancer progression have been developed^[20]. Arpin may be used as a potential biomarker that could provide important information about tumor progress and could even be a possible target for GC therapy.

COMMENTS

Background

Invasion and metastasis are the two main characteristics of malignant tumors. Gastric cancer (GC), the fourth most common malignancy, and is the third leading cause of cancer-related death worldwide. Arpin, a newly found Arp2/3 complex inhibitor reported in Nature, 2013, which was involved in the development and migration of cancer cells for its key role in the filopodia initiation. Thus, further investigations have been carried out about the role of the Arpin protein in GC.

Research frontiers

Arpin, a newly found Arp2/3 complex inhibitor, simultaneously acts on cell speed and directional persistence, which are strongly coupled parameters. Loss of the Arp2/3 inhibitory protein Arpin produces a similar poor outcome in breast cancer as high expression of the NCKAP1 subunit of the Arp2/3 activatory WAVE complex. Moreover, Arpin downregulation may contribute to the initiation and development of breast cancer metastasis. Therefore, as a potential predictive marker, Arpin deserves future studies.

Innovations and breakthroughs

In this study, the authors for the first time investigated the expression of Arpin in GC tissue and its relationship with clinical pathological characteristics and prognosis of GC patients, preliminarily determined the correlations between Arpin and GC, Arpin and Arp2/3 complex, and provided a theoretical and experimental basis for further research of the specific signal transduction mechanism of Arpin and Arp2/3 complex in advanced gastric carcinoma. The authors could not visualize the specific signal transduction and regulation mechanism of Arpin and Arp2/3 complex in GC due to a lack of technical assistance.

Applications

Arpin protein expression was significantly decreased in tumor samples. Low Arpin associates with clinicopathological variables and poor prognosis in GC patients. Arpin may be regarded as potential prognosis indicator of GC. This provides us a more powerful tool and technical guidance to evaluate the clinicopathologic features and prognosis of patients with GC. Further studies are needed to clarify the detailed mechanisms involved.

Peer-review

The authors detected the expression of Arpin, and evaluated its correlation with clinicopathologic characteristics and prognosis of GC patients. They concluded that low Arpin occurs and associates with clinicopathological characteristics and poor prognosis in GC patients. This is a well written interesting study, which is the first report conducted to determine the value of Arpin in patients with GC.

REFERENCES

- 1 **Torre LA**, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin* 2015; **65**: 87-108 [PMID: 25651787 DOI: 10.3322/caac.21262]
- 2 **Mayer RJ**, Venook AP, Schilsky RL. Progress against GI cancer during the American Society of Clinical Oncology's first 50 years. *J Clin Oncol* 2014; **32**: 1521-1530 [PMID: 24752046 DOI: 10.1200/JCO.2014.55.4121]
- 3 **Jemal A**, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011; **61**: 69-90 [PMID: 21296855 DOI: 10.3322/caac.20107]
- 4 **Lee JH**, Kim KM, Cheong JH, Noh SH. Current management and future strategies of gastric cancer. *Yonsei Med J* 2012; **53**: 248-257 [PMID: 22318810 DOI: 10.3349/ymj.2012.53.2.248]
- 5 **Hartgrink HH**, Jansen EP, van Grieken NC, van de Velde CJ. Gastric cancer. *Lancet* 2009; **374**: 477-490 [PMID: 19625077 DOI: 10.1016/S0140-6736(09)60617-6]
- 6 **Hohenberger P**, Gretschel S. Gastric cancer. *Lancet* 2003; **362**: 305-315 [PMID: 12892963 DOI: 10.1016/S0140-6736(03)13975-X]
- 7 **Yamaguchi H**, Condeelis J. Regulation of the actin cytoskeleton in cancer cell migration and invasion. *Biochim Biophys Acta* 2007; **1773**: 642-652 [PMID: 16926057 DOI: 10.1016/j.bbamcr.2006.07.001]
- 8 **Goley ED**, Welch MD. The ARP2/3 complex: an actin nucleator comes of age. *Nat Rev Mol Cell Biol* 2006; **7**: 713-726 [PMID: 16990851 DOI: 10.1038/nrm2026]
- 9 **Rauhala HE**, Teppo S, Niemelä S, Kallioniemi A. Silencing of the ARP2/3 complex disturbs pancreatic cancer cell migration. *Anticancer Res* 2013; **33**: 45-52 [PMID: 23267127]
- 10 **Kaneda A**, Kaminishi M, Sugimura T, Ushijima T. Decreased expression of the seven ARP2/3 complex genes in human gastric cancers. *Cancer Lett* 2004; **212**: 203-210 [PMID: 15279900 DOI: 10.1016/j.canlet.2004.03.020]
- 11 **Otsubo T**, Iwaya K, Mukai Y, Mizokami Y, Serizawa H, Matsuoka T, Mukai K. Involvement of Arp2/3 complex in the process of colorectal carcinogenesis. *Mod Pathol* 2004; **17**: 461-467 [PMID: 14990971 DOI: 10.1038/modpathol.3800062]
- 12 **Kim DH**, Bae J, Lee JW, Kim SY, Kim YH, Bae JY, Yi JK, Yu MH, Noh DY, Lee C. Proteomic analysis of breast cancer tissue reveals upregulation of actin-remodeling proteins and its relevance to cancer invasiveness. *Proteomics Clin Appl* 2009; **3**: 30-40 [PMID: 21136934 DOI: 10.1002/prca.200800167]
- 13 **Veltman D**. Actin dynamics: cell migration takes a new turn with arpin. *Curr Biol* 2014; **24**: R31-R33 [PMID: 24405676 DOI: 10.1016/j.cub.2013.11.022]
- 14 **Dang I**, Gorelik R, Sousa-Blin C, Derivery E, Guérin C, Linkner J, Nemethova M, Dumortier JG, Giger FA, Chipysheva TA, Ermilova VD, Vacher S, Campanacci V, Herrada I, Planson AG, Fetics S, Henriot V, David V, Oguievetskaia K, Lakisic G, Pierre F, Steffen A, Boyreau A, Peyriéras N, Rottner K, Zinn-Justin S, Cherfils J, Bièche I, Alexandrova AY, David NB, Small JV, Faix J, Blanchoin L, Gautreau A. Inhibitory signalling to the Arp2/3 complex steers cell migration. *Nature* 2013; **503**: 281-284 [PMID: 24132237 DOI: 10.1038/nature12611]
- 15 **Zech T**, Calaminus SD, Caswell P, Spence HJ, Carnell M, Insall RH, Norman J, Machesky LM. The Arp2/3 activator WASH regulates $\alpha 5 \beta 1$ -integrin-mediated invasive migration. *J Cell Sci* 2011; **124**: 3753-3759 [PMID: 22114305 DOI: 10.1242/jcs.080986]
- 16 **Yang C**, Svitkina T. Filopodia initiation: focus on the Arp2/3 complex and formins. *Cell Adh Migr* 2011; **5**: 402-408 [PMID: 21975549 DOI: 10.4161/cam.5.5.16971]
- 17 **Krause M**, Gautreau A. Steering cell migration: lamellipodium dynamics and the regulation of directional persistence. *Nat Rev Mol Cell Biol* 2014; **15**: 577-590 [PMID: 25145849 DOI: 10.1038/nrm3861]
- 18 **Liu X**, Zhao B, Wang H, Wang Y, Niu M, Sun M, Zhao Y, Yao R, Qu Z. Aberrant expression of Arpin in human breast cancer and its clinical significance. *J Cell Mol Med* 2016; **20**: 450-458 [PMID: 26811111 DOI: 10.1111/jcmm.12611]

26648569 DOI: 10.1111/jcmm.12740]

- 19 **Mullins RD**, Stafford WF, Pollard TD. Structure, subunit topology, and actin-binding activity of the Arp2/3 complex from *Acanthamoeba*. *J Cell Biol* 1997; **136**: 331-343 [PMID: 9015304

DOI: 10.1083/jcb.136.2.331]

- 20 **Smith AD**, Roda D, Yap TA. Strategies for modern biomarker and drug development in oncology. *J Hematol Oncol* 2014; **7**: 70 [PMID: 25277503 DOI: 10.1186/s13045-014-0070-8]

P- Reviewer: Aoyagi K, Bang YJ, Gotoda T, Raffaniello R, Yashiro M
S- Editor: Gong ZM **L- Editor:** Ma JY **E- Editor:** Zhang FF





Published by **Baishideng Publishing Group Inc**
8226 Regency Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>
<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045