



Patient Empowerment Programme (PEP) and Risk of Microvascular Diseases Among Patients With Type 2 Diabetes in Primary Care: A Population-Based Propensity-Matched Cohort Study

Diabetes Care 2015;38:e116–e117 | DOI: 10.2337/dc14-2213

Carlos K.H. Wong,¹
William C.W. Wong,¹ Y.F. Wan,¹
Anca K.C. Chan,¹ Frank W.K. Chan,²
and Cindy L.K. Lam¹

The benefit of structured diabetes education programs has been confirmed in systematic review (1) and meta-analysis (2). Although the explicit changes in glycemic control after structured education have been well established, limited evidence (3) has demonstrated the effect of structured education on diabetes-related complications. Recent studies (4,5) underwent an investigation of the effects on glycemic control and incidence of cardiovascular complication in a structured diabetes education program, Patient Empowerment Programme (PEP), versus the usual clinical practice in a Hong Kong primary care setting. Detailed description of PEP setting and mode of education delivery have been reported previously (4,5). However, whether the structured education would be associated with a lower risk of diabetic retinopathy, nephropathy, neuropathy, and composite microvascular complication events remains questionable. The aim of this population-based, propensity score–matched cohort study was to evaluate the influence of PEP implemented in primary care as compared with the alternative usual clinical practice.

This study included 13,280 patients with type 2 diabetes mellitus (T2DM) who attended at least one session of PEP from 1 March 2010 to 30 June 2012 and had no prior diagnosis of

microvascular complication before the baseline date. Each patient was observed from baseline date to whichever following event came first until 31 December 2013—the date of an incidence of a microvascular event, date of death, or last follow-up as censoring. To assess the net effect of PEP, an equal number of 13,280 T2DM patients who had not ever participated in PEP on or before 31 December 2013 were matched to PEP subjects on propensity score matching. Four events were outcomes of interest: 1) first microvascular event with one of the following diagnoses, retinopathy, nephropathy, or neuropathy; 2) retinopathy; 3) nephropathy; and 4) neuropathy. Incidence of microvascular events was identified by the ICPC-2 and ICD-9-CM diagnosis coding systems.

In intention-to-treat analysis, multivariable Cox proportional hazards regression was performed to estimate the effect of PEP on the dependent variable of first microvascular event, accounting for all baseline characteristics of patients. Per-protocol analysis was performed using PEP participants who had completed the program and the propensity-matched non-PEP participants. For each model, survival curves were estimated by Kaplan-Meier method.

Kaplan-Meier survival curves are shown in Fig. 1. As observed, PEP participants generally suffered from fewer

cases of composite microvascular disease and diabetic nephropathy. In intention-to-treat analysis, PEP participants were associated with a lower incidence of first microvascular event (HR 0.85, 95% CI 0.78–0.94, $P = 0.001$) and nephropathy (HR 0.71, 95% CI 0.62–0.80, $P < 0.001$) than non-PEP participants, after adjusting for confounding variables. Similar findings were obtained for those PEP participants with the completion of the program in per-protocol analysis.

Enrollment in PEP that enhances both knowledge and self-care of diabetes was associated with decreased microvascular complication events. Program completion was related to the reduction in microvascular complication events, mainly driven by diabetic nephropathy. Findings of this propensity-matched cohort study add to the growing body of literature supporting structured diabetes education as one of the essential and effective components in multifactorial intervention for T2DM in primary care setting.

Acknowledgments. The authors would like to thank the program teams at the Hospital Authority head office (including Margaret Tay, Dr. K.L. Chung, Dr. Alexander Chiu, Dr. Alan Y.L. Kwok, Michelle Wong, and Gary Ching) and all cluster representatives and clinical staff of the PEP for working with the team on this evaluation. The authors also would like to thank

¹Department of Family Medicine and Primary Care, The University of Hong Kong, Ap Lei Chau, Hong Kong

²Integrated Care Programs, Hospital Authority Head Office, Hong Kong Hospital Authority, Kowloon, Hong Kong

Corresponding author: Carlos K.H. Wong, carlosho@hku.hk.

Clinical trial reg. no. NCT01935349, clinicaltrials.gov.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.




Figure 1—Kaplan-Meier survival curves of the composite microvascular complication, retinopathy, nephropathy, and neuropathy. Log-rank test compared the difference between PEP and non-PEP groups.

Dr. S.V. Lo and the staff of the Statistics & Workforce Planning Department (including Edwina Shung, Peggo Lam, Dr. Kelvin Tsoi, Adam Ng, C.F. Yiu, and their teammates) in the Hospital Authority Strategy and Planning Division for their help in coordinating the development of the evaluation frameworks and site visits and facilitating the data collection.

Funding. This study has been funded by the Hong Kong Hospital Authority (ref. no. 8011014157), the Health and Health Services Research Fund, Food and Health Bureau, and HKSAR Commissioned Research on Enhanced Primary Care Study (ref. no. EPC-HKU-2).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. C.K.H.W. wrote the manuscript and researched data. W.C.W.W.

and C.L.K.L. contributed to study design and reviewed and edited the manuscript. Y.F.W. and A.K.C.C. reviewed and edited the manuscript and contributed to statistical analysis and interpretation of results. F.W.K.C. contributed to acquisition of data and reviewed and edited the manuscript. C.K.H.W. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Loveman E, Frampton GK, Clegg AJ. The clinical effectiveness of diabetes education models for type 2 diabetes: a systematic review. *Health Technol Assess* 2008;12: 1–116.iii
2. Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM. Self-management education for

adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care* 2002;25:1159–1171

3. Ko SH, Song KH, Kim SR, et al. Long-term effects of a structured intensive diabetes education programme (SIDEPE) in patients with type 2 diabetes mellitus—a 4-year follow-up study. *Diabet Med* 2007;24:55–62

4. Wong CKH, Wong WCW, Lam CLK, et al. Effects of Patient Empowerment Programme (PEP) on clinical outcomes and health service utilization in type 2 diabetes mellitus in primary care: an observational matched cohort study. *PLoS One* 2014;9:e95328

5. Wong CKH, Wong WCW, Wan YF, et al. Patient Empowerment Programme in primary care reduced all-cause mortality and cardiovascular diseases in patients with type 2 diabetes mellitus: a population-based propensity-matched cohort study. *Diabetes Obes Metab* 2015;17: 128–135