

## Intravenous device associated blood stream staphylococcal infection in paediatric patients

Amita Jain, Astha Agarwal, Raj Kumar Verma, Shally Awasthi\* & K.P. Singh

Departments of Microbiology & \*Pediatrics, Chhatrapati Shahuji Maharaj Medical University, Lucknow, India

Received May 29, 2009

**Background & objectives:** Intravenous device (IVD) associated nosocomial blood stream infections due to staphylococci are major cause of morbidity and mortality. The present study was carried out to assess the frequency of staphylococcal IVD associated infections in a paediatric ward of a tertiary care hospital. Prevalence of resistance to commonly used antimicrobials in hospital acquired staphylococcal isolates was also tested.

**Methods:** Children admitted in paediatric wards with IVD for more than 48 h were enrolled. Blood, IVD tip at the time of removal, skin swab at the site of insertion of IVD and nasal swab were collected and cultured by standard protocol. All staphylococcal isolates from any source were analyzed for antimicrobial susceptibility by disk diffusion method. Genotyping matching of those staphylococcal isolates was done which were isolated from different sites of the same patient, but were phenotypically similar. Genotype of blood isolate was compared with genotype of isolate from nose/IVD/skin.

**Results:** *Staphylococcus aureus* was the most frequent blood isolate (8.7%) followed by *Candida* (2.9%), coagulase negative staphylococci (CoNS 2.6%), *Pseudomonas* spp. (0.4%), *Klebsiella* spp. (0.3%) and *Escherichia coli* (0.1%). Isolation of microorganisms from blood was significantly higher in patients whose skin, IVD and nose were colonized by same microorganism ( $P<0.001$ ). None of the staphylococcal isolate was found to be resistant to glycopeptides (vancomycin and teicoplanin). High penicillin and oxacillin resistance was present in both *S. aureus* (penicillin resistance; 76.8%, oxacillin resistance; 66.7%) and CoNS (penicillin resistance; 73.3%, oxacillin resistance; 60.0%). Among CoNS biotypes, *S. haemolyticus* was commonest blood isolate while *S. epidermidis* was commonest isolate from Skin/nose. Only 33.3 per cent of *S. aureus* blood stream infections and most of *S. epidermidis* and *S. haemolyticus* blood infections were IVD associated.

**Interpretation & conclusions:** Staphylococci were the major causative agent of nosocomial blood stream infections. All episodes of septicaemia due to *S. epidermidis* and *S. haemolyticus* were IVD associated while only 1/3 of *S. aureus* septicaemia was IVD associated.

**Key words** Coagulase negative staphylococci - hospital acquired septicaemia - IVD associated staphylococcal septicaemia - *Staphylococcus aureus*

Nosocomial blood stream infections (BSIs) have been identified as one of the most frequent nosocomial infections in paediatric patients<sup>1-3</sup>. The

risk of nosocomial infections depends on the host characteristics, the number of interventions, invasive procedures, asepsis of techniques, the duration of stay

in the hospital and inappropriate use of antimicrobials. Most often the endogenous flora of the patient, which may be altered because of hospitalization, is responsible for nosocomial infections<sup>4</sup>. Many microorganisms are responsible for nosocomial septicaemia, commonest being Gram-positive cocci primarily staphylococci. These are commensal on human body surfaces and colonize intravenous devices, which become a focus of infection in hospitalized individuals more so in immuno-compromised patients. Colonizing microorganisms may enter into blood stream and cause septicaemia<sup>5</sup>. The emergence of antibiotic resistant strains of staphylococci is considered a major problem in most hospitals. Large proportions of nosocomial staphylococcal strains are resistant to penicillin. Centres for Disease Control and Prevention reported that the methicillin resistant staphylococcal strains are resistant to several other antibiotics in both large and small hospitals<sup>6</sup>.

We have noticed that staphylococci were the commonest cause of nosocomial septicaemia in our paediatric ward (unpublished observation). Therefore, the present investigation was planned to study the frequency of staphylococcal intravenous device (IVD) associated infections in children admitted in paediatric ward of a tertiary care hospital in north India. Percentage of resistance to commonly used antimicrobials in our hospital acquired staphylococcal strains was also tested.

### Material & Methods

The study was done in Chhatrapati Shahuji Maharaj Medical University (CSMMU), a tertiary care hospital at Lucknow, Uttar Pradesh, from January 2003 to January 2006. All consecutive children admitted in paediatric wards with IVD in peripheral vein for more than 48 h, were followed up and if they developed fever (>99 °F) or had >1°F rise in existing body temperature, they were enrolled. From each patient, venous blood, IVD tip at the time of removal, skin swab from the site of IVD insertion and nasal swab were collected for culture. Skin swabs were collected from the site of catheter insertion by rolling the wet swab (dipped in sterile saline) and were transported into screw-capped tube. Nasal swab were collected from anterior nasal chamber by rolling the swab and transported into screw-capped tube. Two ml of venous blood was collected from each child and aseptically transported into screw-capped blood culture bottle containing 10 ml brain heart infusion (BHI) broth with 0.05 per cent sodium polyanethol sulphonate (SPS). Tip of each catheter (1.5

inches) was clipped with a sterile scissor and taken into a sterile screw capped vial. All the samples were sent to the Microbiology laboratory of CSMMU without delay for culture. The study protocol was approved by the ethics committee and informed consent was taken from guardian of each patient.

*Culture of the samples:* Skin and nasal swabs were inoculated on to 5 per cent sheep blood agar (SBA). All the catheter tips were processed on 5 per cent sheep blood agar by the blood agar roll techniques for semi-quantitative culture by the method of Maki *et al*<sup>7</sup>. A colony count of > 15 on SBA was considered significant and further processed. All the inoculated plates were incubated aerobically at 37°C for 24 h; if plates showed no growth after 24 h these were further incubated for 24 h. The blood containing brain heart infusion broth was incubated overnight at 37°C and subcultures were done on 5 per cent sheep blood agar. Inoculated plates were incubated aerobically at 37°C for 24 h. In case first culture was sterile, final subculture was done on appearance of turbidity or on 7<sup>th</sup> day of incubation, whichever was earlier.

*Identification:* Smears were prepared from plates showing growth and stained by Gram's stain. Gram-positive cocci in cluster were tested by modified oxidase, catalase and slide and tube coagulase tests. Oxidase negative, catalase positive and coagulase positive strains were *Staphylococcus aureus*. Oxidase negative, catalase positive and coagulase negative strains were coagulase negative staphylococci (CoNS). CoNS were further biotyped for species identification by using Kloos & Bannerman scheme<sup>8</sup>.

*Antibiotic susceptibility test (AST):* AST of all staphylococcal isolates from all the sources was performed as per Clinical Laboratory Standards Institute (CLSI) guidelines<sup>9</sup>, using Muller-Hinton agar (MHA) (Hi-media, Mumbai, India). The following antibiotic discs, procured from Hi-media (Mumbai) were used; penicillin (10 U), oxacillin (1 µg), vancomycin (30 µg), teikoplanin (30 µg), cefazolin (30 µg), ciprofloxacin (30 µg). A standardized inoculum matching 0.5 McFarland density gradient was swabbed onto the surface of agar plate three times, rotating plate 60 degrees to ensure even distribution. After 5 min of inoculation, discs were applied to plate by sterile forceps and incubated at 37°C for 24 h. A report of resistant, intermediate or sensitive was recorded as per CLSI recommendations<sup>9</sup>. Standard strain of *S. aureus* ATCC 29213 was used as a control strain.

**Genotyping of isolates:** Staphylococcal isolates from those patients whose blood culture was positive for staphylococci were subjected to genotyping<sup>10</sup>. Genotype of blood isolate was compared with genotype of isolate from nose/IVD/skin. Genotyping by random amplification of polymorphic DNA (RAPD)<sup>10</sup> was performed only if blood isolate was phenotypically similar (biotyping, AST pattern) to isolate from any other site (IVD/nose/skin).

**Extraction of DNA:** Briefly, four to five discrete colonies of *Staphylococcus* sp. were suspended in 200 µl lysis buffer (Tris 10 mM, EDTA 2 mM, NaCl 0.4 M and Triton X-100 0.5%) and boiled for 20 min. Ten µg of proteinase K was added. The sample was vortexed and incubated at 56°C for 2 h followed by boiling at 100°C for 10 min to inactivate proteinase K. DNA purification was done by addition of equal volume of phenol: chloroform (24:1) followed by chloroform only. Aqueous phase was finally transferred in 2.5 volume of ethanol and 10 µl sodium acetate (0.3 M final conc.) was added. The tubes were kept at 20°C overnight. The sample was centrifuged at 10,000 g for 10 min and the pellet was washed with chilled 70 per cent ethanol. The pellet was allowed to air dry and finally suspended in 25 µl of sterile triple distilled water for RAPD analysis<sup>11</sup>.

**RAPD analysis:** Three different random amplification primers were used (primer 10265'-TACATTCGAGGACCCCTAAGTG-3', primer 12045'-ATGTAAGCTCCTGGGGATTAC-3' and primer 10145'-AAGTAAGTGACTGGGGTGAGCG-3')<sup>10</sup>. PCR mixture consisted of 10 mM Tris HCl (pH 9.0), 50 mM KCl, 2.5 mM MgCl<sub>2</sub>, 0.01% gelatin, 100 pico moles of each primer and triton X-100 (0.01%). Deoxyribonucleotide triphosphates (Bangalore Genei,

India) were used at a final concentration of 0.2 mM. For each reaction, 0.5 U *Taq* DNA polymerase (Bangalore Genei, India) was added. PCR consisted of 40 cycles of consecutive denaturation, annealing and DNA chain extension (1 min at 94°C, 1 min at 25°C, 2 min at 74°C) in a thermocycler (Techne Progene, USA). Amplicons were analyzed on 1.5 per cent agarose gel electrophoresis and photographed<sup>10</sup>. Amplified products were analyzed on 1.5 per cent agarose gel electrophoresis. Isolates showing 100 per cent similar bands were considered identical.

**Statistical analysis:** Statistical analysis was done by Chi-square test for significance of difference in isolation of microorganisms from blood (septicaemia) of the patients whose skin, IVD and nose was colonized by same microorganism. The data were analyzed by using SPSS software 'version 10' on windows XP. *P*<0.05 was considered significant<sup>12</sup>.

## Results

Total samples collected from 1399 hospitalized children were, 1141 venous blood, 1244 IVD, 1396 skin swabs and 1397 nasal swabs. *S. aureus* was the most frequent isolate; a total of 8.7 per cent venous blood (99/1141), 16.5 per cent nasal swab (231/1397) 13.6 per cent skin swab (190/1396) and 13 per cent catheter tip (162/1244) were positive for *S. aureus* (Table I). Coagulase negative staphylococci (CoNS) were second commonest bacterial isolate; *S. haemolyticus* was the most frequent isolate from blood and IVD while *S. epidermidis* was commonest from skin and nasal swab. The detailed distribution of CoNS species in different samples is shown in Table II. *Candida* was second most frequent isolate followed by CoNS, *Pseudomonas* spp., *Klebsiella* spp., and *Escherichia*

**Table I.** Isolation rate of microorganisms from hospitalized children with clinical suspicion of hospital acquired septicemia

Specimen	Number of specimens collected	Staphylococci			Microorganisms other than staphylococci			
		SA	CoNS	<i>Candida</i>	<i>Acinetobacter</i>	<i>Pseudomonas</i>	<i>Klebsiella</i>	<i>Escherichia coli</i>
Blood	1141	99 (8.7)	30 (2.6)	33 (2.9)	00	3 (0.4)	2 (0.3)	1 (0.1)
IVD	1244	162 (13.0)	98 (7.9)	88 (7.1)	11 (1.2)	8 (0.7)	5 (0.4)	6 (0.2)
Skin swab	1396	190 (13.6)	202 (16.1)	29 (2.1)	20 (1.8)	3 (0.4)	3 (0.2)	1
Nasal swab	1397	231 (16.5)	207 (14.8)	11 (0.8)	8 (0.7)	3 (0.3)	2 (0.1)	00

SA, *Staphylococcus aureus*; CoNS, coagulase negative staphylococci. Values in parenthesis denote percentage

*coli* (Table I). Significance of difference in isolation of microorganisms from blood (septicaemia) of the patients whose skin, IVD and nose was colonized by same microorganism was studied and it was seen that isolation of microorganisms from blood (septicaemia) was significantly higher in those patients whose skin/IVD and/or nose was colonized by same microorganism ( $P < 0.001$ ).

**Antimicrobial resistance (AMR) pattern:** Antimicrobial sensitivity testing of *S. aureus* (n=682) and CoNS (n=537) isolates was done. None of the isolate was found to be resistant to glycopeptides (vancomycin and teicoplanin). Penicillin resistance was high in both *S. aureus* 86.2 per cent (588/682) and CoNS 80.6 per cent (433/537). A total of 76.7 per cent of *S. aureus*

isolates were resistance to oxacillin, 61.5 per cent to cefazolin and 31.6 per cent to ciprofloxacin (Table III). Among CoNS isolates, 63.1 per cent were resistant to oxacillin, 47.8 per cent to cefazolin and 25.5 per cent to ciprofloxacin.

**RAPD analysis:** RAPD assay was performed to analyze the similarity among the staphylococcal isolates from different specimens of the same patient. Staphylococcal isolates (*S. aureus*, *S. epidermidis* and *S. haemolyticus*) from those subjects whose blood cultures were positive for phenotypically similar staphylococci (similar biotype and AMR pattern) were studied by RAPD. Of the 99 patients whose blood culture was positive for *S. aureus*, 34 were also having IVD culture positive for phenotypically similar *S. aureus*, of which 33

**Table II.** Distribution of coagulase negative staphylococci (CoNS) species in different samples collected from children with clinical suspicion of hospital acquired septicaemia

CoNS biotypes	Blood	IVD tip	Skin swab	Nasal swab
<i>Staphylococcus haemolyticus</i>	17 (56.7)	36 (36.7)	33 (16.3)	42 (20.3)
<i>S. epidermidis</i>	5 (16.7)	23 (23.5)	68 (33.7)	69 (33.3)
<i>S. xylosum</i>	4 (13.3)	13 (13.3)	32 (15.8)	19 (9.2)
<i>S. captis</i>	1 (3.3)	12 (12.2)	17 (8.4)	14 (6.8)
<i>S. cohnii</i>	2 (6.7)	8 (8.2)	13 (6.4)	24 (11.6)
<i>S. saprophyticus</i>	0	0	26 (12.9)	18 (8.7)
<i>S. simulans</i>	0	1 (1.0)	0	3 (1.4)
<i>S. hominis</i>	1 (3.3)	3 (3.0)	2 (0.9)	6 (2.9)
<i>S. warneri</i>	0	0	0	1 (0.5)
Non-typable	0	1 (1.0)	11 (5.4)	11 (5.3)
Total isolates/specimen	30/1141	98/1244	202/1396	207/1397

Values are given as no. (%)

**Table III.** Antimicrobial resistance pattern of *S. aureus* and CoNS isolated from different clinical samples from children with clinical suspicion of hospital acquired septicaemia

Antibiotics	% resistance <i>S. aureus</i> (n=682)				
	Venous blood (n=99)	Intravenous device tip (n=162)	Skin swab (n=190)	Nasal swab (n=231)	Total (n=682)
Penicillin	76 (76.8)	141 (87.0)	171 (90.0)	200 (86.6)	588 (86.2)
Oxacillin	66 (66.7)	123 (75.9)	154 (81.0)	180 (77.9)	523 (76.6)
Cefazolin	53 (53.5)	99 (61.1)	121 (63.7)	147 (63.6)	420 (61.5)
Ciprofloxacin	19 (19.2)	52 (32.1)	56 (29.5)	89 (38.5)	216 (31.6)
	% resistance CoNS (n=537)				
	Venous blood (n=30)	Intravenous device tip (n=98)	Skin swab (n=202)	Nasal swab (n=207)	Total (n=537)
Penicillin	22 (73.3)	85 (86.7)	155 (76.7)	171 (82.6)	433 (80.6)
Oxacillin	18 (60.0)	68 (69.4)	116 (57.4)	137 (66.2)	339 (63.1)
Cefazolin	14 (46.7)	44 (44.9)	90 (44.5)	109 (52.7)	257 (47.8)
Ciprofloxacin	9 (30.0)	26 (26.5)	45 (22.3)	57 (27.5)	137 (25.5)

% resistant (including both resistant and intermediate level of resistance). No glycopeptides (vancomycin, teicoplanin) resistance was seen. Value are given as no. (%)

**Table IV.** RAPD analysis result of common isolates from children with clinical suspicion of hospital acquired septicaemia

Organisms (no. of positive blood cultures)	Samples having results similar to blood culture	No. of patients positive for phenotypically similar organism*	No of patients having similar genotype	% similarity among isolates from same patients
<i>S. aureus</i> (n=99)	IVD	34	33	33.3
	SS	29	25	25.2
	NS	35	16	16.1
<i>S. epidermidis</i> (n=5)	IVD	5	5	100
	SS	2	2	40
	NS	1	1	20
<i>S. haemolyticus</i> (n=17)	IVD	17	17	100
	SS	15	12	70.5
	NS	15	6	35.2

IVD, intravenous devices; SS, skin swab; NS, nasal swab. \*Genotype of all these isolates was matched with corresponding blood isolate

blood and IVD isolates were also genetically similar; showing that 33.3 per cent episodes of *S. aureus* blood stream infections are IVD associated. Twenty nine patients were also having skin swabs positive for phenotypically similar *S. aureus*, of which 25 were also genetically similar to blood isolates. Thirty four patients were having nasal swabs positive for phenotypically similar *S. aureus*, of which 16 strains were genotypically identical to blood isolate (Table IV). Five patients had *S. epidermidis* isolated from their blood. All of them had genotypically similar isolate from IVD while only 2 had same isolate from skin and only one had same isolate from nose. Of the 17 patients whose blood culture was positive for *S. haemolyticus*, all had genotypically similar isolate from IVD. Only 12 of them had same isolate from skin and only 6 of them had same isolate from nose (Table IV).

### Discussion

Nosocomial blood stream infections (BSI) are one of the most serious and potentially life threatening infectious diseases in paediatric patients. Early diagnosis and therapy are essential for the prevention of morbidity and mortality<sup>13</sup>. The accurate prediction of likely pathogens and antimicrobial resistance pattern is crucial for successful therapy. In the present study, *S. aureus* was found to be the most frequent blood isolate followed by *Candida*, CoNS, *Pseudomonas* spp., *Klebsiella* spp. and *E. coli*. Among CoNS, *S. haemolyticus* was commonest blood isolate followed by *S. epidermidis*. Berner *et al*<sup>14</sup>, retrospectively analyzed 1037 bacteraemic episodes in children in a German tertiary care center during a 10 years period and noted that Gram-positive bacteria accounted for two third of all bacteraemic episodes in paediatric patients. In

another study<sup>13</sup> 131 episodes of blood stream infection in a paediatric ICU in the UK in a 3 year period were studied. They found that Gram-positive and Gram-negative bacteria accounted for 63 and 31 per cent respectively, 6 per cent were yeast. US medical centers published the data on blood stream infection in 1991 and 1997 focusing on paediatric patients<sup>14</sup>. The key findings were that significant increase in the overall blood stream infection rate was caused by each of the following pathogens groups: CoNS, *S. aureus*, enterococci and *Candida* species. In contrast, the blood stream infection rate caused by Gram-negative bacilli remained stable during the decade. They reported that the greatest increase in blood stream infection rates was observed in coagulase negative staphylococci followed by *Candida* spp. *S. aureus* colonization varied from 8.0 to 71.2 per cent. *Candida* isolation reported from various specimens varies from 3.2 to 26.9 per cent<sup>15,16</sup>. Studies done during 1986 to 2002 reported that the ratio of *S. aureus* to CoNS was around 1:1<sup>17,18</sup> and majority of nosocomial CoNS isolates were obtained from blood samples, catheters and wounds indicating that CoNS have increasingly been recognized as important agents of nosocomial infection. *S. epidermidis* and *S. haemolyticus* were the most responsible CoNS species for septicaemia in paediatric patients<sup>19</sup>, while in another study *S. epidermidis* was reported most frequent blood isolate followed by *S. haemolyticus*<sup>20</sup>.

In our study none of the staphylococcal isolates was found to be resistant to glycopeptides (vancomycin and teicoplanin) while high penicillin and oxacillin resistance was present in both *S. aureus* and CoNS. Silvia *et al*<sup>21</sup> have also reported reduced susceptibility to glycopeptides in 5.4 per cent of CoNS, isolated from blood samples of critically ill haematology patients.

Reduced susceptibility to teicoplanin in blood CoNS isolates was also shown<sup>22</sup>. Resistance to methicillin among *S. aureus* increased from 1.5 per cent in 1986 to 31.2 per cent in 2002<sup>18</sup>. This also implies an increase in resistance to other antimicrobials such as macrolides, lincosamides, aminoglycosides, and quinolones. All *S. aureus* isolates have remained uniformly susceptible to glycopeptides and novel antimicrobials (linezolid, and quinupristin/dalfopristin)<sup>18</sup>.

With regard to CoNS, the increase in resistance to methicillin was even greater, reaching to 61.3%. In most cases, this was associated with resistance to multiple antimicrobials<sup>18</sup>. Antimicrobial resistance pattern of staphylococci varied from study to study. High resistance to penicillin (80 - 90.6% in CNS & 78.6 - 88.9% in *S. aureus*), oxacillin (20.8 - 98.4% in CNS & 24.5 - 87.2% in *S. aureus*), cefazolin (2.7 - 80% in CNS & 1.1 - 23.7% in *S. aureus*) and ciprofloxacin (11 - 66% in CNS 13.1 - 25% in *S. aureus*) was reported<sup>2,23-25</sup>. In the present study, cefazolin resistance was slightly lower than oxacillin resistance. Clinically all oxacillin resistant strains are resistant to all betalactams and should not be treated with penicillins or cephalosporins. All strains showing discrepancy in results were tested by us for detection of *mecA* gene and other phenotypic methods<sup>28</sup>. It was noted that oxacillin disc diffusion test is less specific than *mec A* detection method. Except occasional reports of vancomycin resistant staphylococci, maximum number of studies had reported 0 per cent prevalence of vancomycin resistant staphylococci<sup>23,26,27</sup>. Difference in (i) prescription practices in the hospitals<sup>28</sup>, (ii) geographical locations<sup>2,29,3</sup>, (iii) methods of AMR testing<sup>29,30</sup>, (iv) sample size<sup>24,31</sup>, (v) types of clinical samples are different reasons for these variations<sup>3,31</sup>.

It was found that 1/3 of *S. aureus* blood isolates and 100 per cent of CoNS blood isolates were identical to IVD isolate. Gemme *et al*<sup>32</sup> reported that in young infants 65 per cent of CoNS blood isolates were identical to catheter while Nataro *et al*<sup>33</sup> reported that 42 per cent of the staphylococcal blood isolated from hospitalized infants were genotypically identical to catheter isolates. Another study reported that 64 per cent paediatric patients with CoNS in their blood had same bacteria present on their catheter tips<sup>4</sup>. Similarity of skin and nose isolates to that of blood isolates was not that frequent, varying between 16-70 per cent in our study. This implies that the source of at least some of the blood isolates in hospitalized paediatric patients is not endogenous.

In conclusion, staphylococci were found to be the major causative agent of nosocomial blood stream infections in paediatric population. Penicillin and oxacillin resistance was high in these isolates. Only 33.3 per cent of *S. aureus* blood infections and majority of CoNS (*S. haemolyticus* and *S. epidermidis*) blood infections were IVD associated.

### Acknowledgment

Authors acknowledge the Indian Council of Medical Research, New Delhi, for financial support.

### References

- Piette A, Verschraegen G. Role of coagulase-negative staphylococci in human disease. *Vet Microbiol* 2009; 134 : 45-54.
- Altıparmak MR, Güngör K, Pamuk GE, Pamuk ON, Ozgenç R, Oztürk R. Temporary Catheter infections in hemodialysis patients: results from a single center in Turkey. *Acta Clin Belg* 2003; 58 : 345-9.
- Leibovitch I, Lai TF, Senarath L, Hsuan J, Selva D. Infectious keratitis in South Australia: Emerging resistance to cephazolin. *Eur J Ophthalmol* 2005; 15 : 23-6.
- Lodha R, Natchu UC, Nanda M, Kabra SK. Nosocomial infections in pediatric intensive care units. *Indian J Pediatr* 2001; 68 : 1063-70.
- Jain A, Agarwal A. Biofilm production, a marker of pathogenic potential of colonizing and commensal staphylococci. *J Microbiol Methods* 2009; 76 : 88-92.
- CDC. National Nosocomial Infections Surveillance (NNIS) System Report, Data Summary from January 1990-May 1999, issued June 1999. *Am J Infect Control* 1999; 27 : 520-32.
- Maki DG, Weise CE, Sarafin HW. A semi quantitative culture method for identifying intravenous-catheter-related infection. *N Engl J Med* 1977; 296 : 1305-9.
- Kloos WE, Bannerman YL. Staphylococcus and micrococcus. In: Murry RP, Baron EJ, Pfaller MA, Tenoer MC, Tenover MC, Tenover RH, editors. *Manual of clinical microbiology*, 7<sup>th</sup> ed. Washington, DC: American Society of Microbiology; 1999. p. 262-82.
- Clinical Laboratory Standards Institute (CLSI). *Performance standards for antimicrobial susceptibility testing*. 15<sup>th</sup> Informational Supplement testing M100-S15, Wayne (PA): CLSI; 2005.
- Burnie JP, Naderi-Nasab M, Loudon KW, Matthews RC. An epidemiological study of blood culture isolates of coagulase negative staphylococci demonstrating hospital acquired infection. *J Clin Microbiol* 1997; 35 : 1746-50.
- Jaffe RI, Lane JD, Albury SV, Niemeyer DM. Rapid extraction from and direct identification of clinical samples of methicillin resistant staphylococci using the PCR. *J Clin Microbiol* 2000; 38 : 3407-12.
- Greenwood PE, Nikulin MS. *A guide to chi-squared testing*. New York: Wiley; 1996.
- Gray J, Gossain S, Morris K. Three year survey of bacteraemia and fungemia in a pediatric intensive care unit. *Pediatr Infect Dis J* 2001; 20 : 416-21.

14. Berner R, Sauter S, Duffner U, Brandis M, Niemeyer CM. Bacteremic episodes in pediatric oncologic patients, especially caused by the *Streptococcus viridans* group. *Clin Padiatr* 1998; 210 : 256-60.
15. Kuehnert MJ, Webb RM, Jochimsen EM. *Staphylococcus aureus* blood stream infections among patients undergoing electroconvulsive therapy traced to breaks in infection control and possible extrinsic contamination by propofol. *Anesth Analg* 1997; 85 : 420-5.
16. Glowacki M, Quraishi ZA, Zakhireh B. Risk factors of nosocomial bacteremia associated with pulmonary artery catheters in a critical care unit. *J Am Osteopath Assoc* 1990; 90 : 509-14.
17. Chlebicki MP, Teo EK. Review of peripherally inserted central catheters in the Singapore acute-care hospital. *Singapore Med J* 2003; 44 : 531-5.
18. Cuevas O, Cercenado E, Vindel A, Guinea J, Sanchez-Conde M, Sanchez-Somolinos M, *et al*. Evolution of the antimicrobial resistance of *Staphylococcus* spp. in Spain: Five Nationwide Prevalence Studies 1986 to 2002. *Antimicrob Chemother* 2004; 48: 4240-5.
19. Jog SM, Patole SK. *Staphylococcus warneri* septicemia in preterm neonates-A reminder. *Indian Padiatr* 2002; 39 : 309-10.
20. Mohan U, Jindal N, Aggarwal P. Species distribution and antibiotic sensitivity pattern of coagulase negative staphylococcal isolates from various clinical specimens. *Indian J Med Microbiol* 2002; 20 : 45-6.
21. Silvia N, Carla F, Marco F, Alberto B, Piero TG, Silvia M, *et al*. Characterization of coagulase-negative staphylococcal isolates from blood with reduced susceptibility to glycopeptides and therapeutic options. *BMC Infect Dis* 2009; 9 : 83-6.
22. Del' Alamo L, Cereda RF, Tosin I, Miranda EA, Sader HS. Antimicrobial susceptibility of coagulase-negative staphylococci and characterization of isolates with reduced susceptibility to glycopeptides. *Diagn Microbiol Infect Dis* 1999; 34 : 185-91.
23. Agvvald-Ohman C, Lund B, Edlund C. Multiresistant coagulase negative staphylococci disseminate frequently between intubated patients in a multidisciplinary intensive care unit. *Crit Care* 2004; 8 : R42-7.
24. Mokuolu AO, Jiya N, Adesiyun OO. Neonatal septicaemia in Ilorin: bacterial pathogens and antibiotic sensitivity pattern. *Afr J Med Sci* 2002; 31 : 127-30.
25. Sechi LA, Pinna A, Pusceddu C, Fadda G, Carta C, Zanetti S. Molecular characterization and antibiotic susceptibilities of ocular isolates of *Staphylococcus epidermidis*. *J Clin Microbiol* 1999; 37 : 3031-3.
26. Tenover FC, Biddle JW, Lancaster MV. Increasing resistance to vancomycin and other glycopeptides in *Staphylococcus aureus*. *Emerg Infect Dis* 2001; 7 : 327-32.
27. Ojha N, Deodhar L. Antimicrobial susceptibility pattern of nosocomial pathogens. *Bombay Hosp J* 1997; 39 : 49-51.
28. Jain A, Agarwal A, Verma RK. Cefoxitin disc diffusion test for detection of methicillin-resistant staphylococci. *J Med Microbiol* 2008; 57 : 957-61.
29. Oteo J, Baquero F, Vindel A, Campos J. Spanish members of the European Antimicrobial Resistance Surveillance System. Antibiotic resistance in 3113 blood isolates of *Staphylococcus aureus* in 40 Spanish hospitals participating in the European Antimicrobial Resistance Surveillance System (2001-2002). *J Antimicrob Chemother* 2004; 53 : 1033-8.
30. Jain A, Agarwal A, Verma RK. Cefoxitin disc diffusion test for detection of methicillin-resistant staphylococci. *J Med Microbiol* 2008; 57 : 957-61.
31. Reynolds R, Potz N, Colman M, Williams A, Livermore D, MacGowan A. BSAC extended working party on Bacteraemia Resistance Surveillance. Antimicrobial susceptibility of the pathogens of bacteraemia in the UK and Ireland 2001-2002: the BSCA bacteraemia Resistance Surveillance Program. *J Antimicrob Chemother* 2004; 53 : 1018-32.
32. St Gemme JW, Bell LM, Baumgart S, D'Angio CT, Harris MC. Distinguishing sepsis from blood culture contamination in young infants with blood cultures growing coagulase-negative staphylococci. *Pediatrics* 1990; 86 : 157-62.
33. Nataro JP, Concoran L, Zirin S. Prospective analysis of coagulase-negative staphylococcal infection in hospitalized infants. *J Padiatr* 1994; 125 : 798-804.

Reprint requests: Dr Amita Jain, Department of Microbiology, Chhatrapati Shahuji Maharaj Medical University, Lucknow, UP 226 003, India  
e-mail: amita602002@yahoo.com