

Clinical Article

Effect of Discontinuation of Anticoagulation in Patients with Intracranial Hemorrhage at High Thromboembolic Risk

Hwan-Su Jung, M.D., Ik-Chan Jeon, M.D., Chul-Hoon Chang, M.D., Young-Jin Jung, M.D.

Department of Neurosurgery, Yeungnam University Medical Center, Yeungnam University School of Medicine, Daegu, Korea

Objective : There was no abundance of data on the use of anticoagulant in patients with previous high risk of thromboembolic conditions under a newly developed intracranial hemorrhage in Korean society. The purpose of this study was to evaluate the safety of discontinuance and suggest the proper time period for discontinuance of anticoagulant among these patients.

Methods : We reviewed the medical records of 19 patients who took anticoagulant because of thromboembolic problems and were admitted to our department with newly developed anticoagulation associated intracranial hemorrhage (AAICH), and stopped taking medicine due to concern of re-bleeding from January 2008 to December 2012. Analysis of the incidence of thromboembolic complications and proper withdrawal time of anticoagulant was performed using the Kaplan-Meier method.

Results : Our patients showed high risk for thromboembolic complication. The CHA₂DS₂-VASc score ranged from two to five. Thromboembolic complication occurred in eight (42.1%) out of 19 patients without restarting anticoagulant since the initial hemorrhage. Among them, three patients (37.5%) died from direct thromboembolic complications. Mean time to outbreak of thromboembolic complication was 21.38±14.89 days (range, 8-56 days). The probability of thromboembolic complications at 7, 14, and 30 days since cessation of anticoagulation was 0.00, 10.53, and 38.49%, respectively.

Conclusion : Short term discontinuance of anticoagulant within seven days in patients with AAICH who are at high embolic risk (CHA₂DS₂-VASc score >2) appears to be relatively safe in Korean people. However, prolonged cessation (more than seven days) may result in increased incidence of catastrophic thromboembolic complications.

Key Words : Anticoagulation · Intracerebral hemorrhage · Thromboembolic complication.

INTRODUCTION

Long term use of chronic oral anticoagulants can be complicated by anticoagulation associated intracranial hemorrhage (AAICH), the most devastating complication of anticoagulant therapy. Among patients who received anticoagulant for management of thromboembolic problems, intracranial hemorrhages led to approximately 90% of deaths and major disability in survivors⁵. AAICH is an urgent medical condition requiring prompt reversal of anticoagulation. Rapid reversal is essential for prevention of enlargement of the AAICH, which could result in brain herniation. On the other hand, reversal and discontinuance of anticoagulant can result in development of several kinds of side effects, such as thromboembolic complication. The risk and incidence of thromboembolic complication after cessation

of anticoagulation in Korean society have not been accurately reported; in addition, there is no consensus regarding the proper period for withholding anticoagulant in the case of these patients. Based on results of other studies, discontinuation for seven to 10 days appears to be safe². However, this has limitation as a proper guideline because these data were reported from small western retrospective series^{3,6,11}. In this study, we discussed reasonable access for management of patients with AAICH who are at a high thromboembolic risk in the view of using an anticoagulant.

MATERIALS AND METHODS

Patient selection

From January 2008 to December 2012, 47 patients with war-

• Received : June 11, 2013 • Revised : September 3, 2013 • Accepted : January 10, 2014

• Address for reprints : Young-Jin Jung, M.D.

Department of Neurosurgery, Yeungnam University Medical Center, Yeungnam University School of Medicine, 170 Hyeonchung-ro, Nam-gu, Daegu 705-703, Korea
Tel : +82-53-620-3790, Fax : +82-53-620-3770, E-mail : jyjns@yu.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

farin associated intracranial hemorrhage were admitted to a single neurological institution. Eleven patients (23.4%) died due to an initial massive hematoma and were excluded from this study. Of the remaining 36 patients, 19 patients were not restarted with anticoagulant due to high risk of rebleeding and 17 patients were restarted with anticoagulant at a different time depending on each patient's condition. Restarting of drug was decided after consultation with the doctor who treated the patients due to the underlying disease. However, they did not give a clear answer regarding when the drug should be restarted after hemorrhage. Opinion regarding reuse of anticoagulant differs significantly depending on the physician. If the rebleeding risk was greater than the thromboembolic risk, anticoagulants were not restarted. For these reasons, anticoagulant was not started for 19 patients. Underlying disease to get oral anticoagulant, presenting hemorrhage type, thromboembolic risk factors, event day international normalized ratio (INR), prothrombin time (PT), and thromboembolic complication after cessation of anticoagulation were investigated.

Statistical analysis

Data analysis was performed retrospectively using the Statistical Package for the Social Sciences (SPSS) version 20.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics were performed for determination of the influence of clinical factors on development of thromboembolic events in the non-restarted anticoagulant group. The Kaplan-Meier method was used to assess the incidence and probability of thromboembolic events in patients with passage of time.

CHA₂DS₂-VASc score

Thromboembolic risk is largely dependent on the patient's condition itself. We need to identify the exact degree of thromboembolic risk in the patients' group in order to obtain objective validity when comparing the results with those of other authors. In this study, we used CHA₂DS₂-VASc score. The CHA₂DS₂-VASc [congestive heart failure/left ventricular dysfunction, hypertension, age ≥75 (double), diabetes, stroke (double)-vascular disease, age 65-74, and sex category (female)] has been used in multiple cohorts and is better at identifying patients with true risk. The score is inclusive of the most common risk factors for stroke in everyday clinical practice^{4,8}.

Table 1. Demographic and clinical characteristics of patients without restarting anticoagulant (n=19)

Variables	Mean±SD	Range
Age (years)	66.21±9.18	43-79
PT	37.77±34.08	9.70-150.10
INR	3.38±3.64	0.84-16.72
CHA ₂ DS ₂ -VASc Score (points)	NA	2-5
Discontinuance of warfarin (days)	475.79±682.49	8-1813
Mean follow up period (days)	472.95±575.69	8-1813

PT : prothrombin time, INR : international normalized ratio

RESULTS

The demographic and clinical characteristics of 19 patients in whom anticoagulants were not started are shown in Table 1 and 2. The most common type of AAICH was spontaneous hemorrhage, which presented in eight cases (42.1%). Acute subdural hematoma was the next main type of AAICH in five cases (21.1%). The mean PT was 37.77±34.08 seconds (range, 9.70-150.10 seconds). The thromboembolic risk of patients was calculated by the CHA₂DS₂-VASc score⁸. All patients presented with CHA₂DS₂-VASc scores of 2 or more than 2 points. The mean discontinuance time of the anticoagulant was 475.79±682.49 days (range, 8-1813 days).

There were eight cases (42.1%) with thromboembolic complications, three of which (37.5%) resulted in death. Two deaths were caused by sudden cardiac arrest, which was suspected with acute myocardial infarction, and another patient died of pulmonary thromboembolism. Among the survivors, deep venous thrombosis followed by intravenous filter insertion occurred in two cases and another two patients presented with angina pectoris. And non-fetal pulmonary embolism occurred in one case. The INR score at the time of occurrence of thromboembolic complications was 0.95±0.16. Mean time to occurrence of thromboembolic complication was 21.38±14.89 days (range, 8-56 days). The probabilities of having thromboembolic complications at 7, 14, and 30 days following cessation of warfarin therapy were 0.0% (95% confidence interval, 0.00-20.92%), 10.53% (95% confidence interval, 1.84-34.54%), and 38.49% (95% confidence interval, 18.42-62.85%), respectively (Fig. 1). By seven days, there was no thromboembolic complication or death. After seven days, thromboembolic complications showed a rapid increase. The INR showed a gradual decrease to the normal range, and then thromboembolic complications started to occur after a few days. From the result, we found that all thromboembolic complications and death occurred after seven days in all cases, and most of these thromboembolic complications tended to be concentrated between one and two weeks.

DISCUSSION

This study provided the longest follow up data on discontinuance of anticoagulant in patients presenting with AAICH who have been at high thromboembolic risk estimated by CHA₂DS₂-VASc score in Korean society.

This report presented several important findings. First, discontinuance of the anticoagulant is common in AAICH patients in the actual clinical field. Proper restart of warfarin treatment after AAICH was not prescribed for almost half of patients (47.2%, 17/36). AAICH induced mortality is very high

Table 2. Clinical characteristics of each patient without restarting anticoagulant (n=19)

Case	Age	Sex	Underlying disease	Hemorrhage type	CHA ₂ DS ₂ -VASc Score	Bleed day INR	Bleed day PT	Reversal Vit K (mg)	Event day or 7 days INR	Complications	Event day or Last F/U day	Mortality
1	68	F	AF	S-ICH	2	1.53	18.1	10	1.00	-	275	-
2	60	M	AF	S-ICH	5	2.01	23.5	20	1.14	AMI	21	Death
3	61	M	AF	S-ICH	2	2.06	25.9	20	1.02	-	1809	-
4	56	M	CHF	A-SAH	3	0.94	10.9	-	0.95	AP	18	-
5	66	F	CHF	S-ICH	4	3.72	45	20	1.01	PE	18	-
6	73	F	CHF	S-ICH	3	2.08	26.2	-	2.69	-	1813	-
7	79	M	DVT/PE	A-SDH	-	5.26	62.1	10	1.07	DVT	17	-
8	77	M	DVT/PE	A-SDH	-	1.00	11.6	-	0.60	PE	8	Death
9	78	F	DVT/PE	A-SDH	-	7.21	88.9	20	0.94	DVT	23	-
10	43	M	DVT/PE	C-SDH	-	4.53	53.4	10	0.95	-	169	-
11	66	F	PHV	A-SDH	2	16.72	150.1	30	0.85	AMI	10	Death
12	59	F	PHV	C-SDH	3	2.43	28.4	10	1.29	-	115	-
13	54	F	PHV	S-ICH	3	2.81	33.7	-	1.01	AP	56	-
14	69	M	PHV	T-ICH	3	3.63	43.2	-	1.96	-	11	-
15	64	F	Recurrent stroke	A-SAH	-	0.84	9.7	-	NA	-	439	-
16	70	F	Recurrent stroke	A-SDH	-	1.00	11.6	-	1.00	-	38	-
17	73	M	Recurrent stroke	S-ICH	-	3.31	40.3	10	1.02	-	737	-
18	74	M	Recurrent stroke	S-ICH	-	2.21	23.9	10	1.30	-	1256	-
19	68	M	Recurrent stroke	T-ICH	-	0.96	11.1	-	1.02	-	702	-

AF : atrial fibrillation, CHF : congestive heart failure, DVT/PE : deep vein thrombosis/pulmonary embolism, PHV : prosthetic heart valve, S-ICH : spontaneous intracerebral hemorrhage, A-SAH : aneurysmal subarachnoid hemorrhage, A-SDH : acute subdural hematoma, C-SDH : chronic subdural hematoma, T-ICH : traumatic intracerebral hematoma, AMI : acute myocardial infarction, AP : angina pectoris

and survivors from initial neurological damage are faced with the decision of whether or not to restart anticoagulant^{3,5}. However, proper use of anticoagulant in patients with AAICH has not yet been established, and guidelines could not be obtained from a limited number of previous retrospective series^{3,6,11}. According to American Heart Association recommendations, in patients with a very high risk of thromboembolism in whom restarting warfarin is considered, warfarin therapy may be restarted at seven to 10 days since onset of the original hemorrhage². In this study, only nearly half of patients were prescribed anticoagulant because of hemorrhagic risk. The consensus of the resumption of anticoagulant in AAICH patients is not clear in our institution. Decisions regarding to the starting time of anticoagulant were differed significantly among physicians due to the fact. And available data for determination of the optimal timing and dosage for restarting anticoagulation following an episode of AAICH are inadequate.

Second, in this study, short duration (within 7 days) discontinuance of anticoagulants was relatively safe, similar to the result of a previous study series. Kawamata et al.⁷ found no cases of ischemic stroke when warfarin treatment was discontinued for three days in 27 patients with AAICH. Tinker et al.¹⁰ retrospectively reviewed the risk of discontinuing warfarin therapy in 159 patients with mechanical heart valves undergoing elective surgery; none of the patients had in-hospital thromboembolic complications. Ananthasubramaniam et al.¹¹ also reported that




Fig. 1. Kaplan-Meier thromboembolic complication free survival estimate for patients with anticoagulants associated intracranial hemorrhage without resuming anticoagulants. The probability of thromboembolic complications after cessation of anticoagulation at 7, 14, and 30 days was 0.00, 10.53, and 38.49%, respectively.

no thromboembolic complications occurred in 24 patients with prosthetic heart valves after withholding anticoagulant for an average of 15 days. Like these previous studies, in this study, there was also no occurrence of thromboembolic complications within seven days since AAICH.

The third finding of this study is as follows. Although a temporary withdrawal of anticoagulants within seven days may be safe,

prolonged cessation could lead to catastrophic results. We found that thromboembolic complications begin to occur after seven days and the incidence shows a rapid increase around 14 days later. Two complications occurred within two weeks. However, additional events occurred after two weeks. Overall occurrence of thromboembolic complications was 42.1% (8/19), higher than that reported in previous series¹⁾. The reasons for this phenomenon can be explained as follows. First, the time for withdrawal of anticoagulant was long, compared to the previous reports. In previous studies¹⁾, anticoagulant was restarted within at least two weeks after a hemorrhagic event. The INR in patients with thromboembolic complication showed a gradual reduction and it fell below the normal range on ictus days. Because the thromboembolic risk was accumulated after cessation of anticoagulants, the possibility of thromboembolic event also increased. Second, the patients belonged to a very high risk group for occurrence of thromboembolic events. The CHA₂DS₂-VASc score has been used in multiple cohorts for measurement of severity and for better identification of patients with true risk⁴⁾. The score includes the most common risk factors related to stroke in everyday clinical practice. Among patients with CHA₂DS₂-VASc score=2, the incidence of thromboembolic complication in one year is 2.69% and the rate increased by 3.20% at a score of 3 points. For prevention of thromboembolism, anticoagulants are recommended for patients with a score of more than 2 points⁹⁾. In this study, the CHA₂DS₂-VASc score was more than 2 points for all patients (2 points in three cases, 3 points in five, and 4 and 5 points in one case each), indicating that the patient has a high underlying thromboembolic risk. Otherwise, previous reports did not objectively demonstrate the patients' underlying thromboembolic risk, causing difficulty in comparison and analysis of results between studies.

This study included only 19 cases. The limitation of this study is the retrospective design with a very small number of cases. However, the incidence of patients who did not restart anticoagulation after AAICH was very low. Over the past five years, we were only able to collect 19 cases. Previous studies also included small case series with retrospective review^{3,6,11)}. This study is expected to have worth as the first study of Korean people.

CONCLUSION

Discontinuance of anticoagulant within seven days presented a relatively low incidence of thromboembolic complications in patients with AAICH at a high thromboembolic risk (CHA₂DS₂-VASc score >2), and withdrawal of more than seven days may increase the risk of devastating results. We think that short term discontinuance of anticoagulant limited within seven days ap-

pears to be safe in Korean people. Unfortunately, this study was conducted retrospectively in a small number of cases of AAICH and has several limitations to understanding. For a more reasonable and acceptable guideline on the use of anticoagulant in patients with AAICH, conduct of further studies will be required.

References

1. Ananthasubramaniam K, Beattie JN, Rosman HS, Jayam V, Borzak S : How safely and for how long can warfarin therapy be withheld in prosthetic heart valve patients hospitalized with a major hemorrhage? *Chest* 119 : 478-484, 2001
2. Broderick J, Connolly S, Feldmann E, Hanley D, Kase C, Krieger D, et al. : Guidelines for the Management of Spontaneous Intracerebral Hemorrhage in Adults : 2007 Update : A Guideline From the American Heart Association/American Stroke Association Stroke Council, High Blood Pressure Research Council, and the Quality of Care and Outcomes in Research Interdisciplinary Working Group : The American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. *Circulation* 116 : e391-e413, 2007
3. Butler AC, Tait RC : Restarting anticoagulation in prosthetic heart valve patients after intracranial haemorrhage : a 2-year follow-up. *Br J Haematol* 103 : 1064-1066, 1998
4. Camm AJ, Lip GY, De Caterina R, Savelieva I, Atar D, Hohnloser SH, et al. : 2012 focused update of the ESC Guidelines for the management of atrial fibrillation : an update of the 2010 ESC Guidelines for the management of atrial fibrillation. Developed with the special contribution of the European Heart Rhythm Association. *Eur Heart J* 33 : 2719-2747, 2012
5. Fang MC, Go AS, Chang Y, Hylek EM, Henault LE, Jensvold NG, et al. : Death and disability from warfarin-associated intracranial and extracranial hemorrhages. *Am J Med* 120 : 700-705, 2007
6. Hacke W : The dilemma of reinstating anticoagulation for patients with cardioembolic sources and intracranial hemorrhage : how wide is the strait between Skylla and Karybdis? *Arch Neurol* 57 : 1682-1684, 2000
7. Kawamata T, Takeshita M, Kubo O, Izawa M, Kagawa M, Takakura K : Management of intracranial hemorrhage associated with anticoagulant therapy. *Surg Neurol* 44 : 438-442; discussion 443, 1995
8. Lip GY, Nieuwlaat R, Pisters R, Lane DA, Crijns HJ : Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach : the euro heart survey on atrial fibrillation. *Chest* 137 : 263-272, 2010
9. Olesen JB, Torp-Pedersen C, Hansen ML, Lip GY : The value of the CHA₂DS₂-VASc score for refining stroke risk stratification in patients with atrial fibrillation with a CHADS₂ score 0-1 : a nationwide cohort study. *Thromb Haemost* 107 : 1172-1179, 2012
10. Tinker JH, Tarhan S : Discontinuing anticoagulant therapy in surgical patients with cardiac valve prostheses. Observations in 180 operations. *JAMA* 239 : 738-739, 1978
11. Wijdicks EF, Schievink WI, Brown RD, Mullany CJ : The dilemma of discontinuation of anticoagulation therapy for patients with intracranial hemorrhage and mechanical heart valves. *Neurosurgery* 42 : 769-773, 1998