

TRANSMISSION OF *NEOSPORA CANINUM* BETWEEN WILD AND DOMESTIC ANIMALS

Author(s): L. F P. Gondim, M. M. McAllister, N. E. Mateus-Pinilla, W. C. Pitt, L. D. Mech, and M. E. Nelson

Source: Journal of Parasitology, 90(6):1361-1365.

Published By: American Society of Parasitologists

DOI: <http://dx.doi.org/10.1645/GE-341R>

URL: <http://www.bioone.org/doi/full/10.1645/GE-341R>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

TRANSMISSION OF *NEOSPORA CANINUM* BETWEEN WILD AND DOMESTIC ANIMALS

L. F. P. Gondim, M. M. McAllister, N. E. Mateus-Pinilla*, W. C. Pitt†, L. D. Mech‡, and M. E. Nelson‡

Department of Veterinary Pathobiology, College of Veterinary Medicine, University of Illinois, 2001 South Lincoln Avenue, Urbana, Illinois 61802. e-mail: pita@ufba.br

ABSTRACT: To determine whether deer can transmit *Neospora caninum*, brains of naturally infected white-tailed deer (*Odocoileus virginianus*) were fed to 4 dogs; 2 of these dogs shed oocysts. Oocysts from 1 of the dogs were tested by polymerase chain reaction and found to be positive for *N. caninum* and negative for *Hammondia heydorni*. The internal transcribed spacer 1 sequence of the new strain (designated NC-deer1) was identical to *N. caninum* from domestic animals, indicating that *N. caninum* is transmitted between wild and domestic animals, often enough to prevent divergent evolution of isolated populations of the parasite. NC-deer1 oocysts were administered to a calf that developed a high antibody titer, providing evidence that *N. caninum* from wildlife can infect cattle. In addition, *N. caninum* antibody seroprevalence was detected in 64/164 (39%) free-ranging gray wolves (*Canis lupus*), 12/113 (11%) coyotes (*Canis latrans*), 50/193 (26%) white-tailed deer, and 8/61 (13%) moose (*Alces alces*). These data are consistent with a sylvatic transmission cycle of *N. caninum* between cervids and canids. We speculate that hunting by humans favors the transmission of *N. caninum* from deer to canids, because deer carcasses are usually eviscerated in the field. Infection of canids in turn increases the risk of transmitting the parasite to domestic livestock.

Neospora caninum is a protozoan parasite originally reported in dogs and calves (Bjerkås et al., 1984; O'Toole and Jeffrey, 1987; Parish et al., 1987; Dubey et al., 1988) and commonly diagnosed in bovine abortions worldwide (Dubey, 1999). The parasite has been detected in several mammalian species, i.e., sheep, goats, horses, deer (Dubey and Lindsay, 1996), water buffaloes (Guarino et al., 2000), rhinoceros (Williams et al., 2002), and foxes (Almeria et al., 2002). Dogs were the first-discovered definitive host of *N. caninum* (McAllister et al., 1998); recently, coyotes (*Canis latrans*) were also demonstrated to be definitive hosts of the parasite (Gondim, McAllister et al., 2004). When investigating *N. caninum* in canine feces, it is necessary to avoid confusion with the closely related parasite *Hammondia heydorni*, which has oocysts that appear similar to those of *N. caninum* (Slapeta et al., 2002). There are few confirmed cases of neosporosis in wild animals; however, antibodies against *N. caninum* have been found in several wild mammalian species, suggesting that the parasite may be widely spread among wildlife (for review, see Dubey, 2003).

Natural cases of neosporosis have been reported in different species of wild and captive deer (Woods et al., 1994; Dubey et al., 1996). The seroprevalence of *N. caninum* in free-ranging white-tailed deer (*Odocoileus virginianus*) in some states of the United States has been shown to be between 40 and 50% (Dubey et al., 1999; Lindsay et al., 2002), suggesting that deer may play an important role in the epidemiology of neosporosis.

The aims of this study were to investigate whether naturally infected white-tailed deer tissues can induce dogs to shed *N. caninum* oocysts and to determine the prevalence of antibodies against *N. caninum* in serum samples from gray wolves (*Canis lupus*), coyotes, white-tailed deer, and moose (*Alces alces*). All sera were also tested for *Toxoplasma gondii* antibodies to rule out potential cross-reactivity between the 2 parasites.

MATERIALS AND METHODS

Collection of deer samples

In December 2001, 30 blood samples were collected from the thoracic cavity of hunted white-tailed deer. The samples were selected from hunter-killed deer at the Vermillion County Illinois Department of Natural Resources (IDNR) deer check station in Illinois during the statewide firearm deer harvest season. The sera were tested for *N. caninum* antibodies to estimate the regional seroprevalence. In December 2002, with permission from IDNR, 12 deer heads with matching blood samples were collected from hunter-killed white-tailed deer in the same region. Heads from 4 deer with *N. caninum* antibody titer $\geq 1:800$ by an indirect fluorescent antibody test (IFAT) (described in the Prevalence of *Neospora caninum* and *Toxoplasma gondii* Antibodies Using Wolf, Coyote, White-Tailed Deer, and Moose Sera) were selected and kept at 4 °C for up to 12 hr after collection.

Infection of dogs

Four female mixed-breed hound puppies (A, B, C, and D), 10–14 wk old, purchased from a commercial class A breeder (Covance, Inc., Madison, Wisconsin), were housed individually in indoor kennels. The dogs were fed dry dog food and had never consumed raw meat or meat products. Each pup tested negatively for antibodies against *N. caninum* by IFAT (<1:25) before the experimental challenge.

Fecal samples from each dog were examined for 4 consecutive days before consuming the infected tissues. Within 12 hr after collection of the deer heads, each dog was fed 1 whole brain from a seropositive deer. Fecal samples were examined daily for oocysts, starting on the third day after infection.

Fecal examinations and processing of oocysts

Dog feces were examined by a standard sucrose flotation technique, as described previously (Gondim et al., 2002). Fecal specimens containing oocysts were mixed with 5 volumes of 2% H₂SO₄ and aerated for 3 days to induce sporulation. Sporulated oocysts were stored at 4 °C until used.

DNA extraction from sporulated oocysts

A volume of 140 ml of dog feces diluted in 2% H₂SO₄, containing 700 oocysts, was filtered through a gauze and distributed in 15-ml centrifuge tubes. The tubes were centrifuged at 800 g for 10 min, the supernatant was discarded, and the sediment was suspended with sucrose solution. Coverslips were placed on each tube, and the samples were centrifuged at 800 g for 10 min. The coverslips were removed and washed in water to collect the oocysts. The solution containing the oocysts was centrifuged; the sediment was transferred to a 1.5-ml tube, treated for 5 min with 2.6% sodium hypochlorite, and washed 5 times in water. The sediment was mixed with 700 μ l of digestion buffer (100 mM NaCl, 10 mM Tris-HCl, pH 8.0, 25 mM ethylenediaminetetraacetic acid [EDTA], 0.5% sodium dodecyl sulfate) and 5 μ l of proteinase K (20 mg/ml) and added to a 1.5-ml tube containing 500 μ l of glass beads

Received 20 February 2004; revised 29 March 2004; accepted 30 March 2004.

* Center for Wildlife Ecology, Natural History Survey, Urbana, Illinois 61820.

† USDA-APHIS-WS, National Wildlife Research Center, Hilo Field Station, P.O. Box 10880, Hilo, Hawaii 96721.

‡ U.S. Geological Survey, Northern Prairie Wildlife Research Center, 8711, 37th Street SE, Jamestown, North Dakota 58401-7317.

TABLE I. IFAT for IgG antibodies against *Neospora caninum* and *Toxoplasma gondii* in sera from wildlife from several states of the United States.

Animal species	State	<i>N. caninum</i> IFAT		<i>T. gondii</i> IFAT		IFAT positive for both <i>N. caninum</i> and <i>T. gondii</i>
		Positive	Total (%)	Positive	Total (%)	
Wolf	Minnesota	64/164 (39.0%)	39.0	47/164 (28.7%)	28.7	23/164 (14%)
Coyote	Utah	1/45 (2.2%)		6/45 (13.3%)		
	Colorado	5/28 (17.9%)	10.6	4/28 (14.3%)	31.0	6/113 (5.3%)
	Illinois	6/40 (15.0%)		25/40 (62.5%)		
White-tailed deer	Minnesota	30/150 (20.0%)	25.9	7/150 (4.6%)	12.4	6/193 (3.1%)
	Illinois	20/43 (46.5%)		17/43 (39.5%)		
Moose	Minnesota	8/61 (13.1%)	13.1	1/61 (1.6%)	1.6	0

(0.5 mm in diameter). The suspension was vortexed for 10 min and incubated at 65 C for 2 hr. DNA was extracted using standard phenol-chloroform followed by 2-propanol precipitation.

Polymerase chain reaction assays for *Neospora caninum* and *Hammondia heydorni*

DNA extracted from the oocysts was used in polymerase chain reaction (PCR) assays for *N. caninum* and *H. heydorni*. PCR for *N. caninum* was performed using the species-specific primer pair Np21–Np6 (Yamaga et al., 1996). PCR conditions comprised 40 cycles at 94 C for 1 min, 54 C for 1 min, and 72 C for 1.5 min, with a final extension step at 72 C for 2 min. DNA from *N. caninum* and Vero cells were used as positive and negative controls, respectively. PCR to detect *H. heydorni* was performed using the species-specific primers JS4–JS5 and PCR conditions described by Slapeta et al. (2002); a positive control (*H. heydorni* DNA) and a negative control (Vero cell DNA) were included in the test. Each PCR reaction was performed in a 25- μ l volume containing 30 pmol of each primer, 1.25 U of *Taq* DNA polymerase, 200 μ M of each deoxynucleoside triphosphate, 1 μ l of template, 1.5 mM MgCl₂, 20 mM Tris–HCl (pH 8.4), 50 mM KCl, and sterile ddH₂O up to 25 μ l. PCR products were electrophoresed on 2.0% agarose gel stained with ethidium bromide in a 0.5 \times TBE buffer (45 mM Tris base, 45 mM boric acid, 1 mM EDTA, pH 8.0) at 100 V for 1 hr and observed under ultraviolet (UV) illumination.

Internal transcribed spacer 1 sequencing

Extracted DNA from oocysts was used for amplification and sequencing of the internal transcribed spacer 1 (ITS1) region of the ribosomal DNA (rDNA). PCR for the ITS1 was performed using the primers ITS5–ITS2 (White et al., 1990); PCR conditions consisted of an initial denaturing step at 94 C for 3 min followed by 30 cycles at 94 C for 1 min, 55 C for 1 min, and 72 C for 2 min, with a final extension step at 72 C for 7 min. DNA of *N. caninum* (NC-beef strain) (Gondim, Laski et al., 2004) and Vero cells were included as positive and negative controls, respectively, in the PCR assays. The products were electrophoresed on 1.5% agarose gel stained with ethidium bromide at 100 V for 1 hr and gel purified using the QIAquick Gel Extraction Kit (Qiagen Inc., Valencia, California). Automated sequence reactions for *N. caninum* ITS1 were performed at the Biotechnology Center at the University of Illinois at Urbana–Champaign, using the internal primers PN1–PN2 (Holmdahl and Mattsson, 1996).

PCR-amplified ITS1 using 5% polyacrylamide gel electrophoresis

The amplified ITS1 from the recently isolated oocysts and from the *N. caninum* strain NC-beef (positive control) were electrophoresed on 5% polyacrylamide gel (BioRad, Ready gel, Hercules, California) at 120 V for 2 hr. The gel was stained with ethidium bromide and observed by UV illumination.

Oral inoculation of a calf with oocysts shed by a dog that consumed infected deer brain

A newborn dairy bull calf was purchased from the University of Illinois Dairy. It was housed indoors on a slatted floor, without bedding, and fed milk replacer. The calf was seronegative for *N. caninum* anti-

bodies at 1:25 by IFAT. At 5 days of age, the calf received 2,500 sporulated oocysts shed by dog A, through an esophageal feeder. Blood was collected from the calf 34 days after administration of oocysts, and the serum was separated and kept at –20 C until examined.

Prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies using wolf, coyote, white-tailed deer, and moose sera

Sera from 164 gray wolves, 113 coyotes, 193 white-tailed deer, and 61 moose were examined (Table I). Wolf and moose sera were collected in northeastern Minnesota, coyote sera were collected in Utah, Colorado, and Illinois, and sera from white-tailed deer were collected in northeastern Minnesota and Illinois. All sera had been collected previously for use in other investigations. IFATs were used for the detection of antibodies against *N. caninum* and *T. gondii* using tachyzoites of the strains NC-beef and RH as antigens, respectively. A fluorescein isothiocyanate (FITC)–conjugated anti-dog IgG (Bethyl, Montgomery, Texas), diluted at 1:50 in phosphate-buffered saline (PBS), was used as a secondary antibody for wolf and coyote sera; FITC–conjugated anti-deer IgG (Cappel, Durham, North Carolina), diluted at 1:50 in PBS, was used for white-tailed deer and moose sera. Sera reacting at dilutions \geq 1:50 were considered positive for *T. gondii* antibodies. For *N. caninum* serology, deer and moose sera were considered positive if they reacted at dilutions \geq 1:100, and 1:50 was selected as the cutoff for coyote and wolf sera. The maximum antibody titer was determined for all *N. caninum*–positive samples. Maximum titers for *T. gondii* were determined only for samples that showed positive reactions for both *T. gondii* and *N. caninum*.

RESULTS

Oocyst shedding and PCR from oocyst DNA

Two of 4 dogs (A and B) that ingested naturally infected deer brain shed *N. caninum*–like oocysts in the feces. Dog A shed approximately 12,300 oocysts between 7 and 14 days after infection; dog B shed approximately 500 oocysts on the 11th and 12th days after infection. Oocysts were shed unsporulated and measured 10 μ m in diameter (dog A, n = 10; dog B, n = 5). Sporulation was induced when oocysts were mixed in 2% H₂SO₄ and aerated at room temperature for 3 days. The oocysts shed by dog A tested positively for *N. caninum* by PCR (Fig. 1); in contrast, PCR for *H. heydorni* yielded negative results. DNA could not be extracted from dog B oocysts, which were in low numbers and diluted in a large volume of feces. The 4 dogs remained seronegative for *N. caninum* antibodies at 1:50 by IFAT 28 days after consuming deer brain.

ITS1 sequence of the recently isolated oocysts

The ITS1 sequence obtained from the recently isolated *N. caninum* oocysts (designated NC-deer1) (GenBank AY508811) was identical to the ITS1 sequence of the *N. caninum* strain




FIGURE 1. PCR for *Neospora caninum* using Np21–Np6 primers (Yamaga et al., 1996). (1) A 100-bp DNA ladder; (2) Vero cell DNA (negative control); (3) *N. caninum* DNA (positive control); and (4) a positive reaction for *N. caninum* using DNA extracted from oocysts of the newly described isolate from deer (designated NC-deer1).

NC-Illinois, which had been isolated from a dairy calf in Illinois (Gondim, Laski et al., 2004) (GenBank AY259041), and to other *N. caninum* ITS1 sequences of dogs or cattle (strain/accession number: NC-2/AF249969, NC-5/AF249970, BPA1/AF038860, CN1/AF038861, NC-SweB1/AF029702, NC-beef/AF249968, CZ-4/AF432123). The PCR-amplified product of the NC-deer1 using the primers ITS5–ITS2 yielded a band of approximately 510 base pairs (bp) on the 1.5% agarose gel (Fig. 2A); this band comprises the reported ITS1 sequence for *N. caninum* (421 bp) and part of the 18S and 5.8S rDNA genes. The same PCR-amplified ITS1 of the NC-deer1 isolate was observed as 2 distinct bands on 5% polyacrylamide gel electrophoresis (PAGE), identical to the NC-beef strain that was used as a positive control (Fig. 2B) and similar to other strains examined previously (Gondim, Laski et al., 2004).

Serological response of the calf infected with the newly isolated oocysts

A newborn dairy bull calf developed an antibody titer of 1:800 to *N. caninum* 34 days after oral inoculation with 2,500 oocysts of the newly described isolate (NC-deer1).

Serology of wild animals

Antibodies against *N. caninum* were detected in 39% of wolves, 11% of coyotes, 26% of white-tailed deer, and 13% of moose. Antibodies to *T. gondii* were found in 29% of wolves, 31% of coyotes, 12% of white-tailed deer, and 2% of moose; positive reactions for both *N. caninum* and *T. gondii* were observed in 14% of wolves, 5% of coyotes, 3% of white-tailed deer, and 0% of moose (Table I). The maximum *N. caninum* antibody titer observed in wolf and moose sera was 1:6,400; titers up to 1:3,200 were found in white-tailed deer, and in coy-




FIGURE 2. PCR amplification of the ITS1 region. (1) A 100-bp DNA ladder; (2) Vero cell DNA (negative control); (3) *Neospora caninum* DNA (positive control); and (4) DNA extracted from the NC-deer1 isolate. (A) On a 1.5% agarose gel, a single band of approximately 510 bp was observed for the NC-deer1 isolate and also for the *N. caninum* DNA used as positive control. (B) On a 5% polyacrylamide gel, which has a higher resolution than the 1.5% agarose gel, 2 distinct bands were observed for each PCR-amplified product.

otes the maximum serum titer against the protozoan was 1:800 (Table II). In a total of 133 *N. caninum*-positive serum samples observed for all tested animals, 98 sera reacted solely with *N. caninum* and 35 sera reacted with both *N. caninum* and *T. gondii* (Table I).

DISCUSSION

Two of the 4 dogs shed *N. caninum*-like oocysts after ingesting brain of naturally infected white-tailed deer. Oocysts shed by 1 dog (A) were confirmed to be *N. caninum* using the specific primers Np21–Np6 and by sequencing the ITS1 region of the rDNA. PCR for *H. heydorni* was negative. A calf that ingested these oocysts developed an *N. caninum* antibody titer of 1:800, 34 days after inoculation.

The ITS1 of the rDNA is a region that evolves fast and has frequently been used for phylogenetic comparisons among different organisms (White et al., 1990). The ITS1 sequence of the new *N. caninum* isolate (NC-deer1) is identical to the ITS1 sequence reported for *N. caninum* strains isolated from domestic animals (Gondim, Laski et al., 2004). The separation of the amplified ITS1 into 2 distinct bands on 5% PAGE, as recently reported for 6 different strains of *N. caninum* (Gondim, Laski et al., 2004), was also observed for the ITS1 region of NC-

TABLE II. *Neospora caninum* antibody titers in seropositive wolves, coyotes, white-tailed deer, and moose, tested by IFAT.

Animals	Serum dilution							
	1:50	1:100	1:200	1:400	1:800	1:1,600	1:3,200	1:6,400
Wolves	8	7	9	12	14	10	3	1
Coyotes	3	6	1	1	1	0	0	0
White-tailed deer	ND*	9	10	3	15	7	6	0
Moose	ND*	2	1	1	2	0	1	1

* ND = not determined.




FIGURE 3. Transmission cycles of the protozoan *Neospora caninum* between wild and domestic animals. The domestic cycle, transmission from cattle to coyotes, and transmission from deer to dogs have each been demonstrated. Transmission between coyotes and deer, from coyotes to cattle, and from dogs to deer are probable but as yet untested.

deer1. These results indicate that *N. caninum* infections are horizontally transmitted between wild and domestic animals, often enough to prevent divergent evolution of isolated populations of the parasite.

In this investigation, wild naturally infected deer transmitted *N. caninum* to dogs. In a previous study (Gondim, McAllister et al., 2004), coyotes were shown to be a definitive host of *N. caninum* when they shed oocysts after ingesting tissues from infected cattle. Based on these findings, a crossover cycle of *N. caninum* between wild and domestic animals is proposed and illustrated in Figure 3.

In this study, deer brain was the only tissue fed to dogs because we wanted to avoid shedding of *Sarcocystis* spp. sporocysts in feces, which could have complicated observation and identification of *N. caninum* oocysts. *Sarcocystis* spp. are commonly observed in skeletal muscle and heart of wild deer (Emmett and Huggins, 1982). In a previous study (Gondim et al., 2002), the production of *N. caninum* oocysts in dogs was improved by providing the dogs various tissues from experimentally infected calves, including brain, spinal cord, heart, skeletal muscle, diaphragm, and kidney. Hence, dogs consuming a mixture of different tissues from deer might have shed a greater number of oocysts than that observed in this experiment. The

4 deer brains used in this study were collected from *N. caninum*-seropositive deer; however, it is not known whether all the 4 selected brains contained cysts of the parasite.

White-tailed deer are easily adaptable to a variety of environmental conditions and are widely distributed in North America. In 1998, the number of white-tailed deer in North America was estimated to be almost 29 million in the winter; hunters harvest about 6 million white-tailed deer annually (Crête and Daigle, 1998). When deer are hunted, carcasses are usually field dressed and the offal is left behind. Besides the natural predation of deer by wild canids, during every deer-hunting season the offal is readily available for consumption by wild canids and dogs, which could increase the risk of transmission of *N. caninum* from canids to livestock.

The prevalence of *N. caninum* antibodies found in wolves, coyotes, white-tailed deer, and moose ranged from 11 to 39%. Antibody titers up to 1:6,400 were observed in wolf and moose, titers up to 1:3,200 were found in white-tailed deer, and titers up to 1:800 were found in coyotes; these findings suggest that a wild canid-ruminant life cycle of the parasite is common. All serum samples used in this survey were tested for both *N. caninum* and *T. gondii*; in 133 samples that tested positive for *N. caninum* antibodies, more than 70% of these samples reacted

solely to *N. caninum*, indicating that *N. caninum* titers were not an artifact of cross-reaction with *T. gondii*. The high prevalence (39%) and high titers (up to 1:6,400) of *N. caninum* antibodies observed in wolves are probably related to their diet, which is mostly based on ruminants, including white-tailed deer and moose; wolves may be exposed to *N. caninum*-infected tissues repeatedly throughout their lives. Antibodies to *N. caninum* were found in 11% of coyotes; their diet is more diversified when compared with wolves and they do not primarily rely on ruminants (Arjo et al., 2002). Nevertheless, the prevalence of antibodies found in coyotes (11%) is still higher than that reported for dogs in the United States (7%) (Cheadle et al., 1999).

The isolation of *N. caninum* from wildlife, the serological evidence of infection in several wild animals, and the demonstration that the parasite can cycle between wild and domestic animals pose new questions about the relative importance of transmission of *N. caninum* between wildlife and livestock. Some reasonable control measures may help to prevent transmission of the parasite to domestic ruminants, such as protection of feedstuffs from contamination with canine feces and appropriate disposal of dead livestock to prevent consumption by a definitive host. The large amount of offal from hunted deer carcasses that are field dressed and left available for consumption by a variety of carnivores could pose an added risk of disease transmission each fall.

ACKNOWLEDGMENTS

We thank Mark Lenarz (Minnesota Department of Natural Resources), Michael Schrage (Fond du Lac Band of Lake Superior Chippewa, Minnesota), and Andrew Edwards (1854 Authority, Minnesota) for providing moose sera. We also thank Eric Gese (USDA-APHIS-WS, National Wildlife Research Center, Predator Research and Ecology, Logan, Utah) for providing wolf and coyote sera. This study was supported in part by USDA Hatch funds administered by the University of Illinois Agricultural Experimental Station (ILLU-70-0341) and by the Biological Resources Division, U.S. Geological Survey, and U.S. Department of Agriculture, North Central Research Station. Figure 3 was illustrated by Kerry Helms (College of Veterinary Medicine Biomedical Communication Center, University of Illinois). L.F.P.G. is the recipient of a fellowship from CNPq, Brazil. The *H. heydorni* DNA was kindly provided by John Ellis.

LITERATURE CITED

- ALMERIA, S., D. FERRER, M. PABON, J. CASTELLA, AND S. MANAS. 2002. Red foxes (*Vulpes vulpes*) are a natural intermediate host of *Neospora caninum*. *Veterinary Parasitology* **107**: 287–294.
- ARJO, W. M., D. H. PLETSCHER, AND R. R. REAM. 2002. Dietary overlap between wolves and coyotes in Northwestern Montana. *Journal of Mammalogy* **83**: 754–766.
- BJERKÅS, I., S. F. MOHN, AND J. PRESTHUS. 1984. Unidentified cyst-forming sporozoan causing encephalomyelitis and myositis in dogs. *Zeitschrift für Parasitenkunde* **70**: 271–274.
- CHEADLE, M. A., D. S. LINDSAY, AND B. L. BLAGBURN. 1999. Prevalence of antibodies to *Neospora caninum* in dogs. *Veterinary Parasitology* **85**: 325–330.
- CRÊTE, M., AND C. DAIGLE. 1998. Management of indigenous North American deer at the end of the 20th century in relation to large predators and primary production. *Acta Veterinaria Hungarica* **47**: 1–16.
- DUBEY, J. P. 1999. Recent advances in *Neospora* and neosporosis. *Veterinary Parasitology* **84**: 349–367.
- . 2003. Review of *Neospora caninum* and neosporosis in animals. *Korean Journal of Parasitology* **41**: 1–16.
- , J. L. CARPENTER, C. A. SPEER, M. J. TOPPER, AND A. UGGLA. 1988. Newly recognized fatal protozoan disease of dogs. *Journal of the American Veterinary Medical Association* **192**: 1269–1285.
- , K. HOLLIS, S. ROMAND, P. THULLIEZ, O. C. KWOK, L. HUNGERFORD, C. ANCHOR, AND D. ETTER. 1999. High prevalence of antibodies to *Neospora caninum* in white-tailed deer (*Odocoileus virginianus*). *International Journal for Parasitology* **29**: 1709–1711.
- , AND D. S. LINDSAY. 1996. A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology* **67**: 1–59.
- , J. RIGOLET, P. LAGOURETTE, C. GEORGE, L. LONGEART, AND J. L. LENET. 1996. Fatal transplacental neosporosis in a deer (*Cervus eldi siamensis*). *Journal of Parasitology* **82**: 338–339.
- EMNETT, C. W., AND E. J. HUGGINS. 1982. Sarcocystis of deer in South Dakota. *Journal of Wildlife Diseases* **18**: 187–193.
- GONDIM, L. F. P., L. GAO, AND M. M. MCALLISTER. 2002. Improved production of *Neospora caninum* oocysts, cyclical oral transmission between dogs and cattle, and in vitro isolation from oocysts. *Journal of Parasitology* **88**: 1159–1163.
- , P. LASKI, L. GAO, AND M. M. MCALLISTER. 2004. Variation of the internal transcribed spacer 1 (ITS1) sequence within individual strains and among different strains of *Neospora caninum*. *Journal of Parasitology* **90**: 119–122.
- , M. M. MCALLISTER, W. C. PITT, AND D. E. ZEMLIKA. 2004. Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **34**: 159–161.
- GUARINO, A., G. FUSCO, G. SAVINI, G. DI FRANCESCO, AND G. CRINGOLI. 2000. Neosporosis in water buffalo (*Bubalus bubalis*) in Southern Italy. *Veterinary Parasitology* **91**: 15–21.
- HOLMDAHL, O. J., AND J. G. MATTSSON. 1996. Rapid and sensitive identification of *Neospora caninum* by in vitro amplification of the internal transcribed spacer 1. *Parasitology* **112**: 177–182.
- LINDSAY, D. S., S. E. LITTLE, AND W. R. DAVIDSON. 2002. Prevalence of antibodies to *Neospora caninum* in white-tailed deer, *Odocoileus virginianus*, from the southeastern United States. *Journal of Parasitology* **88**: 415–417.
- MCALLISTER, M. M., J. P. DUBEY, D. S. LINDSAY, W. R. JOLLEY, R. A. WILLS, AND A. M. MCGUIRE. 1998. Dogs are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **28**: 1473–1478.
- O'TOOLE, D., AND M. JEFFREY. 1987. Congenital sporozoan encephalomyelitis in a calf. *Veterinary Record* **121**: 563–566.
- PARISH, S. M., L. MAAG-MILLER, T. E. BESSER, J. P. WEIDNER, T. MCELWAIN, D. P. KNOWLES, AND C. W. LEATHERS. 1987. Myelitis associated with protozoal infection in newborn calves. *Journal of the American Veterinary Medical Association* **191**: 1599–1600.
- SLAPETA, J. R., B. KOUDELA, J. VOTYPKA, D. MODRY, R. HOREJS, AND J. LUKES. 2002. Coprodiagnosis of *Hammondia heydorni* in dogs by PCR based amplification of ITS 1 rRNA: Differentiation from morphologically indistinguishable oocysts of *Neospora caninum*. *Veterinary Journal* **163**: 147–154.
- WHITE, T. J., T. BRUNS, S. LEE, AND J. TAYLOR. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR protocols, a guide to methods and applications*, M. A. Innis, D. J. Gelfand, J. J. Sninsky, and T. J. White (eds.). Academic Press, San Diego, California, p. 315–322.
- WILLIAMS, J. H., I. ESPIE, E. VAN WILPE, AND A. MATTHEE. 2002. Neosporosis in a white rhinoceros (*Ceratotherium simum*) calf. *Journal of the South African Veterinary Association* **73**: 38–43.
- WOODS, L. W., M. L. ANDERSON, P. K. SWIFT, AND K. W. SVERLOW. 1994. Systemic neosporosis in a California black-tailed deer (*Odocoileus hemionus columbianus*). *Journal of Veterinary Diagnostic Investigation* **6**: 508–510.
- YAMAGE, M., O. FLECHTNER, AND B. GOTTSSTEIN. 1996. *Neospora caninum*: Specific oligonucleotide primers for the detection of brain “cyst” DNA of experimentally infected nude mice by the polymerase chain reaction (PCR). *Journal of Parasitology* **82**: 272–279.