



Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Ecology and biogeochemistry of contrasting trophic environments in the South East Pacific by carbon isotope ratios on lipid biomarkers

I. Tolosa¹, J.-C. Miquel¹, B. Gasser¹, P. Raimbault², L. Azouzi³, and H. Claustre⁴

¹International Atomic Energy Agency – Marine Environment Laboratories, 4, quai Antoine 1er, MC 98000, Monaco

²Laboratoire d'Océanographie et de Biogéochimie (UMR 6535 CNRS), Centre d'Océanologie de Marseille, Univ. de la Méditerranée, Campus de Luminy, 13288 Marseille Cx 09, France

³BDSI, Université de Perpignan, 52 avenue Paul Alduy, 66860 Perpignan, France

⁴Laboratoire d'Océanographie de Villefranche, Observatoire Océanologique, INSU/CNRS/UPMC, UMR 7093, BP 08, 06230 Villefranche-sur-Mer, France

Received: 28 November 2007 – Accepted: 28 November 2007 – Published: 18 December 2007

Correspondence to: I. Tolosa (i.tolosa@iaea.org)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The distribution of lipid biomarkers and their carbon isotope composition was investigated on suspended particles from different contrasting trophic environments at six sites in the South East Pacific. High algal biomass with diatom-related lipids was characteristic in the upwelling zone, whereas haptophyte lipids were proportionally most abundant in the nutrient-poor settings of the centre of the South Pacific Gyre and on its easter edge. Dinoflagellate–sterols were minor contributors in all of the studied area and cyanobacteria-hydrocarbons were at maximum in the high nutrient low chlorophyll regime of the subequatorial waters at near the Marquesas archipelago.

The taxonomic and spatial variability of the relationships between carbon photosynthetic fractionation and environmental conditions for four specific algal taxa (diatoms, haptophytes, dinoflagellates and cyanobacteria) was also investigated. The carbon isotope fractionation factor (ε_p) of the diatom marker varied over a range of 16‰ along the different trophic systems. In contrast, ε_p of dinoflagellate, cyanobacteria and alkenone markers varied only by 7–10‰. The low fractionation factors and small variations between the different phytoplankton markers measured in the upwelling area likely reveals uniformly high specific growth rates within the four phytoplankton taxa, and/or that transport of inorganic carbon into phytoplankton cells may not only occur by diffusion but by other carbon concentrating mechanisms (CCM). In contrast, in the oligotrophic zone, i.e. gyre and eastgyre, relatively high ε_p values, especially for the diatom marker, indicate diffusive CO₂ uptake by the eukaryotic phytoplankton. At these nutrient-poor sites, the lowest ε_p values for haptophytes, dinoflagellates and cyanobacteria infer higher growth rates compared to diatoms.

1 Introduction

The sustainability of phytoplankton growth in the world ocean is basically controlled by three factors: nutrient abundance, light availability and the response of phytoplankton

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

EGU

to these sources (Falkowski, 1984; Falkowski et al., 1998; Irwin et al., 2006; Litchman et al., 2006). The variability of these factors and their role in biogeochemical processes emerges from the properties of the surface mixed layer. Hence, eutrophic areas, such as upwellings, with continuous nutrient supply to the euphotic zone differ strongly from areas with a permanently nutrient-depleted surface layer. In these areas, e.g. oceanic gyres, a deep pycnocline prevents surface waters from a supply with deeper, more nutrient-rich waters.

Photosynthesis is a major biogeochemical process where carbon dioxide and water are converted into organic carbon with the presence of light. The fate of this organic carbon is therefore intimately linked to the conditions of its synthesis and depends strongly on the composition of the phytoplankton assemblages. For understanding the global marine carbon cycle, efforts should be made to derive information relative to these ecosystems. In addition to the now widely used chlorophyll and pigment analysis (Barlow et al., 1993; Claustre et al., 2004; Mackey et al., 1996), field data about phytoplankton diversity might also be acquired by the distribution of accessory lipid biomarkers. They provide important information on the phytoplankton composition complementary to that of algal pigments, as well as on the relative importance of carbon sources from heterotrophic bacteria and other zooplankton (Dijkman and Kromkamp, 2006; Pinturier-Geiss et al., 2002; Tolosa et al., 2004).

Isotopic characterization of marine organic matter can provide insight into the conditions under which carbon fixation occurs contributing to the understanding of the global marine carbon cycle. The carbon isotopic composition ($\delta^{13}\text{C}$) of any photosynthetic material and their derived carbon isotope fractionation factor (ε_p) reflects the $\delta^{13}\text{C}$ of the carbon source utilized and the modification by the processes and environmental variables involved in its production: growth rate, temperature, dissolved CO_2 , cell geometry, irradiance, etc. (Burkhardt et al., 1999a; Burkhardt et al., 1999b; Eek et al., 1999; Hayes, 1993; Laws et al., 1997; Popp et al., 1998b; Rau et al., 1996). For example, high growth rates of the phytoplankton producers are linked to high $\delta^{13}\text{C}$ values, i.e. they are enriched in ^{13}C (Bidigare et al., 1999; Laws et al., 1995). However, they

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

 $\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

become depleted in ^{13}C with increasing the concentration of dissolved CO_2 (Burkhardt et al., 1999a; Riebesell et al., 2000). This latter linkage seems to be limited in its extent, and differences of $\delta^{13}\text{C}$ higher than 1–2‰ cannot be explained any more by the change in the CO_2 concentration, but they are rather accounted for by differences in the growth rates of the phytoplankton producers and carbon uptake mechanisms (Bidigare et al., 1997; Burkhardt et al., 1999b, Bentien et al., 2007). Since particulate organic carbon (POC) is a complex mixture of autotrophs, heterotrophs, and detritus, potentially clearer relationships between $\delta^{13}\text{C}$ values and environmental conditions can be obtained using specific biomarkers derived from particular species or taxonomic groups (Bidigare et al., 1999; Pancost et al., 1999; Pancost et al., 1997), compared to the $\delta^{13}\text{C}$ of the bulk POC (Rau et al., 2001; Woodworth et al., 2004). This has become possible with compound-specific isotope-ratio mass spectrometry (Freeman et al., 1990; Hayes et al., 1990).

Our field study uses molecular and isotopic carbon ratios of specific lipid biomarkers to evaluate the sources of the organic matter and to explore variations in the biogeochemistry of the particulate organic matter at the different hydrodynamic and trophic environments from the South East Pacific. These include the eutrophic upwelling area off the Chilean coast, the mesotrophic and HNLC regions south of the equatorial current and the oligotrophic South Pacific Gyre, which was the major focus of this work. The major questions being addressed are:

- 1) What is the spatial distribution of lipid biomarkers of phyto-, zooplankton and bacteria in the contrasting trophic environments of the South East Pacific?
- 2) What is the variability of the carbon isotope fractionation (ε_p) and growth rates of different phytoplankton taxa in relation to the environmental conditions characterized by different nutrient, dissolved inorganic carbon concentrations and productivity regimes? We discuss these results with respect to the processes of carbon uptake and assimilation within the different phytoplankton groups from the South-East Pacific Ocean.

2 Materials and methods

2.1 Sampling

Sampling and hydrographic observations were carried on board R/V "L'Atalante" between October and December 2004 and were organized within the framework of the BIOSOPE (Biogeochemistry & Optics South Pacific Experiment) project which is part of the JGOFS French oceanographic programme PROOF. The main hydrodynamical and trophic features for the different zones are described in Claustre et al. (2007)¹.

Samples were taken at six different sites, so called "long stations", between Tahiti and the Chilean coast (Table 1). The different explored zones exhibited contrasting trophic environments. First, a mesotrophic area downstream of the Marquesas Islands (MAR) and a high nutrient low chlorophyll zone (HNL) upstream of the Islands. Second, an extremely oligotrophic area, very poor in nutrients, located in the centre of the South Pacific Gyre (GYR) and a less oligotrophic site in the east of the gyre (EGY). At the end of the transect, we studied a eutrophic zone highly enriched in nutrients and associated to the upwelling off the Chilean coast (UPW and UPX). UPW station was farther from the coast than UPX and exhibited a more important water stratification.

"Challenger Oceanics" in-situ pumps were used to filter large volumes (400 to 900 l) of water from the upper 300 m of the water column, to collect suspended particles through a Nitex screen of 70 µm and a precombusted (550°C) Microquartz filter (QMF, Sartorius) of 1 µm pore size.

2.2 Bulk measurements

Table 2 summarizes the bulk biochemical parameters of the suspended particulate matter along the transect Marquesas Islands-Chilean coast.

¹ Claustre, H., Sciandra, A., and Vaulot, D.: Introduction to the special section: bio-optical and biogeochemical conditions in the South East Pacific in late 2004 – the BIOSOPE cruise, Biogeosciences Discuss., submitted, 2007.

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

EGU

Total CO₂ (C_T) and total alkalinity (A_T) of water samples were measured by potentiometry (Azouzi et al., 2007) at all sites except UPW. The dissolved CO₂ concentration was calculated from C_T, alkalinity, temperature, salinity and the concentrations of silicate and phosphate using the CO2SYS program developed for CO₂ system (Lewis and Wallace, 1998). This program is based on equations of the seawater CO₂ system (DOE, 1994) and the dissociation constants of Goyet and Poisson (1989). For the UPW site, we assumed the same concentrations as UPX, since vertical profiles for aqueous CO₂ interpolated from the WOCE cruise P06 in GLODAP three-dimensional gridded data at the two stations indicate similar concentrations.

Nutrient concentrations (nitrate, phosphate and silicate) were determined onboard using an autoanalyzer (Raimbault et al., 2007).

Analysis of organic carbon was done with a “Vario EL” elemental analyser (© elementar Analysensysteme GmbH) after acidification of the filter aliquots (Miquel et al., 2007²).

2.3 Lipid extraction

Filters containing the suspended particles were spiked with internal standards (*n*-C₂₄D₅₀, anthracene-*d*₁₀, pyrene-*d*₁₀, perylene-*d*₁₂, friedeline, 5α-androstan-3β-ol and cholic acid), and extracted by microwave oven with 40 ml of a mixture with CH₂Cl₂/MeOH (3:1) at 70°C for 15 min. Isolation of the neutral and acid lipid fractions were done following the method of Tolosa and de Mora (2004). Extractable lipids were saponified using 1 ml KOH 6% in methanol/water (80:20) plus 1 ml of Milli-Q water (80°C, 1 h). Then the neutral fraction was recovered with *n*-hexane and subject to fractionation by HPLC on a normal phase column (Nucleosil column, 20 cm × 0.4 cm i.d. 5 μm) to isolate the aliphatic hydrocarbons (F1), polycyclic aromatic hydrocarbons (F2), ketone compounds (F3) and sterol and alcohol fraction (F4). Saponified solutions

²Miquel, J. C. and Gasser, B.: Downward particulate flux and carbon export in contrasting environments of the South East Pacific, Biogeosciences Discuss., in preparation, 2007.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

were acidified with 1 ml HCl 6 N to pH 2 and the fatty acids obtained by hydrolysis of wax esters, triacylglycerols, steryl esters and phospholipids were extracted with hexane:ethyl acetate 9:1.

2.4 Gas chromatography

- 5 The sterol fraction was treated with *bis*-(trimethylsilyl)-trifluoroacetamide (BSTFA) (200 µl, 70°C, 1 h) to convert the alcohols and sterols to their corresponding trimethylsilyl ethers. The acid fraction was derivatized by transesterifying the lipid extract with 500 µl of 20% BF₃ in methanol at 80°C for 1 h.

Gas chromatography (GC) was performed with a Hewlett Packard HP5890 series
10 II equipped with a flame ionization detector and split/splitless injector. Two fused
silica capillary columns were employed: (A) a DB-5 fused silica capillary column
(30 m × 0.25 mm i.d.; film thickness 0.25 µm) for neutral compounds and fatty acids and
(B) a BPX-70 (SGE, 60 m × 0.32 mm × 0.5 µm) for the fatty acids. Helium was the carrier gas (1.2 ml min⁻¹). The oven temperature for the DB-5 was programmed from
15 60°C (0.5 min hold) to 290°C at 6°C min⁻¹. The GC oven for the BPX-70 column was
programmed from 60°C (0.5 min hold) to 250°C at 6°C min⁻¹. Injector and detector
temperatures were 270°C and 320°C, respectively.

Aliphatic hydrocarbons, ketones, sterols and fatty acids were quantified by internal
standards (*n*-C₂₄D₅₀, friedeline, 5α-androstan-3β-ol, and cholic acid, respectively).
20 Confirmation of peak identity was obtained using GC with mass spectrometric detection
(GC-MS) (Hewlett-Packard 5889B MS “Engine”) operated in the electron impact at
70 eV.

2.5 Compound-specific isotope analysis

The lipid biomarkers were analyzed for their stable carbon isotope composition using
25 an HP 5890 GC equipped with a HP 7673 autoinjector and interfaced through a
combustion furnace with a FINNIGAN MAT Delta C isotope ratio mass spectrometer

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 The GC/C/IRMS was equipped with a 100% methylpolysiloxane fused silica column (Ultra-1, 50 m × 0.32 mm i.d.; 0.5 µm film thickness) pre-connected with a press-fit connector (Supelco, France) to a 0.32 mm i.d. deactivated fused silica capillary retention gap of 5 m. Injections of 2 µl in isoctane were made via an on-column injector. The GC oven was programmed from 60 to 100°C at 10°C min⁻¹, then to 310°C at 4°C min⁻¹ and maintained at 310°C for 40 min. Values reported were determined by at least three replicates to calculate the average and standard deviation. All δ¹³C values are reported in the delta notation relative to the Pee Dee Belemnite (PDB) standard as follows:

$$10 \quad \delta^{13}\text{C} = [(\text{sample}/\text{PDB}) - 1] \times 10^3$$

Corrections for the isotopic change introduced in the derivatisation of sterols, fatty alcohols, and fatty acids were determined through derivatisation of standards of known isotopic composition and applying the equation of Jones et al. (1991). Cholesterol, methanol, 18:0 fatty acid and 18:0 FAME of known isotopic carbon composition (measured by elemental analyser coupled to isotope ratio mass spectrometer), were used to calibrate the GC/C/IRMS and correct the isotopic change introduced by the derivatization. The surrogate standards, 5α-androstan-3β-ol, cholic acid and the GC internal standard friedelin of known isotopic composition served as internal isotopic standards.

The precision (standard deviation) for most analytes with GC-C-IRMS signals higher than 0.5 V (*m/z* 44) was comparable to the instrument specifications (0.5‰).

2.6 Calculations of carbon isotope fractionation (ε_p)

Molecular ε_p was determined following the general equation (1) outlined in Freeman and Hayes (1992):

$$25 \quad \varepsilon_p = [(\delta^{13}\text{CO}_2 + 1000)/(\delta^{13}\text{C}_{pp} + 1000) - 1] \times 10^3 \quad (1)$$

where CO₂ is its dissolved phase in the water column and C_{pp} the primary photosyntheticate.

δ¹³C on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

In this study, direct measurement of $\delta^{13}\text{CO}_2$ was not available. Therefore $\delta^{13}\text{CO}_2$ was calculated according to the Eq. (2) of Mook (1974):

$$\varepsilon_b = [(\delta^{13}\text{CO}_2 + 1000)/(\delta^{13}b + 1000) - 1] \times 10^3 = 24.12 - 9866/T \quad (2)$$

where ε_b is the temperature-dependent carbon isotope fractionation of dissolved CO_2 with respect to bicarbonate, T is the absolute temperature in Kelvin, and the reference value of δ^{13} for bicarbonate (b) in sea surface water was taken as +2.2‰ (Craig, 1970).

$\delta^{13}\text{C}_{pp}$ (primary photosynthate) for eukaryotic organisms was estimated by using a constant isotopic fractionation between photosynthetic lipids and algal biomass, which has been estimated at 4.2‰ for alkenones (Popp et al., 1998a). This value is assumed to be the same for the rest of eukaryotic photosynthetic compounds (Hayes, 1993). For prokaryote, $\delta^{13}\text{C}_{pp}$ was estimated from the n-heptadecane using a constant isotopic fractionation between photosynthetic lipids and algal biomass of 8.4‰ (Sakata et al., 1997).

15 2.7 Estimations of growth rates and intracellular carbon demand in haptophytes

Carbon isotopic fractionation for phytoplankton (ε_p) which obtain CO_2 by passive diffusion is summarized by the expression of Popp et al. (1998b):

$$\varepsilon_p = \varepsilon_f - \beta \frac{\mu(V/S)}{[\text{CO}_2]} \quad (3)$$

where ε_f is the fractionation associated with the enzyme-catalyzed carbon fixation step, β is a constant, μ is the specific growth rate, V and S are the volume and surface area of the alga cells and $[\text{CO}_2]$ is the concentration of dissolved CO_2 external to the algal cell. Since β and (V/S) are practically constant for haptophyte taxa, we can transform this constant and the variable μ into the b-value (‰μmol), which serves as a proxy for

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

growth rate and reflects the intracellular carbon demand. This b-value was calculated following the Eq. (4) of Bidigare et al. (1997):

$$b = (\varepsilon_f - \varepsilon_p) \times [\text{CO}_2]_{\text{aq}} \quad (4)$$

with ε_f values of 25‰ for eukaryotic algae utilizing Rubisco and β -carboxylase enzymes (Bidigare et al., 1997) and $[\text{CO}_2]_{\text{aq}}$ calculated as described in Sect. 2.2.

Specific growth rates (μ , (d⁻¹)) of alkenone producing haptophytes were estimated with the following equation found by Bidigare et al. (1997) in laboratory chemostat culture experiments of *Emiliania huxleyi*:

$$\mu_{cc} = (25 - \varepsilon_p)[\text{CO}_2]/138 \quad (5)$$

and applying the corrections for the effects of day length and respiration on growth rate

$$\mu = [\mu_{cc}/(24/t_p)]0.8 \quad (6)$$

where μ is the 24-h average growth rate, t_p is day length or photoperiod in hours, and the factor 0.8 adjusts the growth rate for dark respiration.

3 Results and discussion

The analytical scheme used in this study identified and quantified ~60 individual compounds in the neutral lipid fraction and ~40 acid compounds in the acid fraction. A summary of selected lipid biomarkers discussed in this study together with their sources is shown in Table 3. Concentrations of the selected lipid biomarkers are summarized in Table 4 and the individual carbon isotope ratio for some of the selected lipid biomarkers are shown in Annex A. The complete data set of concentrations and $\delta^{13}\text{C}$ values for those possible are available on the BIOSOPE Database: <http://www.obs-vlfr.fr/proof/vt/op/ec/biosope/bio.htm>.

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

3.1 Distribution of phytoplankton, zooplankton and bacterial markers in different trophic environments

BGD

4, 4653–4696, 2007

Concentrations of phytol, a non-specific marker for phototrophic organisms, if compared at the depth of chlorophyll maximum, were highest at the upwelling sites with 102 ng l⁻¹ at UPW and 78 ng l⁻¹ at UPX. The mesotrophic sites, MAR and HNL, exhibited intermediate phytol concentrations of 25 to 31 ng l⁻¹. The lowest values were measured at the oligotrophic sites, EGY (16 ng l⁻¹) and in particular at GYR (11 ng l⁻¹), where the chlorophyll maximum was at 175 m depth. Concentrations of diatom biomarkers, e.g. (C₂₈Δ^{5,24(28)} sterol, C₂₅ HBI alkenes, C_{16:4} FA, C_{20:5} FA), haptophytes biomarkers (total alkenones) and dinoflagellates markers (dinosterol) exhibited a similar distribution as phytol concentrations (Table 4), except in the Gyre where alkenones and dinosterol peaked at lower depths than phytol and diatom markers.

Highest concentrations of long-chain C₃₇ and C₃₈ alkenones were measured at the eutrophic UPW site (27 ng l⁻¹) but also at the two gyre sites EGY (20 ng l⁻¹) and GYR (23 ng l⁻¹). These peak values corresponded to the depth of chlorophyll maximum except for the GYR site where the peak was situated above this maximum, at 125 m depth. A much lower concentration (6.5 ng l⁻¹) was recorded at the MAR site. These values are much lower than those reported for suspended particles from the Bering Sea after blooms of *Emiliana Huxleyi* and ranging from 0.15 to 3.12 μg l⁻¹ (Harada et al., 2003), but similar to concentrations observed in suspended particles collected under non-bloom conditions in the surface waters of the North Atlantic and Nordic Sea (Sicre et al., 2002), in the western Sargasso Sea (100 ng l⁻¹) (Conte et al., 2001) and in the oligotrophic North Pacific subtropical gyre (0.5–15 ng l⁻¹) (Prahl et al., 2005).

The C₁₇ n-alkane, which is produced by aerobic photosynthetic bacteria and green algae (Han and Calvin, 1969; Winters et al., 1969) exhibited a maximum concentration of 2.3 ng l⁻¹, below the chlorophyll maximum at the HNL site and of 1.2 ng l⁻¹ at the same depth (70 m) as the chlorophyll maximum at the EGY site. The other sites showed concentration levels below 0.5 ng l⁻¹ (Table 4). These concentrations were

δ¹³C on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

consistent with the abundance distribution of prokaryotic phototrophic organisms, with high abundances at the HNL and EGY sites (Grob et al., 2007).

In all samples, the *n*-alkanols were dominated by the short-chain fatty alcohols of even carbon number (*n*-C14, *n*-C16 and *n*-C18), which are associated to zooplankton markers (Sargent et al., 1977). With the exception of the two gyre sites (GYR and EGY), maximum concentrations of lineal alcohols were found below the depth of chlorophyll maximum, at the upwelling sites (UPW, UPX) at 300 m and at the Marquesas sites (MAR, HNL) at 100 m depth. In the Gyre, maximum concentrations of n-alcohols coincided with the phytol maximum, though another peak (11.6 ng l⁻¹) was registered at the surface of the GYR site. The fatty acids C_{20:1} and C_{22:1}, typical markers of herbivorous mesozooplankton (Lee et al., 2006), exhibited the highest concentrations at the upwelling sites (UPW, UPX) at the depth of chlorophyll maximum, but also below the euphotic zone (300 m) at the MAR and UPX sites.

Similar to the phytol distribution, the concentrations of bacterial biomarkers, such as branched fatty acids, were highest at the UPW site (Table 3), whereas the concentrations of zooplankton markers, such as cholesterol and n-alcohols exhibited higher abundance at UPX. In general, heterotrophic bacterial populations seemed to be associated to diatom biomass, which is supported by the positive correlation between the δ¹³C of the branched fatty acid (*i*-C₁₅ FA) and the δ¹³C of the C_{20:5} FA (*r*=0.81, *p*<0.05).

Some more insight into the phytoplankton distribution may be gained by comparing the relative contribution of the biomarkers within the total neutral lipids, which are illustrated in Fig. 1. The percentage of phototrophic biomarkers generally followed the chlorophyll distribution except at UPX, where phytol and diatoms markers peaked at 100 m depth. Also at HNL, phytol showed relatively high percentages down to 100 m depth. This may be related to the highest relative importance of diatom sterol observed in the euphotic layer (Fig. 1b) and to the presence of diatoms which formed “balls of needles” or clusters (Gómez et al., 2007). In contrast to the diatom biomarkers, the percentage of total alkenones was by far highest in the gyre especially above the chloro-

BGD

4, 4653–4696, 2007

δ¹³C on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

phyll maximum (Fig. 1c), where also concentrations had been among the highest of all sites (Table 4). Prymnesiophytes were likely to be the major constituents of eukaryotic phytoplankton in the mixed surface layer of the gyre, suggesting that haptophytes are well adapted to the low nitrate concentrations prevailing in the oligotrophic zone of the Pacific Gyre. The $C_{28}\Delta^{5,24(28)}$ sterol/alkenones ratio provides us with information about the relative contribution of diatoms to prymnesiophytes. Highest ratios were obtained at the UPX site, and in particular below the euphotic zone. Haptophytes predominated over diatoms at the GYR site, especially above the chlorophyll maximum whereas the diatom signal showed a deeper maximum at 175 m corresponding to the maximum of both phytol and chlorophyll *a*. Dinosterol showed overall low percentages (<3%), which indicated a minor contribution of dinoflagellates in the algal mixture of these Pacific waters, as it was also confirmed by pigment analyses (Ras et al., 2007).

Figure 2 illustrates other diagnostic biomarkers indices to evaluate the relative dominance of zooplankton and bacterial sources within each site as well as the state of the particulate material in the different zones. The cholesterol/phytosterol ratio which indicates dominance of zooplankton over phytoplankton increased with depth at all sites, and in particular at UPX. This was consistent with the substantial concentrations of n-alcohols and zooplanktonic $C_{20:1}$ and $C_{22:1}$ fatty acids found at 300 m depth. At most of the sites, the relative importance of bacterial fatty acids was higher below than at the depth of chlorophyll maximum. A contrasting image was observed at UPX where this relative importance was lower. In fact, high bacterial production and a negative net community production were reported from the euphotic zone of UPX (Van Wambeke et al., 2007), which suggests rather “decomposing” conditions compared to “productive” conditions at UPW. At all sites, the ratio of phytosterols/phytol indicated more degraded phytoplankton material at depths below the chlorophyll maximum, but also above, at the site of the gyre. This ratio showed a slightly more degraded material in the euphotic zone of the UPX site compared to UPW, but fresher material at 300 m depth of UPX.

The polyunsaturation index of C_{16} fatty acids (PUFA % of C_{16}) is an indicator of the ecophysiological state of marine diatom populations because storage lipids, mainly

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$\text{C}_{16:0}$ and $\text{C}_{16:1}$ FA, are synthesized during senescence, rather than during logarithmic growth (Shin et al., 2000). The high indices observed in the euphotic zone of the UPW site and at 300 m depth of UPX suggest that these PUFA originated from diatoms at logarithmic growth. Moreover, the carbon isotope ratios of lipid biomarkers in the euphotic zone were generally more enriched at UPW compared to UPX (Fig. 3), likely indicating higher growth rates at UPW than at UPX. However, below the euphotic zone of UPX, $\delta^{13}\text{C}$ values identified higher growth rates at depth compared to the surface.

- 5 All these parameter point out that post-bloom conditions with high concentrations of animal-derived detritus prevailed at the surface of the UPX site, whereas the important signal of zooplankton and diatom markers below the euphotic layer indicated the presence of zooplankton feeding on phytoplankton produced during bloom conditions. These findings are supported by the highest stock and flux measured at UPX compared to the UPW site (Miquel et al., 2007²) and the high concentration of detritus and senescent colonial diatoms observed by microscope in samples from the euphotic zone 10 at the UPX site (Gómez, personal communication, 2006). These conclusions contrast, however, with those derived from pigment biomarkers (Ras et al., 2007) where UPW site was characteristic of a typically mature bloom of diatoms and the phytoplankton at site UPX was probably at an early stage of development. Also, much higher nutrient concentrations at the surface of UPX site indicated a recent upwelling of deep water. 15 Overall, these contrasted observations might be accounted for by the different turnover and lability between pigment and lipid compounds which represent different pools of the particulate matter. Pigments are relatively more labile and associated with the living material. In contrast, lipids are included in both, the living and detrital particulate pools of the matter. 20

25 3.2 Vertical distribution of biomarkers in the center of the gyre

Depth profiles of selected accessory lipid biomarkers in suspended matter from the center of the Gyre are presented in Fig. 4. Phytol concentrations showed very low surface values and they were increasing progressively with depth, with maximum concen-

tration at 150–200 m depth. Relatively high concentrations of *n*-alcohols, cholesterol and branched fatty acids at the surface indicated an important heterotrophic activity in the upper waters. At higher depths (>75 m), both alcohols and cholesterol showed similar profiles as phytol, likely indicating that here, they were mainly phytoplanktonic-derived or that zooplankton biomass was strongly associated with the phytoplankton abundance. Branched fatty acids which are derived from heterotrophic bacteria exhibited the maximum concentration between 150 and 200 m depth, following the same trend as planktonic biomass. This feature indicates that the bacterial population is associated with the major planktonic biomass.

Other more specific phototrophic biomarkers, such as sterol markers for diatoms exhibited a similar profile as phytol with two maximum at 150 and 200 m depth. Only few macro diatom species, such as *Nitzschia* and *Dactyliosolen* were observed between 200 and 300 m depth and a significant number of *Bacteriastrum* associated to a cyanobacteria symbiont was also observed around 140 m depth (Gómez, personal communication, 2006). In contrast, the sterol marker for dinoflagellates exhibited a uniform distribution from surface to 125 m depth and a deeper maximum concentration at 200 m depth. Small dinoflagellates were observed in the surface waters of the Gyre center (Gómez, personal communication, 2006). The *n*-alkane C₁₇, which is produced by cyanobacteria and other eukaryotic algae, increased with depth showing a maximum concentration between 125 and 175 m. This coincides with the maximum abundances of *Prochlorococcus* and picoeukaryotes recorded between 100 and 200 m depth (Grob et al., 2007).

The C₃₇ alkenones, which are specific markers for some algae of the class *Haptophyceae/Prymnesiophyceae*, including the coccolithophorid species such as *Emiliania huxleyi*, exhibited the maximum concentration above the deep chlorophyll maximum, at 125 m depth. Cell densities of different coccolithophorid taxa showed, however, different depth profiles, with a maximum peak between 150–200 m for *Emiliania huxleyi* whereas other taxa peaked at shallower depths (~100 m) (Beaufort et al., 2007). These findings indicate that (i) other noncalcifying haptophytes might synthesize alkenones at shal-

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

 $\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lower depths, (ii) that alkenones are not associated with the integrity of coccospores and/or (iii) that cellular alkenone concentrations varied with the physiological status and species composition of the coccolithiphorid assemblage. On the other hand, the concentration of alkenones and the accessory carotenoid 19'Hexanoyloxyfucoxanthin (19'HF) (Ras et al., 2007), characteristics of prymnesiophytes, also exhibited different depth distributions (Fig. 5). Such discrepancy reflects that alkenone-producers in these waters are small contributors to the 19'HF stock, and that the habitat of alkenone synthesizers diverges from that of the major phytoplankton taxa contributing to the 19'HF distribution. An analogous feature was observed at station ALOHA from the oligotrophic North Pacific Subtropical Gyre (Prahl et al., 2005) and other studies showed that 19'HF abundance was generally not tightly correlated with that of coccolithophorids (Dandonneau et al., 2006). Figure 6 illustrates the carbon isotope composition of the diunsaturated alkenone together with the total concentrations of C₃₇ alkenones. More enriched $\delta^{13}\text{C}$ values which infer higher growth rates were obtained for those haptophytes inhabiting the depth of the chlorophyll maximum. However, the higher concentrations of alkenones found at 125 m depth were associated to lowest growth rates which indicates a higher concentration of cellular alkenones. Similar observations were reported in nitrate-limited cultures where a small increase in the concentration of cellular alkenones occurred with decreasing growth rates (Conte et al., 1998; Parrish et al., 1998).

The growth unsaturation index ($U_{37}^{K'}$) which is widely used as a proxy of sea surface temperature (Prahl and Wakeham, 1987) was calculated as the relative proportion of di-and triunsaturated C₃₇ alkenones. This index was converted to a measure of temperature by the commonly used empirical calibration equation $T = (U_{37}^{K'} - 0.039)/0.034$ (Prahl et al., 1988). The obtained results (Fig. 7), show that the in situ temperatures of the surface layer were overestimated by 2–3 degrees. Discrepancies between the alkenone-calculated and observed temperatures might be caused by stress due to nutrient limitation and to differences in the stage of the growth cycle (Conte et al., 1998; Epstein et al., 1998; Yamamoto et al., 2000), which result in variable alkenone synthe-

sis. Similar observations were reported in winter at ALOHA station (Prahl et al., 2005). They were explained by simply biogeographical variations observed in the alkenone vs. temperature relationship in natural waters, which may reflect differences in genetic and physiological status of the local alkenone-synthesizing populations. Since 5 haptophytes have a low phosphorous requirement (critical concentrations of $0.2 \mu\text{M}$), nitrogen limitation seems likely since inorganic N concentrations at the Gyre stations were well below the half-saturation constant (K_s , the concentration supporting an uptake rate one-half the maximum rate) determined for *E. Huxley* ($\leq 0.5 \mu\text{M}$) (Eppley et al., 1969). Overall and according to batch cultures of haptophytes (Epstein et al., 1998), 10 the observed increase in the $U_{37}^{K'}$ values with the consequent overestimation of the temperatures, might indicate that this marine phytoplankton taxon is under nutrient-limited “stationary growth” conditions. This overestimation can also be explained by a change 15 in the ecology of alkenone-producing algae (Popp et al., 2006) and by autoxidation of alkenones in these highly irradiated waters (Rontani et al., 2006), especially when residence times of particles are long.

3.3 Biogeochemical implications from carbon isotope fractionation

Stable carbon isotope differences between the inorganic carbon source and that of organic carbon synthesized by autotrophic organisms known as photosynthetic carbon fixation (ε_p), can assist in distinguishing between the different CO_2 fixation pathways 20 (Table 5). Maximum carbon isotope fractionation of photoautotrophic organisms using the Calvin cycle, like micro-algae and cyanobacteria is in the range of 20 to 27‰ (Popp et al., 1998b; Sakata et al., 1997). However, the ε_p expected for biomarkers derived 25 from eukaryotes can vary between 5 and 25‰ depending basically on $[\text{CO}_2]$, growth rate and the ratio of cellular surface area to volume (Bidigare et al., 1997a; Popp et al., 1998). In contrast, ε_p for prokaryotes (cyanobacteria) range between 16 and 22‰ because the large surface-to-volume ratio guarantees a large CO_2 supply relative to the cellular demand. Others pathways, apparently restricted to other bacteria, such as anoxygenic phototrophic bacteria, are the reversed tricarboxylic acid cycle and the

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3-hydroxypropionate pathway, both of which are characterized by significantly smaller isotope effects (ε_p of 2–14‰) (van der Meer et al., 2001).

We observed higher taxonomic variations in ε_p for eukaryotic algae growing in the oligotrophic areas (variations of ~10‰) compared to the eutrophic sites of the upwelling (variations of 3 to 7‰). The variation of the carbon isotope fractionation for the diatom marker covered a range of ~16‰ along the different trophic systems. In contrast, ε_p of dinoflagellate and alkenone markers varied much less ca. 10 and 7‰, respectively.

The carbon isotope fractionation of diatom markers exhibited a significant, negative logarithmic correlation with measured silicate concentrations ($\varepsilon_p = -4.3 \ln[\text{SiOH}_4] + 23.57$, $n=16$, $r=0.96$, $p<0.01$, Fig. 8), whereas lower correlations were found with the other nutrients, nitrates and phosphates. Lower correlations were also obtained with the ε_p of dinoflagellates and no correlation was observed between ε_p of alkenone and nutrients. These findings confirm (Pancost et al., 1999) that silicate is likely one of the essential nutrients which control the ε_p and growth rates of diatoms in the Pacific ocean.

The carbon isotope fractionation of diatom and dinoflagellate markers showed a negative, linear correlation with $[\text{CO}_2]_{\text{aq}}$ ($r_{\text{diatom}}=0.94$ (Fig. 9a), $r_{\text{dinoflagellate}}=0.91$, $n=16$, $p<0.01$). These relationships deviate from the previously reported general oceanic trend (Rau et al., 2001) and culture studies (Burkhardt et al., 1999a) where carbon isotope fractionation increases ($\delta^{13}\text{C}$ decrease) when $[\text{CO}_2]_{\text{aq}}$ increases. However, this apparent deviation has also been observed in Peruvian upwelling waters where it was suggested that a diatom carbon concentrating mechanism (CCM) was likely the cause of the lower ε_p of diatoms in these waters with high $[\text{CO}_2]_{\text{aq}}$. A much lower correlation ($r=0.75$, $n=14$, $p<0.01$) and a slope close to zero was observed for the ε_p of alkenone (Fig. 9b) which agrees with other studies that showed the small effect of $[\text{CO}_2]_{\text{aq}}$ on isotope fractionation in *E. Huxleyi* compared to potential changes of ε_p due to growth rate and carbon uptake mechanisms (Benthien et al., 2007; Bidigare et al., 1997). No correlation was found between $[\text{CO}_2]$ and the ε_p of *n*-C17 ($r_{\text{CO}_2}=0.37$, $n=6$,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$p=0.46$) which is also consistent with a previous work (Popp et al., 1998b) who found for *Synechococcus* that ε_p is independent of the concentration of dissolved CO₂, likely because its cell geometry guarantees a large CO₂ supply.

Overall, the lower isotope fractionation factors and smaller variations between the different eukaryotic markers measured in the upwelling area might indicate uniformly high growth rates for the three phytoplankton taxa, diatoms, haptophytes and dinoflagellates and/or that phytoplankton may employ carbon concentrating mechanisms (CCM) other than diffusion, which actively transport inorganic carbon into cells. Similar findings were reported by other authors (Pancost et al., 1999; Pancost et al., 1997; Rau et al., 2001; Werne and Hollander, 2004) who invoked that an active transport of bicarbonate into the cell may still play a role in the carbon isotope fractionation by phytoplankton in upwelling areas with high concentrations of CO₂.

In contrast, the GYR and EGY sites exhibited the highest carbon isotope fractionation factors for eukaryotic algae and in particular for the diatom marker. Their values reached 25–26‰ which is close to the maximum isotope fractionation of eukaryotic algae utilizing Rubisco and β -carboxylase enzymes (Goericke et al., 1994; Laws et al., 1997). Such high ε_p values cannot be obtained by bicarbonate uptake and are indicative of diffusive CO₂ uptake. Moreover, active uptake of carbon in oligotrophic sites is rather unlikely because of the higher metabolic energy required and CCM may be inhibited by the low oceanic concentrations of certain trace metals (Morel et al., 1994). In these oligotrophic sites, lowest ε_p values were measured for the haptophytes and dinoflagellates, which might infer higher growth rates for these organisms compared to diatoms.

In the mesotrophic areas of the Marquesas Islands (MAR and HNL sites), ε_p values for alkenone producers and dinosterol were also lower than those for diatoms, but still higher than those estimated in the upwelling area. The similar ε_p values for alkenones measured in the poor-nitrated waters of the Gyre and in the high nutrient waters of the HNL site suggest that nitrogen and phosphate are not the limiting nutrients affecting carbon isotope fractionation by the prymnesiophyte algae. This finding contrasts with

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a study from the NE Pacific where nitrogen starvation seemed to affect the ε_p values for alkenones (Eek et al., 1999) and adds further support to the “trace-metal-growth-rate” hypothesis (Bidigare et al., 1997), which suggested that micronutrients control growth regardless of the concentrations of PO_4 . Overall, the relatively low ε_p values for alkenones from the oligo- and mesotrophic waters seem to indicate the use of a CCM other than diffusion. However, recent studies provided clear evidence that haptophytes have developed an inefficient but regulated CCM, with a direct uptake of HCO_3^- (Rost et al., 2003). The highest ε_p for dinoflagellates at the GYR and EGY sites are likely associated to lower growth rates and might be explained by the low N:P ratios since optimum dinoflagellate growth occurs at ratios ranging between 6 and 15 (Hodgkiss and Ho, 1997).

The carbon isotope fractionation derived from n-C₁₇ alkane reached the values of 17–19‰ in the mesotrophic waters of the HNL and MAR sites, which are within the range of 16–22‰ reported for cyanobacteria biomass (Sakata et al., 1997). In contrast, the low carbon isotope fractionations (8 to 12‰) obtained for n-C17 in the oligotrophic waters of the GYR and EGY sites as well as in the eutrophic waters of the upwelling area are suggestive of microorganisms which use CO_2 -concentrating mechanisms. Despite substantial advances over the past few years, in the understanding of the mechanism and genes involved in cyanobacterial CCMs (Badger and Price, 2003), the induction of the CCM mechanism remains to be determined (McGinn et al., 2003; Woodger et al., 2005). On the other hand, since similar ε_p values were measured in the upwelling area for biomarkers derived from eukaryotic plankton, it cannot be ruled out that n-C₁₇ in the upwelling sites is mainly derived from some eukaryotic source.

The b-values and growth rates for alkenone-producing haptophytes varied almost tenfold, ranging from 65 to 500‰ $\mu\text{mol kg}^{-1}$ and from 0.21 to 1.7 d^{-1} , respectively. The highest growth rates were estimated in the waters of the Chilean upwelling, followed by the EGY (0.47 d^{-1}), the GYR (0.35 d^{-1}) and finally the MAR and HNL (0.27 – 0.29 d^{-1}) sites. The lowest values were found in the surface waters of the Gyre (0.21 d^{-1}). Overall, our b-values and growth rates compare to those reported for the Bering Sea, Ara-

bian Sea, Southern Ocean and equatorial Pacific at 140°W (84–136 µmol kg⁻¹ and 0.2–0.4 d⁻¹) (Bidigare et al., 1997; Harada et al., 2003; Laws et al., 2001) but they are slightly higher than those reported from the Peru upwelling zone (197–397 µmol kg⁻¹ and 0.5–1 d⁻¹) (Bidigare et al., 1997). This is probably related to the strength of the upwelling as indicated by the higher nutrient and CO₂ concentrations of our samples.

The b-values for the alkenone synthesizer phytoplankton correlated with concentrations of silicate, nitrate and phosphate concentrations. However, these correlations might not indicate that the nutrients themselves are the growth limiting factor, since a natural correlation between concentrations of dissolved CO₂ and nutrients occurred. Instead, specific growth rate may be controlled by some trace micronutrient that covaries with CO₂, silicate, nitrates and phosphate concentrations (Bidigare et al., 1997; Shaked et al., 2006).

4 Summary and conclusions

Our results along the different trophic systems, showed that source-specific algal biomarkers and compound specific isotope analyses largely responded to the composition of the phytoplankton and to the different processes of carbon acquisition. It has become clear that natural variability in carbon isotope fractionation among algal taxa, is a consequence of multiple factors. Although further studies are needed to clearly define the relationship between ε_p and the specific-growth rates of the different phytoplankton taxa, our field study illustrates the large shift in isotopic fractionation by the different groups of phytoplankton taxa which should reflect the high variability in growth rates and carbon assimilation mechanisms. As a summary, we identified the upwelling zone as the one with highest algal biomass, most developed carbon concentration mechanisms and highest growth rates. In contrast, the oligotrophic area of the Gyre was characterized by the lowest concentration levels of phyto-, zoo- and bacterial markers, extremely deep concentration maxima, relatively high contribution of haptophytes and high carbon isotope fractionation factors indicative of low growth rates and

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

higher diffusive CO₂ uptake.

Acknowledgements. We thank J. Orr and C. Goyet for helpful science discussions. D. Tailliez and C. Bournot are warmly thanked for their efficient help in CTD rosette management and data processing. This is a contribution of the BIOSOPE project of the LEFE-CYBER program.

5 This research was funded by the Centre National de la Recherche Scientifique (CNRS), the Institut des Sciences de l'Univers (INSU), the Centre National d'Etudes Spatiales (CNES), the European Space Agency (ESA), The National Aeronautics and Space Administration (NASA) and the Natural Sciences and Engineering Research Council of Canada (NSERC). The IAEA
10 is grateful for the support provided to its Marine Environment Laboratories by the Government of the Principality of Monaco.

References

- Azouzi, L., Gonçalves Ito, R., Touratier, F., and Goyet, C.: Anthropogenic carbon in the eastern South Pacific Ocean, *Biogeosciences Discuss.*, 4, 1815–1837 2007.
- Badger, M. R. and Price, G. D.: CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution, *J. Exp. Bot.*, 54, 609–622, 2003.
- Baker, E. W. and Louda, J. W.: Thermal aspects in chlorophyll geochemistry, in: *Adv. Org. Geochem.*, 1981, edited by: M. Bjørøy, Wiley, Chichester, 401–421, 1983
- Barlow, R. G., Mantoura, R. F. C., Gough, M. A., and Fileman, T. W.: Pigment signatures of the phytoplankton composition in the northeastern Atlantic during the 1990 spring bloom, *Deep Sea-Res. Pt. II*, 40, 459–477, 1993.
- Beaufort, L., Couapel, M., Buchet, N., and Claustre, H.: Calcite production by Coccolithophores in the South East Pacific Ocean: from desert to jungle, *Biogeosciences Discuss.*, 4, 3267–3299, 2007,
<http://www.biogeosciences-discuss.net/4/3267/2007/>.
- Benthien, A., Zondervan, I., Engel, A., Hefer, J., Terbruggen, A., and Riebesell, U.: Carbon isotopic fractionation during a mesocosm bloom experiment dominated by *Emiliania huxleyi*: Effects of CO₂ concentration and primary production, *Geochim. Cosmochim. Ac.*, 71, 1528–1541, 2007.
- Bidigare, R. R., Fluegge, A., Freeman, K. H., Hanson, K. L., Hayes, J. M., Hollander, D., Jasper, J. P., King, L. L., Laws, E. A., Milder, J., Millero, F. J., Pancost, R. D., Popp, B. N., Steinberg,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- P. A., and Wakeham, S. G.: Consistent fractionation of C in nature and in the laboratory: growth-rate effects in some haptophyte algae, *Global Biogeochem. Cy.*, 11, 279–292, 1997.
- Bidigare, R. R., Hanson, K. L., Buesseler, K. O., Wakeham, S. G., Freeman, K. H., Pancost, R. D., Millero, F. J., Steinberg, P. A., Popp, B. N., Latasa, M., Landry, M. R., and Laws, E. A.: 5 Iron-stimulated changes in ^{13}C fractionation and export by equatorial Pacific phytoplankton: Toward a paleogrowth rate proxy, *Paleoceanography*, 14, 589–595, 1999.
- Burkhardt, S., Riebesell, U., and Zondervan, I.: Effects of growth rate, CO_2 concentration, and cell size on the stable carbon isotope fractionation in marine phytoplankton, *Geochim. Cosmochim. Ac.*, 63, 3729–3741, 1999a.
- Burkhardt, S., Riebesell, U., and Zondervan, I.: Stable carbon isotope fractionation by marine phytoplankton in response to daylength, growth rate, and CO_2 availability, *Mar. Ecol.-Progr. Ser.*, 184, 31–41, 1999b.
- Claustre, H., Hooker, S. B., Van Heukelem, L., Berthon, J.-F., Barlow, R., Ras, J., Sessions, H., Targa, C., Thomas, C. S., van der Linde, D., and Marty, J.-C.: An intercomparison of HPLC 15 phytoplankton pigment methods using in situ samples: application to remote sensing and database activities, *Mar. Chem.*, 85, 41–61, 2004.
- Conte, M. H., Thompson, A., Eglinton, G., and Green, J. C.: Lipid biomarker diversity in the coccolithophorid *Emiliania huxleyi* (*Prymnophyceae*) and the related species *Gephyrocapsa oceanica*, *J. Phycol.*, 31, 272–282, 1995.
- Conte, M. H., Thompson, A., Lesley, D., and Harris, R. P.: Genetic and Physiological Influences 20 on the Alkenone/Alkenoate Versus Growth Temperature Relationship in *Emiliania huxleyi* and *Gephyrocapsa Oceanica*, *Geochim. Cosmochim. Ac.*, 62, 51–68, 1998.
- Conte, M. H., Weber, J. C., King, L. L., and Wakeham, S. G.: The alkenone temperature signal 25 in western North Atlantic surface waters, *Geochim. Cosmochim. Ac.*, 65, 4275–4287, 2001.
- Craig, H.: Abyssal carbon 13 in the South Pacific, *J. Geophys. Res.* 75, 691–695, 1970.
- Dandonneau, Y., Montel, Y., Blanchot, J., Giraudeau, J., and Neveux, J.: Temporal variability 30 in phytoplankton pigments, picoplankton and coccolithophores along a transect through the North Atlantic and tropical southwestern Pacific, *Deep Sea- Res. Pt. I*, 53, 689–712, 2006.
- Dijkman, N. A. and Kromkamp, J. C.: Phospholipid-derived fatty acids as chemotaxonomic markers for phytoplankton: application for inferring phytoplankton composition, *Mar. Ecol.-Progr. Ser.*, 324, 113–125, 2006.
- DOE: Handbook of Methods for Analysis of the Various Parameters of the Carbon Dioxide System in Seawater; version 2, edited by: Dickson, A. G. and Goyet, C., ORNL/CDIAC-74,

1994.

- Eek, M. K., Whiticar, M. J., Bishop, J. K. B., and Wong, C. S.: Influence of nutrients on carbon isotope fractionation by natural populations of Prymnesiophyte algae in NE Pacific, Deep Sea-Res. Pt II, 46, 2863–2876, 1999.
- 5 Eppley, R. W., Rogers, J. N., and McCarthy, J. J.: Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton, Limnol. Oceanogr., 14, 912–920, 1969.
- Epstein, B. L., D'Hondt, S., Quinn, J. G., Zhang, J., and Hargraves, P. E.: An effect of dissolved nutrient concentrations on alkenone-based temperature estimates, Paleoceanography, 13, 122–126, 1998.
- 10 Falkowski, P. G.: Physiological responses of phytoplankton to natural light regimes, J. Plankton Res., 6, 295–307, 1984.
- Falkowski, P. G., Barber, R. T., and Smetacek, V.: Biogeochemical controls and feedbacks on ocean primary production, Science, 281, 200–206, 1998.
- Freeman, K. H. and Hayes, J. M.: Fractionation of carbon isotopes by phytoplankton and estimates of ancient CO₂ levels, Global Biogeochem. Cy., 6, 185–198, 1992.
- 15 Freeman, K. H., Hayes, J. M., Trendel, J. M., and Albrecht, P.: Evidence from carbon isotope measurements for diverse origins of sedimentary hydrocarbons, Nature, 343, 254–256, 1990.
- Goericke, R., Montoya, J. P., and Fry, B.: Physiology of isotopic fractionation in algae and cyanobacteria. In: Stable Isotopes in Ecology and Environmental Science, Blackwell Science Publishers, 187–221, 1994.
- 20 Gómez, F., Claustre, H., Raimbault, P., and Souissi, S.: Two High-Nutrient Low-Chlorophyll phytoplankton assemblages: the tropical central Pacific and the offshore Perú-Chile Current, Biogeosciences Discuss., 4, 1535–1554, 2007,
<http://www.biogeosciences-discuss.net/4/1535/2007/>.
- 25 Goyet, C. and Poisson, A.: New determination of carbonic acid dissociation constants in seawater as a function of temperature and salinity, Deep Sea-Res., 36, 1635–1654, 1989.
- Grob, C., Ulloa, O., Claustre, H., Huot, Y., Alarcón, G., and Marie, D.: Contribution of picoplankton to the total particulate organic carbon concentration in the eastern South Pacific, Biogeosciences, 4, 837–852, 2007,
<http://www.biogeosciences.net/4/837/2007/>.
- 30 Han, J. and Calvin, M.: Hydrocarbon Distribution of Algae and Bacteria, and Microbiological Activity in Sediments, P. Natl. Acad. Sci. USA, 64, 436–443, 1969.

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

- Harada, N., Shin, K. H., Murata, A., Uchida, M., and Nakatani, T.: Characteristics of alkenones synthesized by a bloom of *Emiliania Huxleyi* in the Bering Sea, *Geochim. Cosmochim. Ac.*, 67, 1507–1519, 2003.
- Hayes, J. M.: Factors controlling ^{13}C contents of sedimentary organic compounds: Principles and evidence, *Mar. Geol.*, 113, 111–125, 1993.
- Hayes, J. M., Freeman, K. H., Popp, B. N., and Hoham, C. H.: Compound-specific isotopic analyses: A novel tool for reconstruction of ancient biogeochemical processes, *Org. Geochem.*, 16, 1115–1128, 1990.
- Hodgkiss, I. J. and Ho, K. C.: Are changes in N:P ratios in coastal waters the key to increased red tide blooms?, *Hydrobiologia*, 352, 141–147, 1997.
- Irwin, A. J., Finkel, Z. V., Schofield, O. M. E., and Falkowski, P. G.: Scaling-up from nutrient physiology to the size-structure of phytoplankton communities, *J. Plankton Res.*, 28, 459–471, 2006.
- Jones, D. M., Carter, J. F., Eglinton, G., Jumeau, E. J., and Fenwick, C. S.: Determination of $\delta^{13}\text{C}$ values of sedimentary straight chain and cyclic alcohols by gas chromatography/isotope ratio mass spectrometry, *Biol. Mass Spectrom.*, 20, 641–646, 1991.
- Kaneda, T.: Iso- and anteiso-fatty acids in bacteria: biosynthesis, function, and taxonomic significance. *Microbiol. Rev.*, 55, 288–302, 1991.
- Laws, E. A., Bidigare, R. R., and Popp, B. N.: Effect of growth rate and CO_2 concentration on carbon isotopic fractionation by the marine diatom *Phaeodactylum tricornutum*, *Limnol. Oceanogr.*, 42, 1552–1560, 1997.
- Laws, E. A., Popp, B. N., Bidigare, R. R., Kennicutt, M. C., and Macko, S. A.: Dependence of phytoplankton carbon isotopic composition on growth rate and $[\text{CO}_2]_{\text{aq}}$: Theoretical considerations and experimental results, *Geochim. Cosmochim. Ac.*, 59, 1131–1138, 1995.
- Laws, E. A., Popp, B. N., Bidigare, R. R., Riebesell, U., Burkhardt, S., and Wakeham, S. G.: Controls on the molecular distribution and carbon isotopic composition of alkenones in certain haptophyte algae, *Geochem. Geophys. Geosy.*, 2, 2000GC000057, 2001.
- Lee, R. F., Hagen, W., and Kattner, G.: Lipid storage in marine zooplankton, *Mar. Ecol.-Progr. Ser.*, 307, 273–306, 2006.
- Lewis, E., and Wallace, D. W. R.: Program Developed for CO_2 System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, 1998.
- Litchman, E., Klausmeier, C. A., Miller, J. R., Schofield, O. M., and Falkowski, P. G.: Multi-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nutrient, multi-group model of present and future oceanic phytoplankton communities, *Bio-geosciences*, 3, 585–606, 2006,
<http://www.biogeosciences.net/3/585/2006/>.

- Mackey, M. D., Mackey, D. J., Higgins, H. W., and Wright, S. W.: CHEMTAX - a program for
5 estimating class abundances from chemical markers: application to HPLC measurements of
phytoplankton, *Mar. Ecol.-Progr. Ser.*, 144, 265–283, 1996.
- McGinn, P. J., Price, G. D., Maleszka, R., and Badger, M. R.: Inorganic Carbon Limitation and
Light Control the Expression of Transcripts Related to the CO₂-Concentrating Mechanism
in the Cyanobacterium *Synechocystis* sp. Strain PCC6803, *Plant Physiol.*, 132, 218–229,
10 2003.
- Mook, W. G., Bommerson, J. C., and Staverman, W. H.: Carbon isotope fractionation between
dissolved bicarbonate and gaseous carbon dioxide, *Earth Planet. Sc. Lett.*, 22, 169–176,
1974.
- Morel, F. M. M., Reinfelder, J. R., Roberts, S. B., Chamberlain, C. P., Lee, J. G., and Yee, D.:
15 Zinc and carbon co-limitation of marine phytoplankton, *Nature*, 369, 740–742, 1994.
- Muhlebach, A. and Weber, K.: Origins and fate of dissolved sterols in the Weddell Sea, Antarctica,
Org. Geochem., 29, 1595–1607, 1998.
- Pancost, R. D., Freeman, K. H., and Wakeham, S. G.: Controls on the carbon-isotope compositions
of compounds in Peru surface waters, *Org. Geochem.*, 30, 319–340, 1999.
- Pancost, R. D., Freeman, K. H., Wakeham, S. G., and Robertson, C. Y.: Controls on carbon
20 isotope fractionation by diatoms in the Peru upwelling region, *Geochim. Cosmochim. Ac.*,
61, 4983–4991, 1997.
- Parrish, C. C., Wells, J. S., Yang, Z., and Dabinett, P.: Growth and lipid composition of scallop
juveniles, *Placopecten magellanicus*, fed the flagellate *Isochrysis galbana* with varying lipid
25 composition and the diatom *Chaetoceros muelleri*, *Mar. Biol.*, 133, 461–471, 1998.
- Pinturier-Geiss, L., Mejanelle, L., Dale, B., and Karlsen, D. A.: Lipids as indicators of eutrophication
in marine coastal sediments, *J. Microbiol. Meth.*, 48, 239–257, 2002.
- Popp, B. N., Kenig, F., Wakeham, S. G., Laws, E. A., and Bidigare, R. R.: Does growth rate
affect ketone unsaturation and intracellular carbon isotope variability in *Emiliania huxleyi*?
30 *Paleoceanography*, 13, 35–41, 1998a.
- Popp, B. N., Laws, E. A., Bidigare, R. R., Dore, J. E., Hanson, K. L., and Wakeham, S. G.: Effect
of phytoplankton cell geometry on carbon isotopic fractionation, *Geochim. Cosmochim. Ac.*,
62, 69–77, 1998b.

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid
biomarkers from the
South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Popp, B. N., Prahl, F. G., Wallsgrove, R. J., and Tanimoto, J.: Seasonal patterns of alkenone production in the subtropical oligotrophic North Pacific, *Paleoceanography*, 21, PA1004, doi:10.1029/2005PA001165, 2006.

5 Prahl, F. G., Muehlhausen, L. A., and Zahnle, D. L.: Further evaluation of long-chain alkenones as indicators of paleoceanographic conditions, *Geochim. Cosmochim. Ac.*, 52, 2303–2310, 1988.

Prahl, F. G., Popp, B. N., Karl, D. M., and Sparrow, M. A.: Ecology and biogeochemistry of alkenone production at Station ALOHA, *Deep Sea-Res. Pt I*, 52, 699–719, 2005.

10 Prahl, F. G. and Wakeham, S. G.: Calibration of unsaturation patterns in long-chain ketone compositions for palaeotemperature assessment, *Nature*, 330, 367–369, 1987.

Raimbault, P., Garcia, N. and Cerutti, F.: Distribution of inorganic and organic nutrients in the South Pacific Ocean; -evidence for long-term accumulation of organic matter in nitrogen-depleted waters, *Biogeosciences Discuss.*, 4, 3041–3087, 2007,
<http://www.biogeosciences-discuss.net/4/3041/2007/>.

15 Rau, G. H., Chavez, F. P., and Friederich, G. E.: Plankton $^{13}\text{C}/^{12}\text{C}$ variations in Monterey Bay, California: evidence of non-diffusive inorganic carbon uptake by phytoplankton in an upwelling environment, *Deep Sea-Res. Pt. I*, 48, 79–94, 2001.

Ras, J., Claustre, H. and Uitz, J.: Spatial variability of phytoplankton pigment distributions in the Subtropical South Pacific Ocean: comparison between in situ and predicted data, *Biogeosciences Discuss.*, 4 3409–3451, 2007.

20 Rau, G. H., Riebesell, U., and Wolf-Gladrow, D.: A model of photosynthetic C fractionation by marine phytoplankton based on diffusive molecular CO_2 uptake, *Mar. Ecol.-Progr. Ser.*, 133, 275–285, 1996.

25 Riebesell, U., Revill, A. T., Holdsworth, D. G., and Volkman, J. K.: The effects of varying CO_2 concentration on lipid composition and carbon isotope fractionation in *Emiliania huxleyi*, *Geochim. Cosmochim. Ac.*, 64, 4179–4192, 2000.

Robinson, N., Eglington, G., Brassell, S. C., and Cranwell, P. A.: Dinoflagellate origin for sedimentary 4α methylsteroids and $5\alpha(\text{H})$ -stanols, *Nature*, 308, 419–422, 1984.

30 Rontani, J.-F., Marty, J.-C., Miquel, J.-C., and Volkman, J. K.: Free radical oxidation (autoxidation) of alkenones and other microalgal lipids in seawater, *Org. Geochem.*, 37, 354–368, 2006.

Rost, B., Riebesell, U., Burkhardt, S., and Sultemeyer, D.: Carbon acquisition of bloom-forming marine phytoplankton, *Limnol. Oceanogr.*, 48, 55–67, 2003.

BGD

4, 4653–4696, 2007

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

- Sakata, S., Hayes, J. M., McTaggart, A. R., Evans, R. A., Leckrone, K. J., and Togasaki, R. K.: Carbon isotopic fractionation associated with lipid biosynthesis by a cyanobacterium: Relevance for interpretation of biomarker records, *Geochim. Cosmochim. Ac.*, 61, 5379–5389, 1997.
- 5 Sargent, J. R., Gatten, R. R., and McIntosh, R.: Wax esters in the marine environment - Their occurrence, formation, transformation and ultimate fates, *Mar. Chem.*, 5, 573–584, 1977.
- Shaked, Y., Xu, Y., Leblanc, K., and Morel, F. M. M.: Zinc availability and alkaline phosphatase activity in *Emiliania huxleyi*: Implications for Zn-P co-limitation in the ocean, *Limnol. Oceanogr.*, 51, 299–309, 2006.
- 10 Shin, K. H., Hama, T., Yoshie, N., Noriki, S., and Tsunogai, S.: Dynamics of fatty acids in newly biosynthesized phytoplankton cells and seston during a spring bloom off the west coast of Hokkaido Island, Japan, *Mar. Chem.*, 70, 243–256, 2000.
- Sicre, M.-A., Bard, E., Ezat, U., and Rostek, F.: Alkenone distributions in the North Atlantic and Nordic sea surface waters, *Geochem. Geophys. Geosy.*, 3(2), 2001GC000159, 2002.
- 15 Tolosa, I. and de Mora, S.: Isolation of neutral and acidic lipid biomarker classes for compound-specific-carbon isotope analysis by means of solvent extraction and normal-phase high-performance liquid chromatography, *J. Chromatogr. A*, 1045, 71–84, 2004.
- Tolosa, I., LeBlond, N., Copin-Montegut, C., Marty, J.-C., de Mora, S., and Prieur, L.: Distribution of sterol and fatty alcohol biomarkers in particulate matter from the frontal structure of the Alboran Sea (S.W. Mediterranean Sea), *Mar. Chem.*, 82, 161–183, 2003.
- 20 Tolosa, I., Vescovali, I., LeBlond, N., Marty, J.-C., de Mora, S., and Prieur, L.: Distribution of pigment and fatty acid biomarkers in particulae matter from the frontal structure of the Alboran Sea (S.W. Mediterranean Sea), *Mar. Chem.*, 88, 103–125, 2004.
- van der Meer, M. T. J., Schouten, S., van Dongen, B. E., Rijpstra, W. I. C., Fuchs, G., Damste, J. S. S., de Leeuw, J. W., and Ward, D. M.: Biosynthetic controls on the ^{13}C contents of organic components in the photoautotrophic bacterium *Chloroflexus aurantiacus*, *J. Biol. Chem.*, 276, 10 971–10 976, 2001.
- 25 Van Wambeke, F., Obernosterer, I., Moutin, T., Duhamel, S., Ulloa, O., and Claustre, H.: Heterotrophic bacterial production in the South East Pacific: longitudinal trends and coupling with primary production, *Biogeosciences Discuss.*, 4, 2761–2791, 2007,
<http://www.biogeosciences-discuss.net/4/2761/2007/>.
- 30 Volkman, J. K.: A review of sterol markers for marine and terrigenous organic matter, *Org. Geochem.*, 9, 83–99, 1986.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Volkman, J. K., Barrerr, S. M., Blackburn, S. I., and Sikes, E. L.: Alkenones in *Gephyrocapsa oceanica*: Implications for studies of paleoclimate, Geochim. Cosmochim. Ac., 59, 513–520, 1995.

Volkman, J. K., Barrett, S. M., and Dunstan, G. A.: C₂₅ and C₃₀ highly branched isoprenoid alkenes in laboratory cultures of two marine diatoms, Org. Geochem., 21, 407–414, 1994.

Volkman, J. K. and Hallegraeff, G. M.: Lipids in marine diatoms of the genus *Thalassiosira*: Predominance of 24-methylenecholesterol, Phytochemistry, 27, 1389–1394, 1988.

Volkman, J. K., Jeffrey, S. W., Nichols, P. D., Rogers, G. I., and Garland, C. D.: Fatty acid and lipid composition of 10 species of microalgae used in mariculture, J. Exp. Mar. Biol. Ecol., 128, 219–240, 1989.

Werne, J. P. and Hollander, D. J.: Balancing supply and demand: controls on carbon isotope fractionation in the Cariaco Basin (Venezuela) Younger Dryas to present, Mar. Chem., 92, 275–293, 2004.

Winters, K., Parker, P. L., and Van Baalen, C.: Hydrocarbons of Blue-Green Algae: Geochemical significance, Science, 163, 467–468, 1969.

Woodger, F. J., Badger, M. R., and Price, G. D.: Sensing of inorganic carbon limitation in *Synechococcus* PCC7942 is correlated with the size of the internal inorganic carbon pool and involves oxygen, Plant Physiol., 139, 1959–1969, 2005.

Woodworth, M., Goni, M., Tappa, E., Tedesco, K., Thunell, R., Astor, Y., Varela, R., Rafael Diaz-Ramos, J., and Muller-Karger, F.: Oceanographic controls on the carbon isotopic compositions of sinking particles from the Cariaco Basin, Deep Sea-Res. Pt. I, 51, 1955–1974, 2004.

Yamamoto, M., Shiraiwa, Y., and Inouye, I.: Physiological responses of lipids in *Emiliania huxleyi* and *Gephyrocapsa oceanica* (*Haptophyceae*) to growth status and their implications for alkenone paleothermometry, Org. Geochem., 31, 799–811, 2000.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Table 1. Sampling sites.

DATE	Lat.(° S)	Long. (° W)	ACRONYM	Brief description
28/10/2004	8.4	141.3	MAR	Marquesas Islands characterized by high nutrients high chlorophyll
31/10/2004	9.0	136.8	HNL	High nutrient low chlorophyll area east of the Marquesas Islands
12/11/2004	25.6.	114.0	GYR	Center of the South Pacific Gyre
28/11/2004	31.8	91.4	EGY	Eastern border of the Gyre
6/12/2004	34.0	73.3	UPW	Upwelling area situated above the abyssal plain
10/12/2004	34.5	72.4	UPX	Upwelling area situated above the continental shelf

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Selected environmental parameters from the six sites at the sampled depths.

locations-depth (m)	T °C	density Kg m ⁻³	A _T μmol Kg ⁻¹	C _T μmol Kg ⁻¹	[CO ₂ (aq) μmol Kg ⁻¹]	NO3 μM	PO4 μM	SiOH ₄ μM	POC μM	Chl a μg l ⁻¹	t ³ hours
mar3-50 m	27.7	22.9	2363	2024	11.3	1.59	0.28	1.03	1.96	0.41	12.4
mar3-100 m	26.9	23.2	2356	2052	13.1	3.81	0.39	1.73	0.90	0.21	
mar3-300 m	11.3	26.6	2313	2225	34.7	8.83	0.61	2.66	0.28	0.05	
hn12-75 m	27.3	23.2	2353	2009	11.0	1.69	0.37	1.18	1.60	0.30	12.4
hn11-100 m	26.8	24.5	2382	2097	14.3	1.04	0.50	1.88	0.90	0.26	
hn12-300 m	11.2	26.6	2319	2238	37.2	31.6	2.39	14.45	0.23	0.05	
gyr2-0 m	22.0	24.6	2364	2048	11.9	0.00	0.12	0.88	0.39	0.03	13.3
gyr2-75 m	21.9	25.1	2369	2055	11.8	0.00	0.14	1.04	0.51	0.06	
gyr2-125 m	20.7	25.2	2363	2051	11.7	0.00	0.12	0.96	0.49	0.13	
gyr2-150 m	20.3	25.3	2358	2057	12.1	0.00	0.12	0.71	0.48	0.18	
gyr2-175 m	19.6	25.4	2347	2074	13.3	0.10	0.14	0.71	0.49	0.20	
gyr2-200 m	18.7	25.5	2333	2075	14.1	1.11	0.19	0.79	0.36	0.17	
gyr2-300 m	13.9	26.0	2299	2099	17.2	8.06	0.81	1.93	0.12	0.04	
egy4-70 m	16.5	25.4	2294	2033	12.9	0.53	0.21	1.34	0.95	0.19	13.9
egy2-200 m	14.5	25.8	2271	2071	16.6	2.96	0.41	1.5	0.25	0.04	
egy4-300 m	10.0	26.4	2267	2119	21.8	15.4	1.07	3.99	0.23	0.03	
upw1-40 m	12.8	25.9			37.4	13.0	0.91	8	6.92	2.5	14.3
upw1-100 m	10.8	26.4			52.4	28.1	2.6	20.8	1.83	0.07	
upw2-300 m	8.7	26.8			56.0	39.0	2.8	33.0	1.03	0.08	
upx3-40 m	12.0	26.0	2274	2197	37.4	22.8	2.02	10.7	3.28	0.79	14.4
upx2-100 m	10.5	26.4	2294	2258	52.4	23.2	2.32	24.4	1.25	0.11	
upx2-300 m	10.0	26.7	2307	2281	56.0	35.3	3.5	27.9	0.56	0.07	

¹ Total alkalinity; ² Total CO₂; ³ day length

δ¹³C on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Table 3. Summary of the lipid biomarkers discussed in this study.

ACRONYM	COMPOUND NAME(S)	Main diagnostic (and minor) sources source	References
Phytol	3,7,11,15-tetramethyl-2-hexadecen-1-ol	Phototrophic organisms	(Baker and Louda, 1983)
$\text{C}_{28:4}^{\Delta 5,24(28)}$	24-methylcholesta-5,24(28)-dien-3 β -ol	Diatoms (flagellates)	(Volkman and Hallegraeff, 1988)
C_{25}HBI	Highly branched isoprenoids of C25	Diatoms (flagellates)	(Volkman et al., 1994)
$\text{C}_{16:1}$ FA	6,9,12,15-hexadecatetraenoic acid (C16:4(n-1))	Diatoms	(Dijkman and Kromkamp, 2006)
$\text{C}_{20:5}$ FA	5,8,11,14,17-eicosapentaenoic acid (C20:5(n-3))	Diatoms (flagellates)	(Dijkman and Kromkamp, 2006; Volkman et al., 1989)
Total alkenones	Long-chain (C37-C39) unsaturated ketones	Haptophytes/Prymnesiophycea	(Conte et al., 1995; Volkman et al., 1995)
$\text{C}_{30}\Delta^{22,23}$ -(dinosterol)	4 α -23,24-trimethylcholest-22(E)-en-3 β -ol	Dinoflagellates	(Robinson et al., 1984)
<i>n</i> -C ₁₇	<i>C</i> ₁₇ <i>n</i> -alkane	Cyanobacteria (green algae)	(Han and Calvin, 1969; Winters et al., 1969)
<i>n</i> -alcohols	<i>n</i> -alkanols, mainly <i>n</i> -C14, <i>n</i> -C16 and <i>n</i> -C18	Zooplankton and marine invertebrates (algae)	(Sargent et al., 1977).
$\text{C}_{20:1} + \text{C}_{22:1}$ FA	Long-chain monounsaturated $\text{C}_{20:1}$ and $\text{C}_{22:1}$ FA	Herbivorous mesozooplankton	(Lee et al., 2006)
Branched FA	<i>iso</i> and <i>anteiso</i> branched fatty acids in the carbon number range 15-19.	Heterotrophic bacteria	(Kaneda, 1991)
$\text{C}_{27}\Delta^5$ -(cholesterol)	Cholest-5-en-3 β -ol	Zooplankton (algae)	(Volkman, 1986)
Phytosterols	Includes : 27-nor-24-methylcholesta-5,22(E)-dien-3 β -ol; cholesta-5,22(E)-dien-3 β -ol; 24-methylcholesta-5,22(E)-dien-3 β -ol; $\text{C}_{29}^{\Delta 5,24(28)}$; 24-ethylcholesta-5,22(E)-dien-3 β -ol; 24-ethylcholestan-5-en-3 β -ol and $\text{C}_{30}\Delta^{22}$	Eukaryotic phototrophic organisms	(Muhlebach and Weber, 1998; Tolosa et al., 2003)




Table 4. Selected lipid biomarkers concentrations (ng l^{-1}) in suspended particles from the South Pacific Ocean.

locations-depth (m)	phytol	$C_{28}\Delta^{5,24(28)}$	C_{25}	HBI	$C_{16:4}$	$C_{20:5}$	Total alkenones	$C_{30}\Delta^{22}$	$n\text{-}C_{17}$	$n\text{-}alcohols$	$C_{20:1}+C_{22:1}$	Branched FA	$C_{27}\Delta^5$
mar3-50 m	30.7	20.4	2.32	32.9	97.9	6.5	4.4	0.13	6.7	2.81	21.6	13.0	
mar3-100 m	7.0	4.7	0.02	17.6	36.1	3.7	0.0	0.04	34.5	6.66	31.3	6.7	
mar3-300 m	0.6	3.0	0.00	0.0	9.7	0.0	0.0	0.00	9.5	9.27	2.1	1.1	
hn12-75 m	25.4	29.7	0.80	28.7	61.8	16.0	5.7	0.25	7.3	2.28	19.2	13.0	
hn11-100 m	20.5	15.1	0.07	22.3	54.2	8.8	4.0	2.26	16.5	1.70	3.4	23.5	
hn12-300 m	0.6	1.1	0.00	0.0	5.5	0.0	0.4	0.00	10.0	0.47	1.2	3.8	
gyr2-0 m	1.7	1.9	0.21	1.4	8.2	14.3	1.6	0.00	11.6	0.00	4.6	3.7	
gyr2-75 m	2.3	1.9	0.02	1.4	7.2	18.1	1.7	0.13	4.0	0.52	2.5	2.7	
gyr2-125 m	5.6	3.2	0.03	2.4	16.5	23.4	2.0	0.26	4.2	0.79	3.2	4.5	
gyr2-150 m	9.6	3.8	0.15	5.9	25.4	16.0	1.1	0.29	7.5	0.46	4.6	5.7	
gyr2-175 m	9.9	4.0	0.18	3.8	16.1	13.1	0.8	0.32	7.4	0.33	5.2	3.5	
gyr2-200 m	11.5	6.5	0.41	3.9	20.6	6.9	1.5	0.19	9.8	0.87	5.3	7.0	
gyr2-300 m	0.6	0.3	0.00	0.0	2.2	0.0	0.1	0.00	1.6	0.11	0.7	0.9	
egy4-70 m	16.4	20.1	1.40	17.2	57.1	20.0	2.8	1.15	11.8	1.69	10.1	15.3	
egy2-200 m	2.0	2.8	0.00	0.0	13.0	1.5	0.6	0.67	9.6	0.27	4.7	7.3	
egy4-300 m	0.6	1.4	0.00	0.0	9.4	0.5	0.6	0.00	7.0	0.00	3.1	5.6	
upw1-40 m	102.4	55.4	3.69	139.4	379.1	27.1	9.7	0.23	33.6	15.76	113.7	36.4	
upw1-100 m	23.6	15.2	0.09	15.1	105.7	7.0	3.9	0.28	33.6	2.48	29.6	20.0	
upw2-300 m	2.8	2.5	0.00	0.0	24.2	2.0	0.8	0.08	40.8	1.05	10.1	13.0	
upx3-40 m	78.6	33.0	2.29	25.7	142.8	12.0	6.1	0.18	89.8	7.82	57.0	55.4	
upx2-100 m	26.1	14.8	0.75	8.0	101.0	2.3	3.8	0.16	10.3	1.39	30.0	15.9	
upx2-300 m	5.7	5.2	1.73	16.2	71.7	1.1	1.0	0.00	215.2	2.60	9.4	23.4	

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Carbon isotope fractionation of CO_2 aq with respect to bicarbonate (ε_b), $\delta^{13}\text{CO}_2$ (dissolved in the water column), different carbon isotope fractionation associated with photosynthetic carbon fixation using molecular specific lipid biomarkers (ε_p of biomarkers) and b-value (‰ $\mu\text{mol kg}^{-1}$) and specific growth rate for alkenone synthesizers.

locations-depth (m)	ε_b (‰)	$\delta^{13}\text{CO}_2$ (‰)	ε_p (‰) phytol	ε_p (‰) $\text{C}_{28}\Delta^{5,24(28)}$	ε_p (‰) alkenone	ε_p (‰) $\text{C}_{30}\Delta^{22}$	ε_p (‰) <i>n</i> -C17	b (‰ $\mu\text{mol kg}^{-1}$) alkenone	μ (d ⁻¹) alkenone
mar3-50 m	-8.66	-6.48	18.6	22.4	16.4	16.3	17.2	97.7	0.29
hnl2-75 m	-8.73	-6.55	17.6	24.0	16.7	15.2	19.1	90.9	0.27
hnl1-100 m	-8.79	-6.61	19.8	23.9		16.4			
gyr2-0 m	-9.32	-7.14			19.4			66.8	0.21
gyr2-75 m	-9.33	-7.15			19.4	18.9		65.9	0.21
gyr2-125 m	-9.47	-7.29	18.8	19.3	18.6	18.4		75.3	0.24
gyr2-150 m	-9.52	-7.34		22.8	16.0	18.2	10.2	109.1	0.35
gyr2-175 m	-9.59	-7.41	18.6	26.2	16.4			114.6	0.37
gyr2-200 m	-9.70	-7.52	19.4	24.1	16.5	20.7		119.7	0.38
egy4-70 m	-9.96	-7.78	20.7	26.1	13.5	18.7	8.3	148.5	0.48
egy2-200 m	-10.20	-8.02		22.2	12.9	17.6		201.3	0.58
egy4-300 m	-10.74	-8.57		20.4					
upw1-40 m	-10.38	-8.20	11.1	9.8	11.6	12.6	10.0	501.0	1.73
upw1-100 m	-10.62	-8.44	10.7	7.9	11.4	10.2		712.6	1.72
upw2-300 m	-10.87	-8.69		10.7					
upx3-40 m	-10.50	-8.32	19.5	15.0	12.6	13.4	11.9	461.7	1.61
upx2-100 m	-10.68	-8.50	18.9		12.9	10.7		631.9	1.53
upx2-300 m	-10.74	-8.57				10.7			

Appendix

Table A1. Stable carbon isotopic composition ($\delta^{13}\text{C}(\text{\textperthousand}) \pm$ s.d. of three replicate injections) of selected lipid biomarkers in suspended particles from the South East Pacific Ocean.

locations-depth (m)	phytol	$\text{C}_{28}\Delta^{5,24(28)}$	$\text{C}_{37:2}$ alkenone	$\text{C}_{30}\Delta^{22}$	$n\text{-C}_{17}$	$\text{C}_{20:5}$ FA	$i\text{-C}_{15}^{\text{a}}$ FA
mar3-50 m	-28.8±0.9	-32.4±0.5	-26.7±0.5	-26.6±0.5	-31.7±0.5	-24.0±0.5	-20.7±0.7
hnl2-75 m	-27.9±0.5	-34.0±0.5	-27.1±0.5	-25.7±0.5	-33.6±0.5	-24.8±0.7	-21.6±0.5
hnl1-100 m	-30.1±0.5	-34.0±0.5		-26.9±0.6			
gyr2-0 m			-30.2±0.5			-29.3±0.8	-23.3±0.5
gyr2-75 m			-30.3±0.5	-29.7±0.5		-26.5±0.5	-21.6±0.7
gyr2-125 m	-29.9	-30.3	-29.6±0.5	-29.4±0.5		-26.9±0.5	-20.2±0.5
gyr2-150 m		-33.6	-27.2±0.5	-29.3	-25.8±0.5	-26.8±0.5	-23.4±0.6
gyr2-175 m	-29.8	-37.0	-27.6±0.5			-27.6±0.5	-23.5±0.5
gyr2-200 m	-30.6±1.2	-35.1±1.5	-27.8±0.5	-31.8±0.6		-29.5±0.5	-24.8±0.5
egy4-70 m	-32.1±0.5	-37.3±0.6	-25.2±0.6	-30.2±1.2	-24.4±0.5	-28.8±0.8	-25.5±0.5
egy2-200 m		-3.7±0.5	-24.8±0.6	-29.4±0.8		-28.6±1.6	-25.7±0.7
egy4-300 m		-32.6±0.5				-25.8±1.3	-24.2±0.6
upw1-40 m	-23.3±0.5	-22.0±0.5	-23.8±0.5	-24.7±0.8	-26.5±0.5	-24.7±0.6	-19.8±0.5
upw1-100 m	-23.1	-20.5±0.5	-23.8±0.5	-22.7±0.5		-25.0±0.5	-19.7±0.6
upw2-300 m		-23.4±0.7				-26.1±0.5	-22.1±0.5
upx3-40 m	-31.5±0.5	-27.1±0.9	-24.9±0.5	-25.6±0.5	-28.4±0.5	-28.6±0.5	-21.7±0.5
upx2-100 m	-31.1±2.0	-22.5±1.4	-25.4±0.5	-23.2±0.8		-24.2±0.5	-19.3±0.5
upx2-300 m	-24.1±0.7	-21.3±0.6		-23.3		-24.0±0.5	-20.4±0.5

^a 13-methyl-tetradecanoic acid (*iso*- C_{15})

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 1. Percentage contribution of selected lipid biomarkers and biochemical indices: **(A)**: percentage of phytol relative to total neutral lipid concentrations, **(B)**: percentage of 24-methylcholesta-5,24(28)-dien-3 β -ol relative to total neutral lipids concentrations; **(C)**: percentage of total alkenones relative to total neutral lipid concentrations; **(D)**: Ratio of 24-methylcholesta-5,24(28)-dien-3 β -ol to total alkenones to evaluate the relative contribution of diatoms vs. haptophytes.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 2. Values of selected biochemical indices to elucidate the dominant sources in the suspended particles from the different sites. **(A):** Cholesterol/phytosterols¹ ratio to evaluate the relative contribution of zooplankton vs. algal. **(B):** % Bacterial fatty acid indicator is the sum of all *iso*- and *anteiso*- branched chain fatty acids expressed as percent of total fatty acids. **(C):** Phytosterols/phytol ratio to elucidate the degradation state of the phytoplankton material. **(D):** % C₁₆ PUFA is the polyunsaturation index of C₁₆ fatty acids to evaluate the ecophysiological state of the marine diatoms. 1) Phytosterols are listed in Table 3.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 3. Carbon isotope ratios ($\delta^{13}\text{C}$) for selected lipid biomarkers in the upwelling zone.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 4. Depth distribution of selected lipid biomarkers in the suspended particles from the gyre.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 5. Profiles for total C₃₇ alkenones concentrations (C₃₇ alk) and 19'-hexanoyloxyfucoxanthin pigment (19'HF) with depth in the suspended particles from the gyre.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 6. Profiles for total C_{37} alkenone concentrations and carbon isotope ratio ($\delta^{13}\text{C}$) for the diunsaturated C_{37} alkenone.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 7. Profile for CTD temperature and plot for $U_{37}^{K'}$ -derived water temperature estimates (see text for details) from suspended particles in the gyre.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{13}\text{C}$ on lipid biomarkers from the South East Pacific

I. Tolosa et al.




Fig. 8. Correlation of the carbon isotope fractionation factor of the $\text{C}_{28}\Delta^{5,24(28)}$ diatom marker with measured silicate concentrations across the South Pacific Ocean.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [|◀](#) [▶|](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)




Fig. 9. Correlations of the carbon isotope fractionation factor of diatom (**A**) and alkenone markers (**B**) with CO_2 concentrations.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)