

Comparative sensitivity to methyl eugenol of four putative *Bactrocera dorsalis* complex sibling species – further evidence that they belong to one and the same species *B. dorsalis*

Alvin K.W. Hee¹, Yue-Shin Ooi¹, Suk-Ling Wee^{2,3}, Keng-Hong Tan⁴

1 Department of Biology, Faculty of Science, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia **2** School of Environmental and Natural Resource Sciences **3** Centre of Insect Systematics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia **4** Tan Hak Heng Co., Penang, Malaysia

Corresponding author: Alvin K.W. Hee (alvinhee@upm.edu.my)

Academic editor: A. Clarke | Received 26 June 2015 | Accepted 17 August 2015 | Published 26 November 2015

<http://zoobank.org/66D6227B-27DE-47A4-A349-D09E753D9F1C>

Citation: Hee AKW, Ooi YS, Wee SL, Tan KH (2015) Comparative sensitivity to methyl eugenol of four putative *Bactrocera dorsalis* complex sibling species – further evidence that they belong to one and the same species *B. dorsalis*. In: De Meyer M, Clarke AR, Vera MT, Hendrichs J (Eds) Resolution of Cryptic Species Complexes of Tephritid Pests to Enhance SIT Application and Facilitate International Trade. ZooKeys 540: 313–321. doi: 10.3897/zookeys.540.6099

Abstract

Males of certain species belonging to the *Bactrocera dorsalis* complex are strongly attracted to, and readily feed on methyl eugenol (ME), a plant secondary compound that is found in over 480 plant species worldwide. Amongst those species is one of the world's most severe fruit pests the Oriental fruit fly, *B. dorsalis* s.s., and the former taxonomic species *B. invadens*, *B. papayae* and *B. philippinensis*. The latter species have been recently synonymised with *B. dorsalis* based on their very similar morphology, mating compatibility, molecular genetics and identical sex pheromones following consumption of ME. Previous studies have shown that male fruit fly responsiveness to lures is a unique phenomenon that is dose species-specific, besides showing a close correlation to sexual maturity attainment. This led us to use ME sensitivity as a behavioural parameter to test if *B. dorsalis* and the three former taxonomic species had similar sensitivity towards odours of ME. Using Probit analysis, we estimated the median dose of ME required to elicit species' positive response in 50% of each population tested (ED_{50}). ED_{50} values were compared between *B. dorsalis* and the former species. Our results showed no significant differences between *B. dorsalis* s.s., and the former *B. invadens*, *B. papayae* and *B. philippinensis* in their response to ME. We consider that the

Bactrocera males' sensitivity to ME may be a useful behavioural parameter for species delimitation and, in addition to other integrative taxonomic tools used, provides further supportive evidence that the four taxa belong to one and the same biological species, *B. dorsalis*.

Keywords

Bactrocera dorsalis, *B. invadens*, *B. papayae*, *B. philippinensis*, methyl eugenol, male response, lure sensitivity

Introduction

A number of fruit fly species in the *Bactrocera dorsalis* complex are pests of economic importance. The most notorious is the Oriental fruit fly, *B. dorsalis* (Hendel), a widely distributed and invasive species which includes the recently synonymised *B. invadens* Drew, Tsuruta & White, *B. papayae* Drew & Hancock and *B. philippinensis* Drew & Hancock. The presence of this species in the tropics and subtropics has caused significant damage by rendering infested fruits inedible and prohibiting fruit exports due to strict quarantine restrictions. Direct and indirect losses attributed to pestiferous tephritids are believed to be over US\$2 billion annually (Malavasi 2014). As a case in point, the 2014 ban on the entry of the popular mango (*Mangifera indica*) cv. 'Alphonso' from India to the EU countries (European Commission 2014) due to detection of Oriental fruit fly, made headlines globally and created an uproar in India, which is the largest mango producer in the world. Although the ban was lifted in January 2015 (European Commission 2015), the duration of the ban had adversely affected the lucrative mango export and livelihood of growers.

Further adding to the global fruit fly problem was the incursion into Africa of suspected Oriental fruit fly in 2003 (Lux et al. 2003), subsequently described as a new species, *B. invadens*, two years later (Drew et al. 2005). The *B. invadens*' African invasion was so serious that within three years of first incursion into Kenya, it had already been found to attack over 50 types of fruits in over 20 countries (Ekesi et al. 2006). Whilst *B. invadens*, *B. dorsalis*, *B. papayae* and *B. philippinensis* were known to appear similar morphologically and genetically, could interbreed and produce viable offspring as well as having identical male sex pheromone components (Tan et al. 2011, 2013, Schutze et al. 2013), it was only recently that all four species were synonymized as *B. dorsalis* (Schutze et al. 2015).

Previous studies have shown that male fruit fly responsiveness to the male lure methyl eugenol (ME) is a unique phenomenon that is dose species-specific, besides showing a close correlation to sexual maturity attainment (Wong et al. 1989, 1991, Wee and Tan 2000a). For example, *B. papayae* and *B. dorsalis* demonstrate similar ranges of sensitivity to ME at nanogramme levels, while *B. carambolae* has been shown to be at least 10 times less sensitive to ME compared to the former two species (Wee et al 2002). This led us to hypothesize that if *B. dorsalis*, *B. invadens*, *B. philippinensis* and *B. papayae* are different names for the one biological species, then their sensitivity to ME could be used as a behavioural parameter to confirm, or refute that assumption,

i.e. we evaluated if the sensitivity of those putative species to ME were significantly different (as might be expected if different biological species), or were the same (as might be expected if they are populations of the same biological species). In this paper, the minimum dose of ME needed to elicit species' positive response in 50% of the population tested (ED_{50}) was determined using Probit analysis and compared between the putative species as a species' delimiting tool.

Methods

Insects

Colonies of adult *B. dorsalis* and the former *B. invadens*, *B. philippinensis* and *B. papayae* were maintained in UPM insectary under strict quarantine. *Bactrocera papayae* were raised from locally collected infested starfruits, *Averrhoa carambola* L.; while *B. dorsalis*, *B. invadens*, *B. philippinensis* were obtained as pupae from the FAO/IAEA Insect Pest Control Laboratory in Seibersdorf, Austria in 2010. Pupae were imported into Malaysia using permits issued by the Director-General of Department of Agriculture Malaysia to AKW Hee. The origins of the Seibersdorf cultures are as follows: *B. dorsalis*, Saraburi, Thailand; *B. invadens*, Kenya; and *B. philippinensis*, Guimaras, the Philippines. All adult flies were provided with water and a mixture of sugar and hydrolysed protein (3:1 w/w) *ad libitum*. The flies were bred under conditions of 25–29 °C with 83–90% relative humidity, and a 12 L: 12 D photoperiod. Male flies were separated within three days of emergence (DAE) to prevent mating and were maintained in separate cages (30 cm × 30 cm × 30 cm) until required for bioassay at 19 DAE.

Chemical preparation

Different concentrations of ME (50, 100, 300, 500, 700 and 1000 ng per 5 µl of absolute ethanol, respectively) were prepared by serial dilution from pure ME (>99.8% purity; Merck-Schuchardt, Germany) following preliminary dose-response tests that showed attraction of male flies to ME of between 15–85%, which is in the linear portion of the population response curve (Heong et al. 2013). The use of six dilutions exceeds the minimum of five suggested for studies of this type by Robertson et al. (2007).

Probit regression of male flies' attraction to methyl eugenol

Laboratory bioassays with sexually mature and virgin male flies for their attraction to ME were conducted with slight modifications from the protocol of Wee et al. (2002). Males (30 males per replicate per cage [40 cm × 40 cm × 40 cm]) were acclimatized for 24 h before experimentation, in an isolated indoor area to prevent exposure of ME to

other colonies of flies that were yet to be assayed for ME response. Starting with 50 ng, the lowest concentration, 5 μ l of diluted ME was dispensed using a Hamilton® 5 μ l syringe onto a filter paper disc (3 cm \times 3cm; Whatman® No.1) placed in a disposable Petri dish. During the bioassay, we promptly removed any male that approached the ME spot and attempted to feed. After 5 min, the total number of flies that responded positively was recorded. The same procedure was conducted for new batches of males at higher ME concentrations. There were between 5–9 replicates performed for each dosage and putative species using flies from different cohorts on different days. Absolute ethanol (>99%) was used as a control. All used filter papers and petri dishes were securely disposed in airtight containers to remove traces of ME from the laboratory environment.

The data obtained were pooled and subjected to Probit analyses using the PoloPlus software (LeOra Software 2002) based on Finney (1971). Probit regressions along with ED₅₀ (effective median dose – that dose which elicits a response in 50% of the population tested) values were generated. Regression lines fitted to the dose response curves were subjected to a parallelism test using PoloPlus. This is necessary as relative potency, a measure of the species' responses to ME (based on ratios of ED₅₀), can only be valid when those ME dose-response regression lines are found to be parallel (Robertson et al. 2007; Heong et al. 2013).

Results

The male flies displayed typical behaviour in response to ME i.e. immediate zig-zag flying in locating the source of ME upon sensing the lure, followed by compulsive feeding on ME. When offered to the flies, ME attracted all four putative species at all of the tested doses (Figure 1). Approximately 25% of *B. dorsalis* responded to 50 ng of ME, which was the lowest concentration of ME tested and their responses were observed to gradually increase when subjected to higher dosages (Figure 1). The ED₅₀ response to ME for *B. dorsalis* males was 268 ng (Table 1). Similar behavioural responses were also observed in the males of the other three putative species. At the lowest dose of 50 ng, ME attracted 18%, 22% and 16% of *B. invadens*, *B. philippinensis* and *B. papayae* males, respectively (Figure 1), with their corresponding ED₅₀ values being 222 ng, 256 ng and 247 ng (Table 1). The validity of the ED₅₀ values was confirmed with heterogeneity factors of each species' response to ME showing values < 1 (*B. dorsalis*, 0.484; *B. invadens*, 0.714; *B. philippinensis*, 0.502 and *B. papayae*, 0.859), indicating the data fits well with the model of standardized residuals when plotted against the predicted values. The likelihood ratio test of parallelism revealed that the slopes of the regression lines for *B. dorsalis*, *B. invadens*, *B. philippinensis* and *B. papayae* were not significantly different (Figure 2) ($p > 0.05$; $\lambda^2 = 7.27$; $df = 3$). This allowed us to compare the relative potencies of the four species to ME. The attraction of *B. dorsalis*, *B. invadens*, *B. philippinensis* and *B. papayae* males to ME were found to be not significantly different based on their relative potency values which ranged between 0.9 and 1.1 (Table 1). In all controls, no males were attracted to filter paper containing absolute ethanol.




Figure 1. Dose-sensitivity response curves of *B. dorsalis* and the former taxa *B. papayae*, *B. invadens* and *B. philippinensis* to methyl eugenol at different doses.




Figure 2. Probit lines of *B. dorsalis* and the former taxa *B. papayae*, *B. philippinensis* and *B. invadens* attraction to methyl eugenol.

Table 1. Probit analysis of male attractancy to methyl eugenol for *Bactrocera dorsalis* and the former taxa *B. papayae*, *B. philippinensis* and *B. invadens*.

| Species | <i>n</i> ^a | Regression equation | χ^2 | <i>df</i> | ED ₅₀ ^b (ng) | 95% fiducial limits (ng) | | Relative potency |
|--------------------------|-----------------------|---------------------|----------|-----------|------------------------------------|--------------------------|-------|------------------|
| | | | | | | Lower | Upper | |
| <i>B. papayae</i> | 210 | y=1.108x-2.651 | 3.435 | 4 | 247 | 211 | 287 | 1.0 |
| <i>B. philippinensis</i> | 330 | y=1.043x-2.511 | 2.008 | 4 | 256 | 224 | 290 | 1.0 |
| <i>B. invadens</i> | 270 | y=1.266x-2.970 | 2.857 | 4 | 222 | 195 | 250 | 0.9 |
| <i>B. dorsalis</i> | 210 | y=1.002x-2.433 | 1.936 | 4 | 268 | 226 | 316 | 1.1 |

^a *n* - number of insects tested per dose

^b ED₅₀ (effective median dose) is the dose required to elicit a positive response in 50% of the fruit fly population tested.

Discussion

Bactrocera dorsalis and the three former species *B. invadens*, *B. philippinensis* and *B. papayae* all showed similar sensitivity to the male lure, ME, with non-significant differences in ED₅₀ and potency values. Only marginal difference in the ED₅₀ between *B. papayae* and *B. dorsalis* in the current study is in contrast to our earlier work that showed approximately twice the level of ED₅₀ of *B. papayae* over *B. dorsalis* (Wee et al. 2002). We believe that the discrepancy in those ratios is attributed to the different geographical strains of *B. dorsalis* used, as the strain that was used by Wee et al. originated from Taichung, Taiwan, whilst the current strain used in this study originated from Saraburi, Thailand. Further, it must be noted that there was also no significant difference in the consumption of ME between *B. dorsalis* and *B. papayae* (Wee et al. 2002). The absence of parallelism test in Wee et al. was due to the fact that the inclusion of *B. carambolae* in that study elicited a very high ED₅₀ value (17 and 9 times higher than that of *B. dorsalis* and *B. papayae*) that resulted in the unsuitability of parallelism test of the regression lines for that study.

ME is found naturally in over 480 plant species (Tan and Nishida 2012) and is a pheromone precursor/booster for ME-sensitive *Bactrocera* species (Tan and Nishida 1996, Tan et al. 2014). Synthetic ME is widely used to control pestiferous *Bactrocera* species such as *B. dorsalis* in male annihilation programmes (Steiner et al. 1965, Vargas et al. 2014). However, it was not until the discovery that the responses of male fruit fly species such as *B. dorsalis* and *B. carambolae* to ME was quantifiable (Wee et al 2002), that we have used male ME response as a species' delimitation tool. As an example, the fact that *B. carambolae* is able to interbreed with *B. papayae* and produce viable offspring both in the laboratory and in the wild (Wee and Tan 2000b, 2005, Tan 2003; but see Schutze et al. 2013) suggested that the former might be a sub-species belonging to a single species. However, that male *B. carambolae* possess significantly lower sensitivity to ME compared with that of *B. dorsalis* in cage assays (Wee et al. 2002) further prompted efforts to evaluate the status of *B. carambolae* leading to the recognition that

B. carambolae is currently a distinct, perhaps an incipient, species from that of *B. dorsalis* (Schutze et al. 2015).

An important applied finding of this study, separate to the species delimitation issue, involves the use of ME in the field as a pest control. When ME is applied in the field, the different levels of male attraction to ME between species may impact on chances of male annihilation success. In the case of *B. dorsalis* and the former taxonomic species *B. invadens*, *B. papayae* and *B. philippinensis*, application of male annihilation technique against those flies is not expected to affect the success of the programme given that all four species (now synonymised as *B. dorsalis*) have been proven to have similar sensitivities to ME.

Acknowledgements

The approval granted by the Director-General of Agriculture Malaysia to Alvin K.W. Hee to bring in live pupae shipments of *B. dorsalis*, *B. philippinensis* and *B. invadens* from IPCL, Seibersdorf is much appreciated. The financial support from FAO/IAEA offered to Keng-Hong Tan through IAEA research contract no. 16160 is also gratefully acknowledged. We also thank Todd E. Shelly for his critical comments on an earlier draft of the manuscript.

References

- Drew RAI, Tsuruta K, White IM (2005) A new species of pest fruit fly (Diptera: Tephritidae: Dacinae) from Sri Lanka and Africa. *African Entomology* 13: 149–154.
- Ekesi S, Nderitu PW, Rwomushana I (2006) Field infestation, life history and demographic parameters of the fruit fly *Bactrocera invadens* (Diptera: Tephritidae) in Africa. *Bulletin of Entomological Research* 96: 379–386. doi: 10.1079/BER2006442
- European Commission (2014) EU stops some fruit and vegetable imports from India, to protect EU crops from pests. http://ec.europa.eu/unitedkingdom/press/frontpage/2014/14_36_en.htm [accessed 25 March 2015]
- European Commission (2015) Plant Health: Experts endorse the lift of the import ban of mango fruits from India. http://europa.eu/rapid/press-release_MEX-15-3523_en.htm [accessed 25 March 2015]
- Finney DJ (1971) *Probit Analysis*, Cambridge University Press, Cambridge, 333 pp.
- Heong KL, Tan KH, Garcia CPF, Liu Z, Lu Z (2013) *Research Methods in Toxicology and Insecticide Resistance Monitoring of Rice Planthoppers*, International Rice Research Institute, Los Baños, 145 pp. doi: 10.13140/2.1.4699.5527
- LeOra Software (2002) *PoloPlus*, POLO for Windows LeOra Software.
- Lux SA, Copeland RS, White IM, Manrakhan A, Bilh M (2003) A new invasive fruit fly species from the *Bactrocera dorsalis* (Hendel) group detected in East Africa. *Insect Science and its Application* 23: 355–361. doi: 10.1017/S174275840001242X

- Malavasi A (2014) Introductory remarks. In Shelly TE, Epsky N, Jang EB, Reyes-Flores J, Vargas RI (Eds) (2014) Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies, Springer, Dordrecht, ix.
- Robertson JL, Russell RM, Preisler HK, Savin NE (2007) Bioassays with Arthropods, CRC Press, Boca Raton, 199 pp.
- Schutze MK, Jessup A, Ul-Haq I, Vreysen MJB, Wornoayporn V, Vera MT, Clarke AR (2013) Mating compatibility among four pest members of the *Bactrocera dorsalis* fruit fly species complex (Diptera: Tephritidae). *Journal of Economic Entomology* 106: 695–707. doi: 10.1603/EC12409
- Schutze MK, Aketarawong N, Amornsak W, Armstrong KF, Augustinos A, Barr N, Bo W, Bourtzis K, Boykin LM, Cáceres C, Cameron SL, Chapman TA, Chinvinijkul S, Chomič A, De Meyer M, Drosopoulou ED, Englezou A, Ekesi S, Gariou-Papalexiou A, Hailstones D, Haymer D, Hee AKW, Hendrichs J, Hasanuzzaman M, Jessup A, Khamis FM, Krosch MN, Leblanc L, Mahmood K, Malacrida AR, Mavragani-Tsipidou P, McInnis DO, Mwatawala M, Nishida R, Ono H, Reyes J, Rubinoff DR, San Jose M, Shelly TE, Srikachar S, Tan KH, Thanaphum S, Ul Haq I, Vijaysegaran S, Wee SL, Yesmin F, Zacharopoulou A, Clarke AR (2015) Synonymization of key pest species within the *Bactrocera dorsalis* complex (Diptera: Tephritidae): taxonomic changes based on 20 years of integrative morphological, genetic, behavioural, and chemoecological data. *Systematic Entomology* 40: 456–471. doi: 10.1111/syen.12113
- Steiner LF, Mitchell WC, Harris EJ, Kozuma TT, Fujimoto MS (1965) Oriental fruit fly eradication by male annihilation. *Journal of Economic Entomology* 58: 961–964. doi: 10.1093/jee/58.5.961
- Tan KH (2003) Interbreeding and DNA analysis of sibling species within the *Bactrocera dorsalis* complex. In: Recent Trends on Sterile Insect Technique and Area-Wide Integrated Pest Management- Economic Feasibility, Control Projects, Farmer Organization and *Bactrocera dorsalis* Complex Control Study. Research Institute for Subtropics, Okinawa, 113–122.
- Tan KH, Nishida R (1996) Sex pheromone and mating competition after methyl eugenol consumption in the *Bactrocera dorsalis* complex. In: McPherson BA, Steck GJ (Eds) Fruit Fly Pests. A World Assessment of Their Biology and Management. St. Lucie Press, Florida, 147–153.
- Tan KH, Nishida R (2012) Methyl eugenol - its occurrence, distribution, and role in nature, especially in relation to insect behavior and pollination. *Journal of Insect Science* 20: 74 pp. doi: 10.1673/031.012.5601
- Tan KH, Tokushima I, Ono H, Nishida R (2011) Comparison of phenylpropanoid volatiles in male rectal pheromone gland after methyl eugenol consumption, and molecular phylogenetic relationship of four global pest fruit fly species - *Bactrocera invadens*, *B. dorsalis*, *B. correcta* and *B. zonata*. *Chemoecology* 21: 25–33. doi: 10.1007/s00049-010-0063-1
- Tan KH, Wee SL, Ono H, Nishida R (2013) Comparison of methyl eugenol metabolites, mitochondrial COI, and rDNA sequences of *Bactrocera philippinensis* (Diptera: Tephritidae) with those of three other major pest species within the *dorsalis* complex. *Applied Entomology and Zoology* 48: 275–282. doi: 10.1007/s13355-013-0183-5
- Tan KH, Nishida R, Jang EB, Shelly TE (2014) Pheromones, male lures, and trapping of tephritid fruit flies. In: Shelly TE, Epsky N, Jang EB, Flores-Reyes J, Vargas RI (Eds) Trapping

- and the Detection, Control, and Regulation of Tephritid Fruit Flies, Springer, New York, 15–74. doi: 10.1007/978-94-017-9193-9_2
- Vargas RI, Leblanc L, Piñero JC, Hoffman KM (2014). Male Annihilation, Past, Present, and Future. In: Shelly TE, Epsky N, Jang EB, Flores-Reyes J, Vargas RI (Eds) Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies, Springer, New York, 493–511. doi: 10.1007/978-94-017-9193-9_14
- Wee SL, Tan KH (2000a) Sexual maturity and intraspecific mating success of two sibling species of the *Bactrocera dorsalis* complex. *Entomologia Experimentalis et Applicata* 94: 133–139. doi: 10.1046/j.1570-7458.2000.00613.x
- Wee SL, Tan KH (2000b) Interspecific mating of two sibling species of the *Bactrocera dorsalis* complex in a field cage. In: Tan KH (Ed.) Area-Wide Control of Fruit Flies and Other Insect Pests, Penerbit Universiti Sains Malaysia, Penang, 667–674.
- Wee SL, Hee AKW, Tan KH (2002) Comparative sensitivity to and consumption of methyl eugenol in three *Bactrocera dorsalis* (Diptera: Tephritidae) complex sibling species. *Chemoecology* 12: 193–197. doi: 10.1007/PL00012668
- Wee SL, Tan KH (2005) Evidence of natural hybridization between two sympatric sibling species of *Bactrocera dorsalis* complex based on pheromone analysis. *Journal of Chemical Ecology* 31: 845–858. doi: 10.1007/s10886-005-3548-6
- Wong TTY, McInnis DO, Nishimoto JI (1989) Relationship of sexual maturation rate to response of Oriental fruit fly strains (Diptera: Tephritidae) to methyl eugenol. *Journal of Chemical Ecology* 15: 1399–1405. doi: 10.1007/BF01014839
- Wong TTY, McInnis DO, Ramadan MM, Nishimoto JI (1991) Age-related response of male melon flies *Dacus cucurbitae* (Diptera: Tephritidae) to cue-lure. *Journal of Chemical Ecology* 17: 2481–2487. doi: 10.1007/BF00994596