

ORIGINAL REPORT

## ULTRASONOGRAPHIC MEASUREMENT OF FEMORAL CARTILAGE THICKNESS IN PATIENTS WITH SPINAL CORD INJURY

Murat Kara, MD<sup>1</sup>, Tülay Tiftik, MD<sup>1</sup>, Öznur Öken, MD<sup>1</sup>, Nuray Akkaya, MD<sup>2</sup>, Hakan Tunç, MD<sup>1</sup> and Levent Özçakar, MD<sup>3</sup>

From the <sup>1</sup>Ankara Physical Medicine and Rehabilitation Education and Research Hospital, Ankara, <sup>2</sup>Pamukkale University Medical School, Department of Physical Medicine and Rehabilitation, Denizli, and <sup>3</sup>Hacettepe University Medical School, Department of Physical Medicine and Rehabilitation, Ankara, Turkey

**Objective:** To assess femoral cartilage thickness in patients with spinal cord injury.

**Subjects:** Forty-six patients with SCI (35 men, 11 women; mean age: 33.6 years (standard deviation 8.1) and 46 age-, sex- and body mass index (BMI)-matched healthy subjects were enrolled.

**Methods:** Patients were evaluated with the American Spinal Injury Association (ASIA) Impairment Scale, Modified Ashworth Scale, Walking Index for Spinal Cord Injury, and Functional Independence Measurement. Mid-point ultrasonographic femoral cartilage thickness measurements were taken from the right lateral condyle, right intercondylar area, right medial condyle, left medial condyle, left intercondylar area and left lateral condyle.

**Results:** Ultrasonographic measurements revealed significantly thicker values in the intercondylar areas (bilaterally) and the medial condyle (left knee) of patients with spinal cord injury compared with those of controls. When the subgroups were compared with their paired healthy controls, measurements pertaining to the motor complete group were found to be significantly thicker in the intercondylar areas (bilaterally) and the medial condyle (left knee). Cartilage thickness values correlated negatively with the duration of immobilization (for bilateral intercondylar areas), and with BMI and ASIA level (for bilateral lateral condyles).

**Conclusion:** Femoral cartilage thicknesses were found to change after spinal cord injury, and to have a negative correlation with disease duration and severity. Future studies including histological evaluations may elucidate whether such changes are favourable for the knee joints of patients with spinal cord injury.

**Key words:** spinal cord injury; femoral cartilage; thickness; ultrasound; immobilization.

J Rehabil Med 2013; 45: 145–148

Correspondence address: Murat Kara, Ankara Fizik Tedavi Rehabilitasyon Eğitim ve Araştırma Hastanesi, Sıhhiye, Ankara, Turkey. E-mail: mkarafr@yahoo.com

Submitted December 12, 2011; accepted September 26, 2012

### INTRODUCTION

Spinal cord injury (SCI) is one of the most common causes of long-term unloading and immobilization, whereby changes in the periarticular connective tissue, contractures, periarticular osteoporosis, joint space narrowing and articular cartilage degeneration have been described (1–10).

A literature search into the effects of unloading, due to immobilization, on the joint cartilage, revealed a number of reports showing increased, decreased or unchanged thickness in animal models (11–14). However, there are very few studies in humans that have evaluated the knee cartilage of patients with SCI using magnetic resonance imaging (MRI). The thickness of knee cartilage in patients with SCI has been shown to be decreased, both in cross-sectional comparison with healthy controls (1) and during a 1-year prospective follow-up (2). However, these studies were performed in a very limited number of patients with inhomogeneous disease course. Focussing on the possible impact of cartilage alterations later in their lives, we reasoned that assessment of the femoral cartilage in a larger group of patients with SCI would elucidate the relevant consequences.

The objectives of this study were therefore: (i) to compare the femoral cartilage thicknesses of patients with SCI with those of healthy subjects; (ii) to study the correlation between cartilage thickness and disease-related parameters in these patients. Ultrasonography, which has been shown previously to be a valid and reliable method for evaluating femoral cartilage, was used in this study (15–19).

### METHODS

A total of 46 patients with SCI (35 men, 11 women) and 46 able-bodied age-, sex- and BMI-matched controls (35 men, 11 women) were enrolled in this multi-centre cross-sectional study. Subjects with a history of knee arthritis or surgery before SCI were excluded. All patients were informed about the study procedure and consented to participate. The study protocol was approved by the local ethics committee of one of the attending centres.

Initially, demographic features of the subjects, including age, sex, body weight/height, occupation and other clinical characteristics, were

noted. Patients were asked to evaluate their average daily standing/not standing time since their injury. Immobilization duration was defined as the time since injury for which patients reported daily standing with a standing frame or crutches and braces of less than 1 h daily, or staying in a bed or wheelchair all day.

#### Outcome measures

The functional status of the patients was scored according to the American Spinal Injury Association (ASIA) Impairment Scale (20). Accordingly, the patients were divided into two groups, motor complete (ASIA A and B) and motor incomplete (ASIA C and D).

The Modified Ashworth Scale (MAS) was used to grade the spasticity of the knee flexor and extensor muscles (21). Walking ability was evaluated with the Walking Index for Spinal Cord Injury (WISCI II), which assesses the amount of physical assistance, braces or devices required to walk 10 m. The scores on the WISCI II ranged from 0 (unable to stand and/or participate in walking) to 20 (ambulates with no devices, with brace and no assistance is designated) (22). Motor items of the Turkish version of the Functional Independence Measurement (FIM) scale (scoring from 13 to 91) were used to evaluate motor functioning (23–25).

#### Ultrasonographic cartilage measurements

Ultrasonographic thickness measurements of the cartilage were made with a linear probe (7–12 MHz Logiq P5, GE Medical Systems, California, USA). With subjects sitting in a comfortable position on the examination table with their knees in maximum flexion, the probe was placed in an axial position on the suprapatellar area. The distal femoral cartilage was visualized as a strongly anechoic structure between the sharp bony cortex and the suprapatellar fat. Three (mid-point) measurements were taken from each knee, at the right lateral condyle (RLC), the right intercondylar area (RIA), the right medial condyle (RMC), the left medial condyle (LMC), the left intercondylar area (LIA), and the left lateral condyle (LLC) (Fig. 1).

#### Statistical analysis

Data were expressed as mean (standard deviation (SD)) for nominal variables, and as median (minimum–maximum) for ordinal variables. Paired samples *t*-test was used to compare mean knee cartilage thickness values (SCI vs able-bodied subjects). Correlations between patients' characteristics and femoral cartilage thicknesses were analysed using Pearson's correlation coefficients. Statistical analysis was performed with SPSS 15.0. Statistical significance was set at  $p < 0.05$ .




Fig. 1. Ultrasonographic images (suprapatellar axial view) demonstrating bilateral femoral cartilage measurements of patients with spinal cord injury. RLC: right lateral condyle, RIA: right intercondylar area, RMC: right medial condyle, LMC: left medial condyle, LIA: left intercondylar area, LLC: left lateral condyle, R: right, L: left.

Table I. Clinical characteristics of patients with spinal cord injury (n=46)

	Patients
Age, years, mean (SD)	33.6 (8.1)
Disease duration, months, mean (SD)	26.3 (34.2)
Duration of immobilization, months, mean (SD)	11.5 (16.7)
Level of the lesion, n (%)	
Cervical	12 (26.1)
Thoracic	25 (54.3)
Lumbar	9 (19.6)
ASIA level	
A	17 (37.0)
B	9 (19.6)
C	10 (21.7)
D	10 (21.7)
FIM motor score (median range)	62 (13–91)
WISCI-II (median range)	5 (0–20)
LE-MAS (median range)	0 (0–4)

SD: standard deviation; ASIA: American Spinal Injury Association; WISCI-II: Walking Index for Spinal Cord Injury; LE-MAS: Lower Extremity Modified Ashworth Scale; FIM: Functional Independence Measurement.

## RESULTS

The demographic and clinical characteristics of the patients are shown in Table I. The mean age of the patients and controls was 33.6 years (SD 8.1) (age range 19–47 years). There was no statistical difference between the groups in terms of BMI; the mean BMI of the patients and controls were 23.4 kg/m<sup>2</sup> (SD 3.7) and 24.1 kg/m<sup>2</sup> (SD 2.9), respectively.

Mean femoral cartilage thickness values of the subjects are shown in Table II. Compared with those of the controls', cartilage measurements were significantly thicker at intercondylar areas bilaterally and at the medial condyle on the left knee.

When subgroups were compared with their paired healthy controls, cartilage measurements were significantly thicker at intercondylar areas (right knee  $p = 0.039$ , left knee  $p = 0.015$ ) and at the medial condyle (left knee  $p = 0.028$ ) only in the motor complete group.

Correlation coefficients between cartilage thickness values and patients' characteristics showed that cartilage thickness values were significantly (all  $p < 0.05$ ) negatively correlated with duration of immobilization at bilateral intercondylar areas (right knee  $r = -0.371$ , left knee  $r = -0.349$ ), and with BMI (right knee  $r = -0.380$ , left knee  $r = -0.306$ ) and ASIA level

Table II. Comparison of femoral cartilage thickness measurements (mm)

	Patients Mean (SD)	Controls Mean (SD)	<i>p</i> -value
RLC	2.24 (0.5)	2.22 (0.4)	0.910
RIA	2.54 (0.6)	2.20 (0.5)	0.007
RMC	2.28 (0.5)	2.19 (0.5)	0.417
LMC	2.31 (0.5)	2.10 (0.4)	0.028
LIA	2.60 (0.7)	2.19 (0.5)	0.002
LLC	2.29 (0.6)	2.20 (0.5)	0.399

SD: standard deviation; SCI: spinal cord injury; RLC: right lateral condyle; RIA: right intercondylar area; RMC: right medial condyle; LMC: left medial condyle; LIA: left intercondylar area; LLC: left lateral condyle.

Table III. Correlations between patients' characteristics and cartilage thickness values

	RLC	RIA	RMC	LLC	LIA	LMC
Age	-0.063	-0.064	-0.051	-0.092	-0.061	-0.076
Disease duration	-0.199	-0.163	-0.130	-0.123	-0.173	-0.051
Duration of immobilization	-0.251	-0.371*	-0.115	-0.163	-0.349*	-0.053
BMI	-0.380*	-0.185	-0.130	-0.306*	-0.243	-0.179
ASIA level	-0.408*	-0.137	-0.299	-0.424*	-0.031	-0.203
LE-MAS	0.113	0.197	0.080	-0.081	-0.142	-0.006
FIM motor score	-0.286	-0.270	-0.278	-0.298	-0.260	-0.227
WISCI-II	-0.218	0.023	-0.223	-0.144	-0.027	-0.093

\*Correlation is significant at 0.05 level.

RLC: right lateral condyle; RIA: right intercondylar area; RMC: right medial condyle; LMC: left medial condyle; LIA: left intercondylar area; LLC: left lateral condyle.; BMI: body mass index; ASIA: American Spinal Injury Association; LE-MAS: Lower Extremity Modified Ashworth scale; FIM: Functional Independence Measurement; WISCI-II: Walking Index for Spinal Cord Injury.

(right knee  $r = -0.408$ , left knee  $r = -0.424$ ) at bilateral lateral condyles (Table III).

## DISCUSSION

Femoral cartilage thickness in patients with SCI was evaluated in comparison with healthy subjects. Overall, cartilage measurements were thicker at both intercondylar areas and were negatively correlated with duration of immobilization and severity of injury.

Recent advances in musculoskeletal ultrasonography have enabled quantitative assessment of articular cartilage thickness in different joints. In addition, due to its several advantages (i.e. being a quick, dynamic and cost-effective method), we used ultrasonography as the imaging method in our study.

According to our results, the femoral cartilage was thicker in bilateral intercondylar areas and in left-sided medial femoral condyle. In the relevant literature for different animal models, increased, decreased or unchanged thickness of the articular cartilage has been found with different immobilization positions or with different measurement areas (i.e. different cartilage plates or layers of the knee) (3, 11–14, 26, 27). O'Connor (11) examined knee cartilage in a rat model after unloading combined with non-rigid immobilization in flexion. After 4 weeks, they observed that cartilage thickness had increased by 15–22% at the anterior femoral and by 10% at the anterior tibial regions, with no changes on the posterior sites. In a recent study with a SCI rat model, cartilage thickness has been shown to be increased at unapposed regions and decreased at apposed regions of the flexed knee joint (27). Furthermore, previous studies have also indicated that the contact areas and stresses substantially changed during knee flexion with or without quadriceps contraction (28). Similarly, as patients with SCI spend a lot of time sitting in wheelchairs with their knees in flexion, loading of the knee joint may naturally change after the injury.

In normal knee joints, the maximum femoral cartilage thickness is reported to be located in the middle of the femoral trochlea (29, 30), where high contact pressures are known to act (31, 32). On the other hand, when the immobilization in SCI is taken into account, our aforementioned finding might

seem to be controversial at first glance. However, since previous reports have mentioned additional mechanisms (e.g. sensory denervation) that may play a role in the dynamics of the cartilage milieu (33), we suggest that a biomechanical model alone is not sufficient to explain the whole scenario. Likewise, our findings of thicker cartilage values in patients with SCI (probably due to oedema) might be caused by immobilization and denervation. Moreover, loading and/or long-lasting denervation in the chronic period might cause cartilage degeneration and atrophy, which may explain our finding of negative correlation between cartilage thickness and duration of immobilization. On the other hand, since thicker cartilage values of these patients might not necessarily indicate better/healthier cartilage, our findings need to be evaluated further.

A possible limitation of this study is that we assessed cartilage thickness only, and not volume. However, it has also been shown that differences in cartilage volume may result primarily from a difference in joint size rather than cartilage thickness (34, 35).

We conclude that central portions of the femoral cartilage tend to be thicker and negatively correlated with disease duration and severity after SCI. Future studies, including histological evaluations, could elucidate whether such cartilage changes are favourable for the knee joints of SCI patients, which might be functional thereafter.

*The authors declare no conflicts of interest.*

## REFERENCES

1. Vanwanseele B, Eckstein F, Knecht H, Stüssi E, Spaepen A. Knee cartilage of spinal cord-injured patients displays progressive thinning in the absence of normal joint loading and movement. *Arthritis Rheum* 2002; 46: 2073–2078.
2. Vanwanseele B, Eckstein F, Knecht H, Spaepen A, Stüssi E. Longitudinal analysis of cartilage atrophy in the knees of patients with spinal cord injury. *Arthritis Rheum* 2003; 48: 3377–3381.
3. Moriyama H, Nishihara K, Hosoda M, Saka Y, Kanemura N, Takayanagi K, et al. Contrasting alteration patterns of different cartilage plates in knee articular cartilage after spinal cord injury in rats. *Spinal Cord* 2009; 47: 218–224.
4. Moriyama H, Yoshimura O, Kawamata S, Takemoto H, Saka Y, Tobimatsu Y. Alteration of knee joint connective tissues during contracture formation in spastic rats after an experimentally in-

- duced spinal cord injury. *Connect Tissue Res* 2007; 48: 180–187.
5. Moriyama H, Yoshimura O, Sunahori H, Tobimatsu Y. Comparison of muscular and articular factors in the progression of contractures after spinal cord injury in rats. *Spinal Cord* 2006; 44: 174–181.
  6. Moriyama H, Yoshimura O, Sunahori H, Nitta H, Imakita H, Saka Y, et al. Progression and direction of contractures of knee joints following spinal cord injury in the rat. *Tohoku J Exp Med* 2004; 204: 37–44.
  7. Jurvelin J, Kiviranta I, Tammi M, Helminen JH. Softening of canine articular cartilage after immobilization of the knee joint. *Clin Orthop Relat Res* 1986; 207: 246–252.
  8. Kuettner KE, Cole AA. Cartilage degeneration in different human joints. *Osteoarthritis Cartilage* 2005; 13: 93–103.
  9. Hagiwara Y, Ando A, Chimoto E, Saijo Y, Ohmori-Matsuda K, Itoi E. Changes of articular cartilage after immobilization in a rat knee contracture model. *J Orthop Res* 2009; 27: 236–242.
  10. Ando A, Suda H, Hagiwara Y, Onoda Y, Chimoto E, Saijo Y, et al. Reversibility of immobilization-induced articular cartilage degeneration after remobilization in rat knee joints. *Tohoku J Exp Med* 2011; 224: 77–85.
  11. O'Connor KM. Unweighting accelerates tidemark advancement in articular cartilage at the knee joint of rats. *J Bone Miner Res* 1997; 12: 580–589.
  12. Kiviranta I, Jurvelin J, Tammi M, Saamanen AM, Helminen HJ. Weight bearing controls glycosaminoglycan concentration and articular cartilage thickness in the knee joints of young beagle dogs. *Arthritis Rheum* 1987; 30: 801–809.
  13. Haapala J, Arokoski JP, Hyttinen MM, Lammi M, Tammi M, Kovanen V, et al. Remobilization does not fully restore immobilization induced articular cartilage atrophy. *Clin Orthop Relat Res* 1999; 362: 218–229.
  14. Leroux MA, Cheung HS, Bau JL, Wang JY, Howell DS, Setton LA. Altered mechanics and histomorphometry of canine tibial cartilage following joint immobilization. *Osteoarthritis Cartilage* 2001; 9: 633–640.
  15. Lee CL, Huang MH, Chai CY, Chen CH, Su JY, Tien YC. The validity of in-vivo ultrasonographic grading of osteoarthritic femoral condylar cartilage: a comparison with in vitro ultrasonographic and histologic gradings. *Osteoarthritis Cartilage* 2008; 16: 352–358.
  16. Möller I, Bong D, Naredo E, Filippucci E, Carrasco I, Moragues C, et al. Ultrasound in the study and monitoring of osteoarthritis. *Osteoarthritis Cartilage* 2008; 16: S4–S7.
  17. Yoon CH, Kim HS, Ju JH, Jee WH, Park SH, Kim HY. Validity of the sonographic longitudinal sagittal image for assessment of the cartilage thickness in the knee osteoarthritis. *Clin Rheumatol* 2008; 27: 1507–1516.
  18. Castriota-Scanderberg A, De Micheli V, Scarale MG, Bonetti MG, Cammisà M. Precision of sonographic measurement of articular cartilage: inter- and intraobserver analysis. *Skeletal Radiol* 1996; 25: 545–549.
  19. Mathiesen O, Konradsen L, Torp-Pedersen S, Jorgensen U. Ultrasonography and articular cartilage defects in the knee: an in vitro evaluation of the accuracy of cartilage thickness and defect size assessment. *Knee Surg Sports Traumatol Arthrosc* 2004; 12: 440–443.
  20. Frost FS. Spinal cord injury medicine. In: Braddom RL, editor. *Physical medicine and rehabilitation*. Philadelphia, PA: WB Saunders Co.; 2000, p. 1230–1282.
  21. Bohannon RW, Smith MB. Interrater reliability of a modified Ashworth scale of muscle spasticity. *Phys Ther* 1987; 67: 206–207.
  22. Ditunno PL, Dittuno JF. Walking index for spinal cord injury (WISCI II): scale revision. *Spinal Cord* 2001; 39: 654–656.
  23. Granger CV, Gresham GE. New developments in functional assessment. *Phys Med Rehabil Clin N Am* 1993; 4: 417–499.
  24. Keith RA, Granger CV, Hamilton BB, Sherwin FS. The Functional Independence Measure: a new tool for rehabilitation. *Adv Clin Rehabil* 1987; 1: 6–18.
  25. Kucukdeveci AA, Yavuzer G, Elhan AH, Sonel B, Tennant BA. Adaptation of the Functional Independence Measure for use in Turkey. *Clin Rehabil* 2001; 15: 311–319.
  26. Trudel G, Himori K, Uthoff HK. Contrasting alterations of apposed and unapposed articular cartilage during joint contracture formation. *Arch Phys Med Rehabil* 2005; 86: 90–97.
  27. Moriyama H, Yoshimura O, Kawamata S, Takayanagi K, Kurose T, Kubota A, et al. Alteration in articular cartilage of rat knee joints after spinal cord injury. *Osteoarthritis Cartilage* 2008; 16: 392–398.
  28. Stammberger T, Eckstein F, Michaelis M, Englmeier KH, Reiser M. Interobserver reproducibility of quantitative cartilage measurements: comparison of B-spline snakes and manual segmentation. *Magn Reson Imaging* 1999; 17: 1033–1042.
  29. Adam C, Eckstein F, Milz S, Putz R. The distribution of cartilage thickness within the joints of the lower limb of elderly individuals. *J Anat* 1998; 193: 203–214.
  30. Adam C, Eckstein F, Milz S, Schulte E, Becker C, Putz R. The distribution of cartilage thickness in the knee-joints of old-aged individuals -measurement by A-mode ultrasound. *Clin Biomech (Bristol, Avon)* 1998; 13: 1–10.
  31. Ahmed AM, Burke DL. In-vitro measurement of static pressure distribution in synovial joints-part 1: tibial surface of the knee. *J Biomech Eng* 1983; 105: 216–225.
  32. Ahmed AM, Burke DL, Yu A. In-vitro measurement of static pressure distribution in synovial joints-part 2: Retropatellar surface. *J Biomech Eng* 1983; 105: 226–236.
  33. Finsterbush A, Friedman B. The effect of sensory denervation on rabbits' knee joints. A light and electron microscopic study. *J Bone Joint Surg Am* 1975; 57: 949–956.
  34. Faber SC, Eckstein F, Lukasz S, Mühlbauer R, Hohe J, Englmeier K-H, et al. Gender differences in knee joint cartilage thickness, volume and articular surface areas: assessment with quantitative three-dimensional MR imaging. *Skeletal Radiol* 2001; 30: 144–150.
  35. Tunç H, Oken O, Kara M, Tiftik T, Doğu B, Unlü Z, et al. Ultrasonographic measurement of the femoral cartilage thickness in hemiparetic patients after stroke. *Int J Rehabil Res* 2012; 35: 203–207.