

Effect of Water Temperature and Salinity on the Fertilized Egg Development and Larval Development of Sevenband Grouper, *Epinephelus septemfasciatus*

Young-Bo Song¹, Chi-Hoon Lee¹, Hyeong-Cheol Kang¹, Hyung-Bae Kim² and †Young-Don Lee¹

¹Marine Science Institute, Jeju National University, Jeju 690-968, Republic of Korea

²Dept. of Marine Life Science & Aquaculture, Gangwon Provincial College, Gangwon 210-804, Republic of Korea

ABSTRACT : The fertilized eggs of *E. septemfasciatus* are spherical and transparent with buoyancy at 790 to 890 μm (average $821.8 \pm 2.0 \mu\text{m}$) in diameter with 170 to 230 μm oil globules (average $192.9 \pm 0.93 \mu\text{m}$). Hatching began approximately 46 and 35 hours after fertilization at 22.0°C and 25.0°C water temperature, respectively. The average total length of newly hatched larvae was $1.75 \pm 0.03 \text{ mm}$. Most of the yolk and oil globules were absorbed within 3 to 4 days after hatching. The larvae reached 2.48 to 2.72 mm in total length, and their mouths and anuses opened at 3 to 4 days after hatching. In this time, the mouth diameters of the larvae were 0.209 to 0.238 mm. The larvae reached 3.24 to 4.15 mm in total length at 11 to 17 days after hatching, and began to metamorphose at the time the second dorsal and pelvic spines appeared and elongated. The abdominal cavity was densely lined with melanophores. The larvae reached 5.12 mm in total length at 24 days after hatching.

Key words : Sevenband grouper, *Epinephelus septemfasciatus*, Fertilized egg development, Characteristics of larvae

INTRODUCTION

Sevenband grouper, *Epinephelus septemfasciatus* is a fish genus in the *Epinephelus* subfamily of the family Serranidae, in the order of Perciformes. The subfamily, Epinephelinae contains about 159 species in 15 genera (FAO, 1993). Sevenband groupers, a carnivorous species, mostly inhabit near coral reefs in subtropical or tropical regions and are distributed widely along the southern coast of Korea and Jeju Island. It is also found along the southern coast of Japan, the South China Sea, and as far as the Indian Ocean (Kim et al., 2001). It is a commercially significant product for South Korea, Japan, China, and some Southeast Asian countries. It, along with the longtooth grouper, is considered a local Jeju culinary specialty. However, the sevenband grouper catch has been decreasing

rapidly. In Korea the sevenband grouper spawns during summer season, and the embryogenesis and hatching rate are dependent on water temperature and salinity. However, little data are available regarding fertilized egg, embryonic development and larval stages. The techniques for sevenband grouper seedling production are needed for the Korean aquaculture industry to be more competitive and coastal resources management.

Some research on larval rearing, egg development, and the early life history of Epinephelinae were done in brown spotted grouper (*E. tauvina*) (Hussain & Higuchi, 1980), Nassau grouper (*E. striatus*) (Powell & Tucker, 1992), dusky grouper (*E. marginatus*) (Glamuzina et al., 1998), and longtooth grouper (*E. bruneus*) (Oh et al., 2003) for the aquaculture. Quality issues of cultured fish and advanced commercial production for cultural management

Manuscript received 21 June 2013, Received in revised form 19 August 2013, Accepted 17 November 2013

† Corresponding Author : Young-Don Lee, Marine Science Institute, Jeju National University, 19-5, Hamdeok 5(O)-gil, Jocheon, Jeju Special Self-Governing Province, Jeju 695-965, Republic of Korea. Tel./Fax : +82-64-782-8281, E-mail : leemri@jejunu.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

were studied as well under various conditions such as: water temperature (Watanabe et al., 1995), rearing tank color, density (Duray et al., 1996), aeration, and light intensity (Toledo et al., 2002). This study aims to examine egg development, early larval development under controlled water temperature and salinity for seedling production and feeding management

MATERIALS AND METHODS

1. Egg development and hatching

After artificial fertilization, using eggs selected from mature females and sperm from mature males, we measured the egg size and oil globules and examined their growth and development in elapsed time progress. The experiments were carried out in a thermostatic chamber ($22 \pm 0.5^\circ\text{C}$) and observed with a microscope (Zeiss, Germany) and a profile projector (Mitutoyo, Japan). Developmental progress, from the 8 cell stage, the 16 cell stage, morula stage, blastula stage, gastrula stage, Kupffer's vesicle occurrence, embryo formation, and hatching, was determined when each stage was 50% complete. The fertilized eggs were contained in 1.0 L beakers and experiments were repeated thrice.

2. Egg development under different water temperatures and salinities

To investigate the egg developmental progress depending on the water temperature, the experimental groups were maintained at 22°C and 25°C separately. Each fertilized group had 100 fertilized eggs and was contained in a 250 ml beaker. The time was measured for each developmental stage in the morula stage, blastula stage, gastrula stage, embryo formation, and hatching. The hatching rate for both hatched larvae and failed ones was calculated as day 1 from hatching. Also, results from artificial seawater at the salinity of 34 ppt, 26 ppt, and 18 ppt respectively, were examined to determine characteristics of egg development under controlled salinity. Each group had 50 fertilized eggs and was contained in a 100 ml beaker

respectively, experiments were repeated thrice.

3. Morphological development and growth of larvae

1) Larval rearing

During the experiments, reared larval fish were fed rotifer *Brachionus rotundiformis* (size: 110 to 210 μm) in 15 individuals/ml. They were also fed approximately 500,000 cell/ml of *Nannochloropsis oculata* and *Isochrysis galbana* respectively. Larvae were fed an artificial diet (size: 150 μm . INVE, USA) since the fifteenth day after hatching and *Artemia* (2 to 3 per ml) with rotifer *B. rotundiformis* from the sixteenth day after hatching. The artificial diet's quantity was increased according to the larval development (Fig. 1).

2) Yolk absorption and mouth opening at different water temperatures

The yolk absorption rate and mouth opening time were examined at 22°C and 25°C . The yolk absorption rate and oil globule absorption rate of the larvae hatched from 2,000 to 3,000 reared eggs in a 20 L acrylic tank were investigated. Samples of 5 to 10 larvae were taken daily for measurements of yolk and oil globule. The specimens were anesthetized with MS-222 and then measured on a




Fig. 1. Rearing scheme during the larval rearing of *E. septemfasciatus*.




Fig. 2. Measurement of *E. septemfasciatus* larvae. a, total length; b, anal length; c, major axis of yolk; d, diameter of oil globule; e, minor axis of yolk.

profile projector up to the nearest 1.0 μm (Fig. 2). The yolk and oil globule in each larvae were measured for volume with the Blaxter and Hempel method (1963) as follows:

$$\text{Yolk volume} = \pi/6 \times I h^2$$

(I: major axis of yolk, h: minor axis of yolk)

$$\text{Oil globule volume} = \pi/6 \times d^3$$

(d: diameter of oil globule)

The mouth opening was measured when 80% of mouth opened in the same method as the yolk absorption measurement.

3) Changes in total length and diameter of hatched larvae

The samples of hatched larvae, collected regularly from a 20 L acrylic tank and an indoor concrete water tank (7.0×7.0×1.0 m, effective water capacity of 35 ton) were analyzed to find size changes depending on its rearing condition and time progress. The samples were measured for diameter (d) with a Shirota formula (1970), with $d = \sqrt{2} \times (\text{upper jaw length})$. The total length, upper jaw length, and anal length were measured daily until day 7 after hatching. Samples of 5 to 10 larvae were taken every two to five days as they grew. They were anesthetized with MS-222 and then measured on a profile projector up to the nearest 1.0 μm.

4. Statistical analysis

The data was analyzed for statistical significance of means using the ANOVA-test and Duncan's multiple range test (Duncan, 1955) with SAS software.

RESULTS

1. Egg development

The fertilized eggs, with a diameter from 790 to 890 μm (an average diameter of $821.8 \pm 2.0 \mu\text{m}$), were spherical and transparent with buoyancy (Fig. 3). Oil globules had a diameter range from 170 to 230 μm (an average diameter of $192.9 \pm 0.9 \mu\text{m}$). Table 1 and Fig. 4 detail the phases of embryonic development at 22 °C (Table 1, Fig. 4). Shortly after fertilization, perivitelline space and blastodisc were formed (Fig. 4A). One hour after fertilization, the blastodisc divided into 2 cells for the first cleavage (Fig. 4B). The eggs developed into 4 cells two hours after fertilization (Fig. 4C), 8 cells after 2 hours and 30 minutes (Fig. 4D), and 32 cells after 3 hours 30 minutes (Fig. 4E). Then blastomeres derived during the cleavages and formed the morula after 5 hours (Fig. 4F). The blastula stage started




Fig. 3. Frequency in diameters of fertilized egg (A) and oil globule (B) of *E. septemfasciatus*.

Table 1. Development stages of *E. septemfasciatus* fertilized eggs at 22.0±0.5

Developmental stage	Time after fertilization	Features of eggs and embryos
Fertilized egg	0 h	Spherical & transparent with buoyancy
2 cells	1 h	First cleavage
4 cells	2 hrs	Second cleavage, plane perpendicular to the first
8 cells	2 hrs 30 mins	Third cleavage, plane parallel to the second
32 cells	3 hrs 30 mins	
Morula	5 hrs	
Blastula	12 hrs	Start of blastula stage
Gastrula	16 hrs 30 mins, 23 hrs	Forming a germ ring, blastopore depression
Early embryo formation	27 hrs	Kuffer's vesicle appearance, 7 to 9 myotomes stage
Myotomes formation	29 to 31 hrs	Kuffer's vesicle disappearance, 11 to 12 myotomes stage
Lens & ear vesicle formation	37 hrs	17 to 20 myotomes stage
Heart beat	44 hrs	Embryonic movements
Hatched larval	46 hrs	Hatching begins




Fig. 4. Microscopic view of embryonic development of *E. septemfasciatus*. A, fertilized egg; B, 2 cell; C, 4 cell; D, 8 cell; E, 32 cell; F, morula stage; G, blastula stage; H, I, gastrula stage; J, K, embryo formation; L, myotomes formation; M, lens and ear vesicle formation; N, heart beat; O, hatched larva. Scale bar=400 μ m.

after 12 hours (Fig. 4G), forming the blastoderm. 16 hours after the fertilization, gastrulation started (Fig. 4H), forming a germ ring, as the blastoderm spread around the yolk. Then after 23 hours, embryonic cells began to invaginate, forming embryonic shield (Fig. 4I). 27 hours after the fertilization, the head divided forming optic vesicles, developing 7 to 9 myotomes, and showing Kuffer's vesicles (Fig. 4J, K). 29 to 31 hours after the fertilization, the Kuffer's vesicles disappeared and 11 to 12 myotomes formed (Fig. 4L). After 37 hours, lens and ear vesicles formed and the myotomes increased to 17 to 20 (Fig. 4M). After 44 hours, embryonic movements and heartbeat were observed (Fig. 4N). The hatching began 46 hours after the fertilization (Fig. 4O).

2. Comparison of developing progress in embryogenesis at different water temperatures

Each embryogenesis stage required a different time period at a different water temperature. Developing at the morula stage required an average of 5 hours at 22°C and an average of 4 hours and 30 minutes at 25°C. Blastulation required an average of 12 hours at 22°C and an average of 10 hours and 30 minutes at 25°C. Embryo formation required an average of 23 hours at 22°C and an average of 17 hours at 25°C. Hatching required an average of 46

hours at 22°C and an average of 35 hours at 25°C.

3. Hatching rate at different water temperatures and salinities

Hatching time at salinities of 18 ppt, 26 ppt, and 34 ppt showed no hatching time difference at water temperatures of 22°C and 25°C. However, the different salinities demonstrated a difference in the hatching rates at each temperature. The hatching rate at 22°C was 64.0±6.4% at the salinity of 18 ppt, 89.3±1.8% at 26 ppt, and 99.9% at 34 ppt ($P<0.05$). The hatching rate at 25°C had no difference between the groups of 26 ppt and 34 ppt, showing the percentage 98.7±0.7% and 99.3±0.7% respectively. However, the hatching rate was lower 89.3±3.7% at 18 ppt ($P<0.05$, Fig. 5).

4. The morphological development of larvae

1) The yolk absorption and mouth opening

The yolk volume of the hatched larvae was 0.244±0.020 mm³ (100%). The reared experiment groups at different temperatures showed different results. The yolk absorption was nearly complete, with 83.5% absorption,




Fig. 5. Hatching rates of fertilized eggs of *E. septemfasciatus* at various water temperatures and salinities. Vertical bars denote standard error of means. Different superscripts on the bars are significantly different ($P<0.05$).

Table 2. Time for the embryonic developmental stage at water temperature of 22 and 25 °C for *E. septemfasciatus*

Developmental stage	Water temperature (°C)	
	22°C	25°C
Morula	5.0 hr	4.5 hrs
Blastula	12.0 hrs	10.5 hrs
Gastrula	16.5 hrs	14.0 hrs
Embryo formation	23.0 hrs	17.0 hrs
Hatching	46.0 hrs	35.0 hrs




Fig. 6. Comparison of absorptions of yolk (A) and oil globule (B) in *E. septemfasciatus* larvae at two water temperatures. Vertical bars denote standard error of means.

24 hours after hatching at 25°C and 97.9% 48 hours after hatching. However, the yolk at 22°C absorbed 77.3% 24 hours after hatching, 88.2% 48 hours after hatching, and 93.1% 72 hours after hatching (Fig. 6A). The oil globules in the larvae were measured for volume at 0.0034±0.0002 mm³ (100%). The rates of oil globule absorption at 25°C were 66.0% 24 hours after hatching, 19.7% 48

hours after hatching, and 0.9% 96 hours after hatching while the rates of oil globule absorption at 22°C were 70.5% 24 hours after hatching, 59.8% 48 hours after hatching, 31.8% 96 hours after hatching, and 7.9% 120 hours after hatching (Fig. 6B). The mouth opened in three days after hatching at 25°C, corresponding to the time period of yolk absorption and in four days after hatching at 25°C.

2) The diameter change in hatched larvae mouths

When the mouth opened as larvae three days after hatching at 25°C, the diameter (d) was 0.209 mm. The size, then, increased and was at measured 0.238 mm on the fourth day after hatching and 0.325 mm on the 11th day after hatching. The opening diameter of the hatched larvae ranged from 0.157 mm to 0.244 mm in 0.75d and 0.105 to 0.162 mm in 0.5d (Table 3).

3) Larval morphological development

(1) 0 to 1 day after hatching

The newly hatched larvae ranged from 1.75 to 2.40 mm in total length. The oil globules were located at the posterior edge of the yolk, and the anus was located a little past the body center. The melanophores were distributed on the back of mid tails and the ventral surface. The hatched larvae floated mostly near the water surface (Fig. 7A).

Table 3. Mouth growth of *E. septemfasciatus* larvae at water temperature of 25 °C by days after hatching

Day after hatching	Total length (mm±S.E. ¹⁾	Upper jaw length (mm±S.E.)	Calculated mouth length (mm)		
			d ²⁾	0.75d	0.5d
3	2.48±0.04	0.148±0.009	0.209	0.157	0.105
4	2.72±0.06	0.169±0.006	0.238	0.179	0.119
5	2.75±0.05	0.180±0.002	0.255	0.191	0.127
6	2.78±0.03	0.195±0.004	0.275	0.207	0.138
7	2.62±0.04	0.205±0.017	0.290	0.218	0.145
9	2.94±0.04	0.209±0.005	0.295	0.221	0.148
11	3.24±0.12	0.230±0.005	0.325	0.244	0.162

¹ S.E.: Standard error of means. ²⁾ $\sqrt{2} \times$ upper jaw length

(2) 2 to 3 days after hatching

The larvae ranged from 2.46 to 2.55 mm in total length and still remained yolks and oil globules. For some of the larvae, the mouth opened 3 days after hatching. The head was formed and pectoral fins appeared. Most larvae moved to the tank bottom (Fig. 7B).

(3) 4 to 5 days after hatching

The larvae were from 2.55 to 2.72 mm in total length. Yolk and oil globules absorption was complete. The mouth and anus opened and melanophores were distributed on the lens. The dorsal fins or caudal fins were not formed yet, but the pectoral fins started to develop. The melanophores were scattered on the abdomen along the myotomes (Fig. 7D). The hatched larvae rose to the water surface




Fig. 7. Microscopic photographs of growth of *E. septemfasciatus* larvae. A, newly hatched larva, 1.75 mm TL; B, two days old larva, 2.46 mm TL; C, four days old larva, 2.72 mm TL; D, nine days old larva, 2.94 mm TL, E, twenty four days old larva, 5.12 mm TL. Scale bar=1 mm.

and some rotifer-fed larvae were observed (Fig. 7C).

(4) 7 to 10 days after hatching

The larvae were from 2.62 to 2.94 mm in total length. The primordial fin folds started to disappear and morphological changes to form dorsal fins, caudal fins, and intestine tracts were observed. The melanophores increased on the area of abdomen and primordial fin folds (Fig. 7D).

(5) 17 to 23 days after hatching

The larvae were from 4.15 to 5.12 mm in total length. The second dorsal fins and ventral fins were formed and the layers of melanophore developed on the abdomen, forming caudal fins (Fig. 7E).

DISCUSSION

The size of fertilized eggs is one of the characteristic factors to improve larval viability, forming the total length, starvation tolerance, feeding amount, and the capability to escape from predators (Kayano, 1996). The egg sizes vary depending on the species and even in the same species, the sizes vary according to population, spawning season, and adult ages (Hamamoto et al., 1986). Even within the same subfamily Epinephelinae, location to inhabit determines the differences in the egg size. For example, *E. polyphkadion*, in the Red Sea, measured from 0.71 to 0.79 mm in diameter (Rasem et al., 1997), but the ones inhabiting French Polynesian averaged 0.86 mm (Aquacop et al., 1989) in diameter. *E. coioides*, from the Red Sea, were 0.77 mm (Hussain et al., 1975) in diameter, but the ones along the Philippines coast averaged 0.84 mm in diameter of fertilized egg (Toledo et al., 1993). The above data demonstrates that even for the same species, the larvae inhabiting a higher salinity area (42 to 43 ppt) tend to be smaller in diameter. *E. akaara*, inhabiting along the southern coast of Korea, measured 0.77 mm in diameter (Lee & Hur, 1997). This study on Jeju sevenband groupers showed the fertilized eggs were spherical and transparent with buoyancy. The

eggs ranged from 0.76 to 0.88 mm (an average of 0.82 mm) in diameter and the oil globules were from 0.17 to 0.22 mm in diameter (an average of 0.19 mm) showing similar results from Kitajima et al. (1991). The embryogenesis and hatching rate are dependent on water temperature, salinity, light intensity, and aeration. Hatching for *E. striatus* requires 20 to 25 hours at water temperature ranging from 26 to 30°C (Watanabe et al., 1998). For *E. polyphkadion*, it takes 19 hours at 29 to 30°C (Rasem et al., 1997). The results from this study show that hatching for Jeju sevenband groupers requires 46 hours at 22°C and 35 hours at 25°C. The hatching rate showed no significant difference at different water temperatures ($P>0.05$) but showed low hatching rates at 22°C and 18 ppt ($P<0.05$).

The newly hatched larvae of *E. septemfasciatus* contained yolk and oil globules in the stomach and ranged from 1.75 to 2.40 mm in total length. The anus was located at the edge of the body. The yolks are the major food supplier at the early stage of embryonic development; they contain a lot of energy. Therefore, the larvae with bigger yolks can contain more energy and have more time to be fed (Bagarinao, 1986). Since the newly hatched larvae are relatively bigger than the ones from other developmental stages, the first time feeding improves viability (Hunter, 1981; Quattro & Weeks, 1991). The yolk absorption rate varies according to the rearing conditions. Epinephelinae and Siganidae, tropical marine fish species, contain relatively smaller yolks than temperate fish, but the yolk absorption occurred quicker (Bagarinao, 1986). *E. akaara* is known to absorb 90% of yolks at a water temperature of 25 to 29°C 24 hours after hatching and more than 90% in 64 to 80 hours after hatching (Lee & Hur, 1997). This study showed that sevenband grouper, *E. septemfasciatus*, consumed most of their yolks within 48 hours at the constant water temperature of 25°C but remained partial yolks at 22°C even 72 hours after hatching. The absorption of the oil globules has been reported to progress slower than the yolks (Kuo et al., 1974) due to triglyceride (Fyhn, 1989; Clyde et al., 1992). The absorption

of the oil globules was done before 96 hours after hatching at the constant water temperature of 25°C but remained partially at the constant water temperature of 22°C after the mouth opened. For *E. akaara*, the mouth opening required 42 to 62 hours after hatching at the range of water temperature from 23 to 31°C (Lee & Hur, 1997) and 46 to 84 hours at 20 to 28°C (Kayano, 1988). The results from this study represented that sevenband groupers of mouth open required 72 to 96 hours at 22 to 25°C. The larvae were measured with the Shirota formula (1970) 0.209 mm in diameter and 0.105 mm (0.5d), larger than the minimum diameter of *E. akaara* larvae 0.078 mm in 0.5d for feeding (Kayano, 1988; Lee & Hur, 1997).

The dorsal fins, caudal fins, and the intestinal tracts were formed 7 to 10 days after hatching. 17 to 23 days old larvae grew from 4.15 to 5.12 mm in total length. The second dorsal fins and ventral fins were formed, demonstrating the morphological characteristics of Epinephelinae (Kitajima et al., 1991). When the larvae of sevenband grouper *E. septemfasciatus* grew to approximately 8.0 mm, the total length of second dorsal fins and ventral fins increased 75% and 90% respectively and then grew to adulthood. *C. altivelis* has a different morphological change in second dorsal fins, the total length growing 120% (Sugama & Ikenoue, 1999). *E. tauvina* is also noted to have differential development, growing 48% of their total length (Hussain & Higuchi, 1980). This determines egg development varies according to each species.

REFERENCES

- Aquacop J, Fuchs G, Nedelee, Gasset E (1989) Selection of finfish species as candidates for aquaculture in French Polynesia. In: Barret J (ed.), Advances in Tropical Aquaculture. Actes de Colloques. IFREMER 9:143-156.
- Bagarinao T (1986) Yolk absorption, onset of feeding and survival potential of larvae of three tropical marine fish species reared in the hatchery. Mar Biol 91:449-459.
- Blaxter JHS, Hempel G (1963). The influence of egg size on herring larvae (*Clupea harengus* L.). J Cons int Explor Mer 28:211-244.
- Clyde ST, Harry A, Lee CS (1992) Fatty acid and free amino acid profiles of spawned eggs of striped mullet, *Mugil cephalus*. Aquaculture 105:83-94.
- Duncan DB (1955) Multiple-range test and multiple F test. Biometrics 11:1-42.
- Duray MN, Estudillo CB, Alpasan LG (1996) The effect of background color and rotifer density on rotifer intake, growth and survival of the grouper (*Epinephelus suillus*) larvae. Aquaculture 146:217-224.
- FAO (1993) FAO Species Catalogue Vol. 16. Groupers of the World. FAO, Rome pp 119-120.
- Fyhn HJ (1989) First feeding of marine fish larvae are free amino acids the source of energy. Aquaculture 80:111-120.
- Glamuzina B, Skaramuca B, Glavic N, Kozvul V, Dulcic J, Kraljevic M (1998) Egg and early larval development of laboratory reared dusky grouper, *Epinephelus marginatus* (Lowe, 1834) (Picipes, Serranidae). Sci Mar 62(4):373-378.
- Hamamoto S, Yokogawa K, Tochino M (1986) Several problems on cultivating the parent fish of red spotted grouper, *Epinephelus akaara* (Temminck et Schegel), and judging the qualities of the eggs obtained from them. Bull Kagawa Pref Fish Exp Stn 2:13-22.
- Hunter JR (1981) Feeding ecology and predation of marine fish larvae. In: Lasker RI (ed.), Marine Fish Larvae-Morphology. Ecology and Relation to Fisheries. Univ Washington Press. Seattle and London. pp 33-77.
- Hussain NA, Saif M, Ukawa M (1975) On the culture of *Epinephelus tauvina* (Forsk.). Kuwait Institute for Scientific Research, Kuwait. pp 12.
- Hussain NA, Higuchi M (1980). Larval rearing and development of the brown spotted grouper, *Epinephelus tauvina* (Forsk.). Aquaculture 19:339-350.
- Kayano Y (1988) Development of mouth parts and feeding in the larval and juvenile stages of red spotted grouper, *Epinephelus akaara*. Saibai Giken 3:55-60. (in Japanese)
- Kayano Y (1996) Yearly change in egg production of the

- red spotted grouper, *Epinephelus akaara* in a rearing tank. Saibai Giken 25:47-52. (in Japanese)
- Kim IS, Choi Y, Kim BJ (2001) Percoid fishes of Korea. Korea Research Institute of Bioscience and Biotechnology, Korea. pp 279. (in Korean)
- Kitajima C, Takaya M, Tsukashima Y, Arakawa T (1991) Development of eggs, larvae and juveniles of the grouper, *Epinephelus septemfasciatus*, reared in the laboratory. Jap J Ichthyol 38:47-55. (in Japanese)
- Kuo CM, Nash CE, Shehadeh ZH (1974) A procedural guide to induce spawning in grey mullet (*Mugil cephalus* L.). Aquaculture 3:1-14.
- Lee CK, Hur SB (1997) Yolk absorption, onset of feeding and survival potential of larvae of red spotted grouper, *Epinephelus akaara*. J Aquaculture 10:473-483. (in Korean)
- Oh SR, Kang CH, Lee CH, Hur SW, Lee YD (2003) Sex reversal and masculinization according to growth in longtooth grouper *Epinephelus bruneus*. Dev Reprod 17:79-85.
- Powell AB, Tucker JW (1992) Egg and larval development of laboratory-reared Nassau grouper, *Epinephelus striatus* (Pisces, Serranidae). Bull Mar Sci 50(1):171-185.
- Quattro JM, Weeks SC (1991) Correlations between egg size and egg energetic content within and among biotypes of the genus *Poeciliopsis*. J Fish Biol 38: 331-334.
- Rasem BM, James CM, Al-Thobaiti SA, Carlos MH (1997) Spawning of the camouflage grouper, *Epinephelus plyphekadion* (Bleeker) in the hypersaline waters of Saudi Arabia. Asian Fish Sci 9:251-259.
- Shirota A (1970) Studies on the mouth size of fish larvae. Bull Jpn Soc Sci Fish 36:353-368.
- Sugama K, Ikenoue H (1999) Research and development: The seed production technique of humpback grouper, *Cromileptes altivelis*. JICA. pp 53.
- Toledo JD, Nagi A, Javellana D (1993) Successive spawning of grouper, *Epinephelus suillus* (Valenciennes), in a tank and a floating net cage. Aquaculture 115:361-367.
- Toledo JD, Caberoy NB, Quintio GF, Choresca CH, Nakagawa H (2002) Effects of salinity, aeration and light intensity on oil globule absorption, feeding incidence, growth and survival of early-stage grouper *Epinephelus coioides* larvae. Fish Sci 68:478-483.
- Watanabe WO, Lee CS, Ellis SC, Ellis EP (1995) Hatchery study of the effects of temperature on eggs and yolk sac larvae of the Nassau grouper *Epinephelus striatus*. Aquaculture 136:141-147.
- Watanabe WO, Ellis EP, Ellis SC, Feeley MW (1998) Progress in controlled maturation and spawning of summer flounder *Paralichthys dentatus* broodstock. J World Aqua Soc 29:393-404.