

Human replication protein A unfolds telomeric G-quadruplexes

Tonatiuh Romero Salas, Irina Petrusheva¹, Olga Lavrik¹, Anne Bourdoncle², Jean-Louis Mergny², Alain Favre and Carole Saintomé*

Institut Jacques Monod, CNRS-ParisVI-ParisVII-UMR 7592, 2 place Jussieu, 75251 Paris cedex 05, France, ¹Novosibirsk Institute of Bioorganic Chemistry, Siberian Division of Russian Academy of Sciences, 630090 Novosibirsk, Russia and ²Laboratoire de Biophysique, INSERM U565, CNRS UMR 5153, Muséum National d'Histoire Naturelle USM 503, 43 rue Cuvier, 75005 Paris, France

Received June 15, 2006; Revised July 17, 2006; Accepted July 19, 2006

ABSTRACT

G-quadruplex structures inhibit telomerase activity and must be disrupted for telomere elongation during S phase. It has been suggested that the replication protein A (RPA) could unwind and maintain single-stranded DNA in a state amenable to the binding of telomeric components. We show here that under near-physiological *in vitro* conditions, human RPA is able to bind and unfold G-quadruplex structures formed from a 21mer human telomeric sequence. Analyses by native gel electrophoresis, cross-linking and fluorescence resonance energy transfer indicate the formation of both 1:1 and 2:1 complexes in which G-quadruplexes are unfolded. In addition, quadruplex opening by hRPA is much faster than observed with the complementary DNA, demonstrating that this protein efficiently unfolds G-quartets. A two-step mechanism accounting for the binding of hRPA to G-quadruplexes is proposed. These data point to the involvement of hRPA in regulation of telomere maintenance.

INTRODUCTION

Telomeres are specialized DNA–protein structures that protect the ends of chromosomes and distinguish natural chromosome termini from unnatural breaks produced by DNA damage (1). Alterations in telomere structure are associated with distinct cellular programs including apoptosis and unlimited proliferation indicating that telomeres play an important role in the processes of aging and cancer (2). Telomere DNA length and protein composition vary during the life of a cell, and telomere structures may involve double-stranded DNA-binding proteins (3,4), single-stranded DNA (ssDNA)-binding proteins (5–7) and DNA–DNA interactions (8). Human telomeric DNA, located at the ends of each

chromosome, contains G-rich termini as relatively short single-stranded 3' overhangs designated G-overhangs (9,10). Such single-stranded telomere DNAs have been found in protozoa, yeast and vertebrates (11–13). They are synthesized specifically by a unique ribonucleoprotein reverse transcriptase called telomerase (14,15).

It is well known that clusters of G residues may adopt non-B structures stabilized by interactions between the guanine bases under *in vitro* physiological conditions [for a review see (16)]. These structures are four-stranded DNA complexes where layers of four guanine bases, one from each strand, are bound by Hoogsteen hydrogen bonds, thereby forming stacked G-quartets that hold the chains together. Therefore, it is conceivable that these G-quadruplex structures may occur in living cells (17,18) and affect essential cellular processes, such as recombination and extension of telomeric sequences by the telomerase (19).

It has been suggested that cells might possess a mechanism allowing them to resolve these structures into single strands, thereby providing the best opportunity for telomerase access to the 3' end of a chromosome (14). Replication protein A (RPA) was shown recently to be present at the telomeric ends of chromosomes with maximum association in the S phase and to play an essential role in telomere maintenance (7). Schramke *et al.* (20) proposed that RPA activates the telomerase via its p32 subunit, by maintaining ssDNA in a state amenable to the binding of telomeric components. In addition, Cohen *et al.* (21) showed that in an *in vitro* system, low concentrations of human RPA (hRPA) stimulate extension of G-rich DNA primers by the telomerase (although high concentrations are inhibitory) and proposed a mechanism of unwinding of the unusual structures formed between G residues.

RPA is an ssDNA-binding protein (ssDBP) that is highly conserved in eukaryotes (22,23). hRPA is a heterotrimeric complex consisting of three subunits p70, p32 and p14, named according to their molecular masses of 70, 32 and 14 kDa, respectively. hRPA has four DNA-binding domains (A, B and C in p70 and D in p32), and it binds ssDNA via a multistep pathway (24,25).

*To whom correspondence should be addressed. Tel: +33 1 44 27 40 86; Fax: +33 1 44 27 57 16; Email: saintome@ijm.jussieu.fr

MgCl₂ and 5 mM lithium cacodylate (pH 7.2). Concentrated protein aliquots (0.5 µl) were directly added to the F-htelo-T solution. The spectra were recorded between 490 and 660 nm while exciting at 470 nm, and corrected for background fluorescence, dilution factor and instrument response. The fluorescence intensity of individual fluorophores was estimated by averaging the fluorescence emission intensity in the 522–528 nm region for the donor (FLUO) and 584–590 nm for the acceptor (TAMRA; the low donor emission in that wavelength range was neglected). The ratio P was calculated as $P = I_D/(I_D + I_A)$, where I_D and I_A are the average intensities of the donor and acceptor, respectively.

Fluorescence kinetics

Each solution containing 90 nM F-htelo-T in 2 mM MgCl₂ and 5 mM lithium cacodylate (pH 7.2) was mixed at time zero with a 5× molar excess of its complementary sequence 21C or hRPA. The kinetics were recorded at 20°C in 50 mM KCl. Fluorescence intensity at 516 nm was recorded at regular time intervals (1 s) using band slits of 5 nm. Data fitting was performed as described previously (27).

RESULTS

The model sequence chosen for our study is a 21mer human telomeric repeat sequence (htelo): 5'-GGGTTAGGGTTAGGGTTAGGG-3' which mimics the telomeric G-rich tail. It is well established that telomeric sequences may adopt several quadruplex conformations such as antiparallel (28) and parallel (29) G-quadruplex structures depending on the salt conditions (Figure 1a). We selected a short fragment for our study, because four GGG blocks are sufficient for stable G-quadruplex formation *in vivo* (30).

hRPA binds to G-quadruplexes and forms 1:1 and 2:1 complexes

To determine whether hRPA binds the htelo sequence, we first performed electrophoretic mobility shift assays




Figure 1. Schematic representation of quadruplex and duplex structures. (a) Different possible folding topologies of htelo. Closed circles depict guanines, TTA loops represent single-stranded DNA regions. (b) Schematic representation of a 21 bp duplex formed between F-htelo-T and its complementary sequence 21C where cytosines are depicted by gray circles. FLUO and TAMRA are depicted by closed and open triangles, respectively.

(EMSA). In a standard experiment, 90 nM ³²P-labeled htelo was incubated for 10 min at 20°C with hRPA in the presence of 50 mM of either Na⁺ or K⁺. Each mixture was defined by ' r ', the ratio of hRPA added relative to htelo (expressed as a molar ratio).

Figure 2a illustrates the results of EMSA experiments obtained when the binding reaction was performed in the presence of Na⁺. In the presence of hRPA, one or several retarded bands were obtained, demonstrating that this protein is able to form noncovalent complexes with htelo. Each complex (C) in a lane was defined by its relative mobility m_C taking free htelo as reference, $m_f = 1$ (Figure 2a). At low protein/DNA ratios ($r = 0.5$ –1), a single well-defined complex, designated C_I, with a relative mobility $m_{C_I} = 0.27 \pm 0.01$, was detected. It was accompanied by free htelo and by a fast migrating smear. Increasing r from 2 to 5 progressively reduced the mobility of the main complex and quenched the smear, while free htelo vanished. For r -values >5, most of the htelo (80–90%) concentrated in a low-mobility band, designated C_{II}, with a relative mobility $m_{C_{II}} = 0.16 \pm 0.01$; its mobility remained unchanged upon increasing r up to 14 (data not shown).

To clarify the nature of these retarded complexes, we first determined their hRPA to htelo molar ratios (hRPA/htelo). In this case, after scanning of the radioactivity using a Phosphorimager (Figure 3a), the gels were submitted to Coomassie blue staining; the intensity of the protein-containing bands was quantitatively analyzed with a densitometer (Figure 3b; Materials and Methods). Results showed that free hRPA migrates as a low-mobility band, $m_p = 0.09$, and that hRPA and htelo co-migrate in C complexes. A range of control experiments with Coomassie staining showed that the signal obtained was proportional to the amount of free hRPA loaded on the gel and that the presence of htelo did not alter protein detection and measurement. The data of the ratios obtained are summarized in Table 1. When r varied from 1 to 7, the protein to DNA ratio increased continuously from ~1 to >2, showing that C_I is a 1:1 hRPA–htelo complex, while C_{II} is a 2:1 complex accompanied possibly by higher-order complexes.

To further characterize the complexes at intermediate and high r values, protein–DNA cross-linking experiments with 0.1% glutaraldehyde were performed (31). Reaction mixtures were then loaded onto native gels. As shown in Figure 4a,




Figure 2. Titration of htelo as a function of hRPA concentration. ³²P-htelo (90 nM) was incubated with various amounts of hRPA (from 45 to 630 nM) in the presence of 50 mM NaCl (a) or KCl (b) and separated on a native 5% polyacrylamide gel. C_I and C_{II} represent 1:1 and 2:1 hRPA–htelo complexes, respectively. r is the molar ratio hRPA/htelo.




Figure 3. Determination of the hRPA/htelo ratios in complexes. ^{32}P -htelo (90 nM) was incubated with 90 or 630 nM of hRPA in the presence of 50 mM NaCl and separated by native 5% PAGE. Analysis using a Phosphorimager revealed radioactive htelo (a), and densitometric analysis revealed colored hRPA (b). The boxes show the areas considered to determine the hRPA/htelo ratios.

Table 1. Quantification of hRPA to htelo ratios in complexes C

	$r = 1$	$r = 3$	$r = 5$	$r = 7$
hRPA/htelo ratio	1.04 ± 0.11	1.51 ± 0.1	1.9 ± 0.17	2.27 ± 0.17

For each r -value, the hRPA/htelo ratio found in the retarded band(s) was measured by phosphorimager and densitometric analysis.

glutaraldehyde pretreatment revealed two ($r = 1$) or three well-defined bands ($r = 7$) designated C'_I , C'_{II} , C'_{III} . The covalent complexes C'_I and C'_{II} migrated as their non-cross-linked C_I and C_{II} homologues indicating that C'_I and C'_{II} are 1:1 and 2:1 covalent complexes, respectively. C'_{III} was observed only at $r \geq 3$; it may represent 3:1 hRPA-htelo complex pre-existing in solution, or result from an artifact induced by cross-linking. These cross-linking experiments show clearly that formation of 2:1 complex increases with r and becomes the major species for $r = 7$. Combining these data and the hRPA to htelo ratios found in C complexes (Table 1), we concluded that C complexes observed at intermediate r -values (Figure 2a) that migrate with a relative mobility comprised between C_I and C_{II} are mixtures of 1:1 (C_I) and 2:1 (C_{II}) complexes; its continuously decreasing mobility observed when r increased (Figure 2a) can be explained by the increasing formation of 2:1 complex. Thus, we calculated for each r value the relative weights f_0 , f_1 and f_2 in solution of free htelo, complex C_I and complex C_{II} , respectively (Materials and Methods and Figure 5a).

Parallel experiments were performed in a buffer containing K^+ instead of Na^+ . hRPA binding was qualitatively similar in the presence of either ion (Figures 2b and 5b). However, for the same r value the amount of C complexes was always lower in K^+ than in Na^+ . At $r = 7$, 60% of the htelo was involved in 2:1 complex in K^+ as compared to 90% in Na^+ , and a significant fraction of G-quadruplexes remained unbound (7%). These differences indicate a lower affinity of hRPA for htelo in the presence of K^+ , which is in line with the finding that in the presence of K^+ , G-quadruplex structures are more stable (32).




Figure 4. Comparison of htelo and T21 titrations by hRPA in the absence or presence of glutaraldehyde. ^{32}P -DNA (90 nM) was incubated with various amounts of hRPA (from 90 to 630 nM) in the presence of 50 mM NaCl then cross-linked by the addition of 0.1% glutaraldehyde for 10 min. Individual reaction mixtures were analyzed on a native 5% polyacrylamide gel. (a) EMSA results after incubation of htelo. (b) EMSA results after incubation of T21. r is the molar ratio hRPA/DNA. C_I and C_{II} represent the non-covalent 1:1 and 2:1 hRPA-DNA complexes, respectively. C'_I , C'_{II} and C'_{III} represent the covalent 1:1, 2:1 and putative 3:1 hRPA-DNA complexes, respectively.

The 1:1 complex is predominantly formed with a control oligonucleotide unable to form a G-quadruplex

As a control in these binding experiments, we tested a radio-labeled single-stranded 21mer oligodeoxythymidine T21 (90 nM). Sequential addition of hRPA in the presence of 50 mM Na^+ led to a different EMSA profile (compare Figure 4a and b): (i) the weights of complexes observed on the gel without glutaraldehyde cross-linking at distinct r -values is slightly higher than for htelo, and (ii) the mobility of the C complexes was not reduced as much as for htelo. Glutaraldehyde cross-linking experiments showed that whatever the r -value, the C'_I covalent complex predominates. All these results agree with previously reported data (23) showing that hRPA is an ssDBP which predominantly forms 1:1 complexes with short DNAs. Thus, it may be assumed that formation of major 2:1 complexes with htelo is governed by some peculiar properties of this short structured DNA, which assists in binding of two heterotrimeric molecules.

hRPA binding leads to G-quadruplex unfolding

Fluorescence resonance energy transfer (FRET) experiments were used to study the conformation of htelo bound to hRPA (33). For this purpose, the htelo oligomer was labeled with two fluorophores, fluorescein (FLUO) and tetramethylrhodamine (TAMRA) attached to the 5' and 3' termini, respectively, leading to the dual-labeled oligonucleotide F-htelo-T (Figure 1b). Fluorescence emission of FLUO (the donor, D) is efficiently quenched by TAMRA (the acceptor, A) if the distance D-A is short. If D and A are distant from one another, their respective emissions become independent of each other. Here the extent of energy transfer is represented by an empirical parameter $P = I_D / (I_D + I_A)$, where I_D and I_A are the emission intensities of D and A, respectively (Materials and Methods). For free F-htelo-T, low P -values (P_{Quad}) typical of G-quadruplex structures were obtained with $P_{\text{Quad}} = 0.18 \pm 0.04$ in the presence of Na^+ and $P_{\text{Quad}} = 0.25 \pm 0.05$ in the presence of K^+




Figure 5. Quantification of f_0 , f_1 and f_2 fractions as a function of r . The relative weights in solution of free htelo (f_0 , solid line), complex C_1 (f_1 , long dashed line) and complex C_{II} (f_2 , short dashed line) were quantified for each r -value from hRPA–htelo EMSA (Figure 2) in the presence of Na^+ (a) or K^+ (b). The relative errors vary from 20% for the low r -values ($r = 0.5$ –1) to 10% for the high r -values ($r > 1$). r is the molar ratio hRPA/htelo.




Figure 6. hRPA binding leads to G-quadruplex unfolding. Fluorimetric titration of 90 nM of F-htelo-T by increasing amounts of hRPA was performed in the presence of Na^+ (a) or K^+ (b). The spectra were recorded after 2 min of incubation. (c) Increasing of P as a function of the hRPA/F-htelo-T ratio (r) in the presence of Na^+ (solid line) or K^+ (dashed line) indicates unfolding of F-htelo-T by addition of hRPA. P was calculated from experimental FRET data: $P = I_D / (I_D + I_A)$, where I_D and I_A are the average intensities of the donor (FLUO) and acceptor (TAMRA), respectively. For each P -value, error bar is ± 0.05 . AU, arbitrary units.

(Figure 6) (33). In others words, FRET efficiency in the absence of protein ($r = 0$) is higher in Na^+ than in K^+ , in agreement with recent study (34). A 21mer duplex formed between F-htelo-T and its complementary sequence, designated 21C, served as a control for quadruplex opening

(Figure 1b). It was demonstrated previously that the stability of the F-htelo-T/21C duplex is higher than that of the quadruplex (35). Addition of a 5-fold molar excess of 21C over F-htelo-T (90 nM) led to a high P -value (P_{Duplex}) of 0.83 ± 0.05 after equilibration, whatever the nature of the monovalent

cation. This is consistent with the formation of a B-DNA duplex structure separating D and A by >70 Å (Figure 1b).

Fluorimetric titrations of F-htelo-T by hRPA were performed in the conditions used for hRPA-htelo binding in EMSA experiments. Figure 6a shows the data obtained in the presence of Na^+ . As the ratio r increases, FLUO emission is stimulated while TAMRA fluorescence decreases, indicating that FRET is suppressed and that hRPA is able to unfold quadruplexes. The absence of an isoemissive point (Figure 6a) and the biphasic variation of P as a function of r (Figure 6c) show that the binding process is complex and involves at least three species, in agreement with the formation of the C_I and C_{II} complexes with the free G-quadruplex structures. Thus, we can write (and calculate) that $P = f_0P_0 + f_1P_1 + f_2P_2$, where P_0 , P_1 and P_2 are P values for free htelo, C_I complexes and C_{II} complexes, respectively, and f_0 , f_1 and f_2 the previously calculated fractions of each species (Figure 5a). To fit the experimental (Figure 6c) and calculated P curves, P_1 was considered as an adjustable parameter with $P_0 = 0.18$ and $P_2 = 0.83$. The agreement for both series of data is satisfactory when P_1 is taken as 0.55 ± 0.05 .

Since P_2 is nearly identical to P_{Duplex} , the final conformation of DNA bound with two (or more) hRPA corresponds to an extended structure where both fluorophores are as far away as in the duplex B-DNA structure. The lower P -value for the 1:1 complexes indicates that the distance D–A is smaller in the 1:1 than in the 2:1 complexes. This could be the result of the coming together of the single-stranded tails triggered by the presence of only one bound hRPA.

At first glance, titrations in the presence of K^+ (Figure 6b and c) are similar to those obtained with Na^+ . However, significant differences are observed: (i) for the same r -values, P in the presence of K^+ is always lower than P in the presence of Na^+ in agreement with the binding data discussed above, and (ii) the value of P in the presence of K^+ does not reach the P_{Duplex} value at the highest r -value checked ($r = 7$) as not all htelo is bound (7% remains free) and as a non-negligible amount of 1:1 complexes is still present as shown by the EMSA experiment (Figures 2b and 5b). Taking $P_0 = 0.25$ and $P_2 = 0.83$, the best fit between the experimental (Figure 6c) and calculated data is obtained with $P_1 = 0.6 \pm 0.05$.

One can therefore conclude that in the presence of either Na^+ or K^+ , G-quadruplex structures are unfolded in the hRPA-htelo complexes. In the 1:1 complexes, the oligonucleotide end-to-end distance is intermediate between the one observed in quadruplexes and the larger one ($d \geq 70$ Å) found in the 2:1 complexes.

hRPA efficiently unfolds G-quadruplexes

To gain insight into the opening process of the hRPA-induced G-quadruplex, its kinetics were compared to those obtained during the formation of the duplex F-htelo-T/21C (Figure 1b). Five equivalents of either hRPA or 21C were added to a solution of F-htelo-T and rapidly mixed. The kinetics were followed by measuring the emission spectra at 516 nm (FLUO emission) and are shown in Figure 7. The experiments were performed in K^+ conditions in which the stability of the G-quadruplexes is highest (32). Duplex formation could not be properly fitted with a




Figure 7. Comparison of the F-htelo-T opening by 21C and hRPA. Each solution containing 90 nM F-htelo-T was mixed at time zero with a 5× molar excess of its complementary sequence 21C (black curve) or hRPA (red curve). Fluorescence intensity was recorded at 516 nm in the presence of 50 mM KCl. AU, arbitrary units.

mono-exponential function, and was fitted with a bi-exponential model, as observed previously (27,36), with a fast phase ($k_{\text{fast}} = 0.031 \pm 0.003 \text{ min}^{-1}$) accompanied by a much slower phase ($k_{\text{slow}} = 0.0083 \pm 0.0002 \text{ min}^{-1}$) (data not shown). The fast phase was of the same order of magnitude as reported previously (36,37) for the unfolding step of G-quadruplex structures. A bi-exponential function was also required to fit the binding of hRPA to htelo, but the kinetics were obviously faster (Figure 7), with $k_{\text{fast}} = 0.81 \pm 0.01 \text{ min}^{-1}$ and $k_{\text{slow}} = 0.080 \pm 0.001 \text{ min}^{-1}$. These relatively fast kinetics observed with hRPA as compared to complementary DNA highlight the active role of hRPA in quadruplex opening.

Fluorimetric titrations of the dually labeled single-stranded oligonucleotide control F-T21-T by hRPA were performed in the presence of Na^+ (data not shown). hRPA has a greater affinity for the F-T21-T control sequence compared to F-htelo-T, since FRET titration is complete for $r = 1$ (with P increasing from 0.45 for $r = 0$ to 0.74 for $r \geq 1$). Under the same conditions an important smear was observed in the EMSA experiment when radiolabeled T21 was mixed with one equivalent of hRPA (Figure 4b). This observation indicates that the smear observed in the native gel at low r -values ($r = 0.5$ and 1) represents 1:1 complex dissociation during migration (Figures 2 and 4). In addition, if we compare the P -values of the 1:1 complexes obtained with F-T21-T and F-htelo-T (0.74 ± 0.01 and 0.55 ± 0.05 , respectively), we can conclude that in the 1:1 complexes with hRPA the DNA conformations of T21 and htelo differ. This is in agreement with a different mode of binding of hRPA to the quadruplex-forming substrate, as discussed above. Moreover, kinetic experiments of the dually labeled single-stranded oligonucleotide control F-T21-T by hRPA showed that <30 s were necessary to obtain maximal fluorescence emission (data not shown). This time scale agrees with the one observed previously for hRPA binding to ssDNA (38,39).

DISCUSSION

From the data presented, it can be concluded that hRPA is able to bind 21mer G-quadruplex structures by forming 1:1 complexes and 2:1 complexes, and possibly higher order complexes. The 2:1 complexes are presumably stabilized by cooperative interactions between the two hRPA molecules (26,31) leading to the very stable noncovalent complexes observable by electrophoresis. In comparison with hRPA binding to the single-stranded T21 oligomer, which mainly displays 1:1 complexes, it is clear that the hRPA-binding mode differs depending on the nature of the DNA. It is well known that RPA exhibits relatively low specificity for nucleic acid sequences with a 50-fold preference for polypyrimidine tracks (22–24,40). This sequence preference is similar to those of other nonspecific ssDBP, e.g. the *E.coli* ssDBP (*E.coli* ssDBP) (41). However it was clearly demonstrated recently that there is a general influence of the nucleic acid sequence itself on the binding interactions with ssDBPs: ssDNA binding is influenced by base stacking and the nearest-neighbor (nucleotide sequence in DNA) dependence of this stacking (42). Nevertheless, formation of major stable 2:1 complexes with a 21mer DNA strongly suggests that unlike T21, hRPA binding to G-quadruplex structures is directed by the structure itself with a sequential binding mode. This implies that the first hRPA binds to one extremity of the DNA, allowing binding of the second hRPA. In this case, as htelo is a 21mer, both hRPA should bind htelo by its 8–10 nt binding mode (22–26).

Duplex formation with a G-quadruplex prone sequence is a slow process, especially in the presence of K^+ (35). Opening of the quadruplex is a prerequisite for duplex formation, and the kinetics are dictated by quadruplex unfolding rather than by bimolecular association. In contrast, FRET and kinetics experiments demonstrate that hRPA acts rapidly and efficiently promotes G-quadruplex opening, as a few minutes only are necessary to open the htelo G-quadruplex structure(s). The fast action of hRPA might be required in cases where quadruplex lifetime is long compared to key cellular processes such as replication. Preliminary experiments with an ssDBP such as *E.coli* ssDBP suggest that under similar conditions the binding mode and kinetics of *E.coli* ssDBP are very different compared to hRPA (T. R. Salas, unpublished data), arguing for a specific effect of the hRPA on the G-quadruplex. Other ssDBP or nucleic acids chaperones will be tested to determine if hRPA possesses a unique mode of action on the G-quadruplex.

Taken together, these data lead to the sequential model of hRPA binding summarized in Figure 8. F-htelo-T only displays three short ‘single-stranded’ regions corresponding to the TTA loops (Figure 1a); it is not clear if these short single-stranded regions are sufficient for the initial binding of hRPA. Alternatively, it has been shown that G-quadruplex structures are in equilibrium with partially unfolded G-quadruplex species (F-htelo-T') where some Hoogsteen hydrogen bonds might be transiently opened (43). These single-stranded regions generated at the extremities could be accessible and rapidly trapped by hRPA. This first binding step should destabilize the hydrogen bonds between the remaining Gs of the proximal quartet, permitting the initial conformational change observed by FRET (1:1 complexes). This unfolded




Figure 8. Sequential model of hRPA binding to G-quadruplexes. hRPA binding is directed by single-stranded regions generated at the extremities of the partially unfolded structures as represented by F-htelo-T'. This first binding step destabilizes the G-quadruplex structure to form 1:1 complexes. This unfolded DNA bound to one hRPA molecule at the extremity facilitates binding of a second hRPA molecule to produce 2:1 complexes.

DNA bound with one hRPA molecule could facilitate binding of a second hRPA molecule. Thus hRPA would form two distinct complexes in which DNA is maintained in different unfolded conformations.

The idea that hRPA can remove secondary and tertiary DNA structures by a simple destabilization process was reported previously (44,45). The recent model of a multi-step helix destabilization process by hRPA described by Binz *et al.* (46) supports our results. Unlike hRPA, human helicases such as BLM and WRN, that unwind G-quadruplex structures with a 3'–5' polarity, require ATP, Mg^{2+} and at least one single-stranded 3' tail (47,48). In our system, neither ATP nor 3' tail is present. In addition, even if there is no evidence of polarity of hRPA binding to G-quadruplex structures, it is well known that hRPA binds ssDNA with the opposite 5'–3' polarity (49,50). Clearly, more detailed studies are needed to unravel the mechanism by which hRPA binds and opens G-quadruplex structures. This binding proceeds with high efficiency and significant specificity for G-quadruplexes, suggesting that it has an important biological role in telomere maintenance.

G-quadruplex structures may be important for a number of biological processes and disease-related mechanisms. Particularly, it has been shown that they inhibit telomerase activity by impeding the recruitment and binding of telomerase components to telomeres. We have revealed by this investigation that G-quadruplex structures are specific targets for hRPA and that this protein is able to bind and open G-quadruplex structures much faster than the complementary DNA strand. There is good reason to believe that opening G-quadruplex structures by RPA takes place in order to maintain the telomeric G-overhang in a single-stranded conformation, compatible with telomerase activity. Cohen *et al.* (21) showed that depending on its concentration, hRPA may exert either stimulatory or inhibitory effects on telomerase; it would be important to correlate this observation with the formation of 1:1 or 2:1 complexes. Interestingly, Zaug *et al.* (51) indicated that human POT1 is able to disrupt telomeric G-quadruplexes and that extension by the telomerase depends on the relative position of POT1 on htelo. Thus, the details of the architecture protein–DNA complexes may play an essential role in telomerase activity. We are now

investigating whether similar conclusions may be reached with hRPA.

ACKNOWLEDGEMENTS

We thank Anne-Lise Haenni (IJM, Paris, France) for careful reading of the manuscript. This work was supported by grants from GEFLUC (to C.S.), ARC (#3365 to J.L.M.), EU FP6 'Mol Cancer Med' (# LSHC-CT-2004-502943 to J.L.M.) and RFBR (06-04-48526-a, to I.P., 05-0448319 to O.L. and HSFP-RGP0007/2004-C). We are grateful to CONACYT and SEP Mexico for scholarships (to T.R.S.). Funding to pay the Open Access publication charges for this article was provided by CNRS.

Conflict of interest statement. None declared.

REFERENCES

- Blackburn,E.H. (1991) Structure and function of telomeres. *Nature*, **350**, 569–573.
- Blackburn,E.H. (2000) Telomere states and cell fates. *Nature*, **408**, 53–56.
- Li,B., Oestreich,S. and de Lange,T. (2000) Identification of human Rap1: implications for telomere evolution. *Cell*, **101**, 471–483.
- van Steensel,B., Smogorzewska,A. and de Lange,T. (1998) TRF2 protects human telomeres from end-to-end fusions. *Cell*, **92**, 401–413.
- Price,C.M. and Cech,T.R. (1987) Telomeric DNA–protein interactions of *Oxytricha* macronuclear DNA. *Genes Dev.*, **1**, 783–793.
- Baumann,P. and Cech,T.R. (2001) Pot1, the putative telomere end-binding protein in fission yeast and humans. *Science*, **292**, 1171–1175.
- Smith,J., Zou,H. and Rothstein,R. (2000) Characterization of genetic interactions with RFA1: the role of RPA in DNA replication and telomere maintenance. *Biochimie.*, **82**, 71–78.
- Neidle,S. and Parkinson,G.N. (2003) The structure of telomeric DNA. *Curr. Opin. Struct. Biol.*, **13**, 275–283.
- McEachern,M.J., Krauskopf,A. and Blackburn,E.H. (2000) Telomeres and their control. *Annu. Rev. Genet.*, **34**, 331–358.
- Wright,W.E., Tesmer,V.M., Huffman,K.E., Levene,S.D. and Shay,J.W. (1997) Normal human chromosomes have long G-rich telomeric overhangs at one end. *Genes Dev.*, **11**, 2801–2809.
- Klobutcher,L.A., Swanton,M.T., Donini,P. and Prescott,D.M. (1981) All gene-sized DNA molecules in four species of hypotrichs have the same terminal sequence and an unusual 3' terminus. *Proc. Natl Acad. Sci. USA*, **78**, 3015–3019.
- Henderson,E.R. and Blackburn,E.H. (1989) An overhanging 3' terminus is a conserved feature of telomeres. *Mol. Cell. Biol.*, **9**, 345–348.
- McElligott,R. and Wellinger,R.J. (1997) The terminal DNA structure of mammalian chromosomes. *EMBO J.*, **16**, 3705–3714.
- Collins,K. and Mitchell,J.R. (2002) Telomerase in the human organism. *Oncogene*, **21**, 564–579.
- Reveal,P.M., Henkels,K.M. and Turchi,J.J. (1997) Synthesis of the mammalian telomere lagging strand *in vitro*. *J. Biol. Chem.*, **272**, 11678–11681.
- Williamson,J.R. (1994) G-quartet structures in telomeric DNA. *Annu. Rev. Biophys. Biomol. Struct.*, **23**, 703–730.
- Schaffitzel,C., Berger,I., Postberg,J., Hanes,J., Lipps,H.J. and Plückthun,A. (2001) *In vitro* generated antibodies specific for telomeric guanine-quadruplex DNA react with *Styloynchia lemnae* macronuclei. *Proc. Natl Acad. Sci. USA*, **98**, 8572–8577.
- Duquette,M.L., Handa,P., Vincent,J.A., Taylor,A.F. and Maizels,N. (2004) Intracellular transcription of G-rich DNAs induces formation of G-loops, novel structures containing G4 DNA. *Genes Dev.*, **18**, 1618–1629.
- Zahler,A.M., Williamson,J.R., Cech,T.R. and Prescott,D.M. (1991) Inhibition of telomerase by G-quartet DNA structures. *Nature*, **350**, 718–720.
- Schramke,V., Luciano,P., Brevet,V., Guillot,S., Corda,Y., Longhese,M.P., Gilson,E. and Géli,V. (2004) RPA regulates telomerase action by providing Est1p access to chromosome ends. *Nature Genet.*, **36**, 46–54.
- Cohen,S., Jacob,E. and Manor,H. (2004) Effects of single-stranded DNA binding proteins on primer extension by telomerase. *Biochim. Biophys. Acta*, **1679**, 129–140.
- Wold,M.S. (1997) Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu. Rev. Biochem.*, **66**, 61–92.
- Iftode,C., Daniely,Y. and Borowiec,J.A. (1999) Replication protein A (RPA): the eukaryotic SSB. *Crit. Rev. Biochem. Mol. Biol.*, **34**, 141–180.
- Binz,S.K., Lao,Y., Lowry,D.F. and Wold,M.S. (2003) The phosphorylation domain of the 32-kDa subunit of replication protein A (RPA) modulates RPA–DNA interactions. *J. Biol. Chem.*, **278**, 35584–35591.
- Gomes,X.V., Henriksen,L.A. and Wold,M.S. (1996) Proteolytic mapping of human replication protein A: evidence for multiple structural domains and a conformational change upon interaction with single-stranded DNA. *Biochemistry*, **35**, 5586–5595.
- Lavrik,O.I., Kolpashchikov,D.M., Weissart,K., Nasheuer,H.P., Khodyreva,S.N. and Favre,A. (1999) RPA subunit arrangement near the 3'-end of the primer is modulated by the length of the template strand and cooperative protein interactions. *Nucleic Acids Res.*, **27**, 4235–4240.
- Kumar,N. and Maiti,S. (2005) The effect of osmolytes and small molecule on quadruplex-WC duplex equilibrium: a fluorescence resonance energy transfer study. *Nucleic Acids Res.*, **33**, 6723–6732.
- Wang,Y. and Patel,D.J. (1993) Solution structure of the human telomeric repeat d[AG₃(T₂AG₃)₃] G-tetraplex. *Structure*, **1**, 263–282.
- Parkinson,G.N., Lee,M.P.H. and Neidle,S. (2002) Crystal structure of parallel quadruplexes from human telomeric DNA. *Nature*, **417**, 876–880.
- Vorlickova,M., Chladrkova,J., Kejnovska,I., Fialova,M. and Kyrp,J. (2005) Guanine tetraplex topology of human telomere DNA is governed by the number of (TTAGGG) repeats. *Nucleic Acids Res.*, **33**, 5851–5860.
- Blackwell,L.J., Borowiec,J.A. and Mastrangelo,I.A. (1996) Single-stranded DNA binding alters human replication protein A structure and facilitates interaction with DNA-dependent protein kinase. *Mol. Cell. Biol.*, **16**, 4798–4807.
- Mergny,J.L., Phan,A.T. and Lacroix,L. (1998) Following G-quartet formation by UV-spectroscopy. *FEBS Lett.*, **435**, 74–78.
- Mergny,J.L. and Maurizot,J.C. (2001) Fluorescence resonance energy transfer as a probe for G-quartet formation by a telomeric repeat. *ChemBioChem*, **2**, 124–132.
- Juskowiak,B., Galezowska,E., Zawadzka,A., Gluszynska,A. and Takenaka,S. (2006) Fluorescence anisotropy and FRET studies of G-quadruplex formation in presence of different cations. *Spectrochim. Acta A Mol. Biomol. Spectrosc.*, **64**, 835–843.
- Alberti,P. and Mergny,J.L. (2003) DNA duplex–quadruplex exchange as the basis for a nanomolecular machine. *Proc. Natl Acad. Sci. USA*, **100**, 1569–1573.
- Green,J.J., Ying,L., Klenerman,D. and Balasubramanian,S. (2003) Kinetics of unfolding the human telomeric DNA G-quartet structure using a PNA trap. *J. Am. Chem. Soc.*, **125**, 3763–3767.
- Zhao,Y., Kan,Z.Y., Zeng,Z.X., Hao,Y.H., Chen,H. and Tan,Z. (2004) Determining the folding and unfolding rate constants of nucleic acids by biosensor. *J. Am. Chem. Soc.*, **126**, 13255–13264.
- Wang,M., Mahrenholz,A. and Lee,S.H. (2000) RPA stabilizes the XPA-damaged DNA complex through protein–protein interaction. *Biochemistry*, **39**, 6433–6439.
- Schubert,F., Zettl,H., Häfner,W., Krauss,G. and Krausch,G. (2003) Comparative thermodynamic analysis of DNA–protein interactions using surface plasmon resonance and fluorescence correlation spectroscopy. *Biochemistry*, **42**, 10288–10294.
- Wyka,I.M., Dhar,K., Binz,S.K. and Wold,M.S. (2003) Replication protein A interactions with DNA: differential binding of the core domains and analysis of the DNA interaction surface. *Biochemistry*, **42**, 12909–12918.
- Lohman,T.M. and Ferrari,M.E. (1994) *Escherichia coli* single-stranded DNA-binding protein: multiple DNA-binding modes and cooperativities. *Annu. Rev. Biochem.*, **63**, 527–570.

42. Mou, T.C., Shen, M., Abdalla, S., Delamora, D., Bochkareva, E., Bochkarev, A. and Gray, D.M. (2006) Effects of ssDNA sequences on non-sequence-specific protein binding. *Chirality*, **18**, 370–382.
43. Lee, J.Y., Okumus, B., Kim, D.S. and Ha, T. (2005) Extreme conformational diversity in human telomeric DNA. *Proc. Natl Acad. Sci. USA*, **102**, 18938–18943.
44. Treuner, K., Ramsperger, U. and Knippers, R. (1996) Replication protein A induces the unwinding of long double-stranded DNA regions. *J. Mol. Biol.*, **259**, 104–112.
45. Iftode, C. and Borowiec, J.A. (1998) Unwinding of origin-specific structures by human replication A occurs in a two-step process. *Nucleic Acids Res.*, **26**, 5636–5643.
46. Binz, S.K., Lao, Y., Lowry, D.F. and Wold, M.S. (2003) The phosphorylation domain of the 32-kDa subunit of replication protein A (RPA) modulates RPA–DNA interactions. *J. Biol. Chem.*, **278**, 35584–35591.
47. Huber, M.D., Lee, D.C. and Maizels, N. (2002) G4 DNA unwinding by BLM and Sgs1p: substrate specificity and substrate-specific inhibition. *Nucleic Acids Res.*, **30**, 3954–3961.
48. Mohaghegh, P., Karow, J.K., Brosh, R.M., Bohr, V.A. and Hickson, I.D. (2001) The Bloom's and Werner's syndrome proteins are DNA structure-specific helicases. *Nucleic Acids Res.*, **29**, 2843–28493.
49. Iftode, C. and Borowiec, J.A. (2000) 5'→3' molecular polarity of human replication protein A (hRPA) binding to pseudo-origin DNA substrates. *Biochemistry*, **39**, 11970–1181.
50. Kolpashchikov, D.M., Khodyreva, S.N., Khlimankov, D.Y., Wold, M.S., Favre, A. and Lavrik, O.I. (2001) Polarity of human replication protein A binding to DNA. *Nucleic Acids Res.*, **29**, 373–379.
51. Zaug, A., Podell, E.R. and Cech, T. (2005) Human POT1 disrupts telomeric G-quadruplexes allowing telomerase extension *in vitro*. *Proc. Natl Acad. Sci. USA*, **102**, 10864–10869.