

Retraction Notice

The following article is retracted from publication in Balkan Medical Journal by decision of editors of the journal since it is a duplicate publication, which has been previously published in *Int Arch Med.* 2011 Jan 21;4(1):4.

Additionally, the name of Hafize Öksüz -the second author of this article- was added to this article by the first author without the second author's knowledge. Since this issue is confirmed by the first and second authors and their institutions respectively name of Hafize Öksüz was retracted from this article by the decision of the Editorial Board.

Comparison of Nerve Stimulator and Ultrasonography Application for Brachial Plexus Anesthesia

Beyazıt Zencirci, Hafize Öksüz

Department of Anesthesiology and Reanimation, Faculty of Medicine, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, Turkey

ABSTRACT

Objective: Axillary brachial plexus block can be achieved through various techniques in upper extremity operations. The purpose of our study was to compare the efficacy of axillary brachial plexus block using an ultrasound technique with peripheral nerve stimulation technique.

Material and Methods: Sixty patients for whom elective forearm and hand surgery was planned were included in the study. Group 1 (n=30) was given an axillary block by using ultrasonography, Group 2 (n=30) was given axillary block by using a peripheral nerve stimulator. The quality and time of onset of the sensorial and motor blockade were assessed.

Results: The average time needed to perform the axillary brachial plexus block was similar in both groups. Although not significant statistically, it was observed that the sensory block was achieved earlier in Group 1. However, the degree of motor blockade was more intense in Group 1 than in Group 2.

Conclusion: The ultrasound-guided axillary brachial plexus block is a preferable method with faster onset time and better quality of motor blockade compared to the PNS technique.

Key Words: Nerve stimulator, brachial plexus; ultrasonography, regional anesthesia, axillary nerve block

Received: 19.09.2010

Accepted: 16.12.2010

Introduction

Regional anaesthesia can be defined as removal of nerve conduction and pain in certain parts of the body without causing sensory loss (1). Many other methods have been described up to the present since Hirschel's application of the blinding axillary block in 1911 (2, 3).

The brachial plexus can be blocked through various anatomical approaches such as interscalene, supraclavicular, infraclavicular and axillary approaches. Axillary block techniques can be applied by using transarterial fixation, paresthesia or nerve stimulator (4). Current techniques available for nerve localization mark anatomical indicators for the estimated location of brachial plexus. As well as causing anxiety in the patient and the long application processes, blinding techniques may also cause nerve damage, vein perforations and complications such as systemically local anaesthetic toxic reactions (4). The nerve stimulator technique, however, ensures that the needle is correctly placed without causing paraesthesia. Ultrasonography allows us to display the brachial plexus with a higher quality and helps nerve localization, and these factors can increase the quality of the nerve block. Through ultrasonography (US), peripheral nerves, needle localization and local anaesthetic distribution, which is required for successful conduction of a block, can be directly displayed (5).

In our study, we have aimed to compare the sensory and motor block effects of peripheral nerve stimulation (PNS), which facilitates the application of axillary brachial plexus block (AXB) and increases the prospects, and the technique of US, that has recently been put into use.

Material and Methods

Having obtained the required written consents both from the Ethics Committee and from the 60 ASA I-II patients undergoing elective minor upper-limb surgery, including forearm, wrist, and hand procedures, the patients were prospectively enrolled. Using a computer-generated sequence of random numbers and sealed envelope technique, patients were randomly allocated to receive the axillary brachial plexus block using either ultrasound (US-guided group, n=30) or nerve stimulation (PNS group, n=30) guidance. Those with a history or presence of cardiac, respiratory and/or renal failures or were pregnant were not included in the study. No premedication was administered to the cases.

An intravenous cannula was inserted into the contralateral arm, and a continuous crystalloid solution infusion was started. For the whole procedure, the patients were routinely monitored with an electrocardiogram (ECG), non-invasive blood pressure (NIBP) measurement, and pulse oximetry (SpO₂).

AXB was carried out by abducting the arm that was to be blocked in a position to create a 90° angle with the body and, by flexing and externally rotating the forearm, the hand could be placed right next to the head and the palm positioned facedown. Following the positioning of the cases in both groups, the area on the axillary region to be operated was disinfected.

After the appropriate positioning of the US-guided group-patients, and following the completion of the required preparations, a 22 G insulated needle (Stimuplex® D 50 mm, B.Braun, Germany) was inserted into the axillary region under US guidance (by using Aloka® SSD-4000, Japan, 10 Mhz prob). First the radial, next median, thirdly ulnar and lastly musculocutaneous nerves were identified. After identification of each nerve, 7-10 ml local anesthetic, with a total of 40 ml of 0.75% ropivacaine for the four nerves was injected; until the nerves were completely surrounded.

As for the cases in the PNS Group, following the appropriate positioning and completion of the required preparations, similar to the other group, a total of 40 ml-7-10 ml for each nerve of 0.75% Ropivacaine was injected by using nerve-stimulator-specific, sterile, teflon-isolated needles (22G insulated needle) (Stimuplex® D 50 mm [15°]) in company with the available nerve stimulator (Stimuplex® Dig RC, B.Braun, Melsungen, Germany). At the same time, the motor response elicited by the nerves that form the brachial plexus to nerve stimulation was also considered (radial: arm and finger extension, supination; median: wrist, 2nd and 3rd finger flexion, pronation; ulnar: 4th and 5th finger flexion, thumb adduction, musculocutaneous: arm flexion) (6).

The time included sonographic overview and identification of the targeted structures for US-guided group, identification of the nerves via peripheral nerve stimulator for PNS group, subcutaneous infiltration of the injection site, and application of local anesthetic to the direct vicinity of the four targeted nerves in both of two groups.

At the end of the AXB, the anesthetist performing the block evaluated sensory and motor block as follows: every five minutes and for 30 minutes the innervated areas in each dermatome was evaluated using a pinprick. When the needles were no longer felt, cutaneous anesthesia was considered to be present. The motor block was evaluated once at the end of the 30 minute period. The motor block was estimated as being 0, 33, 66 or 100%: 100%, no movement at all of the upper limb against gravity; 66%, flexion and/or extension movements in the hand but not in the arm; 33%, flexion and/or extension movements in both the hand and the arm against gravity but not against resistance; 0%, flexion and extension movements in both the hand and the arm against resistance (6).

The block was considered to be complete if the dermatomes of the nerves implicated in the surgical site were anaesthetized. All nerves of the surgical site including those of the skin, muscles, and bones were considered. The block was evaluated as incomplete and in need of completion before surgery if one of the nerves of the surgical site was not anesthetized.

Statistical analysis

All data were collected in an Excel®-Sheet for documentation. For statistical analysis, the program SPSS 13.0® for Windows (LEAD Technologies Inc, USA, 2004) was used. The Mann-Whitney U test was used to compare the differences between demographic data of patients such as age, height, weight, and ASA status. The Chi square test was used to compare the differences related to gender. Differences in the onset times and anesthesia between the four nerves were tested using Friedman Repeated Measures Analysis of Variance (ANOVA) on Ranks. Parameters were given as mean±standard deviation. A p value of less than 0.05 was considered statistically significant.

Results

Twenty-nine female and 31 male patients were enrolled in the study. The demographic data and ASA status of the patients are shown in Table 1. No differences between the two groups were found with regard to the demographic data or ASA status.

The average time necessary to perform the AXB was similar in the two groups (p>0.05) (Table 2).

Although not significant statistically, it was observed that the sensory block had formed earlier in US-guided group (7.3±2.6 min in US-guided group, but 6.4±3.9 min in PNS group, p=0.39). The degree of motor blockade was more intense in the US-guided group than in the PNS group (p<0.05) (Table 3). The success rate of the sonographically guided axillary plexus block was 100%.

Table 1. Demographic data and ASA status

	US-guided group	PNS group
Age (year) (mean±SD)	37.07±16.24	39.69±11.27
Gender (M/F)	13/17	18/12
Height (cm) (mean±SD)	167.01±8.69	163.56±7.24
Weight (kg) (mean±SD)	77.41±14.85	74.49±11.26
ASA Status (ASA-1/ASA-2)	14/16	12/18

Table 2. Achievement of sensory block in 4 nerves

	US-guided group	PNS group	p value
10 min	13 (43.33%)	9 (30.00%)	0.29
20 min	24 (80.00%)	17 (56.67%)	0.21
30 min	30 (100.00%)	26 (86.67%)	0.67

Table 3. Frequency distribution of patients in the two groups according to the motor blockade degree or quality

Group	Motor blockade (%)			
	0%	33%	66%*	100%*
US-guided	0	0	0	30
PNS	0	0	7	23

*Significant difference between the two groups (66% and 100%) (p<0.05)

There were neither cardiovascular side effects nor any accidental vascular punctures. No postoperative neurological symptoms were reported.

Discussion

There are various techniques to block the brachial plexus clavicle at different levels from both under and above. Recently, most of the techniques used to inject the local anesthetics have stipulated the use of paraesthesia. However, frequency of neurologic complications that occur following the AXB varies between 0.2 and 19%. This may occur as a result of a direct trauma to the nerve, local anesthetic toxicity, ischemia or a combination of all these factors (7, 8).

The spread of LA around all the nerves is obligatory to achieve complete AXB. Anatomical studies show the neurovascular space to be divided by multiple septae (9). This is the main reason for incomplete AXB. Two different methods are used to solve this problem. One is the use of high LA volumes to achieve a good distribution in the axillary sheath (10). This method has a low risk of nerve damage so the cannula is not redirected in an area already anaesthetised, but incomplete blockades occur in patients with firm tissue surrounding the nerves.

A more effective second method is the multiple approach to terminal nerve branches by using nerve stimulation (11, 12). Nerve stimulators, that were first applied in 1912 but only put into clinical use in 1962, have been an alternative to the technique of paraesthesia. It was believed that the nerve stimulator minimized the possibility of a probable neuropathy that could be caused by direct acute physical contact with the nerve with the paraesthesia technique. However, this method increases the risk of nerve damage by redirecting the cannula in a previously anaesthetized area. Therefore, paraesthesia loses its value as a warning sign (13). Fanelli et al. (11), reported a rate of 1.7% transient neurological complications using a multiple injection technique for peripheral nerve blockade.

The ultrasound approach identifies nerves, vessels, muscles, and septa. One main advantage of the sonographical approach is the ability to monitor the whole procedure of nerve blockade. Damage to important structures like vessels can be avoided during the puncture. We had no accidental vessel puncture in any patient. Therefore, redirecting the cannula can be performed under visual control. The risk of accidental nerve damage can thus possibly be reduced. On the other hand, not only does ultrasonography give us the opportunity to observe the LA solution surrounding the nerve but also it lets us observe the optimal distribution of the injected LA solution around the nerve.

In our study, 86.67% of the cases in the PNS group formed a full sensory block and 76.67% of these formed a full motor block within the first half hour (Table 2 and 3). On the other hand, in US-guided group sensory full block and motor full block rates were 100%. The fact that we obtain better results following the US application is mainly caused by the possibility of observing the nerves forming the brachial plexus and the distribution of local anaesthetic liquid. Whether the consequently applied LA liquid had completely reached the targeted tissues or not can also be monitored.

In addition, ultrasonography can also be used for difficult axillary block applications (14). Li et al. (15) reported that ultrasonography is very useful in terms of application especially for obese cases.

Schwemmer et al. (16, 17) stated that ultrasonography application significantly increases the success rate of axillary blocks and that the starting time of operation following the block is much earlier. Throughout our study, we detected that the sensory block started earlier in the ultrasonography-applied group although this was not significant statistically and on the other hand, that motor block rate in this group was significantly higher in comparison with the other group.

Soeding et al. (18) determined that ultrasonography application significantly reduced the starting time of sensory and motor block and that it significantly increased the block quality. Kefalianakis et al. (19) stated that ultrasonography application decreases the starting time of block. In our study, we have identified that sensory block onset was earlier in the ultrasonography-applied group, although that was not statistically significant.

According to Liu et al. (20), ultrasonography application provides more accomplished sensory and motor blocks. The same researchers also reported that, through ultrasonography, they succeeded in providing a completely adequate analgesia without any complications in sixteen axillary-block applied cases of final-stage renal failures (21). We did not encounter any serious complications in our ultrasonography-applied group throughout the study.

Conclusion

Consequently, we established that sensory block started earlier in the ultrasound-guided AXB, although this was not statistically significant and that the success rate of motor block was higher. We believe that ultrasonography application can be a particularly good alternative, without causing any complications in cases with anatomic complexities.

Conflict of Interest

No conflict of interest was declared by the authors.

References

1. Casati A. Local Anesthetics. In: Chelly JE, Casati A, Fanelli G, editors. *Continuous Peripheral Nerve Blocks: An Illustrated Guide*. Milano: Mosby; 2001. p.37-44.
2. Nielsen ZK. Axillary Brachial Plexus Block. In: Hadzic A, editor. *Textbook of Regional Anesthesia and Acute Pain Management*. New York: The McGraw-Hill; 2007. p.441-51.
3. Wedel DJ, Horlocker TT. Nerve Blocks. In: Miller RD, editor. *Anaesthesia 6th ed. Vol 2*. New York: Churchill Livingstone; 2005. p.1685-95.
4. Urmey WF. Upper Extremity Blocks. In: Brown DL, editor. *Regional Anesthesia and Analgesia*. Philadelphia: W.B.Saunders Company; 1996. p.266-8.
5. Gray AT. Role of ultrasound in startup regional anesthesia practice for outpatients. *Int Anesthesiol Clin* 2005;43:69-78.
6. Lavoie J, Martin R, Tétrault JP, Côté DJ, Colas MJ. Axillary plexus block using a peripheral nerve stimulator: single or multiple injections. *Can J Anaesth* 1992;39:583-6.

7. Franco CD, Vieira ZE. 1,001.subclavian perivascular brachial plexus blocks: success with a nerve stimulator. *Reg Anesth Pain Med* 2000;25:41-6.
8. Cheney FW, Domino KB, Caplan RA, Posner KL. Nerve injury associated with anesthesia: a closed claims analysis. *Anesthesiology* 1999;90:1062-9.
9. Thompson GE, Rorie DK. Functional anatomy of the brachial plexus sheaths. *Anesthesiology* 1983;59:117-22.
10. Jage J, Kossatz W, Biscopig J, Zink KU, Wagner W. Axillary blockade of the brachial plexus using 60 ml prilocaine 0.5% vs. 40 ml prilocaine 1%. A clinical study of 144 patients carried out by the determination of the prilocaine concentration in the central venous blood and by the measurement of the subfascial pressure in the plexus following the injection. *Reg Anaesth* 1990;13:112-7.
11. Fanelli G, Casati A, Garancini P, Torri G. Nerve stimulator and multiple injection technique for upper and lower limb blockade: failure rate, patient acceptance, and neurologic complications. Study Group on Regional Anesthesia. *Anesth Analg* 1999;88:847-52.
12. Coventry DM, Barker KF, Thomson M. Comparison of two neurostimulation techniques for axillary brachial plexus blockade. *Br J Anaesth* 2001;86:80-3.
13. Horlocker TT, Kufner RP, Bishop AT, Maxson PM, Schroeder DR. The risk of persistent paresthesia is not increased with repeated axillary block. *Anesth Analg* 1999;88:382-7.
14. Baumgarten RK, Thompson GE. Is ultrasound necessary for routine axillary block? *Reg Anesth Pain Med* 2006;31:88-9.
15. Li C, McCartney C, Perlas A, Chan V. Successful use of ultrasound guided axillary block in three morbidly obese patients. *Reg Anesth Pain Med* 2004;29(4 Suppl):S52.
16. Schwemmer U, Schleppers A, Markus C, Kredel M, Kirschner S, Roewer N. Operative management in axillary brachial plexus blocks: comparison of ultrasound and nerve stimulation. *Anaesthesist* 2006;55:451-6.
17. Schwemmer U, Markus CK, Greim CA, Brederlau J, Roewer N. Ultrasound-guided anaesthesia of the axillary brachial plexus: efficacy of multiple injection approach. *Ultraschall Med* 2005;26:114-9.
18. Soeding PE, Sha S, Royse CE, Marks P, Hoy G, Royse AG. A randomized trial of ultrasound-guided brachial plexus anaesthesia in upper limb surgery. *Anaesth Intensive Care* 2005;33:719-25.
19. Kefalianakis F, Spohner F. Ultrasound-guided blockade of axillary plexus brachialis for hand surgery. *Handchir Mikrochir Plast Chir* 2005;37:344-8.
20. Liu FC, Liou JT, Tsai YF, Li AH, Day YY, Hui YL, et al. Efficacy of ultrasound-guided axillary brachial plexus block: a comparative study with nerve stimulator-guided method. *Chang Gung Med J* 2005;28:396-402.
21. Liu FC, Lee LI, Liou JT, Hui YL, Lui PW. Ultrasound-guided axillary brachial plexus block in patients with chronic renal failure: report of sixteen cases. *Chang Gung Med J* 2005;28:180-5.