

Poster presentation

Role of the general base Glu268 in nitroglycerin bioactivation and mechanism-based superoxide formation by aldehyde dehydrogenase-2

M Verena Wenzl*¹, Matteo Beretta¹, Antonius CF Gorren¹, Pravas K Baral², Karl Gruber², Michael Russwurm³, Doris Koesling³, Kurt Schmidt¹ and Bernd Mayer¹

Address: ¹Department of Pharmacology and Toxicology, Karl-Franzens-Universität Graz, A-8010 Graz, Austria, ²Department of Molecular Biosciences, Karl-Franzens-Universität Graz, A-8010 Graz, Austria and ³Department of Pharmacology and Toxicology, Ruhr-Universität Bochum, D-44780 Bochum, Germany

Email: M Verena Wenzl* - michaelaverena.wenzl@uni-graz.at

* Corresponding author

from 4th International Conference of cGMP Generators, Effectors and Therapeutic Implications Regensburg, Germany. 19–21 June 2009

Published: 11 August 2009

BMC Pharmacology 2009, 9(Suppl 1):P72 doi:10.1186/1471-2210-9-S1-P72

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S1/P72>

© 2009 Wenzl et al; licensee BioMed Central Ltd.

Background

Mitochondrial aldehyde dehydrogenase (ALDH2) plays an essential role in nitroglycerin (GTN) bioactivation, resulting in formation of nitric oxide (NO) or a related activator of soluble guanylate cyclase (sGC) and consequently in cGMP-mediated vasorelaxation [1]. ALDH2 denitrates GTN to 1,2-glyceryl dinitrate (1,2-GDN) and nitrite but also catalyzes reduction of GTN to nitric oxide (NO) [2]. To elucidate the mechanism of ALDH2-catalyzed GTN bioactivation in relation to the established ALDH2 activities (dehydrogenase, esterase), we compared the function of the wildtype (WT) enzyme with a mutant lacking the general base Glu268 (E268Q).

Results

Despite low dehydrogenase and esterase activities (<3% of WT) the E268Q mutant exhibited virtually unaffected rates of GTN denitration ($133 \pm 11\%$ of WT). The nucleotide cofactor NAD caused a pronounced increase in the rates of 1,2-GDN formation by WT-ALDH2 from 1.21 ± 0.18 to 8.73 ± 0.09 nmol \times min⁻¹ \times mg⁻¹, but inhibited the reaction catalyzed by the E268Q mutant to about 3% of WT. In contrast to WT-ALDH2, the E268Q mutant generated detectable NO measured with a Clark-type electrode

even in the absence of superoxide dismutase (SOD). The apparent initial rate was 2.1 ± 0.31 nmol \times min⁻¹ \times mg⁻¹ and the peak concentration of NO was 0.17 ± 0.03 μ M. Purified sGC was activated by GTN in the presence of increasing amounts of WT-ALDH2, but the effect reached a plateau of about 30% of maximal sGC activity at 50–100 μ g of ALDH2 (9.0 ± 0.38 μ mol cGMP \times min⁻¹ \times mg⁻¹). Superoxide dismutase markedly potentiated the effect of ALDH2, resulting in maximal sGC activation with 25 μ g of protein. With E268Q-ALDH2, maximal sGC activation was observed with 100 μ g of protein even in the absence of SOD. In the presence of SOD, the effect of the mutant was virtually identical to that of WT-ALDH2. Formation of superoxide was confirmed by determination of hydroethidine oxidation that was inhibited by SOD and the ALDH2 inhibitor chloral hydrate. E268Q-ALDH2 exhibited about 50% lower rates of superoxide formation than the WT enzyme.

Conclusion

Our results suggest that E268 is involved in the structural organization of the NAD binding pocket but is not required for GTN denitration. Mechanism-based superoxide formation by ALDH2 may essentially account for oxi-

ductive stress in GTN-exposed blood vessels contributing to nitrate tolerance.

Acknowledgements

We thank Margit Rehn for excellent technical assistance. This work was funded by the Fonds zur Förderung der Wissenschaftlichen Forschung in Austria (W901 DK Molecular Enzymology and P20669) and the Deutsche Forschungsgemeinschaft (KO1157/4-1).

References

1. Chen Z, Foster MW, Zhang J, Mao L, Rockman HA, Kawamoto T, Kitagawa K, Nakayama KI, Hess DT, Stamler JS: **An essential role for mitochondrial aldehyde dehydrogenase in nitroglycerin bioactivation.** *Proc Natl Acad Sci USA* 2005, **102**:12159-12164.
2. Beretta M, Gruber K, Kollau A, Russwurm M, Koesling D, Goessler W, Keung WM, Schmidt K, Mayer B: **Bioactivation of nitroglycerin by purified mitochondrial and cytosolic aldehyde dehydrogenases.** *J Biol Chem* 2008, **283**:17873-17880.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

