

Administration of 3,5-diiodothyronine (3,5-T2) causes central hypothyroidism and stimulates thyroid-sensitive tissues

Alvaro Souto Padron¹, Ruy Andrade Louzada Neto^{1,2}, Thiago Urgal Pantaleão¹, Maria Carolina de Souza dos Santos¹, Renata Lopes Araujo¹, Bruno Moulin de Andrade¹, Monique da Silva Leandro¹, João Pedro Saar Werneck de Castro², Andrea Claudia Freitas Ferreira¹ and Denise Pires de Carvalho¹

¹Laboratório de Fisiologia Endócrina Doris Rosenthal, Instituto de Biofísica Carlos Chagas Filho and Instituto de Pesquisa Translacional em Saúde e Ambiente na Região Amazônica (INPeTAM), CCS-Bloco G- Cidade Universitária, Ilha do Fundo, Rio de Janeiro 21949-900, Brazil

²Laboratório de Biologia do Exercício, Escola de Educação Física e Desportos, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

Correspondence should be addressed to D P de Carvalho
Email
 dencarv@biof.ufrj.br

Abstract

In general, 3,5-diiodothyronine (3,5-T2) increases the resting metabolic rate and oxygen consumption, exerting short-term beneficial metabolic effects on rats subjected to a high-fat diet. Our aim was to evaluate the effects of chronic 3,5-T2 administration on the hypothalamus–pituitary–thyroid axis, body mass gain, adipose tissue mass, and body oxygen consumption in Wistar rats from 3 to 6 months of age. The rats were treated daily with 3,5-T2 (25, 50, or 75 µg/100 g body weight, s.c.) for 90 days between the ages of 3 and 6 months. The administration of 3,5-T2 suppressed thyroid function, reducing not only thyroid iodide uptake but also thyroperoxidase, NADPH oxidase 4 (NOX4), and thyroid type 1 iodothyronine deiodinase (D1 (DIO1)) activities and expression levels, whereas the expression of the TSH receptor and dual oxidase (DUOX) were increased. Serum TSH, 3,3',5-triiodothyronine, and thyroxine were reduced in a 3,5-T2 dose-dependent manner, whereas oxygen consumption increased in these animals, indicating the direct action of 3,5-T2 on this physiological variable. Type 2 deiodinase activity increased in both the hypothalamus and the pituitary, and D1 activities in the liver and kidney were also increased in groups treated with 3,5-T2. Moreover, after 3 months of 3,5-T2 administration, body mass and retroperitoneal fat pad mass were significantly reduced, whereas the heart rate and mass were unchanged. Thus, 3,5-T2 acts as a direct stimulator of energy expenditure and reduces body mass gain; however, TSH suppression may develop secondary to 3,5-T2 administration.

Key Words

- ▶ 3,5-T2
- ▶ TSH
- ▶ thyroid hormone metabolism
- ▶ deiodinase

Journal of Endocrinology
 (2014) 221, 415–427

Introduction

Compound 3,3',5-triiodothyronine (T₃) exerts many important effects on the basal metabolic rate and increases oxygen consumption. Several years ago, it was shown that

3,5-diiodothyronine (3,5-T2) is responsible for certain non-genomic effects of thyroid hormones on the resting metabolic rate (RMR) and oxygen consumption

(Lanni *et al.* 1996, Moreno *et al.* 2002). Compound 3,5-T2 by itself increases the rate of oxygen consumption in perfused rat liver (Horst *et al.* 1989), blood mononuclear cells (Kvetny 1992), and rat liver mitochondria (Lanni *et al.* 1992, 1993, O'Reilly & Murphy 1992) and affects the activity of enzymes involved in energy metabolism, including α -glycerophosphate dehydrogenase (Lombardi *et al.* 2000), cytochrome oxidase (Lanni *et al.* 1992, 1993, Goglia *et al.* 1994, Arnold *et al.* 1998), malate dehydrogenase (Ball *et al.* 1997, Lombardi *et al.* 2000), glucose-6-phosphate dehydrogenase (Lombardi *et al.* 2000), and FOF1-ATP synthase (Mangiullo *et al.* 2010). More recently, it has been shown that the administration of 3,5-T2 to rats fed on a high-fat diet leads to a liver proteome profile similar to that found in non-steatotic livers, which is likely due to the hypolipidemic effect of this thyroid hormone metabolite (Silvestri *et al.* 2010). Because Wistar rats spontaneously develop obesity as they age (Newby *et al.* 1990), chronic 3,5-T2 administration could be beneficial. In these previous studies, pharmacological doses of 3,5-T2 were administered, as Engler & Burger (1984) reported 3,5-T2 physiological serum concentrations to be around 1 ng/dl in men, with a daily production rate estimated to be around 1 μ g (Engler & Burger 1984, Engler *et al.* 1984).

Apart from these unequivocal effects on energy metabolism, 3,5-T2 might also affect the hypothalamus–pituitary–thyroid axis. Horst *et al.* (1995) have shown that 3,5-T2 treatment reduced serum thyroid-stimulating hormone (TSH) and thyroxine (T_4) levels as well as TSH secretion from rat pituitary fragments, and it was also able to decrease the TSH levels of hypothyroid rats (Moreno *et al.* 1998). Moreover, 3,5-T2 increases pituitary type 1 iodothyronine deiodinase (D1) activity and transiently decreases type 2 iodothyronine deiodinase (D2) activity in rats (Baur *et al.* 1997), while in killifish, short-term exposure to 3,5-T2 was shown to decrease D2 activity and *d2* (*dio2*) mRNA levels in the liver (García-G *et al.* 2007).

Despite the possible interference of 3,5-T2 on thyroid function (Horst *et al.* 1995, Baur *et al.* 1997, Moreno *et al.* 1998, García-G *et al.* 2007), to our knowledge, the chronic effects of 3,5-T2 on proteins involved in thyroid hormone biosynthesis (Kopp 2013) and thyroid hormone metabolism have not been assessed thus far. Therefore, our aim was to study the effect of chronic pharmacological 3,5-T2 administration on these parameters. Our goal is based on the fact that 3,5-T2 has been suggested to be a possible therapeutic tool to overcome liver steatosis and insulin resistance; hence, it is tempting to speculate as to whether its chronic administration could lead to thyroid dysfunction, thus affecting other organs that do not respond to 3,5-T2.

In this study, we describe the effects of chronic 3,5-T2 administration on body mass gain and the hypothalamus–pituitary–thyroid axis during aging in male Wistar rats. We show herein that chronic 3,5-T2 administration impairs aging-related body mass gain while suppressing TSH and normal thyroid function.

Materials and methods

Animals

Adult male Wistar rats were housed at a controlled temperature (23 °C) with daily exposure to a 12 h light:12 h darkness cycle and were provided free access to water and standard rat chow. This investigation conforms to the Guidelines for the Care and Use of Laboratory Animals published by the US National Institutes of Health (NIH Publication No.85-23, revised 1996) and was approved (number: IBCCF 070) by the Institutional Animal Care and Use Committee (Comissão de Ética no Uso de Animais do Centro de Ciências da Saúde, CEUA/CCS). All animals were individually housed for a 1-week acclimation period.

Treatment with 3,5-T2

Three-month-old male Wistar rats were divided into the following groups: control (daily s.c. injections of vehicle were administered), 25 μ g 3,5-T2 (daily s.c. injections of 25 μ g 3,5-T2/100 g body weight (BW)), 50 μ g 3,5-T2 (daily s.c. injections of 50 μ g 3,5-T2/100 g BW), and 75 μ g 3,5-T2 (daily s.c. injections of 75 μ g 3,5-T2/100 g BW). The experiments were conducted for 3 months. Compound 3,5-T2 (Sigma) was dissolved in 0.04 M NaOH and subsequently diluted with saline solution (0.04 M NaOH and saline were the vehicle) at the time of administration (0.09% NaCl, 60% v/v; stock solution of 1 μ g 3,5-T2/ μ l), which maintained equal alkalinity in all of the injected solutions. Animals were weighed at the beginning of the treatment and just before killing; food ingestion was determined for 24 h before killing. After the experimental period, animals were killed by decapitation, and blood was collected. The serum was obtained and stored at -20 °C until specific RIA to determine serum TSH, T_3 , and T_4 could be carried out. Hearts, retroperitoneal fat pads (adipose tissue from the perirenal capsules and attached to the dorsal rat body wall, without the adrenal glands that are embedded in the perirenal adipose tissue), and epididymal fat pads were removed and weighed. Liver, kidneys, thyroid glands, hypothalami, and pituitaries were dissected out

and stored at -70°C until processing for enzymatic measurements or for RNA extraction procedure.

Serum insulin, TSH, T_3 , and T_4

The fasting serum insulin concentrations were measured using a commercial RIA kit for rat insulin (MP Biomedicals, LLC, Santa Clara, CA, USA, sensitivity of $5.5\ \mu\text{IU/ml}$), and inter- and intra-assay coefficient of variation (CV) values were 5.0 and 8.9% respectively.

The serum TSH levels were evaluated using a specific RIA obtained from the National Institute of Diabetes, Digestive and Kidney Diseases (NIDDK; Bethesda, MD, USA) and expressed in terms of the reference preparation 2 (RP2). The intra- and inter-assay CV values were 7.7 and 6.5% respectively and the sensitivity was $0.63\ \text{ng/ml}$.

Total serum T_3 and T_4 concentrations were measured using commercial RIA kits based on the presence of specific antibodies adhered to the internal surface of propylene tubes. T_3 : DLS – 3100 Active, sensitivity of $4.3\ \text{ng/dl}$, inter- and intra-assay CV values varied from 4.2 to 6.0% and from 5 to 6.5% respectively; T_4 : DLS – 3200 Active, sensitivity of $0.4\ \mu\text{g/dl}$, inter- and intra-assay CV values varied from 7.1 to 7.4% and from 2.9 to 5.1% respectively (DSL, TX, USA).

All procedures were carried out following the manufacturer's recommendations.

Glucose tolerance test

A glucose tolerance test was carried out before the commencement of 3,5-T2 treatment and after 1, 2, and 3 months of treatment, as described previously (Yagil *et al.* 2005). Briefly, animals were kept on fasting overnight and blood was obtained from the tip of the tail of fully awake, non-anesthetized animals. Blood glucose levels were determined using a glucose reagent strip and a standard automated glucometer (AccuChek Advantage II, Roche). After the fasting blood glucose level was determined, animals were loaded by gavage with $3.5\ \text{g glucose/kg BW}$, and blood glucose levels were measured 30, 60, 120, and 180 min after loading by gavage. The results are expressed as the area under the glycemic curve.

Because a dose of $50\ \mu\text{g 3,5-T2/100 g BW}$ impaired body weight gain and mildly improved glucose tolerance without significantly affecting heart weight, while significantly reducing serum T_3 , T_4 , and TSH levels, we have chosen this daily dose to study the effect of 3,5-T2 administration on thyroid function, thyroid hormone metabolism, and oxygen consumption.

Resting metabolic rate

We evaluated the RMR of rats that received $50\ \mu\text{g 3,5-T2}$ (daily injections of $50\ \mu\text{g 3,5-T2/100 g BW}$) and control rats, as described previously (Araujo *et al.* 2008). The RMR was measured using open-circuit indirect calorimetry for 24 h, after 90 days of treatment. Rats were individually placed in a respiration chamber ($24 \times 21.5 \times 20\ \text{cm}$); and the airflow was maintained constant at $700\ \text{ml/min}$ by a mass flow controller (LE 400; Panlab, Barcelona, Spain). Oxygen was measured using an O_2 analyzer (LE 405; Panlab). Spontaneous activity was measured in the same metabolic cage that contains four sensors in the housing base to detect the movement of the animal. Oxygen consumption and spontaneous activity were recorded at 5-min intervals, and the results were expressed in $\text{ml/min per kg}^{0.75}$ of O_2 and horizontal activity in counts/min respectively. To avoid disruption caused by adaptation to the chamber, the first 3 h (from 1800 to 2100) were excluded from the analyses.

Electrocardiography

Electrocardiogram recording was carried out in conscious animals by the non-invasive method. Electrodes were positioned in specific positions and connected by flexible cables to a differential AC amplifier (model 1700, A-M Systems, Sequim, WA, USA), with a low pass signal filtered at 500 Hz, and digitalized at 1 kHz by a 16-bit A/D converter (AD Instruments, Colorado Springs, CO, USA) using the Lab chart 7.0 software (AD Instruments).

Type 1 and type 2 iodothyronine deiodinase activities

The $5'$ -deiodinase activities in the liver, kidney, thyroid, pituitary, and hypothalamus were evaluated as described previously (Berry *et al.* 1991, Araujo *et al.* 2008, Fortunato *et al.* 2008). Briefly, $25\ \text{mg}$ tissue samples (liver or kidney) or the whole thyroid, pituitary, and hypothalamus were homogenized in 1 ml of 0.1 M sodium phosphate buffer containing 1 mM EDTA (Merck), 0.25 M sucrose (Merck), and 10 mM dithiothreitol (DTT) (USB Products, Cleveland, OH, US) (pH 6.9). For the measurement of type 1 deiodinase activity, duplicate homogenates ($30\ \mu\text{g}$ protein from the liver or kidney and $15\ \mu\text{g}$ protein from the thyroid) were incubated in a water bath for 1 h at 37°C with $1\ \mu\text{M rT}_3$ (Sigma), equal volumes of $[^{125}\text{I}] \text{rT}_3$ (PerkinElmer Life Sciences, Boston, MA, USA) previously purified using Sephadex LH-20, and 10 mM DTT (USB) in 100 mM potassium phosphate buffer containing 1 mM EDTA (pH 6.9). The reaction volume was $300\ \mu\text{l}$, in the presence

or absence of propylthiouracil (PTU) (1 mM) in order to measure specific D1 activity. For the type 2 deiodinase activity assay, duplicate homogenates (30 µg protein from the hypothalamus and pituitary) were incubated in a water bath for 3 h at 37 °C with 1 nM T₄ (Sigma), equal volumes of [¹²⁵I] T₄ (PerkinElmer Life Sciences) previously purified using Sephadex LH-20, 1 mM PTU, and 20 mM DTT (USB) in 100 mM potassium phosphate buffer containing 1 mM EDTA (pH 6.9). The reaction volume was 300 µl, in the presence or absence of excess of T₄ (100 nM) in order to inhibit D2 in the blank assay tube. For D1, blank incubations were carried out in the absence of protein and the presence of 1 mM PTU. For D2, blank incubations were carried out in the presence of 100 nM T₄.

After incubation, the reaction was terminated at 4 °C followed by the addition of 100 µl fetal bovine serum (Cultilab, Campinas, São Paulo, Brazil) and 200 µl trichloroacetic acid (50%, v/v). Samples were centrifuged at 8000 g for 3 min, and the supernatant was collected for the measurement of ¹²⁵I liberated during the deiodination reaction. The protein concentration in the homogenates was measured by the Bradford method (Bradford 1976) after incubation of the homogenates with NaOH (2.5 M).

Thyroid iodide uptake

We had previously demonstrated that the measurement of radioiodide uptake 15 min after ¹²⁵I-NaI administration (short-term iodide uptake) reflects iodide transport through the sodium-iodide symporter (NIS) without the influence of *in vivo* thyroid iodine organification activity (Ferreira *et al.* 2005). Thus, to evaluate the NIS function *in vivo* using thyroid radioiodide uptake measurements, animals received Na-¹²⁵I (3700 Bq, i.p.; Amersham) 15 min before decapitation. The thyroid glands were removed and weighed. The radioactivity of the thyroid glands was measured using a gamma counter (LKB) and expressed as percentage of total ¹²⁵I injected per milligram of thyroid.

Thyroperoxidase preparation and activity

Thyroperoxidase (TPO) extraction and activity measurement were carried out as described previously (Moura *et al.* 1989, Carvalho *et al.* 1994, Ferreira *et al.* 2005). Rat thyroids were minced and homogenized in 0.5 ml of 50 mM Tris-HCl buffer (pH 7.2) containing 1 mM KI using an Ultra-Turrax Homogenizer (IKA, Staufen, Germany). The homogenate was centrifuged at 100 000 g at 4 °C for 1 h. The pellet was suspended in 0.5 ml Triton (0.1%, v/v)

and incubated at 4 °C for 24 h to solubilize TPO. The triton-treated suspension was centrifuged at 100 000 g at 4 °C for 1 h, and the supernatant containing solubilized TPO was used for the assays. The protein content was determined by the Bradford method (Bradford 1976).

To measure TPO iodide oxidation activity, assay mixtures were prepared containing 1.0 ml freshly prepared 50 mM sodium phosphate buffer (pH 7.4), 24 mM KI, 11 mM glucose, and increasing amounts of solubilized TPO. The final volume was adjusted to 2.0 ml with 50 mM sodium phosphate buffer (pH 7.4), and the reaction was initiated by the addition of 10 µl of 0.1% glucose oxidase (Boehringer Grade I). The increase in absorbance at 353 nm (triiodide production) was recorded for 4 min on a Hitachi spectrophotometer (U-3300). The ΔA353 nm/min was determined from the linear portion of the reaction curve and related to protein concentration. One unit of activity corresponds to ΔA353 nm/min = 1.0.

Thyroid H₂O₂ production

H₂O₂ generation was quantified in thyroid particulate fractions by the Amplex red/HRP assay (Molecular Probes, Invitrogen), which detects the accumulation of a fluorescent oxidized product. For particulate preparation, the excised thyroid glands remained at 4 °C for 24 h in 50 mM sodium phosphate buffer (pH 7.2) containing 0.25 M sucrose, 0.5 mM DTT, 1 mM EGTA, 5 mg/ml aprotinin, and 34.8 mg/ml phenylmethylsulfonyl fluoride (PMSF) before homogenization. Then, the homogenate was centrifuged at 100 000 g for 35 min at 4 °C and suspended in 0.25 ml of 50 mM sodium phosphate buffer (pH 7.2) containing 0.25 M sucrose, 2 mM MgCl₂, 5 mg/ml aprotinin, and 34.8 mg/ml PMSF. This particulate fraction was incubated in 150 mM sodium phosphate buffer (pH 7.4) containing 100 U/ml superoxide dismutase (Sigma), 0.5 U/ml HRP (Roche), 50 µM Amplex red (Molecular Probes, Eugene, OR, USA), 1 mM EGTA, with or without 1.5 mM CaCl₂, and the fluorescence was immediately measured in a microplate reader (Victor X4; PerkinElmer, Norwalk, CT, USA) at 30 °C, using wavelength excitation at 530 nm and emission at 595 nm. The H₂O₂ production was quantified using standard calibration curves.

Immunoblotting of NIS, TSHR, dual oxidase, and NADPH oxidase 4

Thyroids were placed in lysis buffer containing 135 mM NaCl, 1 mM MgCl₂, 2.7 mM KCl, 20 mM Tris (pH 8.0), 1% Triton, 10% glycerol, and protease and phosphatase

inhibitors (0.5 mM Na₃VO₄, 10 mM NaF, 1 M leupeptin, 1 M pepstatin, 1 M okadaic acid, and 0.2 mM PMSF) and subsequently homogenized using an Ultra-Turrax Homogenizer (IKA). The samples were then centrifuged at 570 g for 10 min at 4 °C, and the supernatant was collected. An aliquot was used to determine the protein concentration via the BCA protein assay kit (Pierce, Rockford, IL, USA, catalog no. 23 227), following the manufacturer's instructions. The protein samples were then resolved by SDS-PAGE, transferred onto PVDF membranes, and probed with the indicated antibodies. The NIS antibody was kindly provided by Dr Nancy Carrasco, the dual oxidase (DUOX) antibody was kindly provided by Dr Corinne Dupuy, the anti-NADPH oxidase 4 (NOX4) primary antibody (NB 110-58851) was obtained from Novus Biologicals (Littleton, CO, USA), and TSHR antibody was purchased from Santa Cruz Biotechnology. GAPDH was used as an internal control (the antibody was purchased from Millipore Corporation, Billerica, WA, USA), and an anti-rabbit IgG HRP-linked antibody (Cell Signaling Technology, Boston, MA, USA) was used as a secondary antibody. The immunoblots were developed using ECL.

Real-time PCR

Thyroid and hypothalamic total RNAs were extracted using the RNeasy Plus Mini Kit (Qiagen) following the manufacturer's instructions. After DNase treatment, RT was carried out followed by real-time PCR as described previously (Schmittgen *et al.* 2008). Specific oligonucleotides, as described in Supplementary Table 1, see section on supplementary data given at the end of this article, were purchased from Applied Biosystems. For each gene tested, the amplification of different cDNA quantities was linear in the PCR assay conditions used. In this experiment, β -actin was used as an internal control, and the results are expressed as $\Delta\Delta CT$.

Statistical analyses

The results are expressed as the mean \pm S.E.M. Body, heart, and fat mass data were analyzed by one-way ANOVA followed by Dunnett's multiple comparison test. Serum T₄ and T₃ were analyzed by ANOVA followed by the Bonferroni multiple comparison test. Areas under the glycemic curve data were analyzed by two-way ANOVA




Figure 1

Male Wistar rats were treated (or not treated) with 3,5-diiodothyronine (3,5-T2) at doses of 25, 50, and 75 μ g/100 g BW for 90 days. (A) Body mass during 90 days, $n=14-16$ animals per group; $*P<0.05$ control vs 50 μ g 3,5-T2/100 g BW, $^{#}P<0.05$ control vs 75 μ g 3,5-T2/100 g BW. (B) Serum glucose after 90 days of treatment, five to nine animals per group; $*P<0.05$ control vs other groups. (C) Area under the glycemic curve obtained via glucose tolerance testing at different time points, five to nine animals per

group. The test was carried out at the beginning of the treatment with 3,5-T2 (3 months of age) and 1, 2, and 3 months later. (D) Fasting serum glucose, 12-15 animals per group; $*P<0.05$ control (6 months) vs control (2 months). (E) Fasting serum insulin, six to eight animals per group. (F) Heart rate, 10-11 animals per group. The results are expressed as the mean \pm S.E.M. of the area under the glycemic curve. The data were analyzed by two-way ANOVA followed by the Bonferroni multiple comparison test.

Table 1 Body mass of 3-month-old rats at the beginning of a 3-month treatment with 3,5-T2 (25, 50, and 75 µg 3,5-T2/100 g BW, s.c.) and at the end of the treatment. Body mass gain during this period, heart mass, and relative heart mass at the end of the experiment were also recorded. The data are expressed as the mean \pm s.e.m. and were analyzed by one-way ANOVA followed by Dunnett's multiple comparison test. The study included 14–16 animals per group

Group	Body mass (g) (3 months old)	Body mass (g) (6 months old)	Body mass gain (g)	Heart mass (g)	Relative heart mass (g/100 g BW)
Control	291.6 \pm 5.91	401.0 \pm 11.68	105.7 \pm 6.82	1.15 \pm 0.030	0.29 \pm 0.009
25 µg 3,5-T2	293.3 \pm 5.22	372.6 \pm 7.23*	85.1 \pm 4.99*	1.08 \pm 0.034	0.30 \pm 0.009
50 µg 3,5-T2	294.2 \pm 5.12	369.0 \pm 9.11*	78.8 \pm 6.91*	1.16 \pm 0.031	0.32 \pm 0.010
75 µg 3,5-T2	292.0 \pm 5.59	359.4 \pm 10.87*	77.7 \pm 5.62*	1.17 \pm 0.038	0.33 \pm 0.012*

* $P < 0.05$ vs control group.

followed by the Bonferroni post-test. The serum TSH results were analyzed using the non-parametric Kruskal–Wallis test followed by Dunn's multiple comparison test. The area under the curve data for oxygen consumption, food intake, thyroid iodide uptake, TPO activity, deiodinase activities, NIS, and TSHR protein levels as well as *Nis* (*Slc5a5*), *Tpo*, *D1* (*Dio1*) and *Tshr* mRNA levels were analyzed by the unpaired *t*-test. Statistical analyses were carried out using the GraphPad Prism software (version 4; GraphPad Software, Inc., San Diego, CA, USA). Differences were considered to be statistically significant when $P < 0.05$.

Results

Body, heart, and fat mass

After 3 months of 3,5-T2 administration, the final body mass was significantly reduced when compared with the control group independent of the dose (25, 50, or 75 µg of 3,5-T2/100 g BW). Body mass gain during the treatment was \approx 30% lower in rats treated with 3,5-T2 when compared with those in the control group (Fig. 1A). Absolute heart mass did not differ among the groups, whereas relative heart mass was significantly increased in rats treated with the highest dose (75 µg) of

3,5-T2/100 g BW, which was mainly due to decreased body mass rather than increased heart mass (Table 1). Heart rate did not differ among the groups (Fig. 1F), as described previously (Lanni *et al.* 2005). Moreover, although retroperitoneal fat pad mass was significantly reduced in rats treated with 3,5-T2 independent of the 3,5-T2 dose used, the epididymal fat mass remained unchanged (Table 2).

Glucose tolerance test

Control male Wistar rats developed impaired glucose tolerance at 6 months of age (Fig. 1B), and basal fasting serum glucose was significantly higher in the 6-month-old animals when compared with 3,5-T2-treated 6-month-old rats and control 2-month-old rats (Fig. 1D). Treatment with 3,5-T2 improved glucose tolerance at all doses (as shown in Fig. 1B). Animals treated with 3,5-T2 for 3 months have areas under the glycemic curve that are \sim 10% lower than their paired controls (6 months old) (Fig. 1C). The change in the glucose tolerance noted in control aged rats may be related to the higher body mass observed in this group, which was significantly prevented by 3,5-T2 administration to rats. It is noteworthy that serum fasting insulin levels did not differ among the groups (Fig. 1E).

Table 2 Retroperitoneal and epididymal fat pad mass of 6-month-old rats treated with different doses of 3,5-T2 for 3 months (25, 50, and 75 µg 3,5-T2/100 g BW, s.c.). The data are expressed as the mean \pm s.e.m. and were analyzed by one-way ANOVA followed by Dunnett's multiple comparison tests. The study included 14–15 animals per group

Group	Retroperitoneal fat mass (g)	Retroperitoneal fat (g/100 g BW)	Epididymal fat mass (g)	Epididymal fat (g/100 g BW)
Control	7.64 \pm 0.88	1.92 \pm 0.84	7.53 \pm 0.67	1.89 \pm 0.58
25 µg 3,5-T2	5.05 \pm 0.46*	1.52 \pm 0.40	6.12 \pm 0.58	1.68 \pm 0.48
50 µg 3,5-T2	5.67 \pm 0.51*	1.59 \pm 0.42	6.10 \pm 0.53	1.64 \pm 0.46
75 µg 3,5-T2	5.48 \pm 0.58*	1.60 \pm 0.44	6.45 \pm 0.54	1.72 \pm 0.56

* $P < 0.05$ vs control.

Serum TSH, T₃, and T₄

Serum total T₃ and T₄ levels were reduced in rats treated with 3,5-T2, and the reduction was dose dependent (Fig. 2A and B). Despite the decreased serum thyroid




Figure 2

Serum total T₄, T₃, and TSH concentrations in rats treated (or not treated) with 3,5-diiodothyronine (3,5-T2) for 90 days at doses of 25, 50, and 75 µg/100 g BW. Hormone concentrations were determined using specific RIA. (A) Serum total T₄ (13–18 rats per group); (B) serum total T₃ (13–18 rats per group); and (C) serum TSH (11–14 rats per group). The results are expressed as the mean \pm s.e.m. Serum total T₄ and T₃ were analyzed by one-way ANOVA followed by the Bonferroni multiple comparison test, and serum TSH was analyzed by the non-parametric Kruskal–Wallis test followed by Dunn's multiple comparison test. * $P < 0.01$ vs control group; ** $P < 0.001$ vs control group.

hormones, TSH levels remained normal in rats receiving 25 µg of 3,5-T2/100 g BW and were significantly lower in rats that received 50 and 75 µg of 3,5-T2/100 g BW (Fig. 2C), demonstrating that 3,5-T2 by itself might be exerting the negative feedback, as suggested previously (Horst *et al.* 1995, Moreno *et al.* 1998).

Resting metabolic rate

Figure 3A summarizes the oxygen consumption results collected over a 21-h period of time, from 1800 1 day to 2200 the following day. The results are also expressed as the area under the curve of oxygen consumption for 21 h (Fig. 3B). Although 3,5-T2-treated rats had significantly decreased levels of serum T₃ (Fig. 2B), which is the main known regulator of the RMR, oxygen consumption was increased in these animals. Food intake did not significantly differ between the groups (Fig. 3C). Our results reinforce previous data showing that 3,5-T2 can increase the RMR in rats.

Type 1 iodothyronine deiodinase activity

Liver and kidney D1 activities were significantly higher in rats treated with 3,5-T2 (Fig. 4A and B), even in the presence of significantly lower serum T₃, which is the main stimulator of hepatic D1 expression. This result





Figure 3

Resting metabolic rate (RMR) and food intake of control rats and rats treated with 3,5-diiodothyronine (3,5-T2) for 90 days at a dose of 50 µg/100 g BW. The RMR was measured using open-circuit indirect calorimetry for 21 h after 90 days of treatment (five rats per group). (A) Oxygen consumption; (B) area under the curve (AUC) of oxygen consumption (five rats per group); (C) 24 h food intake (nine rats per group); and (D) spontaneous daily activity as counts/min (eight rats per group). The results are expressed as the mean \pm s.e.m. The AUC of oxygen consumption and 24 h food intake were analyzed by the unpaired *t*-test. ** $P < 0.001$ vs control group.

**Figure 4**

Type 1 (D1) and type 2 (D2) iodothyronine deiodinase activities in control rats and rats treated with 3,5-diiodothyronine (3,5-T2) for 90 days at a dose of 50 $\mu\text{g}/100$ g BW. (A) Hepatic D1 (control, $n=12$; 3,5-T2, $n=17$); (B) kidney D1 (control, $n=8$; 3,5-T2, $n=12$); (C) hypothalamic D2 (control, $n=8$; 3,5-T2, $n=11$); and (D) pituitary D2 (control, $n=15$; 3,5-T2, $n=20$) activities. The results are expressed as the mean \pm s.e.m. The data were analyzed by the unpaired t -test. * $P<0.05$ vs control group; ** $P<0.005$ vs control group; *** $P<0.0005$ vs control group.

indicates that 3,5-T2 has a direct stimulatory effect on both kidney and hepatic D1 activities, again simulating the well-described genomic action of T_3 (Araujo *et al.* 2008). These results, together with the data on TSH regulation, strongly support the hypothesis that 3,5-T2 functions as an agonist of the β isoform of thyroid hormone receptor (TR) *in vivo*.

Type 2 iodothyronine deiodinase activity

Treatment with 3,5-T2 is associated with a significant increase in both hypothalamic and pituitary D2 activities (Fig. 4C and D). Presumably, the increase in D2 could be related to the decreased serum T_4 levels and not to 3,5-T2 treatment itself. It is tempting to speculate that the increased hypothalamic and pituitary D2 activities might lead to an increase in local T_3 production and thus could result in inappropriately normal or even decreased TSH in the presence of low serum T_4 .


NIS, TPO, DUOX1, DUOX2, NOX4, D1, TSHR, and TRH expressions

As shown in Fig. 5A, B and C, 3,5-T2 administration significantly reduced thyroid iodide uptake, TPO iodide oxidation activity, and thyroid D1 activity together with the significantly lower *Nis*, *Tpo*, and *D1* mRNA levels


(Fig. 5B, D and F). These results are in accordance with the lower serum thyroid hormone levels and might be secondary to the decreased serum TSH levels. To evaluate whether the reduction in NIS function in 3,5-T2-treated rats was due to decreased NIS protein expression, we evaluated the thyroid NIS content. As shown in Fig. 6A, the NIS protein content was greatly reduced by 3,5-T2 treatment, in accordance with the decreased thyroid iodide uptake and reduced serum TSH.

As TSH is the main regulator of thyroid function, we also evaluated the effect of 3,5-T2 on TSH receptor (TSHR) protein levels. The protein level of both α - and β -TSHR subunits (Fig. 6B) and the *Tshr* mRNA content (Fig. 6C) were significantly higher in the thyroids of 3,5-T2-treated rats, which might at least partially counterbalance the reduced serum TSH levels.

Hypothalamic *Trh* mRNA content did not significantly differ between the groups ($C=1.00\pm 0.14$; 3,5-T2 = 0.80 ± 0.05 , $n=6$).

**Figure 5**

Thyroid iodide uptake, *Nis* mRNA levels, thyroperoxidase (TPO) activity, *Tpo* mRNA levels, thyroid type 1 iodothyronine deiodinase (D1) activity, and thyroid *D1* mRNA levels in control rats and rats treated with 3,5-diiodothyronine (3,5-T2) for 90 days at a dose of 50 $\mu\text{g}/100$ g BW. (A) *In vivo* thyroid iodide uptake (control, $n=11$; 3,5-T2, $n=11$); (B) *Nis* mRNA levels (control, $n=5$; 3,5-T2, $n=6$); (C) *in vitro* TPO iodide oxidation activity (control, $n=12$; 3,5-T2, $n=10$); (D) *Tpo* mRNA levels (control, $n=5$; 3,5-T2, $n=5$); (E) *in vitro* thyroid D1 activity (control, $n=7$; 3,5-T2, $n=10$); and (F) thyroid *D1* mRNA levels (control, $n=5$; 3,5-T2, $n=6$). The results are expressed as the mean \pm s.e.m. The data were analyzed by the unpaired t -test. * $P<0.05$ vs control group; *** $P<0.0005$ vs control group.

**Figure 6**


NIS and TSH receptor (TSHR) protein and *Tshr* mRNA levels in control rats and rats treated with 3,5-diiodothyronine (3,5-T2) for 90 days at a dose of 50 µg/100 g BW. Thyroid lysates were prepared, and western blot analysis was carried out with 30 µg protein using anti-NIS (control, $n=11$; 3,5-T2, $n=11$) (A) and anti-TSHR (control, $n=10$; 3,5-T2, $n=10$) (B) antibodies. GAPDH was used as an internal control. The NIS antibody was diluted 1:1000, the TSHR antibody was diluted 1:500, the GAPDH antibody was diluted 1:4000, and the secondary antibody was diluted 1:1500. (C) Thyroid *Tshr* mRNA levels (control, $n=5$; 3,5-T2, $n=4$). Molecular weights of the proteins: NIS corresponds to 80 kDa, α -TSHR to 62 kDa, β -TSHR to 42 kDa, and GAPDH to 36 kDa. The results are expressed as the mean \pm S.E.M. The data were analyzed by the unpaired *t*-test. * $P<0.05$ vs control group; *** $P<0.001$ vs control group.

In addition, NOX4 that is positively regulated by TSH in the thyroid (Weyemi *et al.* 2010) was significantly decreased, as well as the calcium-independent H_2O_2 generation (Fig. 7B, C and D). By contrast, DUOX calcium-dependent H_2O_2 generation increased due to higher DUOX2 expression (Fig. 7A, C and D).

Discussion

Recent data have demonstrated the beneficial effects of 3,5-T2 in preventing liver steatosis in animals fed on a high-fat diet (Silvestri *et al.* 2010), and these authors claimed that 3,5-T2 exerts non-genomic effects. These interesting results prompted us to evaluate whether the chronic administration of 3,5-T2 to aging Wistar rats could prevent the natural development of overweight that occurs in this rat strain. We were also interested in its possible undesirable effects on the pituitary–thyroid axis.

We show herein that daily 3,5-T2 administration to rats for a period of 90 days leads to a reduction in body mass gain and retroperitoneal fat mass, accompanied by a higher RMR despite the lower serum T_4 and T_3 levels. Regarding body composition, we did not analyze whether the treated animals lost lean body mass; however, because the metabolic rate is corrected by weight, a preservation of the lean body mass could explain the increase in the metabolic rate. Decreased body mass gain in 3,5-T2-treated rats was not dose dependent, suggesting that a maximal effect was already obtained with 25 µg of 3,5-T2/100 g BW; however, it is possible that the decrease in serum T_4 and T_3 levels might in some way compensate for the increase in the 3,5-T2 dose. Horst *et al.* (1995) did not observe any changes in body mass gain in rats treated for 3 months with the same dose of 3,5-T2. A possible explanation for the discrepancy between these results and ours is that these authors administered 3,5-T2 in the

**Figure 7**

Thyroid calcium and NADPH-dependent hydrogen peroxide generation and DUOX and NOX4 protein levels in control rats and rats treated with 3,5-diiodothyronine (3,5-T2) for 90 days at a dose of 50 µg/100 g BW. (A) Calcium-dependent and (B) thyroid NADPH-dependent H_2O_2 -generating activity was measured by the Amplex red method (control, $n=8$; 3,5-T2, $n=8$); (C) thyroid lysates were prepared, and western blot analysis was carried out with 30 µg protein using anti-DUOX (control, $n=4$; 3,5-T2, $n=4$) and anti-NOX4 (control, $n=4$; 3,5-T2, $n=4$) antibodies. GAPDH was used as an internal control. DUOX antibody was diluted 1:2000, NOX4 antibody was diluted 1:1000, β -actin antibody was diluted 1:5000, and the secondary antibody was diluted 1:4000. (D) Thyroid DUOX1, DUOX2 and NOX4 mRNA levels (control $n=5$; 3,5-T2 $n=4$). The results are expressed as mean \pm S.E.M. The data were analyzed by the unpaired *t*-test. * $P<0.05$ vs control group; ** $P<0.005$ vs control group.

drinking water, which could reduce the bioavailability of the drug, whereas in this study, 3,5-T2 was injected subcutaneously. On the other hand, Lanni *et al.* (2005) showed that 25 µg of 3,5-T2/100 g BW were indeed able to reduce the body mass gain and adiposity of rats, whereas no changes in serum free T₄ were detected in animals subjected to a high-fat diet and treated with 3,5-T2. It is important to note that in this previous report, 3,5-T2 was not administered to control animals (Lanni *et al.* 2005). Additionally, Lombardi *et al.* (2009) demonstrated that 3,5-T2 rapidly enhances the fatty acid oxidation rate in skeletal muscle. Apart from the chronic effects of 3,5-T2, acute and most likely non-genomic actions have also been described, such as increased rat liver mitochondrial respiratory rates (Lanni *et al.* 1993), hepatic oxygen consumption (Horst *et al.* 1995), respiration rate in the liver of rats fed on a high-fat diet (Mollica *et al.* 2009), and energy expenditure restoration in hypothyroid rats (Cimmino *et al.* 1996). Thus, these previously described metabolic effects of 3,5-T2 might explain the reduction in body mass gain and retroperitoneal adiposity observed in aged rats treated with 3,5-T2. The epididymal fat pad is important for reproductive function and has been shown to be necessary for the maintenance of spermatogenesis (Chu *et al.* 2010). Furthermore, the retroperitoneal fat pad appears to be more sensitive to thyroid hormone action than epididymal fat, as previously reported in hyperthyroid rats (Blennemann *et al.* 1992); therefore, it is unsurprising that 3,5-T2 treatment did not significantly affect epididymal fat mass in this study.

In addition to the beneficial effects of 3,5-T2 on body mass gain and adiposity, rats treated with 3,5-T2 showed an improvement in the glucose tolerance test by more than 10% when compared with control animals. This result suggests that 3,5-T2 mildly improves glucose tolerance, either directly or due to the decreased adiposity observed in the treated animals. To our knowledge, this is the first report describing the beneficial effect of chronic 3,5-T2 treatment on *in vivo* glucose homeostasis in 6-month-old Wistar rats. These results are in accordance with the finding that 3,5-T2 prevents insulin resistance in the skeletal muscle of rats fed on a high-fat diet and increases GLUT4 (SLC2A4) protein expression and insulin-induced Akt phosphorylation (Moreno *et al.* 2011). In addition to its other effects on the liver, 3,5-T2 also increases nuclear sirtuin 1 and downregulates lipogenic genes (De Lange *et al.* 2011).

In accordance with previous data showing that 3,5-T2 exerts important metabolic effects (Lanni *et al.* 1996, Lombardi *et al.* 2007, 2009), we have found a significant

increase in the oxygen consumption of rats treated with 50 µg of 3,5-T2/100 g BW when compared with control animals. It is important to note that these animals had significantly lower serum T₃ and T₄ levels, which are the main regulators of energy metabolism. It has indeed been shown that 3,5-T2 acutely enhances mitochondrial fatty acid oxidation rates and thermogenesis in rat skeletal muscle (Lombardi *et al.* 2009), increases hepatic lipid β-oxidation (Lanni *et al.* 2005, Lombardi *et al.* 2009), and increases mitochondrial oxygen consumption (Mollica *et al.* 2009) in addition to increasing resting metabolism (Moreno *et al.* 2002). Thus, it is tempting to speculate that 3,5-T2 by itself functions as a chronic stimulator of the RMR, increasing oxygen consumption.

Regardless of the reduced serum T₄ and T₃ levels, TSH was lower in 3,5-T2-treated animals. However, Antonelli *et al.* (2011) reported no changes in serum thyroid hormone levels in two euthyroid subjects treated for 3 weeks with a daily dose of 300 µg of 3,5-T2 (Antonelli *et al.* 2011). Horst *et al.* (1995) showed that 3,5-T2 reduces TSH secretion by rat pituitary fragments stimulated by TRH. These authors also found a reduction in serum T₄ levels after 90 days of treatment with 25 µg of 3,5-T2/100 g BW, which is in accordance with the results obtained herein. Moreover, a single dose of 3,5-T2 was shown to reduce serum *Tsh* and pituitary β-TSH (*Tshb*) mRNA levels (Baur *et al.* 1997). As reviewed by Williams & Bassett (2011), TRβ is the main mediator of thyroid hormone negative feedback. The findings of Ball *et al.* (1997), showing that TRβ2 binds to 3,5-T2 with greater affinity than the other TR isoforms, could well explain the effectiveness of 3,5-T2 for suppressing TSH secretion. Recently, it has also been shown that 3,5-T2 binds to and activates the human THRB isoform of TRs, again showing that this metabolite exerts genomic effects (Mendoza *et al.* 2013). Based on our results, we cannot explain as to why serum thyroid hormone levels were already decreased at a dose of 25 µg 3,5-T2/100 g BW, while serum TSH levels remained in the normal range. It is possible that despite the normal TSH concentration, the hormone could be less bioactive, which could lead to less stimulation of the thyroid and the consequent reduction in T₃ and T₄ production, although the hypothalamic content of *Trh* mRNA did not change in 3,5-T2-treated rats.

Hepatic and kidney D1 activities were significantly increased in rats treated with 3,5-T2 despite significantly reduced serum T₃ levels. As T₃ is the main stimulator of hepatic and kidney D1 activities (Koenig 2005, Gereben *et al.* 2008) through genomic actions, our results reinforce the idea of a T₃-like genomic effect of 3,5-T2 on

D1 regulation (rather than a non-genomic action). The regulation of the *D1* gene by T_3 is mediated by the TR β isoform because T_3 induction of hepatic and renal D1 is severely weakened in *Thrb1* (*Thrb*)-null mice but is normal in *Thra1* (*Thra*)-null mice (Koenig 2005). Thus, 3,5-T2 might activate TR β , leading to increased D1 activity in the liver and kidney once type 1 deiodinase is regulated at the transcriptional level. In accordance with our data, Baur *et al.* (1997) showed that 3,5-T2 stimulates D1 activity in rat anterior pituitaries *in vivo*. By contrast, in the thyroid gland, D1 activity is under the positive control of TSH through the cAMP pathway (St Germain & Galton 1997). As a result, the decreased serum TSH levels in rats receiving 3,5-T2 explain the reduction in their thyroid D1.

On the other hand, D2 activity was increased in both the hypothalamus and pituitary of rats treated with 3,5-T2, despite the thyromimetic effects of 3,5-T2. It has been shown that D2 protein levels and activity are regulated by post-translational mechanisms, especially ubiquitination (Gereben *et al.* 2000), which appears to be induced mainly by T_4 . As T_4 levels are reduced in 3,5-T2-treated rats, this could explain the increase in D2 activity in these animals.

Taken together, our results show that 3,5-T2 significantly downregulates thyroid function. In addition to the decreased levels of thyroid hormones, we have also observed a reduction in the thyroid D1, NIS, and TPO activities as well as their respective mRNA levels. Iodide uptake is a fundamental step in thyroid hormone biosynthesis, and TPO is a key enzyme in thyroid hormone biosynthesis (Dai *et al.* 1996, Carvalho & Ferreira 2007); therefore, the decreased thyroid hormone levels appear to be secondary to the decreased NIS function and TPO activity. The inhibitory effect of 3,5-T2 on thyroid function is at least in part due to the decreased serum TSH levels because TSH is the main stimulator of the thyroid gland. TSHR protein and *Tshr* mRNA levels were increased in the thyroids of 3,5-T2-treated rats. It has been shown that TSH exerts inhibitory effects on *Tshr* promoter activity (Ikuyama *et al.* 1997). Thus, it is conceivable that thyroid TSHR expression was higher in rats treated with 3,5-T2 because the serum TSH levels were significantly reduced. In addition, NOX4 expression is downregulated, while DUOX2 is upregulated, just as previously demonstrated in other models of decreased TSH action on thyrocytes (Milenkovic *et al.* 2007, Weyemi *et al.* 2010).

In conclusion, chronic 3,5-T2 administration reduced body mass gain and retroperitoneal fat mass and increased the RMR regardless of decreased thyroid hormone levels. The reduction in thyroid hormone levels might be

secondary to the decreased serum TSH levels, leading to the reduced activity and expression of NIS, thyroid D1, and TPO. These new data support the idea that exogenous 3,5-T2 causes TSH suppression. Thus, future studies on the possible deleterious effects of hypothyroidism in tissues where 3,5-T2 might not act as a thyromimetic agent are of great importance before this thyroid metabolite can be used as a pharmacological agent.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/JOE-13-0502>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by grants from Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), Programa de Apoio a Núcleos de Excelência (PRONEX/FAPERJ), Instituto de Pesquisa Translacional em Saúde e Ambiente na Região Amazônica (INPeTAm), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). R L A, B M A, and M S L were recipients of CNPq fellowships. Wagner Nunes Bezerra received a fellowship from FAPERJ.

Acknowledgements

The authors are grateful for the technical assistance of Norma Lima de Araújo Faria, Advaldo Nunes Bezerra, Wagner Nunes Bezerra, Walter Nunes Bezerra, and José Humberto Tavares de Abreu.

References

- Antonelli A, Fallahi P, Ferrari SM, Di Domenicantonio A, Moreno M, Lanni A & Goglia F 2011 3,5-diiodo-L-thyronine increases resting metabolic rate and reduces body weight without undesirable side effects. *Journal of Biological Regulators and Homeostatic Agents* **25** 655–660.
- Araujo RL, Andrade BM, Figueiredo AS, Silva ML, Marassi MP, Pereira VS, Bouskela E & Carvalho DP 2008 Low replacement doses of thyroxine during food restriction restores type 1 deiodinase activity in rats and promotes body protein loss. *Journal of Endocrinology* **198** 119–125. (doi:10.1677/JOE-08-0125)
- Arnold S, Goglia F & Kadenbach B 1998 3,5-Diiodothyronine binds to subunit Va of cytochrome-c oxidase and abolishes the allosteric inhibition of respiration by ATP. *European Journal of Biochemistry* **252** 325–330. (doi:10.1046/j.1432-1327.1998.2520325.x)
- Ball SG, Sokolov J & Chin WW 1997 3,5-Diiodo-L-thyronine (T2) has selective thyromimetic effects *in vivo* and *in vitro*. *Journal of Molecular Endocrinology* **19** 137–147. (doi:10.1677/jme.0.0190137)
- Baur A, Bauer K, Jarry H & Köhrle J 1997 3,5-diiodo-L-thyronine stimulates type 1 5' deiodinase activity in rat anterior pituitaries *in vivo* and in reaggregate cultures and GH3 cells *in vitro*. *Endocrinology* **138** 3242–3248. (doi:10.1210/endo.138.8.5333)

- Berry MJ, Kieffer JD, Harnet JW & Larsen PR 1991 Selenocysteine confers the biochemical properties characteristic of the type I iodothyronine deiodinase. *Journal of Biological Chemistry* **266** 14155–14158.
- Blennemann B, Moon YK & Freaque HC 1992 Tissue-specific regulation of fatty acid synthesis by thyroid hormone. *Endocrinology* **130** 637–643. (doi:10.1210/endo.130.2.1733712)
- Bradford MM 1976 A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein dye binding. *Analytical Biochemistry* **72** 248–254. (doi:10.1016/0003-2697(76)90527-3)
- Carvalho DP & Ferreira AC 2007 The importance of sodium/iodide symporter (NIS) for thyroid cancer management. *Arquivos Brasileiros de Endocrinologia e Metabologia* **51** 672–682. (doi:10.1590/S0004-27302007000500004)
- Carvalho DP, Rego KGM & Rosenthal D 1994 Thyroid peroxidase in dys-hormonogenetic goiters with organification and thyroglobulin defects. *Thyroid* **4** 421–426. (doi:10.1089/thy.1994.4.421)
- Chu Y, Huddleston GG, Clancy AN, Harris RB & Bartness TJ 2010 Epididymal fat is necessary for spermatogenesis, but not testosterone production or copulatory behavior. *Endocrinology* **151** 5669–5679. (doi:10.1210/en.2010-0772)
- Cimmino M, Mion F, Goglia F, Minaire Y & Gélœn A 1996 Demonstration of *in vivo* metabolic effects of 3,5-di-iodothyronine. *Journal of Endocrinology* **149** 319–325. (doi:10.1677/joe.0.1490319)
- Dai G, Levy O & Carrasco N 1996 Cloning and characterization of the thyroid iodide transporter. *Nature* **379** 485–460. (doi:10.1038/379458a0)
- Engler D & Burger AG 1984 The deiodination of the iodothyronines and of their derivatives in man. *Endocrine Reviews* **5** 151–184. (doi:10.1210/edrv-5-2-151)
- Engler D, Merkelbach U, Steiger G & Burger AG 1984 The monodeiodination of triiodothyronine and reverse triiodothyronine in man: a quantitative evaluation of the pathway by the use of turnover rate techniques. *Journal of Clinical Endocrinology and Metabolism* **58** 49–61. (doi:10.1210/jcem-58-1-49)
- Ferreira ACF, Lima LP, Araújo RL, Müller G, Rocha RP, Rosenthal D & Carvalho DP 2005 Rapid regulation of the thyroid sodium-iodide symporter (NIS) activity by thyrotropin, iodine and fasting. *Journal of Endocrinology* **184** 69–76. (doi:10.1677/joe.1.05643)
- Fortunato RS, Ignácio DL, Padron AS, Peçanha R, Marassi MP, Rosenthal D, Werneck-de-Castro JPS & Carvalho DP 2008 The effect of acute exercise session on thyroid hormone economy in rats. *Journal of Endocrinology* **198** 347–353. (doi:10.1677/JOE-08-0174)
- García-G C, López-Bojorquez L, Nuñez J, Valverde-R C & Orozco A 2007 3,5-Diiodothyronine *in vivo* maintains euthyroidal expression of type 2 iodothyronine deiodinase, growth hormone, and thyroid hormone receptor β 1 in the killifish. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **293** R877–R883. (doi:10.1152/ajpregu.00101.2007)
- Gereben B, Gonçalves C, Harnet JW, Larsen PR & Bianco AC 2000 Selective proteolysis of human type 2 deiodinase: a novel ubiquitin-proteasomal mediated mechanism for regulation of hormone activation. *Molecular Endocrinology* **14** 1697–1708. (doi:10.1210/mend.14.11.0558)
- Gereben B, Zavacki AM, Ribich S, Kim BW, Huang SA, Simonides WS, Zeöld A & Bianco AC 2008 Cellular and molecular basis of deiodinase-regulated thyroid hormone signaling. *Endocrine Reviews* **29** 898–938. (doi:10.1210/er.2008-0019)
- Goglia F, Lanni A, Barth J & Kadenbach B 1994 Interaction of diiodothyronines with isolated cytochrome *c* oxidase. *FEBS Letters* **346** 295–298. (doi:10.1016/0014-5793(94)00476-5)
- Horst C, Rokos H & Seitz HJ 1989 Rapid stimulation of hepatic oxygen consumption by 3,5-di-iodo-L-thyronine. *Biochemical Journal* **261** 945–950.
- Horst C, Harnet A, Seitz HJ & Rokos H 1995 3,5-Di-iodo-L-thyronine suppresses TSH in rats *in vivo* and in rat pituitary fragments *in vitro*. *Journal of Endocrinology* **145** 291–297. (doi:10.1677/joe.0.1450291)
- Ikuyama S, Ohe K, Takayanagi R, Kohn LD & Nawata H 1997 Cloning and characterization of the 4.2 kb region of the rat thyrotropin receptor promoter. *Endocrine Journal* **44** 247–256. (doi:10.1507/endocrj.44.247)
- Koenig RJ 2005 Regulation of type 1 iodothyronine deiodinase in health and disease. *Thyroid* **15** 835–840. (doi:10.1089/thy.2005.15.835)
- Kopp P 2013 Thyroid hormone synthesis. In *Werner and Ingbar's the Thyroid: a Fundamental and Clinical Text*, 10th edn, pp 48–74. Eds LE Braverman & D Cooper 10th edn. Philadelphia: Lippincott Williams and Wilkins.
- Kvetny J 1992 3,5-T₂ stimulates oxygen consumption, but not glucose uptake in human mononuclear blood cells. *Hormone and Metabolic Research* **24** 322–325. (doi:10.1055/s-2007-1003323)
- de Lange P, Cioffi F, Senese R, Moreno M, Lombardi A, Silvestri E, De Matteis R, Lionetti L, Mollica MP, Goglia F *et al.* 2011 Nonthyrotoxic prevention of diet-induced insulin resistance by 3,5-diiodo-L-thyronine in rats. *Diabetes* **60** 2730–2739. (doi:10.2337/db11-0207)
- Lanni A, Moreno M, Cioffi M & Goglia F 1992 Effect of 3,3'-diiodo-thyronine and 3,5-diiodothyronine on rat liver oxidative capacity. *Molecular and Cellular Endocrinology* **86** 143–148. (doi:10.1016/0303-7207(92)90138-V)
- Lanni A, Moreno M, Cioffi M & Goglia F 1993 Effect of 3,3'-di-iodothyronine and 3,5-di-iodothyronine on rat liver mitochondria. *Journal of Endocrinology* **136** 59–64. (doi:10.1677/joe.0.1360059)
- Lanni A, Moreno M, Lombardi A & Goglia F 1996 Calorigenic effect of diiodothyronines in the rat. *Journal of Physiology* **494** 831–837.
- Lanni A, Moreno M, Lombardi A, de Lange P, Silvestri E, Ragni M, Farina P, Baccari GC, Fallahi P, Antonelli A *et al.* 2005 3,5-diiodo-L-thyronine powerfully reduces adiposity in rats by increasing the burning of fats. *FASEB Journal* **19** 1552–1554. (doi:10.1096/fj.05-3977fje)
- Lombardi A, Beneduce L, Moreno M, Diano S, Colantuoni V, Ursini MV, Lanni A & Goglia F 2000 3,5-diiodo-L-thyronine regulates glucose-6-phosphate dehydrogenase activity in the rat. *Endocrinology* **141** 1729–1734. (doi:10.1210/endo.141.5.7449)
- Lombardi A, Lanni A, de Lange P, Silvestri E, Grasso P, Senese R, Goglia F & Moreno M 2007 Acute administration of 3,5-diiodo-L-thyronine to hypothyroid rats affects bioenergetic parameters in rat skeletal muscle mitochondria. *FEBS Letters* **581** 5911–5916. (doi:10.1016/j.febslet.2007.11.073)
- Lombardi A, de Lange P, Silvestri E, Busiello RA, Lanni A, Goglia F & Moreno M 2009 3,5-Diiodo-L-thyronine rapidly enhances mitochondrial fatty acid oxidation rate and thermogenesis in rat skeletal muscle: AMP-activated protein kinase involvement. *American Journal of Physiology. Endocrinology and Metabolism* **296** E497–E502. (doi:10.1152/ajpendo.90642.2008)
- Mangiullo R, Gnoni A, Damiano F, Siculella L, Zanotti F, Papa S & Gnoni GV 2010 3,5-diiodo-L-thyronine upregulates rat-liver mitochondrial F(o)F(1)-ATP synthase by GA-binding protein/nuclear respiratory factor-2. *Biochimica et Biophysica Acta* **1797** 233–240. (doi:10.1016/j.bbabi.2009.10.009)
- Mendoza A, Navarrete-Ramírez P, Hernández-Puga G, Villalobos P, Holzer G, Renaud JP, Laudet V & Orozco A 2013 3,5-T₂ is an alternative ligand for the thyroid hormone receptor β 1. *Endocrinology* **154** 2948–2958. (doi:10.1210/en.2013-1030)
- Milenkovic M, De Deken X, Jin L, De Felice M, Di Lauro R, Dumont JE, Corvilain B & Miot F 2007 Duox expression and related H₂O₂ measurement in mouse thyroid: onset in embryonic development and regulation by TSH in adult. *Journal of Endocrinology* **192** 615–626. (doi:10.1677/JOE-06-0003)
- Mollica MP, Lionetti L, Moreno M, Lombardi A, De Lange P, Antonelli A, Lanni A, Cavaliere G, Barletta A & Goglia F 2009 3,5-diiodo-L-thyronine, by modulating mitochondrial functions, reverses hepatic fat accumulation in rats fed a high-fat diet. *Journal of Hepatology* **51** 363–370. (doi:10.1016/j.jhep.2009.03.023)
- Moreno M, Lombardi A, Lombardi P, Goglia F & Lanni A 1998 Effect of 3,5-diiodo-L-thyronine on thyroid stimulating hormone and growth hormone serum levels in hypothyroid rats. *Life Sciences* **62** 2369–2377. (doi:10.1016/S0024-3205(98)00219-7)

- Moreno M, Lombardi A, Beneduce L, Silvestri E, Pinna G, Goglia F & Lanni A 2002 Are the effects of T₃ on resting metabolic rate in euthyroid rats entirely caused by T₃ itself? *Endocrinology* **143** 504–510. (doi:10.1210/endo.143.2.8613)
- Moreno M, Silvestri E, De Matteis R, de Lange P, Lombardi A, Glinni D, Senese R, Cioffi F, Salzano AM, Scaloni A *et al.* 2011 3,5-Diiodo-L-thyronine prevents high-fat-diet-induced insulin resistance in rat skeletal muscle through metabolic and structural adaptations. *FASEB Journal* **25** 3312–3324. (doi:10.1096/fj.11-181982)
- Moura EG, Rosenthal D & Carvalho-Guimarães DP 1989 Thyroid peroxidase activity in human nodular goiters. *Brazilian Journal of Medical and Biological Research* **22** 31–39.
- Newby FD, DiGirolamo M, Cotsonis GA & Kutner MH 1990 Model of spontaneous obesity in aging male Wistar rats. *American Journal of Physiology* **259** R1117–R1125.
- O'Reilly I & Murphy MP 1992 Studies on the rapid stimulation of mitochondrial respiration by thyroid hormones. *Acta Endocrinologica* **127** 542–546. (doi:10.1530/acta.0.1270542)
- Schmittgen TD, Lee EJ, Jiang J, Sarkar A, Yang L, Elton TS & Chen C 2008 Real-time PCR quantification of precursor and mature microRNA. *Methods* **44** 31–38. (doi:10.1016/j.ymeth.2007.09.006)
- Silvestri E, Cioffi F, Glinni D, Ceccarelli M, Lombardi A, de Lange P, Chambery A, Severino V, Lanni A, Goglia F *et al.* 2010 Pathways affected by 3,5-diiodo-L-thyronine in liver of high fat-fed rats: evidence from two-dimensional electrophoresis, blue-native PAGE, and mass spectrometry. *Molecular BioSystems* **6** 2256–2271. (doi:10.1039/c0mb00040j)
- St Germain DL & Galton VA 1997 The deiodinase family of selenoproteins. *Thyroid* **7** 655–668. (doi:10.1089/thy.1997.7.655)
- Weyemi U, Caillou B, Talbot M, Ameziane-El-Hassani R, Lacroix L, Laget-Chevallier O, Al Ghuzlan A, Roos D, Bidart JM, Virion A *et al.* 2010 Intracellular expression of reactive oxygen species-generating NADPH oxidase NOX4 in normal and cancer thyroid tissues. *Endocrine-Related Cancer* **17** 27–37. (doi:10.1677/ERC-09-0175)
- Williams GR & Bassett JH 2011 Deiodinases: the balance of thyroid hormone: local control of thyroid hormone action: role of type 2 deiodinase. *Journal of Endocrinology* **209** 261–272. (doi:10.1530/JOE-10-0448)
- Yagil C, Barak A, Ben-Dor D, Rosenmann E, Bernheim J, Rosner M, Segev Y, Weksler-Zangen S, Raz I & Yagil Y 2005 Nonproteinuric diabetes-associated nephropathy in the Cohen rat model of type 2 diabetes. *Diabetes* **54** 1487–1496. (doi:10.2337/diabetes.54.5.1487)

Received in final form 26 March 2014

Accepted 31 March 2014

Accepted Preprint published online 1 April 2014