



# Planting of neonicotinoid-coated corn raises honey bee mortality and sets back colony development

Olivier Samson-Robert<sup>1</sup>, Geneviève Labrie<sup>2</sup>, Madeleine Chagnon<sup>3</sup> and Valérie Fournier<sup>1</sup>

<sup>1</sup> Centre de recherche en innovation sur les végétaux, Université Laval, Québec, Canada

<sup>2</sup> Centre de recherche sur les grains Inc., Saint-Mathieu-de-Beloeil, Québec, Canada

<sup>3</sup> Département des Sciences Biologiques, Université du Québec à Montreal, Montréal, Québec, Canada

## ABSTRACT

Worldwide occurrences of honey bee colony losses have raised concerns about bee health and the sustainability of pollination-dependent crops. While multiple causal factors have been identified, seed coating with insecticides of the neonicotinoid family has been the focus of much discussion and research. Nonetheless, few studies have investigated the impacts of these insecticides under field conditions or in commercial beekeeping operations. Given that corn-seed coating constitutes the largest single use of neonicotinoid, our study compared honey bee mortality from commercial apiaries located in two different agricultural settings, i.e. corn-dominated areas and corn-free environments, during the corn planting season. Data was collected in 2012 and 2013 from 26 bee yards. Dead honey bees from five hives in each apiary were counted and collected, and samples were analyzed using a multi-residue LC-MS/MS method. Long-term effects on colony development were simulated based on a honey bee population dynamic model. Mortality survey showed that colonies located in a corn-dominated area had daily mortality counts 3.51 times those of colonies from corn crop-free sites. Chemical analyses revealed that honey bees were exposed to various agricultural pesticides during the corn planting season, but were primarily subjected to neonicotinoid compounds (54% of analysed samples contained clothianidin, and 31% contained both clothianidin and thiamethoxam). Performance development simulations performed on hive populations' show that increased mortality during the corn planting season sets back colony development and bears contributions to collapse risk but, most of all, reduces the effectiveness and value of colonies for pollination services. Our results also have implications for the numerous large-scale and worldwide-cultivated crops that currently rely on pre-emptive use of neonicotinoid seed treatments.

Submitted 16 May 2017  
Accepted 20 July 2017  
Published 14 August 2017

Corresponding author  
Valérie Fournier,  
valerie.fournier@fsaa.ulaval.ca

Academic editor  
Ilaria Negri

Additional Information and  
Declarations can be found on  
page 17

DOI 10.7717/peerj.3670

© Copyright  
2017 Samson-Robert et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Agricultural Science, Ecology, Entomology, Toxicology, Ecotoxicology

**Keywords** *Apis mellifera*, Clothianidin, Insecticide, Pollinator, Seed treatment, Thiamethoxam, Honey bee, Intoxication, Population dynamic model, Colony health


## INTRODUCTION

In recent years, decline in pollinators, both wild and managed, has garnered much attention, prompting considerable amount of research (see *Lundin et al., 2015*; *Godfray et al., 2015*

for review). In light of these studies, a suite of numerous and interacting factors have been highlighted, including loss of foraging resources due to habitat loss and homogenization, introduction of invasive species, climate change, parasites, pathogens, loss of genetic diversity, beekeeping practices, and exposure to pesticides (Potts *et al.*, 2010; Vanbergen *et al.*, 2013; Goulson *et al.*, 2015; Gill *et al.*, 2016; Klein *et al.*, 2017). The role of pesticides, especially those of the neonicotinoid family, have been of particular concern (Tennekes & Sánchez-Bayo, 2011). While a growing body of evidence has shed light on the multiple routes of exposure to these insecticides and their negative impacts of bee health (Maini, Medrzycki & Porrini, 2010; Pisa *et al.*, 2015; Samson-Robert *et al.*, 2015; Lundin *et al.*, 2015; Tsvetkov *et al.*, 2017; Woodcock *et al.*, 2017) their implications in colony decline are still much debated (Blacquièrre & Van der Steen, 2017). Current gaps in our knowledge, particularly at colony-level (Godfray *et al.*, 2015) and from field-based approaches (Lundin *et al.*, 2015), contribute to uncertainty surrounding these insecticides.

In the province of Quebec alone, prophylactic neonicotinoid seed treatment in corn and soybean represents about 1.5 million acres of land every year (ISQ, 2016), largely located in the southern area of the province. This area is also home to a third of the province's commercial beekeepers, with more than 12,000 colonies (ISQ, 2015). The use of neonicotinoid as seed treatments has diversified and multiplied the routes by which honey bees and other pollinators may be contaminated, consequently increasing the risk of exposure to these systemic insecticides. It is now known that planting of seeds coated with neonicotinoid releases particulate matter contaminated with the insecticide into the environment (Greatti *et al.*, 2003; Greatti *et al.*, 2006; Girolami *et al.*, 2012; Krupke *et al.*, 2012; Sgolastra *et al.*, 2012; Pistorius *et al.*, 2015), and honey bees in agricultural areas are likely to come into contact with clouds of contaminated dust while foraging (Goulson, 2013; Godfray *et al.*, 2015). Airborne particulate matter is also highly susceptible to drifting, settling and thereby contaminating flowers, which increases the risk of intoxication when honey bees collect pollen or nectar from contaminated vegetation (Goulson, 2013; Godfray *et al.*, 2015; Long & Krupke, 2016). It has also been established that only a fraction of the seed coating's active ingredient (1.6–20%) can be absorbed by the crop, thus allowing for the bulk of the remaining insecticide to stay in the soil, leach down and reach surface waters (Sur & Stork, 2003; Goulson, 2013). Ubiquitous contamination of surface water with neonicotinoid insecticides combined with important water requirements for colony development in the spring has also been highlighted as harmful for honey bee health (Samson-Robert *et al.*, 2014; Chrétien *et al.*, 2017). In the province of Quebec and Ontario (eastern Canada), the planting period of treated corn has repeatedly been associated with reported honey bee deaths (PMRA, 2017).

Given the extent to which neonicotinoid seed coating is used and its particular proximity with beekeeping operations, we initiated this work to establish whether large-scale use of treated corn was harmful to honey bee colonies and the beekeeping industry. We conducted a 2-year study using corn fields from commercial growers and commercial apiaries, therefore insuring the most realistic conditions. First, we monitored colony mortality during the corn planting period and analysed dead honey bees for pesticide residues. Second, we used a colony population dynamic model to assess the extent to which



**Figure 1** Map of study sites. Study area in Southern Quebec and locations of commercial apiaries in corn-free areas (open circles) or corn-dominated areas (filled circles). Circle charts describe land cover of each site within a 3 km radius.

honey bee mortality might disrupt colony growth and estimate its resulting impacts on the Southern Quebec's beekeeping industry.

## MATERIALS AND METHODS

### Experimental setting

Field trials took place in the southern part of Quebec, in Eastern Canada in a region with a historically high level of land use for agricultural purposes. This area is Quebec's most productive agricultural region and produces over 60% of the province's corn and soybean crops. Study sites (apiaries) were selected among seasonal commercial bee yards customarily and historically managed by beekeepers. Exposed apiaries ( $n = 9$  for 2012,  $n = 7$  for 2013) were located within a 500 m radius from a treated-corn field, within a broader landscape (3 km radius) dominated by neonicotinoid-treated corn fields (exposed treatment). Control apiaries ( $n = 3$  for 2012,  $n = 7$  for 2013) were located in an area where the main crops are hay, wheat, oats, barley and rye and with minimal distance of 3 km from any crop planted with neonicotinoid seed coating. Minimal distance between same treatment sites was 3 km and 17 km in between exposed-control sites (Fig. 1).

## Honey bee colonies

Commercial honey bee hives were placed at every study site by their respective beekeepers/owners according to the usual seasonal timing of such procedures, either before, or at the very beginning, of large-scale corn planting operations. All colonies remained on site for the duration of the experiment and were then moved for pollination purposes by owners. All study sites were factual customary bee yards locations and colonies were entirely managed by owners. However, colony management was standardised between beekeepers: all colonies (*Apis mellifera* hybrid Italian stock) were reared in Langstroth hives, treated with oxalic acid for varroa mite control and supplied with *ad libitum* pollen supplements. In each apiary, five colonies were randomly selected from among those with similar colony strength at the beginning of the experiment (five frames of bees and four frames of brood). In 2012 and 2013, four beekeepers managed colonies exclusively in exposed apiaries and two beekeepers managed colonies in both exposed and controled apiaries, for a total of six beekeepers per year.

## Mortality index

A one square meter (1 m<sup>2</sup>) piece of pale beige cloth was placed in front of each experimental hive. Every 48 h, dead honey bees found on the cloth were counted to evaluate mortality levels and then removed. In 2012, the sampling period went from May 3rd to June 15th; in 2013, from May 6th to May 24th. By definition, mortality rate is the measure of the precise number of deaths in a known population, per unit of time. Although colony strength can be estimated, the exact number of living bees it comprises cannot be known, especially in daytime when foragers are out of the hive. Similarly, the number of dead bees cannot be established precisely since honey bees exhibit necrophoric behaviour to limit the spread of disease and parasites and to avoid attracting predators ([Visscher, 1983](#)). Furthermore, estimating mortality based on the number of dead bees in front of the hive does not take into account the numerous deaths that might occur away from the hive, for example, when bees die in the field after a lethal exposure to pesticides or when they become disoriented and fail to return to the hive ([Porrini et al., 2002](#)). In fact, most of the mortality happens away from the colony and dead bees found in front of the hive only account for 1 to 20% of all deaths ([Johansen & Mayer, 1990](#)). As such, the estimate of mortality is referred to subsequently as a “mortality index” and acts as the best proxy for the comparison of mortality rates, while still vastly undervaluing honey bee losses. When mortality exceeded 100 dead bees per colony (minimum amount required for chemical analyses), specimens were collected for ulterior residue analyses.


## Chemical analysis

Dead honey bee samples were analyzed using a modified version of the QuEChERS method originally described by [Anastassiades et al. \(2003\)](#). Briefly, 15 ml of acetonitrile was added to 10 mg of a previously ground and homogenized sample of honey bee bodies. Extract was shaken vigorously for 1 min using a Vortex mixer at maximum speed. Afterwards, 0.6 mg of anhydrous magnesium sulfate and 1.5 mg of anhydrous sodium acetate were added to the extract, which was then vortexed for 1 min and centrifuged for 10 min at

3,450 rpm. An aliquot of 1 ml was transferred to a glass vial, and 60  $\mu$ l of methanol and 20  $\mu$ l of isoprocarb standard solution (10 ppm) were added. The solution was then strained through a 0.45  $\mu$ m PTFE filter and 10  $\mu$ l were analyzed by liquid chromatography/mass spectrometry utilizing a Waters Acquity LC interfaced to a Waters Xevo TQ MS (Halo C-18 columns, 4.6 solid core  $\times$  50 mm porous outer shell with 2.7  $\mu$ m particle). The mass spectrometer was positioned in a positive electrospray mode and utilized a different MS/MS scan for each of the pesticides monitored. Liquid chromatography injections were repeated three times. Parent pesticides and metabolites were identified based on comparisons of their chromatographic retention time with known standards and mass abundance ratios to at least two fragment transitions. The ion ratios between the two transitions had to comply with a maximum difference of 20% with the calibration standard. This multi-residue method allows for detection of over 400 agrochemical compounds at parts per billion concentration levels. Analyses were performed by a team of professional chemists from the Laboratory of food analyses at the Quebec Ministry of Agriculture (MAPAQ).

### Intracolony population dynamic model

The commercial beekeeping context in which this study was carried did not allow for controlled conditions after corn planting as colonies were immediately moved for pollination services of different crops (e.g., lowbush blueberry, cranberry), under different climates, at varying distances from our observation locations and for different durations. Therefore, population modeling was used to simulate colony dynamic. The model is based on the mathematical models of *Khoury, Myerscough & Barron (2011)* and operates under different scenarios based on the initial size of the colony's population, queen's egg-laying rate and mortality levels while only considering the population of female worker bees. In all simulations, the evolution of a typical colony was considered during the first three months of the beekeeping seasons and encompasses the corn planting period in our area (roughly the whole month of May) and the blooming period of the two most important pollinator-dependant crops. The simulation-modelling period starts when colonies emerge from their wintering confine, around mid-April. Population size is usually between 10,000 and 14,000 female worker bees, i.e., the average colony size at the beginning of the season in our study area. Forager bees were set to account for 25% of the colony's population (*Fahrbach & Robinson, 1996*). The daily egg-laying rates were set to three realistic levels, namely a rate allowing for fast (1,750 eggs/day), typical (1,500 eggs/day) and slow (1,250 eggs/day) colony development (*Winston, 1991*). Natural forager death rate was set to 0.08 and 0.14 forager bees per day as reported in scientific literature (*Rueppell, Kaftanoglu & Page Jr, 2009; Russell, Barron & Harris, 2013; Betti, Wahl & Zamir, 2014*). The term  $w$  reflects the emergence rate from brood to adult and was set to 16,000 to simulate a fast-growing colony typical of spring and early summer (*Cresswell & Thompson, 2012*). Other dynamic parameters remained unchanged from *Khoury, Myerscough & Barron (2011)* initial model. Simulations were run under the hypotheses of (1) typical and constant forager death rate with no or insignificant foragers' exposure, and (2) forager death rate increased by pesticide exposure during a 30-days corn planting period. In the later configuration,



**Figure 2** Honey bee mortality index. Mean ( $\pm$ CI) number of dead bees collected in front of hives during a 48-hour period over the whole corn planting season ( $F_{1,22} = 19.44$ ;  $p = 0.0002$ ).

exposed foragers death rate was raised by a factor of 3.51 as determined by our mortality survey (see ‘Results’).


### Statistical analyses

The honey bee mortality index data set was analysed using a generalized randomized block design (GRBD) with repeated measures. The experimental units consisted of a group of five colonies located in honey bee yards. GRBD refers to the randomization of study sites within treatments (exposed and control) and within blocks comprised of both study years (2012 and 2013). In this model, treatments and dates were considered as fixed factors, while years and study sites were considered as random factors. Dead honey bees were collected from the same colonies at repeated intervals throughout the corn planting season. Repeated measures design was used to address the longitudinal nature of the data set. The correlation structure for observations within the same experimental unit was selected based on Akaike Information Criterion (AIC). The normality assumption was tested using the Shapiro–Wilk’s statistic, and the homogeneity of variances was verified using the usual residual plots. Box–Cox power transformation was used to remedy violation of the model’s assumptions. Statistical analyses were performed with the `lme()` function in the `nlme` package of R software (V 3.0.2) with the significance level set at 0.05.

## RESULTS

### Honey bee mortality index

Compilation and statistical analysis of numbers of dead honey bees revealed a significant difference ( $F_{1,22} = 19.440897$ ,  $p = 0.0002$ ) in observed mortality during the corn planting period between apiaries from areas with intensive corn cultivation and those from corn-free environment, respectively 32.6 and 9.3 dead bees per colony per 48 h (Fig. 2). Daily mortality was on average 3.51 times higher for colonies located in the vicinity of corn fields planted



**Figure 3** Honey bee mortality in function on planting completion. Mean ( $\pm$ CI) number of dead bees collected in front of hives from apiaries in corn-free environment (A) or near intensive corn cultivation (B) during a 48-hour period. Data shown on the x-axis indicate daily progression of corn planting (0% = planting has not begun, 100% = planting is over) to account for different corn planting duration of each year. Progression of corn planting was determined by estimates provided by agronomists from the government's ministry of agriculture, provincial crop insurance reports and regional fertilizer sales (which immediately follow corn planting).

with neonicotinoid-treated seeds. Mortality levels were not statistically different between sites or years within same treatment, and did not vary as a function of time (Figs. 3A and 3B) during the corn planting season.

### Multi-residue analyses of dead honey bees

Chemical analyses of dead honey bees indicated that they were exposed to various agricultural chemicals throughout the corn planting period. A total of 38 different pesticides and metabolites, 19 insecticides, 11 fungicides, seven herbicides and one synergist, were found in the 74 dead honey bee samples ( $\geq 100$  specimens) that were analysed throughout the study. In the 16 dead honey bee samples collected in control areas (apiaries in corn-free agricultural areas), three pesticides were identified (thiabendazole, clothianidin and thiamethoxam), with some samples containing two pesticides and an average of 1 chemical per sample (Table 1). Of these three pesticides, the fungicide thiabendazole was detected at levels above the limit of quantification, but levels of the neonicotinoid insecticides clothianidin and thiamethoxam were consistently below the limit of quantification, and therefore well below concentrations known to cause acute bee mortality (Lundin et al., 2015). Analysis of the 58 exposed samples collected from apiaries in neonicotinoid-treated corn intensive areas identified 32 pesticides and metabolites, with an average of 3.48 chemicals per sample and up to nine different compounds in a single sample (Table 2).

**Table 1** Summary of multi-residue analyses from control apiaries. Number of detections, concentrations and LD<sub>50</sub> for all actives ingredients identified in samples (16) collected in front of colonies in a neonicotinoid-free agricultural environment during corn planting (LOQ = 1 µg/kg, LOD = 0.1 µg/kg).

Pesticide	Class <sup>a</sup>	Detection	(%)	Concentrations <sup>b</sup> (µg/kg)				LD <sub>50</sub> (ug/bee)	
				Min	Max	Mean	SEM	Contact	Oral
Thibendazole	FUNG	3	18.8	1.00	1.00	1	–	34 <sup>c</sup>	4
Clothianidin	NEO,	0	0.0	–	–	–	–	0.0439	0.00368
	S	(11 < LOQ)	(68.8)	–	–	–	–		
Thiamethoxam	NEO,	0	0.0	–	–	–	–	0.024	0.005
	S	(2 < LOQ)		–	–	–	–		

**Notes.**

<sup>a</sup>Class: FUNG, fungicide, NEO, neonicotinoid, S, systemic.

<sup>b</sup>Concentrations (Min, Max, Mean and SEM) for detections > LOQ.

<sup>c</sup>University of Hertfordshire (2013). The Pesticide Properties DataBase (PPDB) developed by the Agriculture & Environment Research Unit (AERU), University of Hertfordshire, 2006–2013 (Access: <http://sitem.herts.ac.uk/aeru/ppdb/en/index.htm>).

LD50 information from the Pesticide Ecotoxicity Database of the Office of Pesticide Programs, Ecological Fate and Effects Division, of the US Environmental Protection Agency, unless otherwise noted.

Visualisation of pesticide detections in the form of a network highlights that the three most frequent pesticide co-occurrence are clothianidin-thiamethoxam (unsurprisingly, as thiamethoxam is a known precursor of clothianidin), clothianidin-thiabendazole and thiamethoxam-thiabendazole and the most frequent triads of chemicals, which are clothianidin-thiamethoxam-thiabendazole in 30% of samples and atrazine-desethyl atrazine-metolachlor in 15% of samples (Fig. 4). Pesticide risk assessment based on the acute risk quotient (RQ) indicates that the vast majority of compounds score below the threshold level for concern (0.4) (EFED, PMRA & CDP, 2012). RQ is expressed as the ratio of exposure estimate, in this case, concentrations of active ingredient found in dead bees, to point estimates of effects, as established by the acute contact and oral lethal dose to 50% of the organisms tested (LD50). For example, if clothianidin's concentration in a given sample was of 19 µg/kg, its oral LD50 at 24 h is of 0.00368 µg/bee and considering a mean honey bee body weight of 0.128 g (EFED, PMRA & CDP, 2012), then the corresponding RQ value would be of 0.66 ( $19 \times (0.128 \div 1,000) \div 0.00368$ ). The neonicotinoid clothianidin was the only compound whose acute RQ value exceeded the acceptable limit and is therefore an identified source of concerns (Fig. 5).

### Intracolony population dynamic model

Our model reveals that increased mortality levels linked with the planting of neonicotinoid-treated corn has the potential to strongly impact colony dynamics (Fig. 6). Regardless of the queens' egg laying rate and typical mortality levels, populations from exposed colonies would follow a marked decline during the planting period of neonicotinoid-treated corn, after which colony growth would start anew. Simulations show that exposure to planting of neonicotinoid-treated corn by itself should not be sufficient to cause the collapse of exposed colonies. However, the worst-case scenario predicts colony collapse before the end of the corn planting period (Fig. 6F). Also, population sizes of exposed and surviving colonies are predicted to be considerably reduced at key moments during the Quebec's beekeeping season, which are during the lowbush blueberry and cranberry crop pollination

**Table 2** Summary of multi-residue analyses from exposed apiaries. Number of detections, concentrations and LD<sub>50</sub> for all actives ingredients identified in samples ( $N = 58$ ) collected in front of colonies in a neonicotinoid dominated environment during corn planting (LOQ = 1  $\mu\text{g}/\text{kg}$ , LOD = 0.1  $\mu\text{g}/\text{kg}$ ).

Pesticide	Class <sup>a</sup>	Detection	(%)	Concentrations <sup>b</sup> ( $\mu\text{g}/\text{kg}$ )				LD <sub>50</sub> (ug/bee)	
				Min	Max	Mean	SEM	Contact	Oral
Clothianidin	NEO, S	31	53.5	1.00	19	5.81	0.86	0.0439	0.00368
Atrazine	HERB, S	28	48.3	1.00	21.00	4.11	1.07	>97	100 <sup>a</sup>
Metolachlor	HERB, PS	23	39.7	1.00	8.00	2.78	0.42	>110	>110
Thiamethoxam	NEO, S	18 (9 < LOQ)	31.0 (15.5)	1.00	5.00	1.61	0.26	0.024	0.005
Desethyl atrazine	HERB	13	22.4	1.00	33.00	6.69	2.32	>97	ND
Thiabendazole	FUNG	10 (15 < LOQ)	17.2 (25.9)	1.00	4.00	1.50	0.31	34 <sup>c</sup>	4
Boscalid	FUNG, S	10	17.2	5.00	90.00	28.50	8.55	>200	>166
Trifloxystrobin	FUNG, PS	10	17.2	1.00	25.00	7.20	2.65	200	>200 <sup>c</sup>
Azinphos-methyl	OP	9	15.5	1.00	287	98.89	43.62	0.42	0.15
Coumaphos	OP	8	13.8	1.00	3.00	1.63	0.32	20.29	2.99
Pyraclostrobin	FUNG	5	8.6	1.00	5.00	2.40	0.75	>100	>73.1 <sup>c</sup>
Bromoxynil	HERB	4	6.9	3.00	7.00	4.50	0.87	14.5	5
Dimethoate	OP	4	6.9	3.00	12.00	7.50	1.94	0.16	0.056
Iprodione	FUNG	4	6.9	5.00	8.00	6.75	0.75	>120	>25 <sup>c</sup>
Omethoate	OP	3	5.2	5.00	8.00	6.00	1.00	0.48	0,048 <sup>c</sup>
Thiacloprid	NEO, S	3	5.2	1.00	7.00	3.33	1.86	37.83	17.32
Cymoxanil	FUNG	2	3.5	1.00	1.00	1.00	–	>25	>85.3 <sup>c</sup>
Diazinon	OP	2	3.5	1.00	6.00	3.50	2.50	0.22	0.2
Diphenylamine	FUNG	2	3.5	2.00	6.00	4.00	2.00	ND	ND
Kresoxym methyl	FUNG	2	3.5	11.00	18.00	14.50	3.50	>25	>110 <sup>c</sup>
Pendimethalin	HERB	2	3.5	3.00	3.00	3.00	–	49.8	>101.2 <sup>c</sup>
Pyrimethanil	FUNG	2	3.5	1.00	2.00	1.50	0.50	100	100
Aldicarb	CARB, S	1	1.7	1.00	1.00	1.00	–	0.28	ND
Aldicarb sulfone	CARB, S, MET	1	1.7	3.00	3.00	3.00	–	0.28	ND
Aldicarb sulfoxide	CARB, S, MET	1	1.7	4.00	4.00	4.00	–	0.28	ND
Carbaryl	CARB, PS	1	1.7	21.00	21.00	21.00	–	1.1	0.11
Carbendazim	FUNG, S	1	1.7	3.00	3.00	3.00	–	>50	>756 <sup>c</sup>
Dimethenamid	HERB	1	3.4	1.00	1.00	1.00	–	>98	>1,000
Fludioxonil	FUNG	1	1.7	14.00	14.00	14.00	–	>100 <sup>a</sup>	>100
$\lambda$ cyhalothrin	PYR	1	1.7	7.00	7.00	7.00	–	0.038	0.91 <sup>c</sup>
Malathion	OP	1	1.7	2.00	2.00	2.00	–	0.2	0.38
Phosmet	OP	1	1.7	27.00	27.00	27.00	–	1.06	0.37
Piperonyl butoxide	SYN	1	1.7	1.00	1.00	1.00	–	>11	ND
Simazine	HERB, S	1	1.7	366.00	366.00	366.00	–	96.7	ND
Spinetoram A	SPI	1	1.7	11.00	11.00	11.00	–	0.024	0.14 <sup>c</sup>
Spinetoram D	SPI	1	1.7	31.00	31.00	31.00	–	0.024	0.14 <sup>c</sup>

(continued on next page)

Table 2 (continued)

Pesticide	Class <sup>a</sup>	Detection	(%)	Concentrations <sup>b</sup> (µg/kg)				LD <sub>50</sub> (ug/bee)	
				Min	Max	Mean	SEM	Contact	Oral
Spinosad A	SPI	1	1.7	4.00	4.00	4.00	–	0.0029	0.057 <sup>d</sup>
Tau-fluvalinate	PYR	1	1.7	4.00	4.00	4.00	–	0.2	ND

**Notes.**


<sup>a</sup>Class: CARB, carbamate; FUNG, fungicide; HERB, herbicide; MET, metabolite; OP, organophosphate; NEO, neonicotinoid; PS, partially systemic; PYR, pyrethroid; S, systemic; SPI, spinosine; SYN, synergist.

<sup>b</sup>Mean and SEM for detections > LOQ.


<sup>c</sup>University of Hertfordshire (2013). The Pesticide Properties DataBase (PPDB) developed by the Agriculture & Environment Research Unit (AERU), University of Hertfordshire, 2006–2013 (Access: <http://sitem.herts.ac.uk/aeru/ppdb/en/index.htm>).

<sup>d</sup>Miles et al., 2011. Effects of spinosad on honey bees (*Apis mellifera*): findings from over ten years of testing and commercial use. 11th International Symposium of the ICP-BR Bee Protection Group. DOI: 10.5073/jka.2012.437.032. (Access: <http://pub.jki.bund.de/index.php/JKA/article/viewFile/1958/2334>)


<sup>e</sup>LD50 information from the Pesticide Ecotoxicity Database of the Office of Pesticide Programs, Ecological Fate and Effects Division, of the US Environmental Protection Agency, unless otherwise noted.



**Figure 4** Pesticide detection network highlights the combinations of agrochemicals that were detected in a single sample. The wider the circle, the more frequent this compound was detected; the wider the line, the more frequent the combination was. Insecticides are marked in red, fungicides in blue, herbicides in green and synergist in yellow. Coloured areas indicate most common cocktails of chemicals. ALD, Aldicarb; ASO, Aldicarb sulfone; ASX, Aldicarb sulfoxide; ATR, Atrazine; AZI, Azinphos-methyl; BOS, Boscalid; BRO, Bromoxynil; CAR, Carbaryl; CBA, Carbendazim; CLO, Clothianidin; COU, Coumaphos; CYM, Cymoxanil; DAT, Desethyl atrazine; DIA, Diazinon; DIM, Dimethoate; DMA, Dimethenamid; DPA, Diphenylamine; FLU, Fludioxonil; IPR, Iprodione; KME, Kresoxym methyl; LCY, λ cyhalothrin; MAL, Malathion; MET, Metolachlor; OME, Omethoate; PBU, Piperonyl butoxide; PCS, Pyraclostrobin; PEN, Pendimethalin; PHO, Phosmet; PME, Pyrimethanil; SIM, Simazine; SPA, Spinetoram A; SPD, Spinetoram D; SPI, Spinosad A; TBA, Thiabendazole; TCP, Thiacloprid; TFL, Tau-fluvalinate; TFS, Trifloxystrobin; THI, Thiamethoxam.



**Figure 5** Risk assessment based on found concentrations in dead bees and known  $LD^{50}$ . Insecticides are highlighted in red, fungicides in blue, herbicides in green and synergist in yellow. The acceptable limit of concern was set at 0.4 (RQ) based on the historic average dose response relationship for acute studies with bees as described in the White Paper (*EFED, PMRA & CDPR, 2012*). The neonicotinoid clothianidin is the only compound whose RQ exceeds the concern threshold. Hence, it poses the greatest toxicity risk to honey bees relative to other agrochemicals. ALD, Aldicarb; ASO, Aldicarb sulfone; ASX, Aldicarb sulfoxide; ATR, Atrazine; AZI, Azinphos-methyl; BOS, Boscalid; BRO, Bromoxynil; CAR, Carbaryl; CBA, Carbendazim; CLO, Clothianidin; COU, Coumaphos; CYM, Cymoxanil; DAT, Desethyl atrazine; DIA, Diazinon; DIM, Dimethoate; DMA, Dimethenamid; DPA, Diphenylamine; FLU, Fludioxonil; IPR, Iprodione; KME, Kresoxym methyl; LCY,  $\lambda$  cyhalothrin; MAL, Malathion; MET, Metolachlor, OME, Omethoate; PBU, Piperonyl butoxide; PCS, Pyraclostrobin; PEN, Pendimethalin; PHO, Phosmet; PME, Pyrimethanil; SIM, Simazine; SPA, Spinetoram A; SPD, Spinetoram D; SPI, Spinosad A; TBA, Thiabendazole; TCP, Thiacloprid; TFL, Tau-fluvalinate; TFS, Trifloxystrobin, THI, Thiamethoxam.



**Figure 6** Honey bee population dynamics of six simulated colonies. Model is based on the mathematical models of *Khoury, Myerscough & Barron (2011)*. Simulations start at 10,000 and 14,000 bees, which is the average springtime colony size for our study area.  $L$  corresponds to the queen's daily egg-laying rate and varies from 1,750 to 1,250 to account for (A and D) fast, (B and E) regular and (C and F) slow colony growth.  $m$  is the normal level of forager mortality (either 0.08 or 0.14) which is multiplied by a factor of 3.51 for colonies in proximity to neonicotinoid seed-coated corn during planting period. Blue lines for colonies located in corn-free agricultural environment and red lines for colonies in vicinity neonicotinoid seed-coated corn fields. Collapse threshold is set at 5,000 bees (dotted line) based on the European Guidance Document on risk assessment on bees (*EFSA, 2013*).

service periods. For instance, at the beginning of the lowbush blueberry pollination season beginning around early June, exposed colonies are predicted to be at best 2.33 times (*Fig. 6A*) and at worst 3.47 times (*Fig. 6E*) smaller than typical unexposed colonies. As for the cranberry pollination season, usually beginning in early July, colonies are still thought to be affected even if they partially recovered. According to our simulations, predicted populations of exposed colonies would still be 1.65 to 2.16 times smaller (respectively *Figs. 6A* and *6E*) than expected populations of unexposed colonies. After the 100-days period, population sizes of exposed colonies are still predicted to be smaller than expected population of typical colonies by several thousand individuals, between 9,000 (*Fig. 6A*) to 5,600 (*Fig. 6E*) fewer bees.

## DISCUSSION

This study indicates that springtime honey bee mortality is intensified when apiaries are located in the vicinity of neonicotinoid-coated corn fields, impairing colony development and bearing potential contribution to collapse. Our results also confirm that honey bees in Southern Quebec are regularly exposed to combinations of various pesticides, among which clothianidin was the most frequent occurrence and the only active ingredient to exceed threshold for concern. Furthermore, we highlight the special concern posed by co-occurrence of multiple pesticides, especially the clothianidin-thiamethoxam-thiabendazole triad, as it was the most frequent concurrent detection.

### Honey bee mortality index

Throughout the entire corn planting season, honey bee colonies established near neonicotinoid-treated fields showed a 3.51 times higher daily mortality index (Fig. 2). In temperate climates, this increased loss of foragers can be a cause for concern as spring is a particularly challenging time for colonies (Mattila & Otis, 2006; Betti, Wahl & Zamir, 2016). Resumption of intensive brood rearing in late winter and early spring may exhaust the colony's pollen reserve before additional or sufficient resources become available in the environment (Mattila & Otis, 2006). Increased bee mortality would further impair the colony's foraging efficiency and intensify pollen shortages. Should that happen, nutritional deficiencies would trigger a decline in brood production, which could even come to a complete halt (Mattila & Otis, 2006). Declining honey bee populations would result in seasonal vulnerability which would leave the colony more susceptible to other hazards (Betti, Wahl & Zamir, 2016).

As anticipated, mean mortality levels of control colonies were particularly stable and did not differ throughout the corn planting season (Fig. 3A). However, contrary to expectations, worker mortality of exposed colonies did not significantly differ as a function of time (Fig. 3B). We initially believed that mortality levels would be higher at the beginning of corn planting and gradually subside during spring as a similar pattern was observed with acetylcholinesterase expression in bumblebees (Samson-Robert et al., 2015). Our results did suggest that mean mortality levels tended to decrease over the season but the magnitude of variability in mortality between days and colonies of exposed sites did not allow for the detection of a significant effect (Fig. 3B). We hypothesize that weather conditions are a key driver of pesticide related bee deaths which would explain the variability observed in mortality levels between days. Since honey bees do not venture outside when it is raining and corn cannot be planted when heavy or frequent precipitation occurs, both pesticides exposure and honey bee deaths should be minimal. When weather conditions become favourable again, corn planting and honey bee foraging resume and mortality should consequently rise. We also hypothesize that proximity of on-going plantings with honey bee hive and bees foraging areas, rather than regional progression of plantings, play a key role in pesticide exposure and honey bee deaths, which would explain the observed variability in mortality levels between colonies of exposed sites.

## Multi-residue analyses of dead honey bees

Our results also indicate that honey bees from bee yards established in the vicinity of agricultural fields are not only widely exposed to neonicotinoid insecticides during corn planting, but also to a wide array of agricultural pesticides that play an important role in impairing colony health (Tables 1 and 2). Pesticide residue concentrations in dead bees were generally low and seldom exceeded LD<sub>50</sub>. However, many pesticides are rapidly metabolized in live bees, which may explain the general lack of high residue concentrations in recently dead bees. Nicotinic insecticides are particularly problematic in that regard since they are known to be metabolized within a few hours (Suchail, Debrauwer & Belzunces, 2004; Suchail et al., 2004; Cresswell et al., 2014; Blacquièrè & Van der Steen, 2017), making their detection especially tricky. Moreover, metabolites of neonicotinoids, most of which were not targeted by the chemical analyses, have also been shown to be highly toxic for bees (Suchail, Guez & Belzunces, 2001; Nauen et al., 2003; Blacquièrè et al., 2012) increasing the long-term toxicity of these compounds. Most pesticide detections also revealed the co-occurrence of many active ingredients within the same samples (Fig. 4), which would be expected to exhibit enhanced toxicity if they share the same mode of action or target site (Yu, 2014; Zhu et al., 2014) or if compounds compete for a limited pool of detoxification enzyme (Berenbaum & Johnson, 2015; Johnson, 2015). Neonicotinoids thus pose a particular threat, as they are known to be additively toxic when they occur together (Andersch, Jeschke & Thielert, 2010; Pavlaki et al., 2011; Palmer et al., 2013), this was the case for 31% of our samples (Fig. 4). Neonicotinoids are also known to interact additively with various other agrochemicals. As such, combined neonicotinoid exposure to certain pyrethroids, like  $\lambda$ -cyhalothrin, has been related to increased toxicity (Gill, Ramos-Rodriguez & Raine, 2012; Chen et al., 2015). Combined exposure with fungicides like boscalid (Tsvetkov et al., 2017), trifloxystrobin (Wachendorff-Neumann et al., 2012) and ergosterol inhibitor (Iwasa et al., 2004; Thompson et al., 2014; Johnson, 2015; Sgolastra et al., 2017) also result in synergistic toxicity. In the last six years, 12 patent applications have also claimed or demonstrated additive toxicity between the fungicide thiabendazole and thiamethoxam (Donley, 2016), which occurred together in 30% of our samples. Herbicidal compounds have also been shown to be additively toxic when occurring together with neonicotinoids, as was the case with atrazine in 19% of our samples. Finally, combined exposure with in-hive acaricides, such as coumaphos, results in enhanced toxicity in addition with impaired memory and olfactory learning (Williamson & Wright, 2013; Berenbaum & Johnson, 2015; Johnson, 2015). Of all the insecticides identified in dead honey bee bodies, those of the neonicotinoid family, clothianidin and thiamethoxam, were detected far more frequently than others. Furthermore, clothianidin was the only pesticide whose acute RQ exceeded the threshold for concern (Fig. 5) and thereby poses the highest risk for honey bees, which is in accordance with the findings of Sánchez-Bayo & Goka (2014). Taking into account (1) the known high toxicity of nitroguanidine neonicotinoid insecticides (Goulson, 2013); (2) concentration levels found in dead bees in our study; (3) the possible synergistic interactions with other agrochemicals identified in collected dead bees; and (4) the synchronism between the timing of honey bee deaths and concurrence with planting of treated corn in the vicinity,

our study concludes that exposure to neonicotinoid compounds, mostly clothianidin, was the most likely cause of the bee mortality observed at the hives during our study.

### **Impaired colony growth and contributions to collapse**

Unusually high spring forager mortality (Fig. 2) might represent a heavy burden for colonies, assuredly rendering them more vulnerable to other threats and potentially causing their collapse. When implementing the increased in-field mortality into a honey bee population dynamic model, all scenarios exhibited a major deviation from the expected colony dynamic (Fig. 6). When considering the 100-day period, all exposed colonies suffered an important developmental setback, which has the potential to, in extreme scenarios (Fig. 6F), cause the collapse of the colonies. In most cases however, simulations show that colonies should survive the increased mortality and recover over the course of the season, given the proper care and favourable conditions. Nonetheless, colonies seldom encounter such ideal conditions and in reality must usually deal with several pathogens, parasites and predators (Potts *et al.*, 2010; Vanbergen *et al.*, 2013; Goulson *et al.*, 2015; Gill *et al.*, 2016; Klein *et al.*, 2017). In bad-case scenarios (Figs. 6D and 6E), when the corn planting period ends, colonies are expected to dwindle just above 5,000 individuals, which represents the lowest viable colony size (Henry *et al.*, 2012; EFSA, 2013). With such small populations, even a slight disruption of the colony, caused by any type of stress (dietary, parasites, pathogens, weather, transhumance, etc.), might be sufficient to cause its collapse. Recently, Henry *et al.* (2015) confirmed that field exposure to neonicotinoid was indeed associated with increased forager mortality. They also concluded that colonies compensate for the excess mortality without impairing colony growth or honey production. These results appear to be in contradiction with our own conclusions. However, Henry *et al.* colonies were exposed to oilseed rape grown from seeds treated with neonicotinoid and research on honey bees have repeatedly shown little adverse effects at colony-level in this particular crop (Cutler & Scott-Dupree, 2007; Pohorecka *et al.*, 2012; Pilling *et al.*, 2013; Cutler & Scott-dupree, 2014). Being particularly attractive to honey bees, it seems counter intuitive that this crop would not adversely impact colonies. However, oilseed rape produces abundant sugar-rich nectar (Kevan, Lee & Shuel, 1991; Pernal & Currie, 1997) and good quality pollen (Somerville, 2001) allowing for fast and sustained colony development. We believe these benefits outweigh the costs of exposure to neonicotinoid residues from treated seeds.

### **Pollination services**

A recent study by Stanley *et al.* (2015) showed, for the first time, that exposure to neonicotinoid pesticides induced colony-level reduction in pollination services provided by bumblebees to apples. In the case of honey bees, impaired colony development as shown in our population dynamic models may negatively impact both pollination services and honey production, resulting in important economic losses for beekeepers and berry growers in Quebec. For one, to be economically profitable for beekeepers, or even considered for pollination services, honey bee colonies need to meet population size standards in Quebec (FAQ, 2015). Immediately following chronic exposure from corn plantings, predicted

honey bee populations of bad-case scenarios (Figs. 6D and 6E) would not meet the minimal required standard of 7,000 bees and could simply not be rented for pollination services in Quebec. Even mid-summer blooming crops would potentially deal with the after effects of this chronic exposure, as predicted population of colonies would still only be a fraction of the required strength. One common way of dealing with weak colonies is to combine them together so the resulting colonies, although fewer in numbers, would then have the required strength for commercial pollination. As chronic exposure to neonicotinoid compounds reduces the number of honey bee colonies available for rent, pollinator-dependent crops might suffer from this pollination service shortage. For example, lowbush blueberry and cranberry crops are two of Quebec's fastest growing productions and annually monopolise 90% of the colonies on rent throughout the province (ISQ, 2014). Over the last 10 years, honey bee colony rental prices for crop pollination has increased by more than 60% due to the increasing pollination demand from these two crops (Belzile & Li, 2014; ISQ, 2014). Chronic exposures to toxic chemicals thus forces beekeepers to deal with weakened colonies, more vulnerable to parasites and pathogens, worth less for pollination services, sometimes not rentable at all and in all cases producing less honey. Crop growers also bear the consequences as weaker bee colonies are not as efficient at pollinating their crops and rental prices are likely to increase if supply of bee hives becomes a negative issue.

## CONCLUSIONS

Extensive spring planting of neonicotinoid-coated corn resulted in increased daily mortality of individuals from commercial honey bee colonies located in the vicinity (<500 m). Neonicotinoid compounds were the prevailing insecticides detected in dead honey bees from our study with clothianidin being the only compound whose acute RQ exceeded the threshold for concern. The consequence of this type of agrochemical contamination strongly limits honey bee colony development, potentially affecting the number of available foragers, and reducing their colony effectiveness for crop pollination and consequently their monetary value for hive rental. These adverse conditions can likely lead to colony collapse. Additionally, the widespread use of neonicotinoid treated-corn has been repeatedly questioned. Studies from Europe and North America alike have shown that prophylactic use bears insignificant to little increase in yield and that more sustainable pest control alternatives are available (Quesnel, 2005; Goulson, 2013; Furlan & Kreutzweiser, 2014; Labrie et al., 2014; Krupke et al., 2017). Our ISQ provides further evidence of the key role of neonicotinoids in the declining health of honey bee colonies and highlights the potential effects of bee exposure to common combinations of agrochemicals. Policy makers should prioritize the need for future research into pesticide interactions.

## ACKNOWLEDGEMENTS

We would like to thank André Rondeau, André Pettigrew and Ermin Menkovic for their help with site selection. We give special thanks to the CÉROM for fieldwork supplies and technical assistance. We are also grateful to Claude Boucher, Josée Rondeau, and Diane Boucher for assistance with processing chemical analysis of samples during the first year

of the study, Étienne Nadeau for field assistance, Luc Gagnon and his team for conducting chemical analyses, Gaétan Daigle for his help with statistical analyses and André Fortin for his assistance with dynamic population modeling. We are also thankful to anonymous reviewers for their constructive comments on a previous version of the manuscript.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was funded by Prime-Vert sous volet 11,1—Appui à la Stratégie phytosanitaire québécoise en agriculture of the Ministère de l’Agriculture, des Pêcherie et de l’Alimentation (MAPAQ). OS-R received various scholarships from the Quebec’s Centre for Biodiversity Science and NSERC-CANPOLIN. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Prime-Vert sous volet 11,1 —Appui à la Stratégie phytosanitaire québécoise en agriculture of the Ministère de l’Agriculture, des Pêcherie et de l’Alimentation (MAPAQ).  
Quebec’s Centre for Biodiversity Science and NSERC-CANPOLIN.

### Competing Interests

Geneviève Labrie is an employee of CÉROM, Centre de recherche sur les grains Inc. The remaining authors declare there are no competing interests.

### Author Contributions

- Olivier Samson-Robert conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Geneviève Labrie, Madeleine Chagnon and Valérie Fournier conceived and designed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.3670#supplemental-information>.

## REFERENCES

- Anastassiades M, Lehotay SJ, Stajnbaher D, Schenck FJ. 2003.** Fast and easy multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *Journal of AOAC International* **86**:412–431.
- Andersch W, Jeschke P, Thielert W. 2010.** Combination of methiocarb and one or more compounds selected from thiacloprid, thiamethoxam, acetamiprid, nitenpyram,

- and dinotefuran; effective animal pests control and for plant seed dressing. Google patents. Bayer CropScience AG, Monheim am Rhein.
- Belzile L, Li J. 2014.** La croissance de l'industrie apicole québécoise: une fausse joie? In: *Fiche synthèse, IRDA*. 3 p. Available at [https://www.irda.qc.ca/assets/documents/Publications/documents/belzile-li-2014\\_fiche\\_croissance\\_apicole\\_qc.pdf](https://www.irda.qc.ca/assets/documents/Publications/documents/belzile-li-2014_fiche_croissance_apicole_qc.pdf) (accessed on 23 September 2015).
- Berenbaum MR, Johnson RM. 2015.** Xenobiotic detoxification pathways in honey bees. *Current Opinion in Insect Science* **10**:51–58 DOI [10.1016/j.cois.2015.03.005](https://doi.org/10.1016/j.cois.2015.03.005).
- Betti MI, Wahl LM, Zamir M. 2014.** Effects of infection on honey bee population dynamics: a model. *PLOS ONE* **9**:e110237 DOI [10.1371/journal.pone.0110237](https://doi.org/10.1371/journal.pone.0110237).
- Betti MI, Wahl LM, Zamir M. 2016.** Age structure is critical to the population dynamics and survival of honeybee colonies. *R. Soc. Open Sci.* **3**:160444 DOI [10.1098/rsos.160444](https://doi.org/10.1098/rsos.160444).
- Blacquièrè T, Smagghè G, Van Gestel CAM, Mommaerts V. 2012.** Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology* **21**:973–992 DOI [10.1007/s10646-012-0863-x](https://doi.org/10.1007/s10646-012-0863-x).
- Blacquièrè T, van der Steen JJ. 2017.** Three years of banning neonicotinoid insecticides based on sub-lethal effects: can we expect to see effects on bees? *Pest Management Science* **73**:1299–1304 DOI [10.1002/ps.4583](https://doi.org/10.1002/ps.4583).
- Chen C, Wang Y, Qian Y, Zhao X, Wang Q. 2015.** The synergistic toxicity of the multiple chemical mixtures: implications for risk assessment in the terrestrial environment. *Environment International* **77**:95–105 DOI [10.1016/j.envint.2015.01.014](https://doi.org/10.1016/j.envint.2015.01.014).
- Chrétien F, Giroux I, Thériault G, Gagnon P, Corriveau J. 2017.** Surface runoff and subsurface tile drain losses of neonicotinoids and companion herbicides at edge-of-field. *Environmental Pollution* **224**:255–264 DOI [10.1016/j.envpol.2017.02.002](https://doi.org/10.1016/j.envpol.2017.02.002).
- Cresswell JE, Robert F-XL, Florance H, Smirnov N. 2014.** Clearance of ingested neonicotinoid pesticide (imidacloprid) in honey bees (*Apis mellifera*) and bumblebees (*Bombus terrestris*). *Pest Management Science* **70**:332–337 DOI [10.1002/ps.3569](https://doi.org/10.1002/ps.3569).
- Cresswell JE, Thompson HM. 2012.** Comment on “A common pesticide decreases foraging success and survival in honey bees”. *Science* **337**:1453 DOI [10.1126/science.1224618](https://doi.org/10.1126/science.1224618).
- Cutler GC, Scott-Dupree CD. 2007.** Exposure to clothianidin seed-treated canola has no long-term impact on honey bees. *Journal of Economic Entomology* **100**:765–772 DOI [10.1093/jee/100.3.765](https://doi.org/10.1093/jee/100.3.765).
- Cutler GC, Scott-dupree CD. 2014.** A field study examining the effects of exposure to neonicotinoid seed-treated corn on commercial bumble bee colonies. *Ecotoxicology* **23**:1755–1763 DOI [10.1007/s10646-014-1340-5](https://doi.org/10.1007/s10646-014-1340-5).
- Donley N. 2016.** *Toxic concoctions: how the EPA ignores the dangers of pesticide cocktails*. Tucson: Center for Biological Diversity.
- Environmental Fate and Effects Division (EFED), Pest Management Regulatory Agency (PMRA), California Department of Pesticides Regulation (CDPR). 2012.** White paper in support of the proposed risk assessment process for bees. 275 p. Available at [http://www.cdpr.ca.gov/docs/emon/surfwttr/presentations/epa\\_whitepaper.pdf](http://www.cdpr.ca.gov/docs/emon/surfwttr/presentations/epa_whitepaper.pdf) (accessed on 15 October 2015).

- European Food Safety Authority (EFSA). 2013.** Guidance Document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees). *EFSA Journal* 11:3295 DOI 10.2903/j.efsa.2013.3295.
- Fahrbach SE, Robinson GE. 1996.** Juvenile hormone, behavioral maturation, and brain structure in the honey bee. *Developmental Neuroscience* 18:102–114.
- Fédération des Apiculteurs du Québec. 2015.** Contrat de pollinisation–location de ruches. Available at [http://www.apiculteursduquebec.com/documents/74/Contrat\\_type\\_poll\\_revise\\_Fev\\_2015.pdf](http://www.apiculteursduquebec.com/documents/74/Contrat_type_poll_revise_Fev_2015.pdf) (accessed on 5 March 2017).
- Furlan L, Kreuzweiser D. 2014.** Alternatives to neonicotinoid insecticides for pest control: case studies in agriculture and forestry. *Environmental Science and Pollution Research* 22:135–147.
- Gill RJ, Baldock KCR, Brown MJF, Cresswell JE, Dicks LV, Fountain MT, Garratt MPD, Gough LA, Heard MS, Holl JM, Ollerton J, Stone GN, Tang CQ, Vanbergen AJ, Vogler AP, Woodward G, Arce AN, Boatman ND, Brand-Hardy R, Breeze TD, Green M, Hartfield CM, O'Connor RS, Osborne JL, Phillips J, Sutton PB, Potts SG, O'Connor RS, Osborne JL, Phillips J, Sutton PB, Potts SG. 2016.** Chapter four—protecting an ecosystem service: approaches to understanding and mitigating threats to wild insect pollinators. *Advances in Ecological Research* 54:135–206 DOI 10.1016/bs.aecr.2015.10.007.
- Gill RJ, Ramos-Rodriguez O, Raine NE. 2012.** Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature* 491:105–108 DOI 10.1038/nature11585.
- Girolami V, Marzaro M, Vivan L, Mazzon L, Greatti M, Giorio C, Marton D, Tapparo A. 2012.** Fatal powdering of bees in flight with particulates of neonicotinoids seed coating and humidity implication. *Journal of Applied Entomology* 136:17–26 DOI 10.1111/j.1439-0418.2011.01648.x.
- Godfray HCJ, Blacquière T, Field LM, Hails RS, Potts SG, Raine NE, Vanbergen AJ, McLean AR. 2015.** A restatement of recent advances in the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proceedings. Biological Sciences: Biological Sciences* 282:20151821 DOI 10.1098/rspb.2015.1821.
- Goulson D. 2013.** REVIEW: an overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology* 50:977–987 DOI 10.1111/1365-2664.12111.
- Goulson D, Nicholls E, Botias C, Rotheray EL. 2015.** Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 347:1255957 DOI 10.1126/science.1255957.
- Greatti M, Barbattini R, Sabatini AG, Rossi S. 2006.** Presence of the a.i. imidacloprid on vegetation near corn fields sown with Gaucho<sup>®</sup> dressed seeds. *Bulletin of Insectology* 59:99–103.
- Greatti M, Sabatini AG, Barbattini R, Rossi S, Stravisi A. 2003.** Risk of environmental contamination by the active ingredient imidacloprid used for corn seed dressing. Preliminary results. *Bulletin of Insectology* 56:69–72.

- Health Canada's Pest Management Regulatory Agency (PMRA).** 2017. Update on Canadian bee incident reports 2012–2016. Available at [https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/cps-spc/alt\\_formats/pdf/pubs/pest/\\_fact-fiche/bees-incident-abeilles-2012-2016-eng.pdf](https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/cps-spc/alt_formats/pdf/pubs/pest/_fact-fiche/bees-incident-abeilles-2012-2016-eng.pdf) (accessed on 12 February 2017).
- Henry M, Béguin M, Requier F, Rollin O, Odoux J-F, Aupinel P, Aptel J, Tchamitchian S, Decourtye A.** 2012. A common pesticide decreases foraging success and survival in honey bees. *Science* **336**:348–350 DOI [10.1126/science.1215039](https://doi.org/10.1126/science.1215039).
- Henry M, Cerrutti N, Aupinel P, Decourtye A, Gayraud M, Odoux J-F, Pissard A, Rüger C, Bretagnolle V.** 2015. Reconciling laboratory and field assessments of neonicotinoid toxicity to honeybees. *Proceedings of the Royal Society B: Biological Sciences* **282**:20152110 DOI [10.1098/rspb.2015.2110](https://doi.org/10.1098/rspb.2015.2110).
- Institut de la Statistique du Québec (ISQ).** 2014. Statistiques relatives à la location de colonies à des fins de pollinisation selon le type de culture, Québec. Available at [http://www.stat.gouv.qc.ca/statistiques/agriculture/apiculture-miel/h7\\_2014.htm](http://www.stat.gouv.qc.ca/statistiques/agriculture/apiculture-miel/h7_2014.htm) (accessed on 05 March 2017).
- Institut de la Statistique du Québec.** 2015. Statistiques principales relatives au miel, par regroupement de régions administratives, Québec, 2015. Available at [http://www.stat.gouv.qc.ca/statistiques/agriculture/apiculture-miel/h1\\_2015.htm](http://www.stat.gouv.qc.ca/statistiques/agriculture/apiculture-miel/h1_2015.htm) (accessed on 05 March 2017).
- Institut de la Statistique du Québec (ISQ).** 2016. Superficie des grandes cultures, rendement à l'hectare et production, par région administrative. Available at [http://www.stat.gouv.qc.ca/statistiques/agriculture/grandes-cultures/gc\\_2016.htm](http://www.stat.gouv.qc.ca/statistiques/agriculture/grandes-cultures/gc_2016.htm) (accessed on 05 March 2017).
- Iwasa T, Motoyama N, Ambrose JT, Roe RM.** 2004. Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Protection* **23**:371–378 DOI [10.1016/j.cropro.2003.08.018](https://doi.org/10.1016/j.cropro.2003.08.018).
- Johansen CA, Mayer DF.** 1990. *Pollinator protection. A bee and pesticide handbook*. Cheshire: Wicwas Press.
- Johnson RM.** 2015. Honey bee toxicology. *Annual Review of Entomology* **60**:415–434 DOI [10.1146/annurev-ento-011613-162005](https://doi.org/10.1146/annurev-ento-011613-162005).
- Kevan PG, Lee H, Shuel RW.** 1991. Sugar ratios in nectars of varieties of canola (Brassica napus). *Journal of Apicultural Research* **30**:99–102 DOI [10.1080/00218839.1991.11101240](https://doi.org/10.1080/00218839.1991.11101240).
- Khoury DS, Myerscough MR, Barron AB.** 2011. A quantitative model of honey bee colony population dynamics. *PLOS ONE* **6**:e18491 DOI [10.1371/journal.pone.0018491](https://doi.org/10.1371/journal.pone.0018491).
- Klein S, Cabirol A, Devaud J-M, Barron AB, Lihoreau M.** 2017. Why bees are so vulnerable to environmental stressors. *Trends in Ecology & Evolution* DOI [10.1016/j.tree.2016.12.009](https://doi.org/10.1016/j.tree.2016.12.009).
- Krupke CH, Holland JD, Long EY, Eitzer BD.** 2017. Planting of neonicotinoid-treated maize poses risks for honey bees and other non-target organisms over a wide area without consistent crop yield benefit. *Journal of Applied Ecology* DOI [10.1111/1365-2664.12924](https://doi.org/10.1111/1365-2664.12924).

- Krupke CH, Hunt GJ, Eitzer BD, Andino G, Given K. 2012. Multiple routes of pesticide exposure for honey bees living near agricultural fields. *PLOS ONE* 7:e29268 DOI 10.1371/journal.pone.0029268.
- Labrie G, Rondeau A, Faucher Y, Mathieu S, Perreault Y, Tremblay G. 2014. Impact des traitements insecticides de semences sur les insectes ravageurs du sol et sur les paramètres agronomiques dans la culture du maïs grain. Available at [https://www.mapaq.gouv.qc.ca/SiteCollectionDocuments/Agroenvironnement/1582\\_Rapport.pdf](https://www.mapaq.gouv.qc.ca/SiteCollectionDocuments/Agroenvironnement/1582_Rapport.pdf) (accessed on 29 June 2017).
- Long EY, Krupke CH. 2016. Non-cultivated plants present a season-long route of pesticide exposure for honey bees. *Nature Communications* 7:11629 DOI 10.1038/ncomms11629.
- Lundin O, Rundlöf M, Smith HHG, Fries I, Bommarco R, Klein A, Vaissière B, Cane J, Steffan-Dewenter I, Cunningham S, Kremen C, Ollerton J, Winfree R, Tarrant S, Potts S, Biesmeijer J, Kremen C, Neumann P, Schweiger O, Kunin W, Vanbergen A, van der Sluijs J, Simon-Delso N, Goulson D, Maxim L, Bonmatin JJ-M, Belzunces L, Goulson D, Jeschke P, Nauen R, Schindler M, Elbert A, Elbert A, Haas M, Springer B, Thielert W, Nauen R, Jeschke P, Nauen R, Iwasa T, Motoyama N, Ambrose J, Roe R, Nuyttens D, Devarrewaere W, Verboven P, Foqué D, Cresswell J, Krupke C, Hunt G, Eitzer B, Andino G, Given K, Dicks L, Goulson D, Gross M, Cresswell J, Desneux N, Van Engelsdorp D, Maini S, Medrzycki P, Porrini C, Farooqui T, Fairbrother A, Purdy J, Anderson T, Fell R, Smith K, Loh E, Rostal M, Zambrana-Torrel C, Mendiola L, Daszak P, Blacquièrre T, Smagghe G, Van Gestel C, Mommaerts V, Pisa L, Amaral-Rogers V, Belzunces L, Bonmatin JJ-M, Downs C, Goulson D, Godfray H, Blacquièrre T, Field L, Hails R, Petrokofsky G, Potts S, Girolami V, Marzaro M, Vivan L, Mazzon L, Giorio C, Marton D, Rundlöf M, Andersson G, Bommarco R, Fries I, Hederström V, Herbertsson L, Schmuck R, Schöning R, Stork A, Schramel O, Larson J, Redmond C, Potter D, Hoffmann E, Castle S, Stewart S, Lorenz G, Catchot A, Gore J, Cook D, Skinner J, Sechser B, Freuler J, Byrne F, Visscher P, Leimkuehler B, Fischer D, Grafton-Cardwell E, Morse J, Stoner K, Eitzer B, Skerl M, Bolta S, Cesnik H, Gregorc A, Choudhary A, Sharma D, Chen M, Tao L, McLean J, Lu C, Vaikkinen A, Schmidt HH, Kiiski I, Rämö S, Hakala K, Haapala M, Reetz J, Zühlke S, Spittler M, Wallner K, Schreinemachers P, Tipraqsa P, Gallai N, Salles J, Settele J, Vaissière B, Schmuck R, Stadler T, Schmidt HH, Girolami V, Mazzon L, Squartini A, Mori N, Marzaro M, Di Bernardo A, Nguyen B, Saegerman C, Pirard C, Mignon J, Widart J, Thirionet B, Dively G, Kamel A, Gill R, Ramos-Rodriguez O, Raine N, Whitehorn P, O'Connor S, Wackers F, Goulson D, Elston C, Thompson H, Walters K, Laycock I, Cresswell J, Rortais A, Arnold G, Halm M, Touffet-Briens F, Halm M, Rortais A, Arnold G, Tasei J, Rault S, Sanchez-Bayo F, Goka K, Gels J, Held D, Potter D, Larson J, Redmond C, Potter D, Larson J, Redmond C, Potter D, Barbosa W, Smagghe G, Guedes R, Stark JJ. 2015. Neonicotinoid insecticides and their impacts on bees: a systematic review of research approaches and identification of knowledge gaps. *PLOS ONE* 10:e0136928 DOI 10.1371/journal.pone.0136928.

- Maini S, Medrzycki P, Porrini C. 2010.** The puzzle of honey bee lossess: a brief review. *Bulletin of Insectology* **63**:150–160.
- Mattila HR, Otis GW. 2006.** Influence of pollen diet in spring on development of honey bee (Hymenoptera: Apidae) Colonies. *Journal of Economic Entomology* **99**:604–613 DOI [10.1093/jee/99.3.604](https://doi.org/10.1093/jee/99.3.604).
- Nauen R, Ebbinghaus-Kintscher U, Salgado VL, Kaussmann M. 2003.** Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pesticide Biochemistry and Physiology* **76**:55–69 DOI [10.1016/S0048-3575\(03\)00065-8](https://doi.org/10.1016/S0048-3575(03)00065-8).
- Palmer MJ, Moffat C, Saranzewa N, Harvey J, Wright GA, Connolly CN. 2013.** Cholinergic pesticides cause mushroom body neuronal inactivation in honeybees. *Nature Communications* **4**:1634 DOI [10.1038/ncomms2648](https://doi.org/10.1038/ncomms2648).
- Pavlaki MD, Pereira R, Loureiro S, Soares AMVM. 2011.** Effects of binary mixtures on the life traits of *Daphnia magna*. *Ecotoxicology and Environmental Safety* **74**:99–110 DOI [10.1016/j.ecoenv.2010.07.010](https://doi.org/10.1016/j.ecoenv.2010.07.010).
- Pernal SF, Currie RW. 1997.** Nectar quality in open-pollinated, pol CMS hybrid, and dominant SI hybrid oilseed summer rape. *Canadian Journal Plant Sci* **78**:79–89 DOI [10.4141/P97-071](https://doi.org/10.4141/P97-071).
- Pilling E, Campbell P, Coulson M, Ruddle N, Tornier I. 2013.** A four-year field program investigating long-term effects of repeated exposure of honey bee colonies to flowering crops treated with thiamethoxam. *PLOS ONE* **8**:e77193 DOI [10.1371/journal.pone.0077193](https://doi.org/10.1371/journal.pone.0077193).
- Pisa LW, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Downs CA, Goulson D, Kreuzweiser DP, Krupke C, Liess M, McField M, Morrissey CA, Noome DA, Settele J, Simon-Delso N, Stark JD, van der Sluijs JP, Van Dyck H, Wiemers M. 2015.** Effects of neonicotinoids and fipronil on non-target invertebrates. *Environmental Science and Pollution Research* **22**:68–102 DOI [10.1007/s11356-014-3471-x](https://doi.org/10.1007/s11356-014-3471-x).
- Pistorius J, Wehner A, Kriszan M, Bargaen H, Knäbe S, Klein O, Frommberger M, Stähler M, Heimbach U. 2015.** Application of predefined doses of neonicotinoid containing dusts in field trials and acute effects on honey bees. *Bulletin of Insectology* **68**:161–172.
- Pohorecka K, Skubida P, Miszczak A, Semkiw P, Sikorski P, Zagibajlo K, Teper D, Koltowski Z, Skubida M, Zdanska D, Bober A. 2012.** Residues of neonicotinoid insecticides in bee collected plant materials from oilseed rape crops and their effect on bee colonies. *Journal of Apicultural Science* **56**:115–134 DOI [10.2478/v10289-012-0029-3](https://doi.org/10.2478/v10289-012-0029-3).
- Porrini C, Ghini S, Girotti S, Sabatini AG, Gattavecchia E, Celli G. 2002.** Use of honey bees as bioindicators of environmental pollution in Italy. In: Devillers J, Pham-Delegue M-H, eds. *Honey bees: estimating the environmental impact of chemicals*. Florance: CRC Press, 186–247.
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010.** Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* **25**:345–353 DOI [10.1016/j.tree.2010.01.007](https://doi.org/10.1016/j.tree.2010.01.007).

- Quesnel G. 2005.** Corn seed treatment to control seed and seedling insect pests. Crop Advances: OMAFRA Field Crop Project Reports. Available at <http://www.ontariosoilcrop.org/wp-content/uploads/2015/07/V2Cor13.pdf> (accessed on 29 June 2017).
- Rueppell O, Kaftanoglu O, Page Jr RE. 2009.** Honey bee (*Apis mellifera*) workers live longer in small than in large colonies. *Experimental Gerontology* **44**:447–452 DOI [10.1016/j.exger.2009.04.003](https://doi.org/10.1016/j.exger.2009.04.003).
- Russell S, Barron AB, Harris D. 2013.** Dynamic modelling of honey bee (*Apis mellifera*) colony growth and failure. *Ecological Modelling* **265**:158–169 DOI [10.1016/j.ecolmodel.2013.06.005](https://doi.org/10.1016/j.ecolmodel.2013.06.005).
- Samson-Robert O, Labrie G, Chagnon M, Fournier V. 2014.** Neonicotinoid-contaminated puddles of water represent a risk of intoxication for honey bees. *PLOS ONE* **9**:e108443 DOI [10.1371/journal.pone.0108443](https://doi.org/10.1371/journal.pone.0108443).
- Samson-Robert O, Labrie G, Mercier P-L, Chagnon M, Derome N, Fournier V. 2015.** Increased acetylcholinesterase expression in bumble bees during neonicotinoid-coated corn sowing. *Scientific Reports* **5**:12636 DOI [10.1038/srep12636](https://doi.org/10.1038/srep12636).
- Sánchez-Bayo F, Goka K. 2014.** Pesticide residues and bees—a risk assessment. *PLOS ONE* **9**:e94482 DOI [10.1371/journal.pone.0094482](https://doi.org/10.1371/journal.pone.0094482).
- Sgolastra F, Medrzycki P, Bortolotti L, Renzi MT, Tosi S, Bogo G, Teper D, Porrini C, Molowny-Horas R, Bosch J. 2017.** Synergistic mortality between a neonicotinoid insecticide and an ergosterol-biosynthesis-inhibiting fungicide in three bee species. *Pest Management Science* **73**:1236–1243 DOI [10.1002/ps.4449](https://doi.org/10.1002/ps.4449).
- Sgolastra F, Renzi T, Draghetti S, Medrzycki P, Lodesani M, Maini S, Porrini C. 2012.** Effects of neonicotinoid dust from maize seed-dressing on honey bees. *Bulletin of Insectology* **65**:273–280.
- Somerville DC. 2001.** Nutritional value of bee collected pollens. A report for the Rural Industries Research and Development Corporation, Publication No. 01/047; Project No. DAN-134A. NSW Agriculture Goulburn, NSW Kustry, 57–58. Available at <https://rirdc.infoservices.com.au/downloads/01-047> (accessed on 23 December 2016).
- Stanley DA, Garratt MPD, Wickens JB, Wickens VJ, Potts SG, Raine NE. 2015.** Neonicotinoid pesticide exposure impairs crop pollination services provided by bumblebees. *Nature* **528**:548–550 DOI [10.1038/nature16167](https://doi.org/10.1038/nature16167).
- Suchail S, Debrauwer L, Belzunces LP. 2004.** Metabolism of imidacloprid in *Apis mellifera*. *Pest Management Science* **60**:291–296 DOI [10.1002/ps.772](https://doi.org/10.1002/ps.772).
- Suchail S, Guez D, Belzunces LP. 2001.** Discrepancy between acute and chronic toxicity induced by imidacloprid and its metabolites in *Apis mellifera*. *Environmental Toxicology and Chemistry* **20**:2482–2486 DOI [10.1002/etc.5620201113](https://doi.org/10.1002/etc.5620201113).
- Suchail S, De Sousa G, Rahmani R, Belzunces LP. 2004.** *In vivo* distribution and metabolism of <sup>14</sup>C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Management Science* **60**:1056–62 DOI [10.1002/ps.895](https://doi.org/10.1002/ps.895).
- Sur R, Stork A. 2003.** Uptake, translocation and metabolism of imidacloprid in plants. *Bulletin of Insectology* **56**:35–40.

- Tennekes HA, Sánchez-Bayo F. 2011.** Time-Dependent toxicity of neonicotinoids and other toxicants: implications for a new approach to risk assessment. *Journal of Environmental & Analytical Toxicology* **S4**:001 DOI [10.4172/2161-0525.S4-001](https://doi.org/10.4172/2161-0525.S4-001).
- Thompson HM, Fryday SL, Harkin S, Milner S. 2014.** Potential impacts of synergism in honeybees (*Apis mellifera*) of exposure to neonicotinoids and sprayed fungicides in crops. *Apidologie* **45**:545–553 DOI [10.1007/s13592-014-0273-6](https://doi.org/10.1007/s13592-014-0273-6).
- Tsvetkov N, Samson-Robert O, Sood K, Patel HS, Malena DA, Gajiwala PH, Maciukiewicz P, Fournier V, Zayed A. 2017.** Chronic exposure to neonicotinoids reduces honey bee health near corn crops. *Science* **356**:1395–1397 DOI [10.1126/science.aam7470](https://doi.org/10.1126/science.aam7470).
- Vanbergen AJ, Baude M, Biesmeijer JC, Britton NF, Brown MJF, Brown M, Bryden J, Budge GE, Carvell C, Challinor AJ, Connolly CN, Evans DJ, Feil EJ, Garratt MP, Greco MK, Heard MS, Jansen VAA, Keeling MJ, Kunin WE, Marris GC, Memmott J, Murray JT, Nicolson SW, Osborne JL, Paxton RJ, Pirk CWW, Polce C, Potts SG, Priest NK, Raine NE, Roberts S, Ryabov EV, Shafir S, Shirley MDF, Simpson SJ, Stevenson PC, Stone GN, Termansen M, Wright GA. 2013.** Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment* **11**:251–259 DOI [10.1890/120126](https://doi.org/10.1890/120126).
- Visscher PK. 1983.** The honey bee way of death: necrophoric behaviour in *Apis mellifera* colonies. *Animal Behaviour* **31**:1070–1076.
- Wachendorff-Neumann U, Mauler-Machnik A, Erdelen C, Ohtake H. 2012.** Synergistic mixture of trifloxystrobin and imidacloprid. Google patents. Bayer CropScience AG, United States.
- Williamson SM, Wright GA. 2013.** Exposure to multiple cholinergic pesticides impairs olfactory learning and memory in honeybees. *The Journal of Experimental Biology* **216**:1799–1807 DOI [10.1242/jeb.083931](https://doi.org/10.1242/jeb.083931).
- Winston ML. 1991.** *The biology of the honey bee*. Cambridge: Harvard University Press.
- Woodcock BA, Bullock JM, Shore RF, Heard MS, Pereira MG, Redhead J, Ridding L, Dean H, Sleep D, Henrys P, Peyton J, Hulmes S, Hulmes L, Sárospataki M, Saure C, Edwards M, Genersch E, Knäbe S, Pywell RF. 2017.** Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science* **356**:1393–1395 DOI [10.1126/science.aaa1190](https://doi.org/10.1126/science.aaa1190).
- Yu SJ. 2014.** *The toxicology and biochemistry of insecticides*. Boca Raton: CRC Press.
- Zhu W, Schmehl DR, Mullin CA, Frazier JL. 2014.** Four common pesticides, their mixtures and a formulation solvent in the hive environment have high oral toxicity to honey bee larvae. *PLOS ONE* **9**:e77547 DOI [10.1371/journal.pone.0077547](https://doi.org/10.1371/journal.pone.0077547).