

# Inferring Past Environments from Ancient Epigenomes

David Gokhman,<sup>†,1</sup> Anat Malul,<sup>†,1</sup> and Liran Carmel<sup>\*,1</sup>

<sup>1</sup>Department of Genetics, The Alexander Silberman Institute of Life Sciences, Faculty of Science, The Hebrew University of Jerusalem Edmond J. Safra Campus, Givat Ram, Jerusalem, Israel

<sup>†</sup>These authors contributed equally to this work.

\*Corresponding author: E-mail: liran.carmel@huji.ac.il.

Associate editor: Connie Mulligan

## Abstract

Analyzing the conditions in which past individuals lived is key to understanding the environments and cultural transitions to which humans had to adapt. Here, we suggest a methodology to probe into past environments, using reconstructed premortem DNA methylation maps of ancient individuals. We review a large body of research showing that differential DNA methylation is associated with changes in various external and internal factors, and propose that loci whose DNA methylation level is environmentally responsive could serve as markers to infer about ancient daily life, diseases, nutrition, exposure to toxins, and more. We demonstrate this approach by showing that hunger-related DNA methylation changes are found in ancient hunter-gatherers. The strategy we present here opens a window to reconstruct previously inaccessible aspects of the lives of past individuals.

**Key words:** ancient DNA, ancient epigenetics, DNA methylation, environmental paleoepigenetics, environmental epigenetics, paleoepigenetics.

## Introduction

In the past tens of thousands of years humans went through dramatic shifts in their life style and environment. Examples include the exodus from Africa, which exposed humans to new climatic conditions, pathogens, and nutritional sources; the transition from hunting and gathering to farming, which altered nutritional composition, culture, and social structure; and the advent of technology (Ermini et al. 2015). Thus, the environment of a human individual thousands of years ago was very different from that of a present-day human. Here, we use the term environment to describe the host of extrinsic and intrinsic conditions affecting an individual, such as nutrition, diseases, social interactions, psychological state, physical activities, ambient temperature, altitude, and exposure to toxins. In order to understand the nature of the challenges that faced ancient humans and the changes and adaptations that followed them, it is critical to develop means to study the environment in which these past individuals lived.

Until recently, methods to investigate the history and life style of an individual were mainly archaeological and paleo-anthropological (Klein 2000). Through findings such as bones, ornaments, and tools, researchers have reconstructed aspects of ancient human daily life (Klein 2000). For example, it was shown that Neanderthals used manganese oxides to lower the auto-ignition temperature of wood for fire-making (Heyes et al. 2016); and bone fossils and lithic findings in excavations in Mount Carmel shed light on tool production technology and burial rituals (Bar-Yosef et al. 1992). Another approach which allows inference on ancient life is the study of present-day hunter-gatherers. Often criticized, such studies rely on the idea that some ancient practices survived to this day, or have

otherwise been developed in parallel in these populations (Hawkes et al. 1997; Barnard 1998). Though valuable and informative in many ways, the above approaches have some known limitations (Justeson 1973). For example, archaeological analyses rely on preserved artifacts, with a bias towards those which tend to be preserved better. Moreover, the interpretation of findings is sometimes subjective and done through modern eyes (Justeson 1973). Therefore, the study of past environments could benefit from the introduction of new and complementary strategies that will be integrated with the collection of existing methods.

Progress in ancient DNA sequencing technology (Orlando et al. 2015; Krause and Paabo 2016) allowed the development of new ways to study ancient environments. For example, shotgun sequencing of ancient DNA from the calcified dental plaque (calculus) of five Neanderthals allowed inference about variation in Neanderthal diet, health, and microbiome (Weyrich et al. 2017). Similarly, sequencing of 34 early European calculi revealed how the shift from hunting and gathering to farming manifested in alterations in the oral microbiome (Adler et al. 2013). Sequencing of ancient DNA from the *Yersinia pestis* bacteria, which caused recurrent plague pandemics such as the Black Death (14–17th centuries AD), revealed the evolutionary dynamics and epidemiology of these plagues (Schuenemann et al. 2011; Wagner et al. 2014; Rasmussen et al. 2015).

Technological developments in ancient DNA sequencing were also used to genotype, or even fully sequence, the genomes of the ancient individuals themselves, using DNA extracted from bones, hair, and teeth. As of today, DNA was successfully sequenced from hundreds of individuals that

lived from medieval times to tens of thousands of years ago (for example see Fu et al. 2014; Gamba et al. 2014; Raghavan et al. 2014; Seguin-Orlando et al. 2014; Allentoft et al. 2015; Haak et al. 2015; Mathieson et al. 2015; Raghavan et al. 2015; Gallego-Llorente et al. 2016; Lazaridis et al. 2016; Skoglund et al. 2016; González-Forbes et al. 2017). Several sequenced individuals belong to extinct human groups such as the Neanderthal (Green et al. 2010; Prüfer et al. 2014) and the Denisovan (Reich et al. 2010; Meyer et al. 2012), see reviews in (Orlando et al. 2015; Krause and Paabo 2016; Skoglund and Reich 2016; Llamas et al. 2017; Nielsen et al. 2017). Using these genomes, researchers were able to reveal historical events affecting human demographics, such as patterns of migrations, the spread of diseases, population admixtures, and replacements, and adaptation to agricultural lifestyle, colder climates, and high altitudes (Huerta-Sánchez et al. 2014; Allentoft et al. 2015; Mathieson et al. 2015; Lazaridis et al. 2016; Racimo et al. 2017).

Yet, the genetic makeup of populations changes slowly, over many generations, and hence tracking down DNA changes may provide information on long-term processes, but can rarely be used as a tool to study responses to new environments (Yona et al. 2015). In contrast, epigenetics—defined as somatically heritable chemical modifications of the DNA that do not entail alterations to the sequence itself—is plastic and responsive to the environment. As epigenetic layers react to external and internal conditions through modulation of their pattern of chemical modifications, they are sometimes referred to as “middleman” between the environment and the DNA (Feil and Fraga 2012; Marsit 2015; Rubin 2015; Yona et al. 2015; Etchegaray and Mostoslavsky 2016). Therefore, identifying environmentally responsive loci (ERLs) of epigenetic layers in ancient individuals has the potential to serve as a novel means to investigate the conditions in which they lived. Here, we define an ERL as any locus whose epigenetic pattern carries information on environmental cues, regardless of whether this locus responds directly to the environment, or is simply associated with it.

Epigenetic patterns play a pivotal role in determining and marking the activity of genes. The term “Epigenetics” refers to several regulatory layers, including histone modifications, chromatin remodeling, nucleosome positioning, and DNA methylation. In this paper, we focus on DNA methylation—the addition of a methyl group to a cytosine nucleotide. In mammals, DNA methylation usually occurs in the context of a cytosine that is followed by a guanine, thus named CpG position. Non-CpG (i.e., CT, CA, or CC) methylation also occurs in mammals, mainly in pluripotent and brain cells, but is otherwise rare (Ziller et al. 2011; Varley et al. 2013). The role of DNA methylation in gene regulation depends on its position and context. In promoters it is generally associated with silencing of genes, whereas in other regions it plays a role in modulating enhancer activity, X-chromosome inactivation, splicing, transposable element silencing, and more (Feil and Fraga 2012; Jones 2012).

Methylation patterns along a genome are determined by two main factors. The first is genetic; much of the methylome is determined directly by the DNA through the binding of

methylation-altering factors to specific sequences (Lienert et al. 2011; Teh et al. 2014). Being chiefly dictated by DNA sequence, such methylation patterns persist across generations, and are, by large, less responsive to environmental changes within the lifetime of an individual (Feil and Fraga 2012; Jones 2012; Yuan 2012; Whitaker et al. 2015).

The second factor is external and internal environmental signals that alter the methylome, such as changes in caloric intake, psychological trauma, and *in utero* exposure to toxins. The mechanisms mediating environmental cues and methylation changes are poorly understood, but they are often associated with DNA-binding activity, or affect one-carbon metabolic pathways that dictate the availability of methyl donors (Jaenisch and Bird 2003; Feil and Fraga 2012). Regardless of the underlying mechanisms, the accumulation of whole-genome methylation maps, combined with an increasing interest in the relationship between the environment and the methylome, lead to the identification of a growing number of ERLs and the factors that shape them (Bollati and Baccarelli 2010; Feil and Fraga 2012).

Therefore, given a sufficiently long list of ERLs, and their methylation levels in ancient individuals, we could identify “environmental signatures” that would shed light on the conditions in which these individuals lived. Such inference requires precise knowledge of ERLs and the environmental changes they are associated with (environmental epigenetics), as well as an ability to infer accurate epigenetic patterns from ancient DNA (paleoepigenetics). As of today, both of these fields are in their infancy. Environmental epigenetics still lacks sufficient knowledge about the mechanisms and factors that shape the methylome, and faces challenges in designing well-controlled experiments. Paleoepigenetics still relies on a small set of high-quality methylomes that come from a limited repertoire of tissues. However, as both fields are rapidly expanding, we see great value in combining them into what we term environmental paleoepigenetics. The development of environmental paleoepigenetics is a forward-looking research goal that could contribute to the study of past environments by integrating multiple lines of evidence from archeological, paleontological, microbial, and paleoepigenetic approaches.

In this review, we examine the use of paleoepigenetic techniques to reconstruct premortem methylation maps from ancient DNA. We then offer guidelines how to identify ERLs that could be useful for environmental paleoepigenetics. Next, we survey works that reported such potentially useful ERLs, mainly in response to diseases, changes in nutrition, and exposure to toxins. Finally, we point out the presence of hunger-related ERLs in some ancient individuals, and discuss how this may be interpreted.

## Reconstructing Ancient Methylation Maps

The spontaneous deamination of cytosines is a central chemical degradation process of ancient DNA. This process produces distinct degradation signatures for methylated and for unmethylated cytosines, whereby methylated cytosines are deaminated into thymines, whereas unmethylated cytosines

are deaminated into uracils. During library preparation uracils are removed, leading to enrichment of thymines in premortem methylated positions. Using this signal, premortem methylation maps of ancient individuals can be reconstructed (Briggs et al. 2010; Gokhman et al. 2014; Pedersen et al. 2014; Hanghøj et al. 2016). We have previously used this idea to reconstruct the full methylomes of our closest extinct relatives—the Neanderthal and the Denisovan (Gokhman et al. 2014). Comparing these maps to a modern bone methylation map, we identified ~2,000 differentially methylated regions (DMRs), providing information on the genes that are differentially regulated between these human groups.

### Which ERLs Could Be Useful in Environmental Paleoepigenetics?

Using ancient DNA methylation patterns to infer about the environment of an individual that died thousands of years ago must address the fact that DNA methylation is tissue-specific. DNA does not survive long in most tissues, and the majority of ancient DNA material comes from bone, hair, and teeth (Orlando et al. 2015; Gokhman et al. 2016). This means that we can reconstruct the methylomes of a limited number of tissues, which are not necessarily those whose methylation had changed in response to an environmental cue. For instance, an individual exposed to dietary stress may exhibit DNA methylation alterations in the brain or adipose tissues, but not necessarily in bone tissue.

At first glance, this renders the use of bones, teeth, and hair for inference about environmental signatures in other tissues inadvisable. However, it has been shown that some genomic regions establish their methylation patterns early in embryogenesis (Roemer et al. 1997; Rakyán et al. 2002; Dolinoy et al. 2007). Therefore, if an environmentally induced methylation change happened early during embryogenesis, prior to the differentiation of tissues, then this change would be carried on to the daughter cells and tissues (Rakyán et al. 2002; Dolinoy et al. 2007; Gokhman et al. 2016). Thus, timing is critical to the information-content of methylation in ERLs: the earlier in development an epigenetic alteration occurred, the more cell types it is expected to affect.

A special case of methylation changes following early exposure is that of metastable epialleles. These are loci that are differentially methylated between genetically identical individuals, but show low variability between tissues of the same individual. These properties suggest that their methylation state is established early in development, and that they are particularly influenced by environmental conditions during early pregnancy. Metastable epialleles may be identified by comparing DNA methylation maps in different tissues across individuals, for example comparing monozygotic twins or comparing individuals from different controlled environments. The methylation state of metastable epialleles was described as an environmentally dependent stochastic event that results in cell variegation, and in some cases affects gene expression and phenotypes. Metastable epialleles have been shown to affect coat color, transgene expression in myocytes,

neural tube development, and more (Rakyán et al. 2002; Dolinoy et al. 2007; Waterland and Michels 2007; Alan Harris et al. 2013). Regardless of their phenotypic effect, the combination of high susceptibility to environmental conditions, low intertissue variability, and lifelong stability makes metastable epialleles good candidates to serve as loci whose methylation state in bones, teeth, and hair could be a proxy to the conditions during early embryogenesis of an archaic individual.


Importantly, the idea of using epigenetics to infer about past environments does not rely on whether the epigenetic modifications are heritable across generations. The extent to which DNA methylation in mammals is heritable across generations is debated (Heard and Martienssen 2014), but an ERL could serve as an environmental marker regardless of whether it emerged during the lifetime of an individual or passed to him/her from past generations. It is important to emphasize that if an ERL was shown to be transgenerationally inherited, it could probably serve as a good marker for environmental paleoepigenetics, because its methylation state exists in the predifferentiated cells of the early embryo and is therefore likely to be found in bones, teeth, and hair as well.

### The Effects of Nutrition, Toxins, and Diseases on the Methylome

Environmental epigenetics investigates how different factors such as diseases, exposure to toxins, and nutrition shape the epigenome. We suggest environmental paleoepigenetics build upon this growing body of knowledge to address the reverse process—analyzing patterns of ancient epigenomes and deducing what environmental factors might underlie them (fig. 1).

Using information embedded in ancient methylation maps to infer on nonenvironmental factors has already been achieved with regard to age and tissue type. Pedersen et al. (2014) used the reconstructed methylation map of an ancient Saqqaq Eskimo in order to infer his age at death, and we have previously used the reconstructed Neanderthal and Denisovan methylomes in order to show that their DNA came from bone cells (Gokhman et al. 2014).

Below, we survey studies in environmental epigenetics that reported links between DNA methylation changes and environmental factors. We use two criteria to determine the scope of studies we cover. First, we focus on nutrition, toxins, and diseases, as these are extensively studied factors which are particularly relevant to environmental reconstruction. Second, we spotlight ERLs that might be useful in environmental paleoepigenetics, namely ERLs that are either 1) established early in development; 2) transmitted transgenerationally; or 3) influenced by environmental factors in tissues from which ancient DNA usually originates. As transgenerational inheritance of DNA methylation in mammals is still contested, and as environmental epigenetics rarely looks at tissues such as bones, teeth, and hair, the vast majority of ERLs reported here stem from environmental changes that affect the early embryo. In addition, we include studies involving nonhuman mammals, especially rodents and primates.



**Fig. 1.** Environmental paleoepigenetics builds on environmental epigenetics and paleoepigenetics. In environmental epigenetics, researchers study how extrinsic and intrinsic factors affect the epigenome (blue arrows). Paleoepigenetics harnesses degradation signals in ancient DNA to reconstruct premortem DNA methylation maps (green arrows). Environmental paleoepigenetics would use the reconstructed methylation maps of ancient individuals to infer on the unknown extrinsic and intrinsic factors that shaped them (orange arrows).

Although it is still to be determined which of these nonhuman ERLs also exist in humans, they provide a more comprehensive view of the way environmental factors shape the methylome. Additionally, these nonhuman ERLs could be used in the future to study the environments of additional ancient organisms, as nonhuman methylation data (Seguin-Orlando et al. 2015) and high-quality nonhuman ancient genomes (Palkopoulou et al. 2015) are starting to emerge.

Most studies to date examined the nutritional factor. One of the most well-known examples in humans described how periconceptional exposure to famine (the Dutch Hunger Winter of 1944–1945) was associated with hypomethylation of the 5′-UTR of the *IGF2* gene (Heijmans et al. 2008). Later, it was shown that many other loci change their methylation in response to famine, including regions in the *INSIGF*, *IL10*, *ABCA1*, *GNASAS*, *MEG3*, and *LEP* genes (Tobi et al. 2009). On the other end, high-fat diet (HFD) was found to significantly increase methylation levels in mice at 30 CpGs in intron 2 of *igf2r* (Gallou-Kabani et al. 2010). In rats, paternal HFD drove hypermethylation of a CpG site upstream of *Il13ra2* in female offspring (Ng et al. 2010). As may be expected, the level of methyl-donor compounds in the diet was also shown to affect methylation: Waterland et al. supplemented the diet of pregnant female mice with extra folic acid, vitamin B<sub>12</sub>, choline, and betaine—all methyl-donor-rich

foods—resulting in hypermethylation of several loci in the offspring (Waterland and Jirtle 2003; Waterland et al. 2006). See supplementary table 1, Supplementary Material online, for additional studies of nutrition–methylation relationship.

A recent study separated the effects of genetic background and environment on the methylome of human populations, by comparing Central African populations that differ in genetic background, lifestyle, or both (Fagny et al. 2015). The study looked at the genotypes and whole blood methylomes of hundreds of individuals from five populations. Among the investigated populations were West African rainforest hunter-gatherers, West African agrarians, and West African agrarians that readopted life in the rainforest. Although the first and second populations differ by both ancestry and lifestyle, the second and third populations differ mostly by lifestyle. The authors identified >3,000 differentially methylated sites that could be attributed to the recent change of habitat, and showed that they are overrepresented in genes involved in immune response, host–pathogen interactions, and various cellular processes. The developmental stage at which these environmental cues leave a mark on the methylome is still to be determined.

Another factor that was shown to alter DNA methylation at various sites was exposure to toxins and pollutants. *In utero* exposure to smoking was associated with alterations in



methylation across various genes, including hypermethylation in *BDNF* (Toledo-Rodriguez et al. 2010), *AXL*, and *PTPRO* (Breton et al. 2009) and hypomethylation in *CYP1A1* (Suter et al. 2010) and in the repetitive element *AluYb8* (Breton et al. 2009). Other compounds that triggered local changes in methylation include ethanol (Kim et al. 2014), arsenic (Intarasunanont et al. 2012), and cadmium (Vidal et al. 2015).

The last group of factors that interacts with DNA methylation is diseases and disorders. In this regard, the interaction between methylation and diseases is bidirectional. On one hand, many studies have shown how changes in methylation mark or even drive a wide variety of disorders (Robertson 2005; Bergman and Cedar 2013; Chen et al. 2016; Li et al. 2016; Martino et al. 2016; Reynard 2016; Wüllner et al. 2016). On the other hand, it was shown how maternal diseases leave epigenetic marks in the offspring. In the latter case, most studies have focused on diabetes and showed localized changes in specific genes, as well as genome-wide differences affecting thousands of genes. For example, children of type 2 diabetic mothers had over 4,000 differentially methylated regions (Del Rosario et al. 2014), and offspring to mothers with gestational diabetes mellitus had lower methylation levels in several CpG sites in the lipoprotein lipase gene (*LPL*) (Houde et al. 2014), hypermethylation in leptin promoters (Lesseur et al. 2014), and in over 1,000 additional genes (Ruchat et al. 2013). Regardless of the directionality of effect (methylation changes that drive diseases or diseases that drive methylation changes), these studies open a window to investigate the health profile of past individuals.

Perhaps the most intriguing factors that affect DNA methylation are psychological and social. Most studies that looked into these factors examined methylation patterns associated with behavioral, psychological, or social events at late stages of life (supplementary table 1, Supplementary Material online). However, studies investigating prenatal maternal stress, such as the Great Ice Storm of 1998 (Cao-Lei et al. 2014), war stress in the Democratic Republic of Congo (Mulligan et al. 2012), intimate partner violence (Radtke et al. 2011), and depression during pregnancy (Liu et al. 2012), provide insight into methylation changes that are associated with stress exposure in early stages of development. Finally, physical exercise, as well as external conditions such as temperature, humidity, and altitude, has been repeatedly associated with various ERLs. However, to our knowledge, these studies have all been conducted on adults, and on tissues other than teeth, hair, and bone, and thus, their relevance environmental paleoepigenetics is still to be determined.

Additional studies that investigated ERLs and effects of the environment on global DNA methylation level are summarized in supplementary table 1, Supplementary Material online.


### Methylation Patterns of Hunter-Gatherers and Archaic Humans Are Consistent with Low Caloric Intake

Environmental epigenetics is still a young field, and as of today the number of ERLs that can be used for inference on past

environments is very small. One of the goals of this review is to encourage researchers in the field to dedicate efforts to identifying ERLs suitable to environmental paleoepigenetics. As a further encouragement, we provide here a small-scale demonstration of the potential of environmental paleoepigenetics to enrich our understanding of the conditions under which humans lived thousands of years ago.

In a study by Dominguez-Salas et al., researchers focused on a rural Gambian population where caloric intake varies considerably according to season, with a “hungry season” followed by a “harvest season.” The “hungry season” is the rainy season, characterized by restricted protein-energy availability, whereas the dry “harvest season” does not hold any nutritional stress. The oscillations in nutrient availability have been shown to affect *in utero* development and growth (Rayco-Solon et al. 2005). These studies found that children conceived during the “hungry season” showed hypermethylation at six metastable epialleles, residing near or within the following genes: *LOC654433*, *EXD3*, *RBM46*, *BOLA3*, *ZNF678*, and *ZFYVE28* (Waterland et al. 2010; Dominguez-Salas et al. 2014). In follow-up work on these data, we crossed these six ERLs with a list of DMRs found between archaic humans (Neanderthal and Denisovan) and 21 present-day humans (Gokhman et al. 2014). We found that three of the ERLs partially or completely overlap DMRs (in *EXD3*, *RBM46*, and *ZNF678*) that are significantly hypermethylated in the Neanderthal, the Denisovan, or both (fig. 2a–c). The observed overlap is significantly higher than would be expected by chance, even when controlling for GC content ( $P < 10^{-6}$ , randomization test, 1,000,000 iterations). Also, these methylation differences are unlikely to be driven by sequence changes, as the closest single nucleotide change or indel that differentiate the hominin groups are found tens of thousands of bases away (23 kb for *RBM46*, 84 kb for *EXD3*, and 125 kb for *ZNF678*), whereas methylation-affecting polymorphisms tend to be much closer to DMRs (with a peak enrichment at 45 bp) (Gibbs et al. 2010). Since the ERLs were found within metastable epialleles, they are likely to reflect not only the methylation state in the tissues where they were measured, but rather across tissues, including bones and teeth (Rakyan et al. 2002; Dolinoy et al. 2007; Waterland and Michels 2007; Waterland et al. 2010; Alan Harris et al. 2013; Dominguez-Salas et al. 2014). It is important to note that the present-day methylation maps used in order to identify these DMRs (Gokhman et al. 2014) were produced by reduced representation bisulfite sequencing (RRBS), and thus provide information for only ~10% of CpGs in the genome. Therefore, further research is needed in order to fully characterize the extent of these DMRs.

To explore whether the differences between archaic and present-day humans could reflect the transition from hunting and gathering to farming, we looked at additional published DNA methylation maps of modern and ancient individuals, totaling five hunter-gatherers and four sedentary individuals (Gokhman et al. 2017). We found that these ERLs are hypermethylated by 7% on average in skeletal tissues of hunter-gatherers ( $P = 7.3 \times 10^{-5}$ , *t*-test), similarly to the effect size detected in the Gambian population study



**Fig. 2.** Methylation patterns in the Neanderthal and the Denisovan point to a low-calorie diet. Methylation maps are shown for a present-day human, a Neanderthal, and a Denisovan. Each line represents a CpG position. Methylation levels are color-coded from green (unmethylated) to red (methylated). Present-day human maps are partial because the protocol used to produce the maps was reduced representation bisulfite sequencing (RRBS), which provides information for  $\sim 10\%$  of CpG positions. Reconstructed ancient methylation and DMRs were taken from (Gokhman et al. 2014). ERLs were taken from Dominguez-Salas et al. (2014). (a) Archaic humans are hypermethylated in the *EXD3* gene compared with the present-day human. The DMR completely overlaps the ERL, where hypermethylation is associated with low-calorie diet. (b) The Neanderthal is hypermethylated in the *RBM46* gene compared with the present-day human. The DMR partially overlaps with the ERL, where hypermethylation is associated with low-calorie diet. (c) The Denisovan is hypermethylated upstream of the *ZNF678* gene compared with the present-day human. The DMR completely overlaps the ERL, where hypermethylation is associated with low-calorie diet. (d) Box plots of methylation levels of hunter-gatherers and sedentary individuals within the six hunger-related ERLs. In *LOC654433*, *RBM46*, and *EXD3* hunter-gatherers are significantly hypermethylated compared with sedentary individuals, reflecting possible low caloric intake. Within *BOLA3*, however, hunter-gatherers are hypomethylated.

(Dominguez-Salas et al. 2014). Particularly, we found that *LOC654433*, *RBM46*, and *EXD3* are significantly hypermethylated in hunter-gatherers compared with sedentary individuals, compatible with their hypermethylation in Gambian individuals conceived during the hungry season, whereas *BOLA3* presents the opposite trend, incompatible with hunter state (fig. 2d).

Though further research is needed in order to fully understand the precise factors that trigger such changes in methylation, these preliminary findings point to the possibility that *in utero*, archaic humans, and possibly hunter-gatherers in general, experienced conditions that are somewhat similar to those experienced by present-day individuals conceived during a hungry season. More importantly, this highlights

the potential of using ERLs that are established during early development to investigate the environment in which ancient individuals lived.

## Summary

Although some ERL studies analyzed metastable epialleles or ERLs that are established early in development, many others focused on a specific tissue of interest and on regions that have not been tested for stability across tissues. In this review, we have focused on studies that identified ERLs in early stages of development, and are therefore likely to be shared by many tissues. However, in order to understand the extent to which these ERLs persist in different cell types, and specifically, to be able to test them in tissues from which ancient DNA samples are collected, it is imperative to conduct further studies which cover a large array of tissues and individuals.

Importantly, although the mechanisms that drive the changes in methylation and their resulting phenotypic effects are of great interest, they are practically irrelevant for environmental reconstruction. These ERLs serve only as markers for the conditions to which an individual was exposed. Thus, in this regard, it is irrelevant whether they are only correlated with specific environmental conditions, or directly driven by them, and whether they have phenotypic consequences, or are completely neutral.

A single ERL may respond to more than one environmental cue. For example, *NR3C1*, *SLC6A4*, and *IGF2* have all been shown to be influenced by multiple factors (supplementary table 1, Supplementary Material online). It is therefore critical to identify a full complement of ERLs that respond to each extrinsic or intrinsic factor, so as to obtain an “ERL signature” to each factor. Such signatures would allow reconstruction of past environments, whereas taking into account pleiotropic effects as well as correlation between various factors affecting the same locus. In this way, each ERL may be viewed as a broadly tuned environmental sensor, and each environmental cue may be associated with a specific pattern of ERL responses.

It is also important to consider effect size. Although some studies report considerable effect sizes, for example 24.4% methylation change in *AHRR* (Zeilinger et al. 2013), and 30% change in *GF11* (Elliott et al. 2014), both in response to smoking, most effect sizes are smaller, sometimes as little as 1% (Fagny et al. 2015). This poses a particular challenge when analyzing reconstructed methylation maps, where signals are smoothed using a sliding window, and where the variance in predicting methylation level depends on how well we can estimate the deamination rate of methylated cytosines (Gokhman et al. 2016). For this reason, environmental paleoepigenetics should focus on higher effect sizes or longer ERLs where the smoothing effect is mitigated. Environmental inference where effect sizes are small would probably require the use of methylation maps from many individuals, which would increase statistical power.

DNA sequence can directly affect methylation states through factors such as CpG density and the activity of binding proteins (Lienert et al. 2011). Genetic changes affecting

DNA methylation patterns may be detected through various means, for example by identifying methylation quantitative trait loci (meQTLs) (ZhaNg et al. 2010). The genetic background of an individual can therefore affect the way his/her ERLs respond to environmental cues (Teh et al. 2014). Genotype–methylation–environment interplay is well demonstrated in the *FKBP5* gene, where the risk for childhood trauma-dependent demethylation is increased by a polymorphism that affects the chromatin interaction between the enhancers of *FKBP5* and its transcription start site (Klengel et al. 2012). Therefore, methylation-based environmental inferences, and specifically those that include samples from different ancestries, should consider potential genetic effects, which tend to reside close to the site of differential methylation (Gibbs et al. 2010).

Our focus in this perspective was on humans. However, the use of ERLs in order to reconstruct past environments could be implemented on additional methylation maps, such as those of nonhuman mammals (Llamas et al. 2012) and even plants (Smith et al. 2014). Many studies have demonstrated how environmental factors, such as temperature, alter DNA methylation patterns in plants (Feil and Fraga 2012). Thus, we anticipate that ancient plant methylation maps (Smith et al. 2014), as well as nonhuman mammalian methylation maps (Seguin-Orlando et al. 2015), could be analyzed to infer the environmental conditions to which they were exposed thousands of years ago.

The motivation behind many of the above studies was mainly medical—to understand how different environmental factors affect our physical and mental health. This field is growing rapidly, and our understanding of how the environment shapes our epigenome is becoming deeper and more complex. In parallel, the fields of paleogenetics and paleoepigenetics are gaining momentum (Hanghøj et al. 2016), with ancient DNA samples being sequenced in increasing numbers, to higher quality and from older periods (Orlando et al. 2015; Krause and Paabo 2016). We believe that the integration of these two fields will have a synergistic effect, opening new angles to explore how the lives of archaic individuals looked like. Once enough knowledge is gained in both fields, we believe that the epigenomes of ancient individuals could reveal aspects of their daily life, diseases from which they suffered, substances to which they were exposed, and possibly even their psychosocial state.

## Supplementary Material

Supplementary data are available at *Molecular Biology and Evolution* online.

## Acknowledgements

We thank Shiran Bar for useful advice. The work was supported by the Israel Science Foundation FIRST individual grant (ISF 1430/13 to L.C.). D.G. is supported by the Clore Israel Foundation.



## References

- Adler CJ, Dobney K, Weyrich LS, Kaidonis J, Walker AW, Haak W, Bradshaw CJA, Townsend G, Sołtysiak A, Alt KW. 2013. Sequencing ancient calcified dental plaque shows changes in oral microbiota with dietary shifts of the Neolithic and industrial revolutions. *Nat Genet.* 45(4): 450–455.
- Alan Harris R, Nagy-Szakal D, Kellermaier R. 2013. Human metastable epiallele candidates link to common disorders. *Epigenetics* 8(2): 157–163.
- Allentoft ME, Sikora M, Sjögren K-G, Rasmussen S, Rasmussen M, Stenderup J, Damgaard PB, Schroeder H, Ahlström T, Vinner L. 2015. Population genomics of Bronze Age Eurasia. *Nature* 522(7555): 167–172.
- Barnard A. 1998. The foraging spectrum: diversity in hunter-gatherer lifeways. *Am Ethnologist* 25: 36–37.
- Bar-Yosef O, Vandermeersch B, Arensburg B, Belfer-Cohen A, Goldberg P, Laville H, Meignen L, Rak Y, Speth JD, Tchernov E, et al. 1992. The excavations in Kebara Cave, Mt. Carmel. *Curr Anthropol.* 33: 497.
- Bergman Y, Cedar H. 2013. DNA methylation dynamics in health and disease. *Nat Struct Mol Biol.* 20(3): 274–281.
- Bollati V, Baccarelli A. 2010. Environmental epigenetics. *Heredity* 105(1): 105–112.
- Breton CV, Byun HM, Wenten M, Pan F, Yang A, Gilliland FD. 2009. Prenatal tobacco smoke exposure affects global and gene-specific DNA methylation. *Am J Respir Crit Care Med.* 180(5): 462–467.
- Briggs AW, Stenzel U, Meyer M, Krause J, Kircher M, Pääbo S. 2010. Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res.* 6: e87.
- Cao-Lei L, Massart R, Suderman MJ, Machnes Z, Elgbeili G, Laplante DP, Szyf M, King S, Iwamoto K. 2014. DNA methylation signatures triggered by prenatal maternal stress exposure to a natural disaster: project ice storm. *PLoS One* 9(9): e107653.
- Chen L, Ge B, Casale FP, Vasquez L, Kwan T, Garrido-Martín D, Watt S, Yan Y, Kundu K, Ecker S, et al. 2016. Genetic drivers of epigenetic and transcriptional variation in human immune cells. *Cell* 167(5): 1398–1414.e24.
- Del Rosario MC, Ossowski V, Knowler WC, Bogardus C, Baier LJ, Hanson RL. 2014. Potential epigenetic dysregulation of genes associated with MODY and type 2 diabetes in humans exposed to a diabetic intra-uterine environment: an analysis of genome-wide DNA methylation. *Metab Clin Exp.* 63: 654–660.
- Dolinoy DC, Das R, Weidman JR, Jirtle RL. 2007. Metastable epialleles, imprinting, and the fetal origins of adult diseases. *Pediatr Res.* 61(5 Part 2): 30R.
- Dominguez-Salas P, Moore SE, Baker MS, Bergen AW, Cox SE, Dyer RA, Fulford AJ, Guan Y, Laritsky E, Silver MJ, et al. 2014. Maternal nutrition at conception modulates DNA methylation of human metastable epialleles. *Nat Commun.* 5: 3746.
- Elliott HR, Tillin T, McArdle WL, Ho K, Duggirala A, Frayling TM, Davey Smith G, Hughes AD, Chaturvedi N, Relton CL. 2014. Differences in smoking associated DNA methylation patterns in South Asians and Europeans. *Clin Epigenetics* 6(1): 4.
- Ermimi L, Der Sarkissian C, Willerslev E, Orlando L. 2015. Major transitions in human evolution revisited: a tribute to ancientDNA. *J Hum Evol.* 79: 4–20.
- EtcheGARAY J, Mostoslavsky R. 2016. Interplay between metabolism and epigenetics: a nuclear adaptation to environmental changes. *Mol Cell* 62(5): 695–711.
- Fagny M, Patin E, Maclsaac JL, Rotival M, Flutre T, Jones MJ, Siddle KJ, Quach H, Harmant C, McEwen LM, et al. 2015. The epigenomic landscape of African rainforest hunter-gatherers and farmers. *Nat Commun.* 6: 10047.
- Feil R, Fraga MF. 2012. Epigenetics and the environment: emerging patterns and implications. *Nat Rev Genet.* 13(2): 97–109.
- Fu Q, Li H, Moorjani P, Jay F, Slepchenko SM, Bondarev AA, Johnson PLF, Aximu-Petri A, Prüfer K, de Filippo C, et al. 2014. Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* 514(7523): 445–449.
- Gallego-Llorente M, Connell S, Jones ER, Merrett DC, Jeon Y, Eriksson A, Siska V, Gamba C, Meiklejohn C, Beyer R, et al. 2016. The genetics of an early Neolithic pastoralist from the Zagros, Iran. *Sci Rep.* 6: 31326.
- Gallou-Kabani C, Gabory A, Tost J, Karimi M, Mayeur S, Lesage J, Boudadi E, Gross MS, Taurelle J, Vigé A, et al. 2010. Sex- and diet-specific changes of imprinted gene expression and dna methylation in mouse placenta under a high-fat diet. *PLoS One* 5(12): e14398.
- Gamba C, Jones ER, Teasdale MD, McLaughlin RL, Gonzalez-Fortes G, Mattiangeli V, Domboróczki L, Kóvári I, Pap I, Anders A, et al. 2014. Genome flux and stasis in a five millennium transect of European prehistory. *Nat Commun.* 5: 5257.
- Gibbs JR, van der Brug MP, Hernandez DG, Traynor BJ, Nalls MA, Lai S-L, Arepalli S, Dillman A, Rafferty IP, Troncoso J, et al. 2010. Abundant quantitative trait loci exist for DNA methylation and gene expression in human brain. *PLoS Genet.* 6(5): e1000952.
- Gokhman D, Agranat-Tamir L, Housman G, García-Pérez R, Nissim-Rafinia M, Mallick S, Nieves-Colón MA, Gu H, Ferrando-Bernal M, Gelabert P, et al. 2017. Recent regulatory changes shaped human facial and vocal anatomy. bioRxiv 106955. doi: <https://doi.org/10.1101/106955>.
- Gokhman D, Lavi E, Prüfer K, Fraga MF, Riancho JA, Kelso J, Pääbo S, Meshorer E, Carmel L. 2014. Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science* 344(6183): 523–527.
- Gokhman D, Meshorer E, Carmel L. 2016. Epigenetics: it's getting old. Past meets future in Paleoeugenetics. *Trends Ecol Evol.* 31(4): 290.
- González-Fortes G, Jones ER, Lightfoot E, Bonsall C, Lazar C, Grandal-d'Anglade A, Garralda MD, Drak L, Siska V, Simalcsik A, et al. 2017. Paleogenomic evidence for multi-generational mixing between neolithic farmers and mesolithic hunter-gatherers in the lower Danube Basin. *Curr Biol.* 27(12): 1801–1810.e10.
- Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz MH-Y, et al. 2010. A draft sequence of the Neandertal genome. *Science* 328(5979): 710–722.
- Haak W, Lazaridis I, Patterson N, Rohland N, Mallick S, Llamas B, Brandt G, Nordenfelt S, Harney E, Stewardson K, et al. 2015. Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* 522(7555): 207–211.
- Hanghøj K, Seguin-Orlando A, Schubert M, Madsen T, Pedersen JS, Willerslev EOL. 2016. Fast, accurate and automatic ancient nucleosome and methylation maps with epiPALEOMIX. *Mol Biol Evol* 33(12): 3284–3298.
- Hawkes K, O'Connell JF, Rogers L. 1997. The behavioral ecology of modern hunter-gatherers, and human evolution. *Trends Ecol Evol.* 12(1): 29–32.
- Heard E, Martienssen RA. 2014. Transgenerational epigenetic inheritance: myths and mechanisms. *Cell* 157(1): 95–109.
- Heijmans BT, Tobin EW, Stein AD, Putter H, Blauw GJ, Susser ES, Slagboom PE, Lumey LH. 2008. Persistent epigenetic differences associated with prenatal exposure to famine in humans. *Proc Natl Acad Sci U S A.* 105(44): 17046–17049.
- Heyes PJ, Anastasakis K, de Jong W, van Hoesel A, Roebroeks W, Soressi M. 2016. Selection and use of manganese dioxide by Neanderthals. *Sci Rep.* 6: 22159.
- Houde AA, St-Pierre J, Hivert MF, Baillargeon JP, Perron P, Gaudet D, Brisson D, Bouchard L. 2014. Placental lipoprotein lipase DNA methylation levels are associated with gestational diabetes mellitus and maternal and cord blood lipid profiles. *J Dev Orig Health Dis.* 5(2): 132–141.
- Huerta-Sánchez E, Jin X, Asan, Bianba Z, Peter BM, Vinckenbosch N, Liang Y, Yi X, He M, Somel M. 2014. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature* 512: 194–197.
- Intarasunanont P, Navasumrit P, Waraprasit S, Chaisatra K, Suk WA, Mahidol C, Ruchirawat M. 2012. Effects of arsenic exposure on DNA methylation in cord blood samples from newborn babies and in a human lymphoblast cell line. *Environ Health* 11(1): 31.



- Jaenisch R, Bird A. 2003. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet.* 33(3s): 245–254.
- Jones PA. 2012. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat Rev Genet.* 13(7): 484–492.
- Justeson JS. 1973. Limitations of archaeological inference: an information-theoretic approach with applications in methodology. *Am Antiq.* 38(02): 131.
- Kim P, Choi CS, Park JH, Joo SH, Kim SY, Ko HM, Kim KC, Jeon SJ, Park SH, Han SH, et al. 2014. Chronic exposure to ethanol of male mice before mating produces attention deficit hyperactivity disorder-like phenotype along with epigenetic dysregulation of dopamine transporter expression in mouse offspring. *J Neurosci Res.* 92(5): 658–670.
- Klein RG. 2000. Archeology and the evolution of human behavior. *Evol Anthropol.* 9(1): 17–36.
- Klengel T, Mehta D, Anacker C, Rex-Haffner M, Pruessner JC, Pariante CM, Pace TWW, Mercer KB, Mayberg HS, Bradley B, et al. 2012. Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat Neurosci.* 16(1): 33–41.
- Krause J, Paabo S. 2016. Genetic time travel. *Genetics* 203(1): 9–12.
- Lazaridis I, Nadel D, Rollefson G, Merrett DC, Rohland N, Mallick S, Fernandes D, Novak M, Gamarra B, Sirak K, et al. 2016. Genomic insights into the origin of farming in the ancient Near East. *Nature* 536(7617): 1–22.
- Lesseur C, Armstrong DA, Paquette AG, Li Z, Padbury JF, Marsit CJ. 2014. Maternal obesity and gestational diabetes are associated with placental leptin DNA methylation. *Am J Obstet Gynecol.* 211(6): 654.e1–654.e9.
- Li X, Xiao B, Chen X-S. 2016. DNA methylation: a new player in multiple sclerosis. *Mol Neurobiol* 54(6): 4049–4059.
- Lienert F, Wirbelauer C, Som I, Dean A, Mohn F, Schübeler D. 2011. Identification of genetic elements that autonomously determine DNA methylation states. *Nat Genet.* 43(11): 1091–1097.
- Liu Y, Murphy SK, Murtha AP, Fuemmeler BF, Schildkraut J, Huang Z, Overcash F, Kurtzberg J, Jirtle R, Iversen ES, et al. 2012. Depression in pregnancy, infant birth weight and DNA methylation of imprint regulatory elements. *Epigenetics* 7(7): 735–746.
- Llamas B, Holland ML, Chen K, Cropley JE, Cooper A, Suter CM, Lalueza-Fox C. 2012. High-resolution analysis of cytosine methylation in ancient DNA. *PLoS One* 7(1): e30226.
- Llamas B, Willerslev E, Orlando L. 2017. Human evolution: a tale from ancient genomes. *Philos Trans R Soc B Biol Sci.* 372(1713): 20150484.
- Marsit CJ. 2015. Influence of environmental exposure on human epigenetic regulation. *J Exp Biol.* 218(Pt 1): 71–79.
- Martino DJ, Saffery R, Allen KJ, Prescott SL. 2016. Epigenetic modifications: mechanisms of disease and biomarkers of food allergy. *Curr Opin Immunol.* 42: 9–15.
- Mathieson I, Lazaridis I, Rohland N, Mallick S, Patterson N, Roodenberg SA, Harney E, Stewardson K, Fernandes D, Novak M, et al. 2015. Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* 528(7583): 499–503.
- Meyer M, Kircher M, Gansauge M, Li H, Racimo F, Siebauer M, Green RE, Bryc K, Briggs AW, Stenzel U, et al. 2012. A high-coverage genome sequence from an archaic denisovan individual a high-coverage genome sequence from an archaic denisovan individual a high-coverage genome sequence from an archaic denisovan individual. *Science* 338(6104): 1–14.
- Mulligan CJ, D'Errico NC, Stees J, Hughes DA. 2012. Methylation changes at NR3C1 in newborns associate with maternal prenatal stress exposure and newborn birth weight. *Epigenetics* 7(8): 853–857.
- Ng S-F, Lin RCY, Laybutt DR, Barres R, Owens JA, Morris MJ. 2010. Chronic high-fat diet in fathers programs  $\beta$ -cell dysfunction in female rat offspring. *Nature* 467(7318): 963–966.
- Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E. 2017. Tracing the peopling of the world through genomics. *Nature* 541(7637): 302–310.
- Orlando L, Gilbert MTP, Willerslev E. 2015. Reconstructing ancient genomes and epigenomes. *Nat Rev Genet.* 16(7): 395–408.
- Palkopoulou E, Mallick S, Skoglund P, Enk J, Rohland N, Li H, Omrak A, Vartanyan S, Poinar H, Götherström A, et al. 2015. Complete genomes reveal signatures of demographic and genetic declines in the woolly mammoth. *Curr Biol.* 25(10): 1395–1400.
- Pedersen JS, Valen E, Velazquez AMV, Parker BJ, Rasmussen M, Lindgreen S, Lilje B, Tobin DJ, Kelly TK, Vang S, et al. 2014. Genome-wide nucleosome map and cytosine methylation levels of an ancient human genome. *Genome Res.* 24(3): 454–466.
- Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud G, Sudmant PH, de Filippo C, et al. 2014. The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* 505(7481): 43–49.
- Racimo F, Gokhman D, Fumagalli M, Ko A, Hansen T, Moltke I, Albrechtsen A, Carmel L, Huerta-Sánchez E, Nielsen R. 2017. Archaic adaptive introgression in TBX15/WARS2. *Mol Biol Evol.* 34(3): 509–524.
- Radtke KM, Ruf M, Gunter HM, Dohrmann K, Schauer M, Meyer A, Elbert T. 2011. Transgenerational impact of intimate partner violence on methylation in the promoter of the glucocorticoid receptor. *Transl Psychiatry* 1(7): e21.
- Raghavan M, DeGiorgio M, Albrechtsen A, Moltke I, Skoglund P, Korneliussen TS, Gronnow B, Appelt M, Gullov HC, Friesen TM, et al. 2014. The genetic prehistory of the New World Arctic. *Science* 345: 1255832–1255832.
- Raghavan M, Steinrücken M, Harris K, Schiffels S, Rasmussen S, DeGiorgio M, Albrechtsen A, Valdiosera C, Ávila-Arcos MC, Malaspina A-S, et al. 2015. Genomic evidence for the Pleistocene and recent population history of Native Americans. *Science* 349(6250): aab3884.
- Rakyan VK, Blewitt ME, Druker R, Preis JI, Whitelaw E. 2002. Metastable epialleles in mammals. *Trends Genet.* 18(7): 348–351.
- Rasmussen S, Allentoft ME, Nielsen K, Orlando L, Sikora M, Sjögren K-G, Pedersen AG, Schubert M, Van Dam A, Kapel CMO, et al. 2015. Early divergent strains of *Yersinia pestis* in Eurasia 5, 000 years ago. *Cell* 163(3): 571–582.
- Rayco-Solon P, Fulford AJ, Prentice AM. 2005. Differential effects of seasonality on preterm birth and intrauterine growth restriction in rural Africans. *Am J Clin Nutr.* 81(1): 134–139.
- Reich D, Green RE, Kircher M, Krause J, Patterson N, Durand EY, Viola B, Briggs AW, Stenzel U, Johnson PLF, et al. 2010. Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature* 468(7327): 1053–1060.
- Reynard LN. 2016. Analysis of genetics and DNA methylation in osteoarthritis: what have we learnt about the disease? *Semin Cell Dev Biol* 62: 57–66.
- Robertson KD. 2005. DNA methylation and human disease. *Nat Rev Genet.* 6(8): 597–610.
- Roemer I, Reik W, Dean W, Klose J. 1997. Epigenetic inheritance in the mouse. *Curr Biol.* 7(4): 277–280.
- Rubin LP. 2015. Maternal and pediatric health and disease: integrating biopsychosocial models and epigenetics. *Pediatr Res* 79(1-2): 127–35.
- Ruchat SM, Houde AA, Voisin G, St-Pierre J, Perron P, Baillargeon JP, Gaudet D, Hivert MF, Brisson D, Bouchard L. 2013. Gestational diabetes mellitus epigenetically affects genes predominantly involved in metabolic diseases. *Epigenetics* 8(9): 935–943.
- Schuenemann VJ, Bos K, DeWitte S, Schmedes S, Jamieson J, Mittnik A, Forrest S, Coombes BK, Wood JW, Earn DJD, et al. 2011. Targeted enrichment of ancient pathogens yielding the pPCP1 plasmid of *Yersinia pestis* from victims of the Black Death. *Proc Natl Acad Sci.* 108(38): E746–E752.
- Seguin-Orlando A, Gamba C, Sarkissian C, Der Ermini L, Louvel G, Boulygina E, Sokolov A, Nedoluzhko A, Lorenzen ED, et al. 2015. Pros and cons of methylation-based enrichment methods for ancient DNA. *Sci Rep.* 5(1): 11826.
- Seguin-Orlando A, Korneliussen TS, Sikora M, Malaspina A-S, Manica A, Moltke I, Albrechtsen A, Ko A, Margaryan A, Moiseyev V, et al. 2014.

- Genomic structure in Europeans dating back at least 36, 200 years. *Science* 346(6213): 1113–1118.
- Skoglund P, Posth C, Sirak K, Spriggs M, Valentin F, Bedford S, Clark GR, Reepmeyer C, Petchey F, Fernandes D, et al. 2016. Genomic insights into the peopling of the Southwest Pacific. *Nature* 538(7626): 510–513.
- Skoglund P, Reich D. 2016. A genomic view of the peopling of the Americas. *Curr Opin Genet Dev.* 41: 27–35.
- Smith O, Clapham AJ, Rose P, Liu Y, Wang J, Allaby RG. 2014. Genomic methylation patterns in archaeological barley show de-methylation as a time-dependent diagenetic process. *Sci Rep.* 4: 5559.
- Suter M, Abramovici A, Showalter L, Hu M, Shope C, Do Varner M, Aagaard-Tillery K. 2010. In utero tobacco exposure epigenetically modifies placental CYP1A1 expression. *Metab Clin Exp.* 59: 1481–1490.
- Teh AL, Pan H, Chen L, Ong ML, Dogra S, Wong J, MacIsaac JL, Mah SM, McEwen LM, Saw SM, et al. 2014. The effect of genotype and in utero environment on interindividual variation in neonate DNA methylomes. *Genome Res.* 24(7): 1064–1074.
- Tobi EW, Lumey LH, Talens RP, Kremer D, Putter H, Stein AD, Slagboom PE, Heijmans BT. 2009. DNA methylation differences after exposure to prenatal famine are common and timing- and sex-specific. *Hum Mol Genet.* 18(21): 4046–4053.
- Toledo-Rodriguez M, Lotfipour S, Leonard G, Perron M, Richer L, Veillette S, Pausova Z, Paus T. 2010. Maternal smoking during pregnancy is associated with epigenetic modifications of the brain-derived neurotrophic factor-6 exon in adolescent offspring. *Am J Med Genet B Neuropsychiatr Genet.* 153B(7): 1350–1354.
- Varley KE, Gertz J, Bowling KM, Parker SL, Reddy TE, Pauli-Behn F, Cross MK, Williams BA, Stamatoyannopoulos JA, Crawford GE, et al. 2013. Dynamic DNA methylation across diverse human cell lines and tissues. *Genome Res.* 23(3): 555–567.
- Vidal AC, Semenova V, Darrah T, Vengosh A, Huang Z, King K, Nye MD, Fry R, Skaar D, Maguire R, et al. 2015. Maternal cadmium, iron and zinc levels, DNA methylation and birth weight. *BMC Pharmacol Toxicol.* 16: 20.
- Wagner DM, Klunk J, Harbeck M, Devault A, Waglechner N, Sahl JW, Enk J, Birdsell DN, Kuch M, Lumibao C, et al. 2014. *Yersinia pestis* and the Plague of Justinian 541–543 AD: a genomic analysis. *Lancet Infect Dis.* 14(4): 319–326.
- Waterland RA, Michels KB. 2007. Epigenetic epidemiology of the developmental origins hypothesis. *Annu Rev Nutr.* 27: 363–388.
- Waterland RA, Dolinoy DC, Lin JR, Smith CA, Shi X, Tahiliani KG. 2006. Maternal methyl supplements increase offspring DNA methylation at Axin fused. *Genesis* 44(9): 401–406.
- Waterland RA, Jirtle RL. 2003. Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol Cell Biol.* 23(15): 5293–5300.
- Waterland RA, Kellermayer R, Laritsky E, Rayco-Solon P, Harris RA, Travisano M, Zhang W, Torskaya MS, Zhang J, Shen L, et al. 2010. Season of conception in rural gambia affects DNA methylation at putative human metastable epialleles. *PLoS Genet.* 6(12): 1–10.
- Weyrich LS, Duchene S, Soubrier J, Arriola L, Llamas B, Breen J, Morris AG, Alt KW, Caramelli D, Dresely V, et al. 2017. Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature Genesis* 544(7650): 357–361.
- Whitaker JW, Chen Z, Wang W. 2015. Predicting the human epigenome from DNA motifs. *Nat Methods* 12(3): 265–272.
- Wüllner U, Kaut O, DeBoni L, Piston D, Schmitt I. 2016. DNA methylation in Parkinson's disease. *J Neurochem.* 139 Suppl 1: 1–13.
- Yona AH, Frumkin I, Pilpel Y. 2015. A relay race on the evolutionary adaptation spectrum. *Cell* 163(3): 549–559.
- Yuan GC. 2012. Linking genome to epigenome. *Wiley Interdiscip Rev Syst Biol Med.* 4(3): 297–309.
- Zeilinger S, Kühnel B, Klopp N, Baurecht H, Kleinschmidt A, Gieger C, Weidinger S, Lattka E, Adamski J, Peters A, et al. 2013. Tobacco smoking leads to extensive genome-wide changes in DNA methylation. *PLoS One* 8(5): e63812.
- Zhang D, Cheng L, Badner JA, Chen C, Chen Q, Luo W, Craig DW, Redman M, Gershon ES, Liu C. 2010. Genetic control of individual differences in gene-specific methylation in human brain. *Am J Hum Genet.* 86(3): 411–419.
- Ziller MJ, Müller F, Liao J, Zhang Y, Gu H, Bock C, Boyle P, Epstein CB, Bernstein BE, Lengauer T, et al. 2011. Genomic distribution and inter-sample variation of non-CpG methylation across human cell types. *PLoS Genet.* 7(12): e1002389.