

Pathogenesis of Maternal-Fetal Syphilis Revisited

Victoria Wicher and Konrad Wicher

Wadsworth Center for Laboratories and Research, New York State Department of Health, Albany, New York

Although congenital syphilis has been recognized for several centuries and an efficient treatment with penicillin became available more than a half-century ago, the disease is still with us. Inability to culture *in vitro* the causative agent, *Treponema pallidum*, and the lack of an adequate animal model have prevented exploration of the various immunopathological events affecting the natural course of congenital infection. The purpose of this review is to analyze the disease in the context of recent knowledge acquired from human and experimental animals, particularly from the guinea pig model of congenital and neonatal syphilis, and to describe how the infection interacts with the maternal-fetal unit and how it is further modulated by the conceptus' ontogenic development. We also attempt to elucidate several old immunologic concepts and misconceptions that have remained unchallenged for too long.

Congenital syphilis results from transplacental infection with the spirochete *Treponema pallidum* subspecies *pallidum*. The disease has been recognized for ~500 years. Descriptions of the devastating early and late clinical and pathological manifestations of the disease have been reported in excellent treatises in the older literature [1–3]. A milder, less aggressive clinical presentation of the condition observed in the postpenicillin era is also the subject of several reports in the modern literature [4–7]. Despite the current availability and application of modern methodologies (reviewed in [8]) for evaluation of the maternal and neonatal humoral response, serodiagnosis continues to be a problem, because >50% of infected infants are asymptomatic at birth and syphilitic babies may even be born to seronegative mothers [9, 10].

The fact is that the nature and role of the maternal-fetal immune response, their mutual interaction, and their effect on the outcome of pregnancy and ontogenic

development of the conceptus remain poorly understood. The scarcity of relevant animal models (reviewed in [11, 12]) has thwarted our understanding of the immunopathological events affecting the various presentations and the natural course of the disease. It has also prevented experimental verification and challenge of old immunologic concepts (quoted in [6, 7]), born mostly from empirical observations that were not (and still are not) clearly understood.

EFFECT OF SYPHILIS ON PREGNANCY OUTCOME

Congenital syphilis may occur when an infected woman becomes pregnant or when a pregnant woman becomes infected. For the sake of clarity, these sequences of events will be called A and B, respectively (figure 1). The human period of gestation normally is 9 months, which is approximately the duration of the most active primary, secondary, and early latent stages of maternal syphilis. Therefore, several combinations between stages of pregnancy and infection—and consequently, different outcomes—can be expected.

Sequence A. It has long been recognized that transmission of the disease to the fetus is largely dependent on the duration of the disease in the mother [3, 13–16], although there are controversial reports regarding the incidence of fetal morbidity and mortality (figure 1).

Received 5 July 2000; revised 9 January 2001; electronically published 25 June 2001.

Financial support: National Institute of Allergy and Infectious Diseases (grant AI 21833), US Public Health Service, Bethesda, MD.

Reprints or correspondence: Drs. Victoria Wicher/Konrad Wicher, Wadsworth Center for Laboratories and Research, New York State Dept. of Health, Empire State Plaza, PO Box 509, Albany, NY 12201-0509 (WicherK@wadsworth.org).

Clinical Infectious Diseases 2001;33:354–63

© 2001 by the Infectious Diseases Society of America. All rights reserved.
1058-4838/2001/3303-0014\$03.00




Figure 1. In congenital syphilis, the outcome of pregnancy depends on the stage of maternal syphilis at time of conception (*sequence A*) or the stage of gestation at the time of infection (*sequence B*).

In 1937, Paley [17] reported that the pregnancies of mothers with untreated syphilis of <2 years' duration resulted in ~50% of infected and 50% of uninfected infants. Fiumara et al. [15] reported that pregnancy of untreated women with primary or secondary syphilis will result in infection of all conceptuses. Approximately 50% will be dead in utero, prematurely born, stillborn, or dead shortly after delivery. The longer the mother has had the disease at the time of pregnancy, the chance of fetal infection decreases, to ~40% in early latency and 10% in late latency. This has been recently confirmed by Sanchez et al. [16].

Sequence B. In utero transmission may occur at any stage of pregnancy [18, 19], although the highest rate of fetal mortality and morbidity occur with untreated first-trimester and second-trimester infection, as seen in sequence A (figure 1); a higher percentage of asymptomatic disease occurs in third-trimester infection [4, 15]. Likewise, when congenital rubella and toxoplasmosis are acquired during the last trimester, the incidence of clinical manifestations in the infected infant is lower than when microbial infection occurs during the first and second trimesters [20, 21].

It should be noted that in the literature, the description of the temporal correlation between the stages of syphilis and gestation is frequently unclear and misleading. Therefore, to explain the predictable effect of maternal syphilis on the outcome of pregnancy, it is frequently stated that the most affected infants are those born to mothers with primary and secondary syphilis, whereas those born to mothers in the early-late or late stages of the disease are less affected [4, 7, 22]. In fact, the most affected infants are those conceived in, not born to, mothers

with primary or secondary syphilis, whereas the less affected are those conceived in mothers in the early-late or late stage of the disease [15]. Both situations relate exclusively to sequence A (figure 1).

On the basis of empirical observations in humans [2, 3, 23], it was long believed that reduction in transplacental infection with progression of maternal syphilis was related to evolution of maternal immunity, also known as Kassowitz's law [23]. However, more recent observations in humans [2, 24] have demonstrated that the possibility of fetal infection is never eliminated. A classic example was provided by Fiumara in 1965 [24], in which an untreated syphilitic mother transmitted the disease to her children, although with decreasing frequency, during a period of 10 years. The most puzzling and as yet unexplained finding in this report, as in other reports [2], was the intermittence of the transmission, because normal offspring could be preceded and followed by an infected infant [24].

These findings are interesting in light of recent studies of the guinea pig model of congenital syphilis [25]. In the latter studies, controlled by more specific and sensitive techniques than those used by Fiumara in the human study, it was demonstrated that untreated syphilitic sows could consistently transmit the infection to their young; this infection would perpetuate itself through multiple pregnancies and up to the fourth generation. Whereas congenital infection in guinea pigs is asymptomatic, infection of most littermates examined was demonstrated by use of ≥ 1 specific and sensitive methods: the IgM test, PCR, and the rabbit infectivity test (RIT) [25].

It is interesting that in the animal studies, what seemed to decrease in successive progeny was not the maternal transmission but the load of transmitted microorganisms and/or their virulence. This assumption was in accordance with the expression of a milder humoral response in consecutive progeny and the extremely low infectivity displayed by their organs when injected into rabbit testes. Neither obvious local inflammation nor systemic seroconversion was observed, and yet *T. pallidum* DNA was repeatedly detected in the rabbit testes by PCR [25]. PCR detects *T. pallidum* regardless of its virulence or viability.

Changes in pathogenicity are not foreign to treponemal diseases and are due to multiple factors: nutritional, environmental, behavioral, and immunologic [26]. Furthermore, as demonstrated in the guinea pig model of neonatal and congenital syphilis [25, 27, 28], the pathogen's virulence may be modulated by the maternal immune response but also by the conceptus' genetic background, which is at least as important as, if not more important than, the traits of the bacterium [29]. This could explain why, in many situations, congenitally infected infants are asymptomatic at birth and remain so for variable periods of time, whereas other newborns present with full-blown disease [30]. In the untreated congenitally asymptomatic infant, the organisms apparently vegetate in a com-

mensal state until appropriate biological conditions that promote virulence and pathogenicity develop. It is possible, therefore, that “normal” infants preceded and followed by congenitally infected siblings [24] represent instances where the commensal state remained unaltered for life, undetectable by old, insensitive serological methods.

Also uncertain, and rather unchallenged, are several immunopathological concepts that were advanced by earlier investigators to explain the maternal and fetal response when infection occurs at various stages of gestation (sequence B). The rare frequency of organisms found by Harter and Bernischke [18] in aborted fetuses in early pregnancy and the absence of local inflammation or necrosis surrounding the organisms led the authors to postulate that “the spirochetes rarely, if ever, overwhelm the early fetus, causing intrauterine death and expulsion, as is common in the last trimester.”

There are 2 problems with this postulate. The first is the implicit assumption that the risk of fetal infection, morbidity, and mortality increases as the state of pregnancy advances. The second is the exclusive association of that postulate, made by clinicians and researchers in the field, with the notion advanced by Silverstein [31] and Silverstein and Lukes [32], who suggested that fetal infection becomes evident only after a functional immune system develops. In fact, Harter and Bernischke’s postulate is counter to the recognition that the highest incidence of fetal death occurs when a woman is infected shortly before or after conception, when the fetal immune system is nonexistent or underdeveloped. The higher percentage of asymptomatic disease is frequently associated with third-trimester infection [15], when there is a higher degree of fetal immunocompetence and an increased chance of finding treponemes in tissues [18] (figure 1). Certainly, both postulates are at odds with the fact that many congenitally infected newborns do not have any obvious clinical manifestation of the disease, and a good percentage of them may remain unaffected for life [6, 15, 33].

The rarity of spirochetes in early as opposed to later gestation has been recently confirmed in our laboratory with use of 2 strains of guinea pigs, one that was susceptible (C4-deficient [C4D]) and one that was resistant (Albany [Alb]) to cutaneous infection with *T. pallidum*. In the first trimester (20–25 days’ gestation), only 3 (13%) of 22 fetuses and 3 (23%) of 13 examined within 24–48 h of intravenous infection of C4D and Alb dams, respectively, were positive by PCR or RIT. However, 8 of 8 fetuses obtained 24 h after maternal infection of 3 C4D dams in the third trimester (at 50–60 days) showed that ≥ 1 of the organs examined (lymph nodes, brain, spleen, placenta, and umbilical cord) were positive by PCR (authors’ unpublished observations). Following a recent human study, Nathan et al. [19] reported similar findings: *T. pallidum* (PCR positive and RIT positive) was detected in only 4 (36%) of 11 amniotic

fluid specimens extracted from women 14–19 weeks pregnant who had secondary and early latent syphilis, at the time when increased treponemia and transmission could be expected.

A more plausible explanation of these apparently paradoxical situations may be afforded by recent findings concerning the immune interaction at the fetal-maternal interface and new concepts of *T. pallidum* biology. The low number of treponemas found in early macerated fetuses [18] could result from an overwhelming cell-killing (apoptosis) of maternal or fetal infected cells in placenta. It has been shown that bacterial lipoproteins from *T. pallidum* and *Borrelia burgdorferi* can induce the synthesis in murine macrophages of biologically active TNF- α [34, 35], which has the potential for cell activation and apoptosis [36]. Apoptosis has been implicated in immune defense against several pathogens, including some of those that cause congenital infection [37, 38], although the full mechanism remains controversial and not clearly understood [39]. To our knowledge, apoptosis in congenital syphilis has not been explored.

Alternatively, the completion of the sequence of the *T. pallidum* genome [40] has provided clues regarding the pathogen’s limited biosynthetic capabilities, which explain its full dependence on the host for survival, propagation, and dissemination by adhesion to cells, including endothelial cells and extracellular matrix components [41–44]. Indeed, the survival of the pathogen should be highly restricted in the incipient embryo-fetal microenvironment (figure 1). This could also explain why treatment initiated early in pregnancy is highly effective.

In light of these new findings, how then can we explain the high incidence of fetal death associated with pregnancy that occurs immediately before or after infection? As indicated above, one possibility is that resorption or fetal demise early during pregnancy is caused by the infected fetal-maternal unit’s immune reaction, due to an intense release of inflammatory cytokines from activated leukocytes infiltrating the infected membranes [45]. In recent years, extensive evidence has been accumulated regarding the bidirectional interaction between mother and fetus [46]. In fact, during gestation, cytokine production not only precedes but is an active part of the process of implantation, growth, and survival of the fetus. Cytokines also contribute to the process of fetal lymphogenesis and organogenesis [47, 48].

More important, these cytokines are produced locally by lymphoid and resident nonlymphoid cells of the decidua, chorion, and trophoblasts [45, 47, 49]. Their main role is to discriminate between 2 apparently contradictory requirements, to protect the utero from invading infectious agents while preventing the rejection of the allotransplant. Under normal circumstances the fetoplacental unit itself spontaneously secretes the anti-inflammatory cytokines IL-10, IL-4, and transforming growth factor (TGF)- β to fend off any harmful effect of sys-

temic or local inflammation [50–52]. Increased production of the inflammatory cytokines IL-2, IFN- γ , and TNF- α and prostaglandins (Th1) induced by uterine infection has been associated with fetal demise, growth retardation, and preterm delivery at any stage of gestation [52–55], whereas a predominant Th2 response, although it will not prevent fetal infection, is unlikely to cause fetal damage [50, 51]. One may reason, therefore, that a detrimental Th1 immune response must be prevalent when a recently infected mother becomes pregnant or when maternal syphilis occurs in very early pregnancy, causing resorption or fetal death.

Although it is not totally eliminated, the risk of fetal loss should progressively diminish when infection occurs in later stages of gestation and ontogenic development, when the fetal-placental unit immune response is tilted toward preservation of the conceptus [47, 48, 50]. The report of Desmonts and Couvreur [56] on a prospective study of congenital toxoplasmosis is worth considering. The majority of congenital infections (65%) occurred following third-trimester maternal infection. However, the vast majority of the infants were asymptomatic. In contrast, only 17% of neonates who were born to mothers infected during the first trimester were congenitally infected, but the majority were severely damaged.

Dramatically depressed cellular responses to cytomegalovirus and herpes simplex infection, potential sources of gestational morbidity, have been observed during second-trimester and third-trimester human pregnancies [57, 58]. The relatively large percentage of congenitally infected but asymptomatic newborns with syphilis is consistent with the above hypothesis. None of the above immune mechanisms have been explored in cases of natural congenital syphilis.

EFFECT OF PREGNANCY ON THE COURSE OF MATERNAL SYPHILIS

Clinical and experimental studies have shown that several hormonal and immunologic changes that take place during pregnancy [47, 52, 59–61] are potential modulators of maternal susceptibility to infection. Various hormones made by the trophoblast have been shown to interfere with the induction of the immune response. These include progesterone and estrogen, which inhibit cytotoxic T cells and natural killer (NK) cells [60–62]. Progesterone is essential for the maintenance of pregnancy in several mammalian species, apparently because it affects the Th1/Th2 ratio and decreases cell-mediated responses [60]. NK activity is significantly lower during healthy human pregnancy than in nonpregnancy, whereas it is substantially increased in patients who experience spontaneous abortion or preceding parturition [63]. Parker and Wendel [64] reported that reduction of estrogen production may be associated with severe intrauterine stress in pregnancies complicated by syphilis.

In the prepenicillin era, when expression of the disease was

much more dramatic [1–3] than it is today [4–7] and only few serological tests (of low sensitivity and specificity) were available, observations made by clinicians in the field of syphilis indicated that pregnancy affected the natural course of syphilis [65–68]. As early as 1922, Moore [67] wrote: “The fact that women who bear syphilitic children often give no history of syphilis and present no signs of the disease (except a positive Wasserman reaction) has long been a common knowledge.” and “a woman infected at or shortly after the time of conception usually does not develop a chancre or secondary syphilis. When infection takes place late in pregnancy, on the other hand, the usual course of events follow[s], but is often much delayed.” These observations, reassessed and confirmed 20 years later by Moore et al. [68], were consistent with those made earlier by Brown and Pearce [65] with regard to the rabbit model.

Although the aforementioned concepts are not shared by modern clinicians [4, 5, 69], one cannot but recognize how the prevailing diagnosis of congenital syphilis relies so heavily on maternal and neonatal serological tests (VDRL, RPR, FTA-Abs) and molecular tests (*T. pallidum* DNA), on neonatal clinical symptoms, and on radiological findings, because maternal clinical symptoms are often absent in pregnancy, even during active disease [10, 70]. In a retrospective study, Reyes et al. [33] reported that of 148 serologically (RPR, FTA-Abs) positive mothers who delivered either congenitally infected stillborn or liveborn infants, only 6 (4%) had a history of primary or secondary syphilis. The remaining 142 infected mothers (96%) were asymptomatic.

If implantation and survival of the fetus is associated with an altered Th1/Th2 balance, one might predict that the humoral response in pregnancy is increased, whereas cell-mediated responses are either ameliorated or seriously impaired. In fact, some clinical and experimental data [61, 71] support the concept that healthy pregnancy is characterized by a Th2-like phenomenon. Systemic lupus erythematosus, an antibody-mediated autoimmune disease, tends to flare up, especially in women with recently active disease, before conception [72, 73]. A noticeable temporary abatement of the symptoms of rheumatoid arthritis (a cell-mediated disorder) has been consistently observed in pregnancy [74], whereas a strong Th1 response against intracellular pathogens is known to compromise pregnancy, as demonstrated in the murine model of *Leishmania* infection [59] and infections with other intracellular pathogens such as *Toxoplasma* and *Plasmodium*, as well as HIV-associated infections (reviewed in [46]).

Fortuitously, we have observed what seems to be a Th2 phenomenon during the course of recent experimental studies directed toward elucidation of maternal transmission of syphilis through several litters and generations of guinea pigs [25]. In those studies, *T. pallidum*-infected multiparous females were bled for serological examination approximately every 23 days




Figure 2. Kinetics of humoral response in multiparous female guinea pig with acquired syphilis resembles that of multiparous congenitally infected sow shown in figure 3A. O.D., optical density. Figure is reproduced from [25] with permission.

(early, middle, and late gestation). Analysis of the kinetics of the maternal humoral response showed a consistent pattern of fluctuations throughout consecutive pregnancies (figures 2 and 3A). Early in pregnancy the levels of maternal IgM and IgG antitreponemal antibodies increased; this was followed by a substantial drop in the titers before and immediately after each delivery. Apparently, upregulation of the Th2-like maternal immune response, although it prevents fetal demise, facilitates the growth and circulation (depressed Th1?) of *T. pallidum*, as shown by the obvious transmissibility to the fetuses (IgM positive, PCR positive, and RIT positive). It could also account for the high levels of maternally transmitted IgG antibodies and the fetal production of circulating immune complexes and IgM rheumatoid factor commonly observed in congenitally infected infants [75] and guinea pigs [11].

These unexpected findings were observed in 2 multiparous does with acquired syphilis, which were the source of 3 (figure 2) and 5 consecutive litters, respectively, and in 1 multiparous dam that was congenitally infected (figure 3A). This pattern of humoral response was not observed in nulliparous syphilitic females tested for a similar 12-month period (figure 3B) [25]. Although the number of infected pregnant females that were serologically studied is low and more studies are needed to confirm these results, the consistency of the findings in a total of 13 pregnancies is not trivial.

Indeed, it is interesting how often the clinician is confronted with negative serological results of maternal blood tests performed just before or after delivery [9, 10, 70, 76], which cannot always be explained by recent infection or the prozone effect [70].

CONGENITAL SYPHILIS

Perinatal infection with *T. pallidum* exhibits a wide range of clinical manifestations. The general assumption that the outcome of pregnancy is determined by the time of gestational infection [15] and by fetal immunogenesis [31, 32] has been insufficient to define the various modalities and presentations of the disease. It cannot explain why the majority of congenitally infected newborns are free of symptoms at birth or why some of them develop early or late manifestations, whereas others remain asymptomatic for life. Indeed, there is a paucity of information about congenital syphilis with regard to the immunomodulatory role played by the fetal-maternal relationship, and changes associated with fetal ontogenic development and genetic background.

During pregnancy, the human placenta undergoes critical structural changes, which are synchronized with the development of the embryonic/fetal and maternal compartments [77]. Normally, the placenta is a main source of anti-inflammatory cytokines (IL-10, IL-4) and a variety of growth factors, including TGF- β . These factors not only stimulate or inhibit fetal growth, but they also participate in cellular and tissue differentiation, programmed cell death, metabolism, nutrient uptake, and angiogenesis [48, 77]. In the fetus, TGF- β plays a critical role in the differentiation of oligodendrocytes [78], in the integrity of the musculoskeletal system, and in the remodeling and mineralization of bone tissue [77]. Together with IL-10 and IL-4, the growth factors protect the fetus from systemic or local inflammation by inhibiting lymphokine-activated killer cells [79], allospecific cytotoxic T cells [80], antigen-presenting cells, and macrophage activation [81]. An additional property of IL-10 is its potent inhibition of osteoclast formation and prevention of bone resorption [82].

Thus, one may speculate that factors known to promote placental and fetal growth, as well as fetal survival as an allograft, may incidentally benefit and protect an invasive pathogen, such as *T. pallidum*, from an immune attack, while promoting asymptomatic infection. Following the same line of reasoning, *T. pallidum*-associated temporary or permanent damage to a variety of embryonic/fetal tissues and organs, including skin, brain, bones, liver, and pancreas, may be regarded as a failure of the fetal-maternal unit to prevent an inflammatory response triggered by the pathogen in a relatively immunocompetent fetal microenvironment.

As early as 18–20 weeks' gestation, the fetus has the potential to respond to infection with use of nonspecific host-resistant factors, such as phagocytic cells and NK cells, or with a relatively low level of specificity provided by the $\gamma\delta$ T cells and B1 CD5⁺ B cells [83]. The mammalian immune system, however, is not completely functional during fetal life or even at birth but undergoes a gradual maturation, which may be completed after

the neonatal or even adolescent life [84]. Although the normal newborn produces a large number of B cells with different specificities, the full array of antibody response is reached only postnatally by the gradual exposure to different bacteria and viruses and the acquisition of memory cells [83, 85, 86].

Furthermore, the immune response to a particular antigen remains constant for the individual animal within a particular strain but shows marked differences between different strains [87, 88], which suggests an important role for genetic factors in the evolution of the host's immunocompetence. Several reports have shown genetic differences among mouse strains in their susceptibility to bacterial [88], viral [89], and protozoal infection [90]. It has also been shown that expression of resistance genes may not be equally distributed in different organs [91, 92] or at all stages of ontogenic development [27, 93]. We have shown strain-associated and age-associated differences in lymphocyte phenotypes and immune responsiveness to alloantigens and mitogens in 2 strains of guinea pigs: (1) C4D genetically associated with inbred strain 13 and (2) the Alb line haplotype identical to inbred strain 2 [94].

More interesting is that the ontogeny of the humoral response and cutaneous reaction to *T. pallidum* infection in these

2 strains follow a different time course in neonates and in adults [27]. C4D neonates (1–3 days old) are temporarily resistant (2–3 months) to intradermal infection but are highly susceptible as adults. The opposite occurs with Alb strain neonates. Ontogenic changes in susceptibility to cutaneous challenge with *T. pallidum* has also been demonstrated in the rabbit model [95] and in infection of mice with another pathogenic spirochete, *B. burgdorferi* [96].

There is little information about the role of the cellular arm of the immune response in congenital syphilis. Friedmann, in 1977 [97], reported that peripheral blood lymphocytes from 7 babies aged 2 months to 2 years with active disease responded to *T. pallidum* antigen with stimulation ratios significantly higher than those of normal age-matched control cells, whereas the cellular response of 16 asymptomatic, serologically positive newborns (24–48 h old) was not significantly different from that of controls. Samsom et al. [98] did not find any significant differences between 17 newborns (1–7 days old) with early congenital syphilis and 17 age-matched controls in response to phytohemagglutinin antigen (PHA). These authors found significantly increased levels of B cells and production of IgM and circulating immune complexes (CICs). Fitzgerald and Froberg [99] showed that congenitally infected rabbits have an elevated T cell proliferative response to mitogen and B cell function involving increased levels of antitreponemal antibodies.

We have shown that transplacental infection with *T. pallidum* in C4D and Alb dams occurs irrespective of their different genetic backgrounds and their susceptibility to cutaneous infection [11]. In these 2 strains, in utero infection elicits early and extended (2–4 months of age) production of IgM anti-treponemal antibodies, IgM–rheumatoid factor, IgM–CIC, and a delayed switch to IgG antibodies [28]. Compared to those of normal control animals, spleen cells from congenitally infected pups aged 1–20 weeks showed a remarkably prolonged naïve-type of immune response, as reflected in the significantly higher response to both T cell mitogens concanavalin-A (ConA) and PHA and a weaker response to B cell mitogen.

Several studies [100–102] have indicated that T lymphocytes can be phenotypically divided into 2 different subsets of CD4⁺ T cells: naïve (CD45R^{hi}) and memory (CD45R^{lo}) T cells. Naïve T cells represent the suppressor-inducer subset [100]; they display an increased lymphoproliferative response to T cell mitogens but react poorly to antigens and are associated with a primary IgM immune response and induction of suppression of antigen-driven IgG production. Memory CD4⁺ T cells, on the other hand, represent the helper-inducer subpopulation for antigenic and mitogenic responses of B cells [101]. They have a relatively lower proliferative response to T cell mitogens but react strongly to antigens. Both subpopulations have different activation requirements and secretion of cytokines [102]. These




Figure 3. Kinetics of humoral response to *Treponema pallidum* in a congenitally infected guinea pig mother (generation II) during 5 consecutive pregnancies (A). For comparison, kinetics of humoral response of nulliparous congenitally infected female (B). O.D., optical density. Figure is reproduced from [25] with permission.

characteristics of naïve and memory T cells are consistent with the elevated response to T cell mitogens in asymptomatic congenitally infected rabbits [99] and guinea pigs [103] and the lower response to *T. pallidum* antigen in asymptomatic infants [97]. They also correlate with the abundant production of IgM polyclonal antibodies and IgM–rheumatoid factor, reflecting the activation of naïve B1-CD5⁺ B cells [28, 104, 105]. Indeed, delayed maturation of T cells and B cells will contribute to the dysfunction of the immune response with decreasing resistance to pathogens and increased production of autoantibodies, particularly of the IgM type [28, 75, 103, 105].

Whether the prolonged naïve-type of humoral immune response in asymptomatic congenital syphilis is associated with some mechanism of immunologic tolerance at the cellular level, as occurred in children congenitally infected with *Toxoplasma gondii* [21] and lymphocytic choriomeningitis virus [106], has remained unexplored. The presence of the pathogen at the critical stage of development could result in a lymphocyte population unable to distinguish between the microorganism and self. However, immunologic tolerance that results from an efficient process of clonal deletion of T cells capable of mounting an aggressive response to “self”-MHC molecules [107] may not play a major role in congenital syphilis, because such type of tolerance leaves the organism unable to induce an autoimmune response, a process commonly seen in patients with syphilis [108]. High levels of circulating immune complexes containing *T. pallidum*, creatine kinase (CK), fibronectin (Fn), and IgG1 and IgG3, in addition to IgM, have been found in serum samples of congenitally infected babies [104, 105]. Antibodies to host antigens, such as cardiolipin, CK, collagen, laminin, and Fn, are often present in serum samples obtained from *T. pallidum*-infected human and experimental animals (see review of literature in [104]). We are more inclined to believe that a process akin to clonal inactivation (1) by antigen presentation in the absence of costimulatory signals, due to the dominance of immature T and B cells early in ontogeny, or (2) by a state of “immunological ignorance,” where T cells that are able to recognize the antigen will be present in the mature host but will not normally be activated [109], may have some relevance in congenital syphilis.

Apparently, failure to maintain an appropriate balance between self-reactive naïve CD45R^{hi} T cells and normally present antagonistic mature CD45R^{lo} T cells may result in an autoimmune response [110] ranging from a lethal wasting disease in the newborn, as shown in the rat model [111], to irreversible tissue damage. Indeed, a subtle but insidious inflammatory autoimmune response could explain why late manifestations of congenital syphilis, such as interstitial keratitis, Clutton’s joints, and sensorineural deafness, may respond to steroid treatment rather than to antibiotics.

THE GUINEA PIG MODEL

Although congenital syphilis causes no apparent disease in guinea pigs, this animal has been the source of valuable information on various immunologic and clinical aspects of perinatal syphilis that could not be explored in humans or other animal models. The guinea pig host has been instrumental in the addressing of 3 important but controversial clinical puzzles. The first relates to the possibility of transmission of congenital syphilis through several generations, also known as tertiary syphilis, which has been suspected since the end of the 19th century [1, 2]. The animal model demonstrated not only third-generation and fourth-generation syphilis but also an association between congenital transmission and pregnancy-related changes in the maternal humoral response and a decline in immune response in successive congenitally infected progeny [25].

Second, we have conclusively shown that despite its close genetic relationship with *T. pallidum* subspecies *pallidum*, *T. pallidum* subspecies *pertenue*, the causative agent of yaws, shows distinctive pathogenic properties, such as its inability to cross the placenta and cause congenital infection [112]. Third, we addressed the question of whether infection occurring during delivery or immediately after birth differs from infections occurring during gestation. The results of these studies indicated that, as opposed to congenital infection, intracutaneous inoculation of 1–3-day-old neonates with *T. pallidum* did not induce IgM rheumatoid factor or IgM-CIC, which reflect a host’s naïve B1 (CD5⁺) B cell response to maternally transmitted IgG. The inclusion of both strains of guinea pigs in these studies of neonatal infection also served to demonstrate age-associated and genetically associated differences in the cutaneous response to *T. pallidum* [27].

CONCLUSION

Although in the past decade the number of new cases of congenital syphilis in the United States has substantially decreased, from a peak of 107 cases per 100,000 live-born infants in 1991 to 30 per 100,000 in 1996 [113], isolated outbreaks of infection with up to 10-fold higher rates have been reported [113–115]. This suggests that eradication of the disease cannot be taken for granted. The failure to eradicate the disease now and in the past should in part be attributed to our insufficient understanding of the immunologic mechanisms involved in the maternal-fetal interface that cause early fetal death, stillbirth, and early and late manifestations of congenital syphilis. Adequate knowledge of immunity is essential to comprehend the natural history of the disease and how it relates to latency, clinical progression, and relapse and to properly evaluate therapeutic results.

Indeed, there is a need for new epidemiological, immuno-

logic [116], and clinical strategies, if not to fully eliminate the pathogen, then to at least keep it under control.

Acknowledgments

We gratefully acknowledge Drs. Daniel Musher and Robert Baughn at the Veterans Affairs Medical Center, Houston, for critical evaluation of the manuscript, and Dr. Katherine Henrikson, for her editorial help.

References

1. Hutchinson J. Syphilis. 2d ed. New York: Funk & Wagnalls, 1910:1–571.
2. Nabarro D. Congenital syphilis. London: Edward Arnold Publishers, 1954:1–461.
3. Diday PE. A treatise on syphilis in newborn children and infants at the breast. London: The New Sydenham Society, 1959.
4. Wendel GD. Gestational and congenital syphilis. Clin Perinatol 1988; 15:287–303.
5. Zenker PN, Berman SM. Congenital syphilis: trends and recommendations for evaluation management. Pediatr Infect Dis J 1991; 10: 516–22.
6. Ingall D, Sanchez PJ, Musher DM. Syphilis. In: Remington JS, Klein JO, eds. Infectious diseases of the fetus and newborn infant. 4th ed. Philadelphia: WB Saunders, 1995:529–64.
7. Radolf JD, Sanchez PJ, Schulz KF, Murphy KF. Congenital syphilis. In: Holmes KK, Mårdh P-A, Sparling PF, et al., eds. Sexually transmitted diseases. 3d ed. New York: McGraw-Hill, 1999:1165–89.
8. Wicher K, Horowitz HW, Wicher V. Laboratory methods of diagnosis of syphilis for the beginning of the third millennium: a review. Microbes and Infection 1999; 1:1035–49.
9. Dorfman DH, Glaser JH. Congenital syphilis presenting in infants after the newborn period. N Engl J Med 1990; 323:1299–302.
10. Sanchez PJ, Wendel GD, Norgard MV. Congenital syphilis associated with negative results of maternal serologic tests at delivery. Am J Dis Child 1991; 145:967–9.
11. Wicher K, Baughn RE, Wicher V, Nakeeb S. Experimental congenital syphilis: guinea pig model. Infect Immun 1992; 60:271–7.
12. Wicher V, Wicher K. Guinea pig model of acquired and congenital syphilis. In: Zak O, Sande M, eds. Handbook of animal models of infection. New York: Academic Press, 1999:291–301.
13. Hutchinson J. A clinical memoir on certain diseases of the eye and ear consequent on inherited syphilis. London: John Churchill, 1863.
14. Ingraham NR. The value of penicillin alone in the prevention and treatment of congenital syphilis. Acta Derm Venereol (Stockh) 1951; 31(Suppl 24):60–87.
15. Fiumara NJ, Fleming WL, Downing JG, Good F. The incidence of prenatal syphilis at the Boston City Hospital. N Engl J Med 1951; 245:634–40.
16. Sanchez PJ, Wendel GD, Grimprel E, et al. Evaluation of molecular methodologies and rabbit infectivity testing for the diagnosis of congenital syphilis and central nervous system invasion by *Treponema pallidum*. J Infect Dis 1993; 167:148–57.
17. Paley SS. Syphilis and pregnancy. NY State J Med 1937; 37:585–90.
18. Harter CA, Bernischke K. Fetal syphilis in the first trimester. Am J Obstet Gynecol 1976; 7:705–11.
19. Nathan L, Bohman Van R, Sanchez PJ, et al. In utero infection with *Treponema pallidum* in early pregnancy. Prenat Diagn 1997; 17: 119–23.
20. Buimovici-Klein E, Lang PB, Ziring PR, Cooper LZ. Impaired cell-mediated immune response in patients with congenital rubella: correlation with gestational age at time of infection. Pediatrics 1979; 64: 620–6.
21. McLeod R, Mack DG, Boyer K, et al. Phenotypes and functions of lymphocytes in congenital toxoplasmosis. J Lab Clin Med 1990; 116: 623–35.
22. Sanchez PJ. Congenital syphilis. In: Aronoff SC, ed. Advances in pediatric infectious diseases. St. Louis: Mosby Year Book 1992; 7:161–80.
23. Kassowitz M. Die Vererbung der Syphilis. Wien: Med Jahrbücher, Wilhem Braumüller, 1876.
24. Fiumara NJ. A legacy of syphilis. Arch Dermat 1965; 92:676–78.
25. Wicher K, Baughn RE, Abbruscato F, Wicher V. Vertical transmission of *Treponema pallidum* to various litters and generations of guinea pigs. J Infect Dis 1999; 179:1206–12.
26. Willcox RR. Changing patterns of treponemal disease. Br J Vener Dis 1974; 50:169–78.
27. Wicher V, Wicher K, Rudofsky U, et al. Experimental neonatal syphilis in a susceptible (C4D) and resistant (Albany) strain of guinea pig. Clin Immunol Immunopathol 1990; 55:23–40.
28. Wicher V, Baughn RE, Wicher K. Congenital and neonatal syphilis in guinea pig shows a different pattern of immune response. Immunology 1994; 82:404–9.
29. Salyers AA, Whitt DD. Virulence factors that promote colonization. In: Bacterial pathogenesis: a molecular approach. Washington, DC: American Society for Microbiology Press, 1994:30–46.
30. Fogel BJ, Tamer AM, Fojaco RT, et al. Immunologic response of the fetus in congenital syphilis. J Florida M A 1969; 56:777–9.
31. Silverstein AM. Congenital syphilis and the timing of immunogenesis in the human fetus. Nature 1962; 194:196–7.
32. Silverstein AM, Lukes RJ. Fetal response to antigenic stimulus. I. Plasmacellular and lymphoid reactions in the human fetus to intra-uterine infection. Lab Invest 1962; 11:918–32.
33. Reyes MP, Hunt N, Ostrea EM, George D. Maternal/congenital syphilis in a large tertiary-care urban hospital. Clin Infect Dis 1993; 17: 1041–6.
34. Radolf JD, Norgard MV, Brandt ME, et al. Lipoproteins of *Borrelia burgdorferi* and *Treponema pallidum* activate cachectin/tumor necrosis factor synthesis: analysis using a CAT reporter construct. J Immunol 1991; 147:1968–74.
35. Radolf JD, Arndt LL, Akins DR, et al. *Treponema pallidum* and *Borrelia burgdorferi* lipoproteins and synthetic lipopeptides activate monocytes/macrophages. J Immunol 1995; 154:2866–77.
36. Aliprantis AO, Yang RB, Mark MR, et al. Cell activation and apoptosis by bacterial lipoproteins through toll-like receptors-2. Science 1999; 285:736–9.
37. Economides A, Schmid I, Anisman-Posner DJ, et al. Apoptosis in cord blood T lymphocytes from infants of human immunodeficiency virus-infected mothers. Clin Diagn Lab Immunol 1998; 5:230–4.
38. Fujino T, Iwamoto I, Otsuka H, et al. Apoptosis in placentas from human T-lymphotropic virus type I-seropositive pregnant women: a possible defense mechanism against transmission from mother to fetus. Obstet Gynecol 1999; 94:279–83.
39. Hsu C-D, Hong S-F, Harirah H, et al. Amniotic fluid soluble Fas levels in intra-amniotic infection. Obstet Gynecol 2000; 95:667–70.
40. Fraser CM, Norris SJ, Weinstock GM, et al. Complete genome sequence of *Treponema pallidum*, the syphilis spirochete. Science 1998; 281:375–88.
41. Fitzgerald TJ, Repesh LA, Blanco DR, Miller JN. Attachment of *Treponema pallidum* to fibronectin, laminin, collagen IV, and collagen I, and blockage of attachment by immune rabbit IgG. Br J Vener Dis 1984; 60:357–63.
42. Thomas DD, Baseman JB, Alderete J. Fibronectin mediates *Treponema pallidum* cytoadherence through recognition of fibronectin cell binding domain. J Exp Med 1985; 161:514–25.
43. Steiner BM, Sell S, Schell R. *Treponema pallidum* attachment to surface and matrix proteins of cultured rabbit epithelial cells. J Infect Dis 1987; 155:742–8.
44. Burgess AW, Paradise LJ, Hilbelink D, Friedman H. Adherence of

- Treponema pallidum* subsp. *pallidum* in the rabbit placenta. Proc Soc Exp Biol Med **1994**; 207:180–5.
45. Keelan JA, Marvin KW, Sato TA, et al. Cytokine abundance in placental tissues: evidence of inflammatory activation in gestational membranes with term and preterm parturition. Am J Obstet Gynecol **1999**; 181:1530–6.
 46. Wegmann TG, Lin H, Guilbert L, Mosmann TR. Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a Th2 phenomenon? Immunol Today **1993**; 14:353–6.
 47. Robertson SA, Seemark RF, Guilbert LJ, Wegmann TG. The role of cytokines in gestation. Clin Rev Immunol **1994**; 14:239–92.
 48. Chaouat G, Menu E, Athanassakis I, Wegmann TG. Maternal T cells regulate placental size and fetal survival. Regional immunology. New York: John Wiley and Sons, **1988**; 1:143–8.
 49. Steinborn A, Niederhut A, Solbach C, et al. Cytokine release from placental endothelial cells, a process associated with preterm labour in the absence of intrauterine infection. Cytokine **1999**; 11:66–73.
 50. Lin H, Mosmann TR, Guilbert L, et al. Synthesis of T helper 2-type cytokines at the maternal-fetal interface. J Immunol **1993**; 151:4562–7.
 51. Chaouat G, Meliani AA, Martal J, et al. IL-10 prevents naturally occurring fetal loss in the CBA x DBA/2 mating combination, and the local defect in IL-10 production in this abortion-prone combination is corrected by in vivo injection of IFN α . J Immunol **1995**; 154:4261–8.
 52. Chaouat G, Menu E, Szekeres-Bartho J, et al. Immunological and endocrinological factors that contribute to successful pregnancy. In: Wegmann G, Gill TG, Nisbet-Broem R, eds. Molecular and cellular immunology of the fetal maternal interface. New York: Oxford University Press, **1991**:277–86.
 53. Heyborne JD, Witkin SS, McGregor JA. Tumor necrosis factor- α in midtrimester amniotic fluid is associated with impaired intrauterine fetal growth. Am J Obstet Gynecol **1992**; 167:920–5.
 54. Hill JA, Polgar K, Anderson DJ. T-helper 1-type immunity to trophoblast in women with recurrent spontaneous abortion. JAMA **1995**; 273:1933–6.
 55. Gorivodsky M, Zemlyak I, Orenstein H, et al. TNF- α messenger RNA and protein expression in the uteroplacental unit of mice with pregnancy loss. J Immunol **1998**; 160:4280–8.
 56. Desmots G, Couvreur J. Toxoplasmosis: epidemiologic and serological aspects of perinatal infections. In: Krugman S, Gershon A, eds. Infections of the fetus and newborn infants. New York: Alan R Liss, **1975**:115–32.
 57. Gehrz RC, Christianson WR, Linner KM, et al. Cytomegalovirus-specific humoral and cellular responses in human pregnancy. J Infect Dis **1981**; 143:391–5.
 58. Kumar A, Madden DL, Nankervis GA. Humoral and cell mediated immune responses to herpesvirus antigens during pregnancy: a longitudinal study. J Clin Immunol **1984**; 4:12–7.
 59. Krishnan L, Guilbert LJ, Russell AS, et al. Pregnancy impairs resistance of C57Bl/6 mice to *Leishmania major* infection and caused decreased antigen-specific IFN α responses and increased production of T helper 2 cytokines. J Immunol **1996**; 156:644–52.
 60. Szekeres-Bartho J, Wegmann TG. A progesterone-dependent immunomodulatory protein alters the Th1/Th2 balance. J Reprod Immunol **1996**; 31:81–95.
 61. Myers MJ, Petersen BH. Estradiol induced alterations of the immune system. I. Enhancement of IgM production. Int J Immunopharmacol **1985**; 7:207–313.
 62. Szekeres-Bartho J, Kinski R, Chaouat G. The effect of progesterone induced immunologic blocking factor on NK mediated resorption. Am J Reprod Immunol **1990**; 24:105–7.
 63. Szekeres-Bartho J, Varga P, Pacsa AS. Immunological factors contributing to initiation of labour-lymphocyte reactivity in term labour and threatened pre-term delivery. Am J Obstet Gynecol **1986**; 155:108–13.
 64. Parker CR, Wendel GD. The effect of syphilis on endocrine function of the fetoplacental unit. Am J Obstet Gynecol **1988**; 159:1327–31.
 65. Brown WH, Pearce L. On the reaction of pregnant and lactating females to inoculation with *Treponema pallidum*: a preliminary note. Am J Syph **1920**; 4:593–7.
 66. Stokes JH. The treatment of late syphilis and of syphilis in mother and child. Arch Dermatol Syphilol **1921**; 4:778–95.
 67. Moore JE. Studies in asymptomatic neurosyphilis. III. The apparent influence of pregnancy on the incidence of neurosyphilis in women. Arch Intern Med **1922**; 30:548–54.
 68. Moore JE, Kemp J, Eagle H, et al. The treatment of syphilis complicated by pregnancy: the antenatal prevention of congenital syphilis. In: Moore JE, ed. Modern treatment of syphilis. 2d ed. Springfield, Illinois: Charles C Thomas, **1941**:474–501.
 69. Ingall D, Norins L. Syphilis. In: Remington JE, Klein JO, eds. Infectious diseases of the fetus and newborn infant. Philadelphia: WB Saunders, **1976**:414–63.
 70. Berkowitz KM, Stampf K, Baxi L, Fox HE. False negative screening test for syphilis in pregnant women [letter]. N Engl J Med **1990**; 322:270–1.
 71. Dresser DE. The potentiating effect of pregnancy on the humoral immune responses of mice. J Reprod Immunol **1991**; 20:253–66.
 72. Varner MW. Autoimmune disorders and pregnancy. Semin Perinatol **1991**; 15:238–50.
 73. Hayslett JP, Lynn RI. Effect of pregnancy in patients with lupus nephropathy. Kidney Int **1980**; 18:207–20.
 74. Da Silva JA, Spector TD. The role of pregnancy in the course and aetiology of rheumatoid arthritis. Clin Rheumatol **1992**; 11:189–92.
 75. Dobson SRM, Taber LH, Baughn RE. Recognition of *Treponema pallidum* antigens by IgM and IgG antibodies in congenitally infected newborn and their mothers. J Infect Dis **1988**; 157:903–10.
 76. Spangler AS, Jackson JH, Fiumara NJ, Warthin TA. Syphilis with a negative blood test reaction. JAMA **1964**; 189:113–6.
 77. Han VKM. Growth factors in placental growth and development. In: Rice GE, Shain P, Brenneke Y, eds. Molecular aspects of placental and fetal membrane autacoids. Miami: CRC Press, **1993**:395–445.
 78. McKinnon RD, Piras G, Ida JA, Dubois-Dalcq M. A role for TGF- α in oligodendrocyte differentiation. J Cell Biol **1993**; 121:1397–407.
 79. Rook AH, Kehrl JH, Wakefield LM, et al. Effects of transforming growth factor- β on the functions of natural killer cells: depressed cytolytic activity and blunting of interferon responsiveness. J Immunol **1986**; 136:3916–20.
 80. Mule JJ, Schwartz SL, Roberts AB, et al. Transforming growth factor- β inhibits the in vitro generation of lymphokine-activated killer cells and cytotoxic T cells. Cancer Immunol Immunother **1988**; 26:95–100.
 81. Czarniecki CW, Chin HH, Wong GH, et al. Transforming growth factor- β 1 modulates the expression of class II histocompatibility antigens on human cells. J Immunol **1988**; 140:4217–23.
 82. Owens JM, Gallagher AC, Chambers TJ. IL-10 modulates formation of osteoclasts in murine hemopoietic cultures. J Immunol **1996**; 157:936–40.
 83. Adinolfi M. Ontogeny of human natural and acquired immunity. In: Olding LB, ed. Reproduction and immunology. New York: Springer, **1997**:67–102.
 84. Silverstein AM. Ontogeny of the immune response: a perspective. In: Cooper MD, Dayton DH, eds. Development of host defense. New York: Raven Press, **1977**:1–10.
 85. Adinolfi M. The development of lymphoid tissues and immunity. In: Davis JA, Dobbing J, eds. Scientific foundation of pediatrics. New York: Heineman, **1974**:333–48.
 86. Sherwin WK, Rowlands DT. Determinants of the hierarchy of humoral immune responsiveness during ontogeny. J Immunol **1975**; 115:1549–54.
 87. Playfair JHL. Strain differences in the immune response of mice: the neonatal response to sheep red blood cells. Immunology **1968**; 15:35–50.
 88. Yang L, Weis JH, Eichwald E, et al. Heritable susceptibility to severe *Borrelia burgdorferi*-induced arthritis is dominant and is associated

- with persistence of large numbers of spirochetes in tissues. *Infect Immun* **1994**; 62:492–500.
89. Chalmers JE, Mackenzie JS, Stanley NF. Resistance to murine cytomegalovirus linked to the major histocompatibility complex of the mouse. *J Gen Virol* **1977**; 37:107–14.
 90. Trishmann TM, Bloom RR. Genetics of murine resistance to *Trypanosoma cruzi*. *Infect Immun* **1982**; 35:546–51.
 91. Wicher K, Abbruscato F, Wicher V, et al. Target organs of infection in guinea pig with acquired and congenital syphilis. *Infect Immun* **1996**; 64:3174–9.
 92. Orme IM, Collins FM. Demonstration of acquired resistance in Bcg^r inbred mouse strain infected with a low dose of BCG Montreal. *Clin Exp Immunol* **1984**; 56:81–8.
 93. Gros P, Skamene E, Forget A. Cellular mechanisms of genetically controlled host resistance to *Mycobacterium bovis* (BCG). *J Immunol* **1983**; 131:1966–72.
 94. Zhao J, Wicher V, Burger R, Schafer H, Wicher K. Strain and age-associated differences in lymphocyte phenotype and immune responsiveness in C4-deficient and Albany strains of guinea pigs. *Immunology* **1992**; 77:165–70.
 95. Gamboa D, Miller JN. Experimental neonatal syphilis. I. Evidence of resistance to symptomatic infection in neonatal rabbits following intradermal inoculation with *Treponema pallidum* (Nichols strain). *Pediatr Res* **1984**; 18:965–71.
 96. Barthold SW, Beck DS, Hansen GM, et al. Lyme borreliosis in selected strains and ages of laboratory mice. *J Infect Dis* **1990**; 162:133–8.
 97. Friedmann PS. Cell mediated immunological reactivity in neonates and infants with congenital syphilis. *Clin Exp Immunol* **1977**; 30: 271–6.
 98. Samsom GR, Beatty DW, Malan AF. Immune studies in infants with congenital syphilis. *Clin Exp Immunol* **1990**; 81:315–8.
 99. Fitzgerald TJ, Froberg MK. Congenital syphilis in newborn rabbits: immune functions and susceptibility to challenge infection at 2 and 5 weeks of age. *Infect Immun* **1991**; 59:1869–71.
 100. Morimoto C, Letvin NL, Distaso JA, et al. The isolation and characterization of the human suppressor inducer T cells subset. *J Immunol* **1985**; 134:1508–15.
 101. Morimoto C, Letvin NL, Boyd AW, et al. The isolation and characterization of the human helper-inducer T cell subset. *J Immunol* **1985**; 134:3762–9.
 102. Sanders ME, Makgoba MW, Shaw S. Human naïve and memory T cells: reinterpretation of helper-inducer and suppressor-inducer subsets. *Immunol Today* **1988**; 9:195–8.
 103. Wicher V, Zhao J, Dilwith R, et al. Immune abnormalities in guinea pigs with asymptomatic congenital syphilis. *Pediatr Res* **1997**; 42:794–8.
 104. Baughn RE. Triggering of autoimmune antibody response in syphilis. In: *Microorganisms and autoimmune diseases*. Friedman H, Rose N, Bendinelli M, eds. New York: Plenum Press, **1996**:79–103.
 105. Dobson SRM, Taber LH, Baughn RE. Characterization of the components in circulating immune complexes from infants with congenital syphilis. *J Infect Dis* **1988**; 158:940–7.
 106. Jamieson BD, Ahmed R. T-cell tolerance: exposure to virus in utero does not cause a permanent deletion of specific T cells. *Proc Natl Acad Sci USA* **1988**; 85:2265–8.
 107. Nossal GJV. Cellular mechanisms of immunological tolerance. *Annu Rev Immunol* **1983**; 1:33–62.
 108. Wicher K, Wicher V. Autoimmunity in syphilis. In: *Organ-specific autoimmunity*. Bigazzi PE, Wick G, Wicher K, eds. New York: Marcel Dekker, **1990**:101–24.
 109. Janeway CA, Travers P. Tolerance and response to self and non-self tissues. In: *Janeway CA, Travers P, eds. Immunobiology. The Immune System in Health and Disease*. 3d ed. New York, London: Garland Publishing, **1997**:12.26–12.33.
 110. Fawell D, McKnight AJ, Powrie F, et al. Subsets of CD4⁺ T cells and their roles in the induction and prevention of autoimmunity. *Immunol Rev* **1991**; 123:37–64.
 111. Powrie F, Mason D. OX-22^{high} CD4⁺ T cells induce wasting disease with multiple organ pathology: prevention by the OX-22^{low} subset. *J Exp Med* **1990**; 172:1701–8.
 112. Wicher K, Wicher V, Abbruscato F, Baughn RE. Experimental yaws: Guinea pig model. *Treponema pallidum* subsp. *pertenue* displays pathogenic properties different from those of *T. pallidum* subsp. *pallidum*. *Infect Immun* **2000**; 68:3219–25.
 113. Centers for Disease Control and Prevention. Primary and secondary syphilis: United States, 1997. *MMWR Morb Mortal Wkly Rep* **1998**; 47:493–7.
 114. Centers for Disease Control and Prevention. Epidemic of congenital syphilis: 1996–1997. *MMWR Morb Mortal Wkly Rep* **1998**; 47:904–7.
 115. St. Louis ME, Farley TA, Aral SO. Untangling the persistence of syphilis in the South. *Sex Transm Dis* **1996**; 23:1–4.
 116. Scarozza AM, Ramsingh AI, Wicher V, Wicher K. Spontaneous cytokine gene expression in normal guinea pig blood and tissues. *Cytokine* **1998**; 10:851–9.