



NCBI



www.pubmed.gov

A service of the U.S. National Library of Medicine  
and the National Institutes of Health

My NCBI 

[\[Sign In\]](#) [\[Register\]](#)

---

All Databases
PubMed
Nucleotide
Protein
Genome
Structure
OMIM
PMC
Journals
Books

Search  for    [Advanced Search](#)

About Entrez


Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy 

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health


Clinical Alerts

ClinicalTrials.gov


PubMed Central

Limits
Preview/Index
History
Clipboard
Details

Display
Abstract
Show
20
Sort By
Send to

All: 1
Review: 0


1: [J Neurosurg.](#) 2008 Dec 12. [Epub ahead of print] [Related Articles](#), [Links](#)



### Reduced local recurrence of a single brain metastasis through microscopic total resection.

[Yoo H.](#), [Kim YZ.](#), [Nam BH.](#), [Shin SH.](#), [Yang HS.](#), [Lee JS.](#), [Zo JI.](#), [Lee SH.](#)

1 Neuro-Oncology Clinic, Center for Specific Organs Cancer., 2 Cancer Registration and Biostatistics Branch; and, 3 Center for Lung Cancer, National Cancer Center, Goyang, Republic of Korea.

Object The goal of this study was to evaluate the therapeutic impact of the resection of metastatic brain tumor cells infiltrating adjacent brain parenchyma. Methods Between July 2001 and February 2007, 94 patients (67 males and 27 females, with a mean age of 55.0 +/-12.0 years) underwent resection of a single brain metastasis, followed by systemic chemotherapy with or without radiotherapy. In 43 patients with tumors located in noneloquent areas, the authors performed microscopic total resections (MTRs) that included tumor cells infiltrating adjacent brain parenchyma, and they pathologically confirmed during surgery that the resection margins were free of tumor cells (MTR group). In 51 patients with lesions in eloquent locations, gross-total resections (GTRs) were performed without the removal of neighboring brain parenchyma (GTR group). The 2 groups were then compared for local recurrence and survival. Results The MTR group had better local control of the tumor than did the GTR group; 10 (23.3%) of 43 patients in the MTR group and 22 (43.1%) of 51 patients in the GTR group had a local recurrence (p = 0.04). The median time to tumor progression in the MTR group could not be calculated using the Kaplan-Meier method, whereas it was 11.4 months in the GTR group. The 1- and 2-year respective local recurrence rates were 29.1 and 29.1% in the MTR group and 58.6 and 63.2% in the GTR group (p = 0.01). Multivariate analysis showed that the MTR procedure was associated with a decreased risk of local recurrence (p = 0.003). A Cox regression analysis revealed that the hazard ratio for a local recurrence in the MTR group versus the GTR group was 3.14 (95% CI 1.47-6.72, p = 0.003). There was no significant difference in the local recurrence rate between the MTR group without radiotherapy (10 [30.3%] of 33) and the GTR group with postoperative radiotherapy (5 [26.3%] of 19). Conclusions The results in this study suggest that MTRs including tumor cells infiltrating adjacent brain parenchyma for a single brain metastasis provide better local tumor control.

PMID: 19072310 [PubMed - as supplied by publisher]

Display
Abstract
Show
20
Sort By
Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)