

genome consensus sequence data provides information that represents the dominant virus subtype. It does not provide sufficient information to resolve transmission events particularly for rapidly spreading viruses. However, changes in the composition of minor variants between hosts and the pattern of minor variants fixation during outbreaks, could provide additional high-resolution data on who is infecting whom. The same data could also potentially inform the extent of within-host virus diversity as well as the proportion of diversity that is transmitted between individuals. We have developed a reproducible semi-automated whole genome variant calling pipeline to explore the role of minority variants in resolving transmission patterns and within host viral evolution. The pipeline is available as modular Bash scripts that run on a Linux cluster environment.

A28 Frequent co-infection among human group A rotaviruses in Thailand

M.A. Rabaa,^{1,2} T.T.N. Dung,¹ P.T. Duy,¹ L. Bodhidatta,³ B. Swierczewski,⁴ O.M. Sessions,⁴ C.J. Mason,³ S. Baker,^{1,2,5}

¹The Hospital for Tropical Diseases, Wellcome, Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam, ²Centre for Tropical Medicine, Oxford University, Oxford, UK, ³U.S. Armed Forces Research Institute of Medical Science, Bangkok, Thailand, ⁴Program in Emerging Infectious Diseases, Duke-National University of Singapore Graduate Medical School, Singapore, Singapore and ⁵The London School of Hygiene and Tropical Medicine, London, UK

Rotavirus (RoV) is a non-enveloped dsRNA virus in the Reoviridae family, with a 18.5-kb genome of 11 segments encoding six structural (VP1–4, VP6 and VP7) and five or six non-structural proteins (NSP1–NSP5/6). Reassortment between human and/or animal RoVs plays an important role in the generation of genetic diversity in these viruses, and is presumed to result from co-infection in human or animal reservoirs. However, co-infection with heterologous RoV has rarely been documented, in part due to inadequate detection methods and a lack of large-scale genomic investigations. Despite the availability of an efficacious vaccine, the burden of rotaviral diarrhea remains high in many developing countries, with rotavirus infection detected in 40–50% of all pediatric patients hospitalized with diarrhea. In addition to its cost, reduced vaccine effectiveness in developing country settings has contributed to its low uptake and the lack of government support for vaccination programs across Southeast Asia. The genetics and dynamics of rotavirus (RoV) have rarely been systematically investigated in these settings. The government of Thailand is preparing to add the rotavirus vaccine to its immunization program but has expressed concern regarding its effectiveness in the population and its long-term impact on rotavirus diversity and disease burden. To investigate the diversity of Group A RoV prior to the initiation of immunization programs, we performed full genome sequencing of 200 RoV from infected children across the country from 2004 to 2010 using novel in-house rotavirus capture and sequencing methods. Whilst the majority of samples (76%) showed infection with RoV of the common G1-P[8]-I1-C1-M1-A1-N1-T1-E1-H1 (Wa-like) genotype constellation (representing VP7-VP4-VP6-VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5, respectively), 41% of all samples additionally showed heterologous rotavirus co-infection, containing one or more segments representing an additional, less common genotype. Co-infection of G1-P[8] Wa-like viruses together with segments representing G9-P[19]-I5 lineages was particularly common and phylogenetic analysis suggests that these lineages may have spread through one region of Thailand for three years via serial co-transmission. G1-P[8]

Wa-like infection with additional G3 or G9 VP7 segments was also common across the country. Despite the high frequency of RoV co-infection in this setting, reassortant viruses were rarely observed (in ~8% of cases), and none of these appeared to represent reassortment among the common co-infecting viral segments detected within this study. The impacts of co-infection and reassortment on transmission fitness, the epidemiology of rotaviral diarrhea, and vaccine efficacy and long-term viral diversity warrant further investigation via full genome sequencing following the large-scale introduction of immunization programs in Southeast Asia.

A29 Wolbachia for dengue control; will dengue viruses evolve resistance?

T.T. Vi,¹ D.T. Kien,¹ B. Wills,^{1,2} C.P. Simmons,^{1,2,3}

¹Oxford University Clinical Research Unit, Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam, ²Centre for Tropical Medicine & Global Health, Oxford University, Oxford, UK and ³Dept. of Microbiology and Immunology, University of Melbourne, Parkville, Victoria, Australia

Dengue viruses (DENV) are major human pathogens, transmitted between people mainly by *Aedes aegypti* mosquitoes. Recently, the endosymbiont bacteria *Wolbachia* was horizontally introduced into *Ae. Aegypti*; *Wolbachia* infection non-specifically impairs DENV replication in these mosquitoes opening up a potential new strategy for dengue control. This project will test the hypothesis that DENV cannot easily develop resistance to the mechanisms by which *Wolbachia* blocks virus replication. Using contemporary viruses currently circulating in Vietnam, we will perform 30 cycles of repeated passage of DENV in wild-type and *Wolbachia*-infected *Ae. Aegypti* (wMel strain). This experimental system maximizes the possibility of observing selective pressure by wMel on DENV populations in mosquito tissues. We will use intermittent quantitative virology and virus genome sequencing to detect changes over time in the genotype and phenotype of DENV in *Ae. aegypti*. The results will provide valuable insights into how easily DENV adapt to *Wolbachia* in a controlled system. If *Wolbachia* specific evolution is observed, then we will test the infectiousness and replication profile of such viruses in cultured human cells. The results will deliver the first evidence of the likelihood of DENV “escape” from wMel, with implications for future and ongoing field releases of wMel in dengue-endemic countries.

A30 Recombination & evolution in two viral families: effective steps or a random walk?

R. Bingham,^{1,*} G. Leonov,¹ N. Patel,² E. Weiss,¹ S. White,² E. Dykeman,¹ P. Stockley,² R. Twarock,¹

¹York Centre for Complex Systems Analysis, University of York, York YO10 5GE, UK and ²Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds LS2 9JT, UK

Recombination or reassortment occurs in many different viral families and is understood as one of the predominant mechanisms of evolution. We perform a broad analysis of the mechanisms and recombinants of two viral families, Picornaviridae and Hepadnaviridae, which exhibit very different viral behaviour. Combining information about recombinants with knowledge of the evolutionary selection pressures on the genome and insights from the conservation of functional aspects of the genome, we find that viruses can dynamically exploit recombination in different ways, either to evade antigens and prolong infection, or to enhance viral diversity and allow the virus to infect new hosts. Better understanding of the mechanisms of