

Recruitment of Osteoclast Precursors by Purified Bone Matrix Constituents

J. D. MALONE, S. L. TEITELBAUM, G. L. GRIFFIN, R. M. SENIOR, and A. J. KAHN

Departments of Medicine, Pathology, and Biomedical Science, Washington University Schools of Medicine and Dental Medicine, and The Jewish Hospital of St. Louis, St. Louis, Missouri 63110

ABSTRACT The osteoclast, the multinucleated giant cell of bone, is derived from circulating blood cells, most likely monocytes. Evidence has accrued that is consistent with the hypothesis that the recruitment of monocytes for osteoclast development occurs by chemotaxis. In the present study, we have examined the chemotactic response of human peripheral blood monocytes and related polymorphonuclear leucocytes to three constituents of bone matrix: peptides from Type I collagen, α_2 HS glycoprotein, and osteocalcin (bone *gla* protein). The latter two substances are among the major noncollagenous proteins of bone and are uniquely associated with calcified connective tissue.

In chemotaxis assays using modified Boyden chambers, Type I collagen peptides, α_2 HS glycoprotein, and osteocalcin evoke a dose-dependent chemotactic response in human monocytes. No chemotaxis is observed in PMNs despite their ontogenetic relationship to monocytes and their documented sensitivity to a broad range of other chemical substances. Our observations are consistent with the view that osteoclast precursors (monocytes) are mobilized by chemotaxis, and suggest that the chemoattractants responsible for this activity are derived from the bone matrix or, in the case of collagen and osteocalcin, directly from the osteoblasts which produce them.

Osteoclasts are multinucleated giant cells that originate through the fusion of uninucleated precursors (1–4). Within the past five years, substantial evidence has accumulated which indicates that these precursors are derived from hematopoietic stem cells located in bone marrow, and that they are carried to sites of incipient or active bone resorption by the blood vascular system (5–7). The available evidence strongly suggests that this precursor cell either is a mononuclear phagocyte (i.e., a macrophage or monocyte) or is derived from the same progenitor which gives rise to mononuclear phagocytes (8–11).

It has been proposed that monocytes are recruited from the circulation to sites of osteoclast differentiation by chemotaxis (12–14). This suggestion follows from the observation that mononuclear phagocytes exhibit directional migration when placed in a gradient of conditioned medium collected from cultures of resorbing bone rudiments (15) and are chemotactically attracted to a partially purified factor isolated from neonatal mouse bone (16). Clearly, both conditioned media and bone extracts are likely to contain a number of factors capable of eliciting a chemotactic response in cells as broadly sensitive to environmental stimulation as mononuclear phagocytes (17), including some which are probably not present under normal conditions of bone turnover. However, the osteoclast is a cell specifically associated with bone and, consequently, it seems reasonable to suppose that, if chemotaxis is

involved in mobilizing osteoclast precursors, the chemoattractants originate from the cells or the extracellular matrix which characterize this tissue.

Mature bone matrix is a complex, highly mineralized (~65% dry weight) material built upon a structural framework of Type I collagen (~25–30%) and containing a comparatively small number of noncollagenous proteins and glycoproteins (<5%) (18). Two of the better characterized of the latter proteins are α_2 HS glycoprotein and osteocalcin. α_2 HS glycoprotein is synthesized in the liver and is selectively accumulated, relative to other plasma proteins, in bone and other mineralized tissues (19, 20). Osteocalcin (bone *gla* protein) is similarly associated with a range of mineralized tissues but, at least in bone, appears to be synthesized locally, probably by osteoblasts (21–23). Both osteocalcin and α_2 HS glycoprotein have been implicated in the mineralization process (18, 24).

In the present study, we have attempted to establish whether α_2 HS glycoprotein, osteocalcin, and mammalian collagenase-produced fragments of Type I collagen might serve as the chemoattractive agents involved in the recruitment of osteoclast precursors. It has already been shown that peripheral monocytes respond to bacterial enzyme digests of Type I collagen (17) and, in fact, to dipeptides and tripeptides containing glycine and proline, amino acids closely identified with the primary structure of collagen (25). Our data demonstrate that

human peripheral monocytes respond chemotactically to osteocalcin and α_2 HS glycoprotein as well as to collagen peptides derived through the action of mammalian collagenase. Polymorphonuclear leucocytes (PMNs) do not exhibit these responses despite their known sensitivity to other chemotactic stimuli (26) and their ontogenetic relationship to monocytes (27-29).

MATERIALS AND METHODS

Cell Isolation

Mononuclear cells and PMNs were separated from freshly drawn, heparinized human blood according to the Ficol-Hypaque density gradient technique (30). The cells were washed in Eagle's minimal essential medium with L-glutamine (α -MEM [alpha modification]; KC Biological, Lenexa, Kans.) and were suspended to a final concentration of 2.5×10^6 mononuclear cells or 1.5×10^6 PMNs per milliliter in α -MEM or in α -MEM supplemented with 10% heat-inactivated (60°C for 45 min) fetal calf serum (α -10 HI). 0.7 ml of cell suspension was then pipetted into the upper compartment of modified Boyden chambers (AHLCO Mfg. Co., Southington, Conn.).

Chemotaxis Assay

The chemotactic activities of osteocalcin, α_2 HS glycoprotein, and Type I collagen peptides were determined by use of the double-membrane technique developed by Campbell (31). The lower membrane was cellulose acetate, pore size $0.45 \mu\text{m}$ (Millipore Corp., Bedford, Mass.); the upper membrane (facing the cells) was polycarbonate (Nucleopore Corp., Pleasanton, Calif.), pore size either $2 \mu\text{m}$ for PMN assays or $5 \mu\text{m}$ for monocytes. The chambers were prepared in triplicate for each variable and were incubated in humidified air at 37°C. After 60 min (for PMNs) or 120 min (for monocytes), the membrane pairs were fixed in isopropanol, stained with hematoxylin, and mounted on slides. The slides were then coded, randomized, and read without reference to the code. Chemotactic activity was quantified on the basis of the number of cells located at the interface between the two membranes and falling within the boundaries of an eyepiece grid at $\times 400$ magnification. Five random grids were read for each membrane pair. The results were expressed as the mean number of cells/grid \pm SE, for each set of triplicate samples, and were corrected for random cell migration as determined from scoring control chambers having only medium in the lower compartment. Statistical analysis was made using Student's *t* test of means.

The positive control in all experiments was C_{5a} -derived chemotactic activity (C_{5a}), prepared from zymosan-activated human serum (a gift from Dr. D. Kreutzer, University of Connecticut, Farmington, Conn.) and standardized for 50% maximum chemotactic activity (ED_{50}) against human monocytes (32, 33). Twice the ED_{50} , in $25 \mu\text{l}$, was added to the lower compartment of each positive control chamber. Both the test substances and the C_{5a} were diluted to the desired concentration in either α -MEM or α -10 HI. These media were supplemented with 100,000 U/liter penicillin and 100 mg/liter streptomycin, and were filter-sterilized before use.

Preparation of Test Materials

Type I collagen peptides ($\sim 7,500$ mean mol wt) were produced by digesting native guinea pig skin collagen with purified mammalian collagenase and were generously provided by Dr. J. Jeffrey (Washington University School of Medicine, St. Louis, Mo.). Human α_2 HS glycoprotein (49,000 mol wt) was isolated from human serum using the purification sequence described by Triffitt et al. (20) and was kindly prepared for us by Dr. Triffitt (Medical Research Council Bone Research Laboratory, Oxford, England). Osteocalcin (5,670 mol wt) was extracted from lyophilized chicken bone with EDTA and purified by gel filtration and ion-exchange chromatography according to the methods described by Hauschka and collaborators (21). This substance, which was determined to be free of collagen contamination, was provided by Dr. Hauschka (Harvard University School of Dental Medicine, Cambridge, Mass.).

RESULTS

Human peripheral monocytes respond chemotactically to collagen peptides produced by the action of mammalian collagenase. The magnitude of the response is dose dependent within the range tested (6.25-100 $\mu\text{g/ml}$) and is optimal at 12.5-25 $\mu\text{g/ml}$ (1.7 - 3.3×10^{-6} M) (Fig. 1a). At these latter concentrations, collagen peptides are 40-50% as effective as the positive




FIGURE 1 (a and b) The chemotactic response of human peripheral monocytes to peptides from Type 1 collagen. Note that monocytes are optimally responsive to collagen peptides at concentrations of 12.5 and 25 $\mu\text{g/ml}$ (1.7 - 3.3×10^{-6} M) (a) and that the response is abolished in the absence of a gradient (b). The "fractions" used in labeling the abscissa indicate the location of the test substance in the Boyden chamber (top [upper] or bottom [lower] compartment) and its concentration in $\mu\text{g/ml}$. Striped bars denote chambers containing test material; open bars, chambers containing the positive control substance, C_{5a} . Values are expressed as net cells/HPF (high-power field) and are plotted \pm SEM.

control substance, C_{5a} . Net migration is not observed when the gradient is abolished by placing collagen peptides in both compartments of the Boyden chamber or exclusively in the cell-containing, upper compartment (Fig. 1b). Similarly, no chemotaxis is observed when mammalian collagenase alone is used as a test substance (data not shown).

Purified osteocalcin also promotes a dose-dependent chemotactic response in monocytes (Fig. 2a). This response, elicited over the range of 6.25-100 $\mu\text{g/ml}$, is optimal at 12.5-25 $\mu\text{g/ml}$ (2.2 - 4.3×10^{-6} M). Chemotaxis fails to occur if the osteocalcin is placed in both compartments of the Boyden chamber or if the protein is present only in the upper compartment (Fig. 2b). (In fact, in the particular experiment illustrated in Fig. 2b the presence of osteocalcin in the upper compartment significantly inhibited ($P < 0.01$) the net migration of cells into the membrane). Neither osteocalcin nor collagen peptides used over the same concentration range evoke a chemotactic response in PMNs (data not shown).




FIGURE 2 (a and b) The chemotactic response of human peripheral monocytes to osteocalcin. Note that the cells show significant directed migration to osteocalcin at concentrations ranging from 6.25 to 100 $\mu\text{g/ml}$ and that they are optimally responsive at 12.5 and 25 $\mu\text{g/ml}$ ($2.2\text{--}4.3 \times 10^{-6}$ M). Note also that a net positive migration is only observed when the chemoattractant is present in the lower (bottom) compartment of the Boyden chamber (b).

The limited availability of purified $\alpha_2\text{HS}$ glycoprotein precluded our fully assessing the chemotactic nature of this compound. However, the results, to date, demonstrate that the glycoprotein does elicit the increased migration of monocytes under conditions identical to those employed in collagen peptide and osteocalcin assays, and is effective at comparable concentrations, i.e., between 6.25 and 50 $\mu\text{g/ml}$ (1.3×10^{-7} to 1.02×10^{-6} M) (Fig. 3).

DISCUSSION

Osteoclast development occurs under physiological conditions at sites within the skeleton which are undergoing (or about to undergo) bone resorption. The first step in this process, the mobilization of precursor cells, appears to result from the chemotaxis of mononuclear phagocytes (15, 16), cells known to exhibit directional migration in response to a variety of chemical agents (17). Since osteoclast development is a tissue-specific phenomenon and is generally linked to other events associated with normal bone turnover, it follows that the chemotactic factor (or factors) responsible for mobilizing the precursor cells should exhibit a similar tissue specificity and become available as a consequence of the processes of bone formation or resorption.

Because the organic constituents of bone matrix largely




FIGURE 3 The in vitro migration of human peripheral monocytes in a gradient of $\alpha_2\text{HS}$ glycoprotein. Note that the magnitude of the response to this serum glycoprotein and the effective concentration (1.3×10^{-7} to 1.02×10^{-6} M) are similar to those observed with collagen peptides and osteocalcin.

define the microenvironment in which osteoclasts are formed, these constituents would clearly fulfill the "geographic" requirement for chemoattractants which might be involved in the recruitment of osteoclast precursor cells. In addition, such substances would become available as recruiting agents when bone matrix is broken down during osteolysis or, in the case of matrical components produced *in situ*, as a direct result of secretion by osteoblasts lining the bone surface. In fact, as documented below, the bone literature contains reports consistent with both modes of chemoattractant release.

When bone is stimulated to resorb, by any of a number of chemical agents, the osteoclast population characteristically responds with a burst of resorptive activity by extant cells followed, after a lag period of several hours, by the generation of new, and often larger, osteoclasts (34–37). During this lag period, the mineral component of the bone matrix is mobilized and, along with it, $\alpha_2\text{HS}$ glycoprotein, osteocalcin, and, probably, collagen. These substances would then serve to attract monocytes (precursor cells) to sites of osteoclast differentiation. In support of this possible mechanism of precursor recruitment, we note that demineralization is the first phase of bone resorption (38), and that both $\alpha_2\text{HS}$ glycoprotein and osteocalcin are released from bone when the mineral component of the matrix is removed (19, 24). We also note that urinary excretion of hydroxyproline (presumably derived from the degradation of bone collagen) is elevated during periods of bone resorption (39) and that plasma levels of osteocalcin (bone *gla* protein) are increased in bone disease characterized by high bone turnover (40).

There are other circumstances, such as those encountered in embryonic and fetal life, where osteoclasts appear *de novo*, in the absence of any apparent, prior resorptive activity. Here, it seems reasonable to speculate that the secretion of soluble chemoattractive matrix constituents by osteoblasts is responsible for attracting osteoclast precursor cells. In support of this proposition, we note that osteoblast-derived osteogenic sarcoma cells synthesize and release osteocalcin (bone *gla* protein) *in vitro* (41), and that the level of synthesis of this compound can be significantly increased by exposing the cells to parathyroid hormone (PTH) (42). In addition, the serum level of bone *gla* protein is increased in rats after the injection of $1,25(\text{OH})_2\text{D}_3$ (43). Finally, we call attention to the growing

body of evidence indicating that PTH acts to enhanced resorption in vitro via receptors located on osteoblasts rather than osteoclasts (44).

The results of the present study demonstrate that monocytes respond chemotactically to osteocalcin, to α_2 HS glycoprotein, and to Type I collagen peptides produced through the action of a purified mammalian collagenase. By contrast, PMNs do not respond to either osteocalcin or collagen peptides despite the fact that they are derived from the same hematopoietic stem cell as monocytes (27–29) and are chemotactically sensitive to a broad spectrum of other substances (25). These findings support the hypothesis that chemotaxis plays a major role in the recruitment of osteoclast precursor cells, and suggest that the likely sources for the relevant chemoattractive agents are the bone matrix and the osteoblast.

We thank Drs. Kreuzer, Jeffrey, Triffitt, and Hauschka for their generosity in providing the purified bone-matrix-associated macromolecules and the C_{5a} used in these experiments, Mr. Michael Richards for his technical assistance in performing the chemotaxis assays, and Ms. K. Farris for typing the manuscript.

This research was supported, in part, by funds from the Jewish Hospital of St. Louis and by grants DE-04629-04, HL-16118, and DE-05413-01 from the National Institutes of Health.

Received for publication 18 August 1981, and in revised form 8 October 1981.

REFERENCES

- Hancox, N. M. 1949. The osteoclast. *Biol. Rev. Camb. Philos. Soc.* 24:448–469.
- Hall, B. K. 1975. The origin and fate of osteoclasts. *Anat. Rec.* 183:1–12.
- Owen, M. 1978. Histogenesis of bone cells. *Calcif. Tissue Int.* 25:205–207.
- Chambers, T. J. 1978. Multinucleate giant cells. *J. Pathol.* 126:125–148.
- Kahn, A. J. and D. J. Simmons. 1975. Investigation of cell lineage in bone using a chimaera of chick and quail embryonic tissue. *Nature (Lond.)* 258:325–327.
- Coccia, P. F., W. Krivit, J. Cervinka, C. C. Clawson, J. H. Kersey, T. H. Kim, M. E. Nesbit, N. K. C. Ramsay, P. I. Warkentin, S. L. Teitelbaum, A. J. Kahn, and D. M. Brown. 1980. Successful bone marrow transplantation for infantile malignant osteopetrosis. *N. Engl. J. Med.* 302:701–708.
- Ash, P., J. F. Loutit, and K. M. S. Townsend. 1980. Osteoclasts derived from haemopoietic stem cells. *Nature (Lond.)* 283:669–670.
- Fischman, D. A., and E. D. Hay. 1962. Origin of osteoclasts from mononuclear leukocytes in regenerating new limbs. *Anat. Rec.* 143:329–337.
- Jee, W. S. S., and P. D. Nolan. 1963. Origin of osteoclasts from the fusion of phagocytes. *Nature (Lond.)* 200:225–226.
- Buring, K. 1975. On the origin of cells in heterotopic bone formation. *Clin. Orthop. Relat. Res.* 110:293–302.
- Gothlin, G., and J. L. E. Ericsson. 1976. The osteoclast. Review of ultrastructure. *Clin. Orthop. Relat. Res.* 120:201–231.
- Minkin, C., and G. Trump. 1977. Mononuclear phagocytes and bone remodeling: bone mediated macrophage chemotaxis. In *Mechanisms of Localized Bone Loss*. J. E. Horton, T. M. Tarpley, and W. F. Davis, editors. Informational Retrieval, Inc., Washington, D.C. 438 (Abstr.).
- Mundy, G. R., and T. Yoneda. 1981. Monocytes: interactions with bone cells and lymphocytes during bone resorption. In *Hormonal Control of Calcium Metabolism*. D. V. Cohn, R. V. Talmage, and J. L. Matthews, editors. Excerpta Medica, Amsterdam. 178–181.
- Kahn, A. J., J. D. Malone, and S. L. Teitelbaum. 1981. Mononuclear phagocytes respond to "bone seeking" hormones and bone matrix constituents. In *Hormonal Control of Calcium Metabolism*. D. V. Cohn, R. V. Talmage, and J. L. Matthews, editors. Excerpta Medica, Amsterdam. 182–189.
- Mundy, G. R., J. Varani, W. Orr, M. D. Gondek, and P. A. Ward. 1978. Resorbing bone is chemotactic for monocytes. *Nature (Lond.)* 275:132–135.
- Minkin, C. R. Posek, and J. Newbrey. 1981. Mononuclear phagocytes and bone resorption: identification and preliminary characterization of a bone-derived macrophage chemotactic factor. *Metabolic and Bone Disease and Related Research*. 2:363–369.
- Postlethwaite, A. E., and A. H. Kang. 1976. Collagen- and collagen peptide-induced chemotaxis of human blood monocytes. *J. Exp. Med.* 143:1299–1307.
- Triffitt, J. T. 1980. Organic matrix of bone tissue. In *Fundamental and Clinical Bone Physiology*. M. R. Urist, editor. J. B. Lippincott and Co., New York. 45–82.
- Ashton, B. A., H. J. Hohling, and J. T. Triffitt. 1976. Plasma proteins present in human cortical bone: enrichment of the α_2 HS-glycoprotein. *Calcif. Tissue Res.* 22:27–33.
- Triffitt, J. T., M. E. Owen, B. A. Ashton, and J. M. Wilson. 1978. Plasma disappearance of rabbit α_2 HS-glycoprotein and its uptake by bone tissue. *Calcif. Tissue Res.* 26:155–161.
- Hauschka, P. V., J. B. Lian, and P. M. Gallop. 1975. Direct identification of the calcium-binding amino acid, γ -carboxyglutamate, in mineralized tissue. *Proc. Natl. Acad. Sci. U. S. A.* 72:3925–3929.
- Price, P. A., A. S. Otsuka, J. W. Poser, J. Kristaponis, and N. Raman. 1976. Characterization of a γ -carboxyglutamic acid-containing protein from bone. *Proc. Natl. Acad. Sci. U. S. A.* 73:1447–1451.
- Hauschka, P. V., and M. L. Reid. 1978. Vitamin K dependence of a calcium-binding protein containing γ -carboxyglutamic acid in chicken bone. *J. Biol. Chem.* 253:9063–9068.
- Hauschka, P. V., and M. L. Reid. 1978. Timed appearance of a calcium-binding protein containing γ -carboxyglutamic acid in developing chicken bone. *Dev. Biol.* 65:426–434.
- Zigmond, S. H. 1978. Chemotaxis by polymorphonuclear leukocytes. *J. Cell Biol.* 77:269–287.
- Wilkinson, P. C. 1974. Chemotaxis and Inflammation. Churchill Livingstone, Edinburgh.
- Bradley, T. R. and D. Metcalf. 1966. The growth of mouse bone marrow cells in vitro. *Aust. J. Exp. Biol. Med. Sci.* 44:287–300.
- Metcalf, D., and M. A. S. Moore. 1971. Haemopoietic cells. *Front. Biol.* Vol. 24.
- Williams, N., and R. R. Eger. 1978. Purification and characterization of clonable murine granulocyte-macrophage precursor cell populations. In *Hematopoietic Cell Differentiation*. D. W. Golde, M. J. Cline, D. Metcalf, and C. F. Fox, editors. Academic Press, Inc., New York. 385–398.
- Boym, A. 1968. Isolation of mononuclear cells and granulocytes from human blood. *Scand. J. Clin. Lab. Invest.* 21(Suppl. 97):77–89.
- Campbell, P. B. 1977. An improved method for the in vitro examination of monocyte leukotaxis. *J. Lab. Clin. Med.* 90:381–388.
- O'Flaherty, J. T., C. L. Kreuzer, and P. A. Ward. 1977. Neutrophil aggregation and swelling induced by chemotactic agents. *J. Immunol.* 119:232–239.
- Kreuzer, D. L., and W. D. Claypool, M. L. Jones, and P. A. Ward. 1979. Isolation by hydrophobic chromatography of the chemotactic factor inactivators from human serum. *Clin. Immunol. Immunopathol.* 12:162–176.
- Holtrop, M. R., L. G. Raisz, and H. A. Simmons. 1974. The effects of parathyroid hormone, calcichine, and calcitonin on the ultrastructure and the activity of osteoclasts in organ culture. *J. Cell Biol.* 60:346–355.
- Müller, S. C. 1978. Rapid activation of the medullary bone osteoclast cell surface by parathyroid hormone. *J. Cell Biol.* 76:615–618.
- Holtrop, M. E., and L. G. Raisz. 1979. Comparison of the effects of 1,25-dihydroxycholecalciferol, prostaglandin E_2 , and osteoclast-activating factor with parathyroid hormone on the ultrastructure of osteoclasts in cultured long bones of fetal rats. *Calcif. Tissue Int.* 29:201–205.
- Rowe, D. J., and E. Hausmann. 1977. Quantitative analyses of osteoclast changes in resorbing bone organ cultures. *Calcif. Tissue Res.* 23:283–289.
- Bonucci, E. 1974. The organic-inorganic relationships in bone matrix undergoing osteoclastic resorption. *Calcif. Tissue Res.* 16:13–36.
- Smith, R. 1980. Collagen and disorders of bone. *Clin. Sci.* 215–223.
- Price, P. A., J. G. Parthemore, and L. J. Deftos. 1980. New biochemical marker for bone metabolism. *J. Clin. Invest.* 66:878–883.
- Nishimoto, S. K., and P. A. Price. 1980. Secretion of the vitamin-K-dependent protein of bone by rat osteosarcoma cells. *J. Biol. Chem.* 255:6579–6583.
- Rodan, G. A., R. J. Majeska, S. B. Rodan, and J. J. Egan. 1981. The use of osteosarcoma clonal variants in the study of PTH effects on osteoblastlike cells. In *Hormonal Control of Calcium Metabolism*. D. V. Cohn, R. V. Talmage, and J. L. Matthews, editors. Excerpta Medica, Amsterdam. 191 (Abstr.).
- Price, P. A., and S. A. Baukol. 1981. 1,25-dihydroxyvitamin D_3 increases serum levels of the vitamin K-dependent bone protein. *Biochem. Biophys. Res. Commun.* 99:928–935.
- Rodan, G. A., and T. J. Martin. 1981. The role of osteoblasts in hormonal control of bone resorption—a hypothesis. *Calcif. Tissue Int.* 33:349–351.