

Guanosine 5'-monophosphate-chelated calcium and iron feed additives maintains egg production and prevents *Salmonella Gallinarum* in experimentally infected layers

Hye-Ji Noh¹, HeeKyong Kim¹, Su Jeong Heo², Hyang Hyun Cho³, Hong Bum Koh^{2,*}

¹Medinutrol Co., Ltd., Yeonggwang 57024, Korea

²College of Veterinary Medicine, Chonnam National University, Gwangju 61186, Korea

³Hanilbiomed Inc., Gwangju 62363, Korea

We evaluated the effects of guanosine 5'-monophosphate (GMP)-chelated calcium and iron (CaFe-GMP) on health and egg quality in layers experimentally infected with *Salmonella Gallinarum*. In this study, a CaFe-GMP feed additive was added to a commercial layer feed and fed to layers over a four-week period. All were inoculated with *Salmonella Gallinarum*. Body weight, mortality, clinical symptoms, and poultry production including feed intake, egg production, egg loss, and feed conversion rate were observed, and *Salmonella Gallinarum* was re-isolated from the liver, spleen, and cecum of the layers. All tested internal organs for the CaFe-GMP additive group exhibited significantly lower re-isolation numbers of *Salmonella Gallinarum* and less severe pathological changes than those in the control group, indicating that the CaFe-GMP feed supplement induced bacterial clearance and increased resistance to *Salmonella Gallinarum*. Additionally, due to the inhibitory action of CaFe-GMP on the growth of *Salmonella Gallinarum*, the CaFe-GMP additive group exhibited better egg production, including a higher laying rate and fewer broken eggs. The results suggest that a 0.16% CaFe-GMP additive may help prevent salmonellosis in the poultry industry.

Keywords: *Salmonella Gallinarum*, feed additives, layer, organic mineral

Introduction

Fowl typhoid is caused by *Salmonella Gallinarum*, the transmission route of which is horizontal, *i.e.*, from an infected ovary to the egg and the progeny [3]. The disease can be either acute or chronic and usually results in a high mortality rate (60%) and increased morbidity [2,29]. Although *Salmonella Gallinarum* has adapted to its avian host and rarely induces food poisoning in humans, fowl typhoid outbreaks cause severe economic losses worldwide [36].

The worldwide increase in multi-antibiotic resistance in bacteria is a problem caused by major drug resistant nosocomial pathogens. As a result of multiple antibiotic inefficiencies, the mortality rate caused by *Salmonella* has increased [12]. Recently, several attempts have been made to prevent salmonellosis by using alternatives to antibiotics, such as vaccination, bacteriophages, probiotics, and herbs [10,13,15]. However, these methods have demonstrated limited efficiency. Therefore,

the development of efficacious feed additives to prevent poultry salmonellosis is a major research goal [1,20].

Formulation of balanced diets is fundamental to economical poultry production, and this process depends on knowledge of the nutrient requirements of poultry and the nutritional attributes of various feed sources. Minerals are naturally occurring inorganic solids and dietary nutrients essential in the maintenance of homeostasis for laying hens [22]. In addition, minerals have important roles in pathogen virulence and host antimicrobial resistance [27,34]. Calcium enhances intestinal resistance to infectious diseases and protects against *Salmonella*- and enterotoxigenic *Escherichia coli* (ETEC)-associated diarrhea [5,6]. Iron deficiency is associated with lowering resistance to infection, whereas iron supplements reverse this effect [21,23,26].

Recently, organic minerals (chelated minerals), because of their potential for higher bioavailability than inorganic minerals, have been studied by several researchers [19]. Organic minerals include any mineral bound such that it is chelated or bonded to

Received 24 Aug. 2016, Revised 12 Sep. 2016, Accepted 23 Nov. 2016

*Corresponding author: Tel: +82-62-530-2849; Fax: +82-62-530-2809; E-mail: hbkoh@chonnam.ac.kr

Journal of Veterinary Science · © 2017 The Korean Society of Veterinary Science. All Rights Reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pISSN 1229-845X

eISSN 1976-555X

organic molecules, such as an amino acid, thereby forming chemical structures with unique characteristics such as stability and high mineral bioavailability. Several reports have indicated that organic sources of trace minerals have a higher bioavailability than that of inorganic forms [4,19,37]. After absorption, organic minerals may induce physiological effects, which could improve specific metabolic responses, such as the immune response. Many studies have demonstrated the benefits of metal-amino acid chelates on animal metabolism [33].

In this study, we evaluated the effects of the feed additive guanosine 5'-monophosphate (GMP)-chelated calcium and iron (CaFe-GMP), a metal-nucleotide chelate, against salmonellosis in layers.

Materials and Methods

Preparation of organic mineral feed additives

The CaFe-GMP was provided by Medinutrol (Korea). CaFe-GMP, an organic formulation in which calcium and iron are provided as GMP chelates was used as a supplement to the layers basal diet. A commercial, nutritionally complete, antibiotic-free, general layer feed was used as a basal diet to meet the nutrient requirements for layers in accordance with Nutrient Requirements of Poultry recommendations [22]. The basal diet was formulated to contain a given chemical composition (Table 1).

In our previous study, the potential effect of prebiotics on the growth of *Lactobacillus* spp. in layers using a feed supplement of 0.16% CaFe-GMP, which contained 83 ppm calcium (Ca) and 55 ppm iron (Fe) (unpublished data) was investigated. In the present experiment, for the CaFe-GMP additive group, CaFe-GMP was completely mixed into the basal diet to form 0.16% CaFe-GMP containing 83 ppm Ca and 55 ppm Fe (Table 1).

Animal experiment

Hy-line brown layer chickens (34 weeks old) with no history of salmonellosis were obtained from a local farm. Microbiological testing did not detect *Salmonella* Gallinarum in the feces of the chickens. All chickens were maintained at the Chonnam National University College of Veterinary Medicine (Chonnam, Korea) in separate air-controlled rooms. Chickens were allowed free access to tap water, and their respective diets were supplied three times per day. All experimental procedures were approved by the Institutional Animal Care and Use Committee of the Chonnam National University (approval No. CNU IACUC-YB-2015-59).

After a 7-day acclimation period, the chickens were divided into three groups with six chickens in each group. The groups were as follows: normal group, fed with a normal commercial layer feed without *Salmonella* Gallinarum inoculation; control group, fed with a normal commercial layer feed but with a *Salmonella* Gallinarum inoculation; additive group, fed with a normal commercial layer feed supplemented by the addition of

CaFe-GMP and with *Salmonella* Gallinarum inoculation. All chickens were fed their respective diet for 4 weeks, and all but the normal group were inoculated with *Salmonella* Gallinarum. The experiment was conducted in triplicate.

Preparation of *Salmonella* Gallinarum (inoculum)

Salmonella Gallinarum (SG3001) for experimental inoculation was provided by the Animal and Plant Quarantine Agency (Korea). The SG3001 was cultured on MacConkey agar plates overnight. One colony was selected and inoculated in Brain Heart Infusion (Difco Laboratories, USA) broth and incubated at 10 × g for 6 h at 37°C. Viable counts of the broth

Table 1. Composition of the basal diet

Ingredients (%)	Layer feed (30–39 weeks of age)	
Yellow corn	48.3	
Wheat	4.39	
Soybean meal	13	
Distiller's dried grains with solubles	13.2	
Canola meal	5	
Sesame oil meal	2	
Corn gluten	1.5	
Oil	1	
Salt	0.25	
Calcium carbonate (CaCO ₃)	10.5	
Mono-dicalcium phosphate	0.5	
Dicalcium phosphate	0.1	
Choline chloride	0.06	
Vitamin premix*	0.1	
Trace mineral premix [†]	0.1	
Calculated chemical composition	Normal/control group	CaFe-GMP group
ME (kcal/kg)	2,800	2,800
Crude protein (%)	17	17
Crude ash (%)	15.5	15.5
Calcium (%)	3.9	3.9
Available phosphorus (%)	0.35	0.35
Methionine (%)	0.67	0.67
Methionine + Cystine (%)		
Calcium (ppm)	–	83 [‡]
Iron (ppm)	–	55 [§]

*Supplied per kg of diet: vitamin A 13,000,000 IU, vitamin D3 4,000,000 IU, vitamin B1 3,000 mg, vitamin B2 7,000 mg, vitamin B6 5,000 mg, vitamin B12 20 mg, lysin 3,000 mg, methionine 1,000 mg, tryptophane 700 mg, pantothenic acid 12,000 mg, niacin 50,000 mg, biotin 150 mg, and folic acid 1,500 mg. [†]Supplied per kg of diet: Iron 46 mg, zinc 78 mg, copper 7 mg, Selenium 0.1 mg, Manganese 87 mg, iodine 1.3 mg, and chromium 0.12 mg. [‡]Guanosine 5'-monophosphate (GMP)-chelated calcium (organic mineral form). [§]Guanosine 5'-monophosphate (GMP)-chelated iron (organic mineral form). ME, metabolizable energy.

culture were obtained by using the spread plate method. Inoculated broth was diluted with phosphate buffer saline (PBS; Sigma-Aldrich, USA) to obtain an infective dose of 1×10^8 CFU/mL (CFU, colony-forming unit) (the optimal dose was determined in preliminary experiments). Experimental inoculations were administered orally.

Body weight, clinical symptoms, and pathological changes

Body weights of chickens were measured at 0 and 5 days post-infection (dpi). The chickens were monitored daily for clinical signs, including anorexia, diarrhea, and depression, among others. Postmortem examinations of all dead and sacrificed chickens were conducted and any pathological changes associated with fowl typhoid were noted. Additionally, the mortality rate in each group during the experimental period was recorded.

Experimental parameters measured of poultry production

Egg production (eggs/chicken/day, %), egg loss (broken and cracked eggs, %), and egg weight (g) were recorded daily. Feed intake (g/chicken/day) was calculated by weighting the initial feed input against the uneaten food over the experimental period. Feed conversion rate (FCR; g feed/g egg) were determined on a daily basis. FCR was expressed as kilograms of feed consumed per kilogram of egg produced. Mortality rate (%) was recorded daily as it occurred.

Re-isolation of *Salmonella Gallinarum* from internal organs

Re-isolation numbers for *Salmonella Gallinarum* was determined for liver, spleen, and feces collected from each sacrificed chicken at 5 dpi. In addition, feces were retrieved from sterilized paper that was used to line the chicken's housing unit. For non-selective pre-enrichment processing, 0.1 g of tissue and feces from the cecum of each sample were aseptically collected and added to 10 mL of sterile Buffered Peptone Water (Sigma-Aldrich), homogenized completely, and incubated at 37°C for 24 h. Then, the homogenate was transferred from 1 mL of the pre-enrichment broth to 10 mL Tetrathionate Broth (TTB; Difco Laboratories) and incubated at 37°C for 24 h for selective enrichment processing. The TTB was serially diluted 10-fold in PBS, and 50 µL of each dilution was spread onto a *Salmonella Shigella* agar (Difco Laboratories) plate and incubated at 37°C for 24 h. The resulting characteristic black-colored colonies were counted and expressed as CFU/0.1 g tissue, but only for those plates with counts of 30 to 300 colonies per plate.

Statistical analysis

The significance of treatment effects was analyzed by comparing mean \pm SEM values. Statistical analysis was performed by using analysis of variance (ANOVA) and Duncan's multiple-range tests. Differences were considered significant at $p < 0.05$.

Results

Changes in body weight and clinical symptoms after *Salmonella Gallinarum* infection

The adverse effects of the *Salmonella Gallinarum* infection induced in the present study were assessed by examining body weight and clinical symptoms. For each group, body weight change was compared between mean weight at 0 dpi for all chickens in a group to that at 5 dpi in the surviving chickens. The body weights of each group at 0 and 5 dpi are shown in Table 2. The non-inoculated, normal group did not exhibit a change in body weight, whereas the inoculated, control group (1,741.0 g \pm 125.0 g) had significantly lower body weight at 5 dpi than at 0 dpi (1,905.7 g \pm 155.3 g), a loss of approximately 8.6% ($p < 0.07$). However, the inoculated, CaFe-GMP additive group exhibited only slightly lower body weight at 5 dpi (1,493.6 g \pm 59.4 g) than at 0 dpi (1,598.0 g \pm 34.3 g); the change was not significant. The percentage decline in body weight for the CaFe-GMP additive group was less than that of the control group. Clinical signs (anorexia, diarrhea, and depression) were observed in the control group, but the CaFe-GMP additive group showed very few typical clinical signs of *Salmonella Gallinarum*.

Changes in mortality rate of layers experimentally infected with *Salmonella Gallinarum*

Mortality rate is a major evaluation marker used in treatment toxicity analysis for preclinical studies. Mortality rates in the three trial groups are summarized in Table 3. Mortality was first observed at 4 dpi in the control group, whereas there was no mortality in the CaFe-GMP additive group or the normal group during the experimental period. Although low levels of clinical symptoms of *Salmonella Gallinarum* were observed in the CaFe-GMP additive group, the group suffered no mortality from the infection.

Changes in feed intake rate in layers experimentally infected with *Salmonella Gallinarum*

For each group, feed intake was compared at 0 dpi for all

Table 2. Effect of dietary supplementation of organic minerals on the body weight in layer groups, post-challenge

Group	Mean body weight per chicken (g)	
	0 dpi	5 dpi
Normal	1,522 \pm 167	1,544 \pm 174
Control	1,905 \pm 155	1,712 \pm 115
CaFe-GMP	1,598 \pm 34	1,493 \pm 59

dpi, days post-infection.

Table 3. Effect of dietary supplementation of organic minerals on performance parameters and egg production characteristics in layer groups, post-challenge

Group	Mean parameters per chicken (g)					
	0 dpi			5 dpi		
	Normal	Control	CaFe-GMP	Normal	Control	CaFe-GMP
Feed intake (g/chicken/d)	93.9 ± 12.9	110.7 ± 4.5	100.8 ± 7.0	99.5 ± 8.1	91.4 ± 29.9*	102.1 ± 9.5
Egg production (%)	77.8	87.5	83.3	86.7	80	84
Cracked/broken egg (%)	0	0	0	0	3.3	0
Egg weight (g)	56.65 ± 2.50	61.09 ± 5.30	57.26 ± 1.95	54.99 ± 5.35	61.32 ± 3.64	57.54 ± 1.50
FCR (g feed/g egg)	1.36 ± 0.20	1.55 ± 0.17	1.45 ± 0.10	1.51 ± 0.13	1.33 ± 0.30	1.48 ± 0.12
Mortality rate (%)	0	0	0	0	20	0

FCR, feed conversion rate. * $p < 0.05$ compared with 0 dpi.

Table 4. Effect of dietary supplementation of organic minerals on the weight of spleen in layer groups, post-challenge


Group	Weight of spleen (g)
Normal	1.6 ± 0.3
Control	4.7 ± 1.5*
CaFe-GMP	2.6 ± 1.6 [†]

* $p < 0.05$ compared with normal group. [†] $p < 0.05$ compared with control group.

chickens and at 5 dpi for all surviving chickens. The mean feed intake for chickens in the normal, CaFe-GMP additive, and control groups at 0 and 5 dpi are shown in Table 3. Compared to 0 dpi, feed intake in the control group decreased significantly by approximately 17.44% at 5 dpi, whereas in the CaFe-GMP additive group it increased slightly (1.2%) at 5 dpi; a non-significant change. The slight increase in feed intake of the CaFe-GMP additive group was less than that in the normal group, although neither increase was significant.

Evaluation of egg production and quality

Average egg production rate for the CaFe-GMP additive group was not significantly different from that of the control group before the *Salmonella Gallinarum* infection period (control group: 87.5%, CaFe-GMP group: 83.3%). After *Salmonella Gallinarum* infection, the CaFe-GMP additive group maintained a similar laying rate, whereas the laying rate for the control group decreased by 7.5% from that in the uninfected experimental period. The percentage of cracked or broken eggs for the control group after *Salmonella Gallinarum* infection was 3.3% at 5dpi, whereas no broken eggs occurred in the CaFe-GMP additive group during the experimental period (Table 3).

**Fig. 1.** Effect of dietary supplementation of organic minerals on the macroscopic change in liver in layer groups post-challenge. Representative images from the (A) normal group, (B) control group and (C) CaFe-GMP group. Scale bars = 5 cm.

Macroscopic changes of spleen and liver

As shown in Table 4, the control group had significantly heavier spleens than those in the normal group. The CaFe-GMP additive group had a spleen weight significantly lower than that in the control group ($p < 0.05$); moreover, spleen size was similar to that in the normal group (data not shown). In addition, hepatomegaly was reduced in the CaFe-GMP additive group (Fig. 1). Furthermore, necropsy revealed that the CaFe-GMP additive group exhibited milder and less severe gross pathological abnormal changes when compared with those in the control group.

Bacterial clearance in internal organs

Surviving chickens from each group were sacrificed at 5 dpi. The liver, spleen, and feces from the cecum were aseptically collected and tested to determine whether *Salmonella Gallinarum* used for the challenge could be re-isolated from organs of the experimental chickens. The liver, spleen, and feces samples were negative for *Salmonella Gallinarum* in the normal group. In the control group, bacteria were re-isolated at 5 dpi from the liver, spleen, and cecum, indicating a systemic infection had

Table 5. Effect of dietary supplementation of organic minerals on the re-isolation number of *Salmonella* in layer groups, post-challenge

Group	Mean CFU per chicken at 5 dpi (10 ⁶ CFU/0.1 g)			
	Liver	Spleen	Feces in cecum	Paper litter from housing
Normal	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0
Control	9,751.15 ± 3,863.05*	791.07 ± 126.67*	1,833.70 ± 4,483.24*	1,571.34 ± 358.10*
CaFe-GMP	5,751.30 ± 3,389.94	336.890 ± 230.78 [†]	0.08 ± 0.17	1.50 ± 0.17 [†]

CFU, colony-forming unit. * $p < 0.05$ compared with normal group. [†] $p < 0.05$ compared with control group.

occurred. As shown in Table 5, the numbers of *Salmonella* Gallinarum colony forming units in the CaFe-GMP additive group were considerably lower at 5 dpi in the liver (45.7%, $p < 0.06$), spleen (57.4%, $p < 0.006$), and feces from the cecum (96.6%, $p < 0.057$) than those in the control group. In particular, *Salmonella* Gallinarum detection was significantly cleared in excreted feces on sterilized paper litter from the housing units of the CaFe-GMP additive group (99.9%, $p < 0.001\%$) at 5 dpi when compared with that of the control group.

Discussion

Avian systemic salmonellosis is primarily caused by the *Salmonella* Gallinarum strain and results in the development of fowl typhoid disease [7]. Fowl typhoid can be acute or chronic and usually results in a high mortality rate [2]. Symptoms include diarrhea, intestinal and womb bleeding, and enlargement of the liver and spleen.

Antibiotics may be used as feed supplements to improve physical performance by suppressing subclinical disease challenge in industrial animals [8]. However, there is worldwide concern regarding the overuse of antibiotics because of the development of multi-antibiotic resistant bacteria and resistance genes and their spread from animals to humans through consumption of food products that contain antibiotic residues [16]. Hence, there is an urgent need to identify natural-origin feed supplements that can reduce antibiotic use. Several studies have demonstrated that prevention of *Salmonella* colonization in chickens can be achieved by feeding with prebiotics, such as saccharides, organic acid, and inorganic minerals [14,24,25,28,37,38].

Mineral additives in poultry feed are needed for bird survival and for efficient poultry production. The mineral requirements of laying hens are well reported [22], but information on trace mineral requirements and their related effects on various aspects such as immune function, resistance to disease, growth, and carcass composition is limited. In this study, we evaluated the effect of a CaFe-GMP feed supplement on the health and egg quality in layers experimentally infected with *Salmonella* Gallinarum.

Our results demonstrated that administering 0.16% CaFe-GMP

as a feed additive had a beneficial effect on body weight in layers after *Salmonella* Gallinarum infection. It has been reported that mice on an iron-deficient diet gained less weight and ate less [32], suggesting that chickens fed a diet with an iron additive may exhibit less weight loss during intestinal inflammation by *Salmonella* Gallinarum infection.

Salmonella Gallinarum causes a high mortality rate in fowl [9]. However, in the present study, mortality did not occur in the *Salmonella* Gallinarum-infected CaFe-GMP additive group during the experimental period. Moreover, clinical signs of an infection were delayed, and the severity of related symptoms was lower in the CaFe-GMP additive group than in the control group. In the control group, there was severe, creamy whitish diarrhea observed from 4 dpi onwards, whereas in the CaFe-GMP additive group there were few typical clinical symptoms of *Salmonella* Gallinarum infection observed. In addition, the CaFe-GMP additive group had no fatalities. These results are similar to previous results showing that parenteral administration of complexed iron can enhance survival in chicks infected with *Salmonella* Gallinarum endotoxin (LPS) [30].

With regard to feed intake, the amount of feed consumed in the control group abruptly decreased, supporting results reported by Kim *et al.* [17]. In the *Salmonella* Gallinarum-infected CaFe-GMP additive group, there were no significant changes in feed consumption during the experimental period. Moreover, our results showed that the CaFe-GMP additive group displayed no significant change in growth performance.

Infection of fowl by *Salmonella* Gallinarum results in a decline in egg production rate and broken eggs [11,31]. Interestingly, the infected CaFe-GMP additive group retained a normal egg production rate and broken eggs were not observed during the experimental period. Similarly, others have shown that dietary supplementation with a Fe-soy proteinate effectively improved egg quality [28], and dietary supplementation of an organic mineral mixture increased the bioavailability of minerals compared to that from inorganic sources [37]. CaFe-GMP supplementation may protect against the penetration of *Salmonella* spp. into the oviduct of layers and, therefore, could increase economic profits by maintaining egg production and

hatchability in *Salmonella*-infected layers.

Salmonella infection in chickens occurs at all ages and is characterized by severe hepatomegaly and splenomegaly accompanied by anemia, septicemia, and a liver with a bronzing aspect [29]. Gross changes were observed in the organs of the control group, mainly in the liver, which was green-yellowish in color, larger, and more friable than that in normal layers. There was also spleen congestion and splenomegaly observed in the control group. In the CaFe-GMP additive group, all live chickens exhibited normal organ appearance.

We also examined whether *Salmonella* Gallinarum could be re-isolated, post-challenge, from organs of the experimental chickens. In the control group, bacteria were re-isolated at 5 dpi from the liver, spleen, and feces from the cecum, which indicated the occurrence of an infection. The numbers of *Salmonella* Gallinarum in all tested tissues of the CaFe-GMP additive group were significantly lower than those in the control group. Moreover, mortality in the CaFe-GMP additive group was delayed compared to that in the control group. Several reports have demonstrated that dietary iron intake lessens intestinal inflammatory responses and the pathology of enteric infection by foodborne bacterial pathogens, such as the *Salmonella enterica* serovar Typhimurium in mice [18,34]. In addition, iron deficiency has been associated with an impaired immune response, but it can also substantially increase resistance against intracellular pathogens, probably due to an increase in nutritional immunity [35]. Because dietary iron status can affect the immune response, it is likely that the array of antimicrobial defenses secreted from the intestinal mucosa is also affected. Additionally, in the present study, necropsy revealed that the CaFe-GMP additive group had milder and less severe abnormal gross lesions and pathological changes than those observed in the control group, indicating that the CaFe-GMP feed supplement produced bacterial clearance as well as resistance against *Salmonella* Gallinarum.

Our findings suggest that adding 0.16% CaFe-GMP to the diet of layers can improve clinical symptoms of *Salmonella* Gallinarum, reduce diarrhea, and result in mild and muted abnormal gross pathological changes in liver and spleen. In addition, CaFe-GMP feed supplementation enabled bacterial clearance from internal organs and maintained egg laying production in layers experimentally infected with *Salmonella* Gallinarum. Consequently, a CaFe-GMP feed supplement may be useful in preventing salmonellosis in the poultry industry.

Acknowledgments

This research was supported by the High Value-added Food Technology Development Program (113024-3), Ministry of Agriculture, Food and Rural Affairs, Republic of Korea.

Conflict of Interest

The authors declare no conflict of interests.

References

1. **Barrow PA.** Salmonella infections: immune and non-immune protection with vaccines. *Avian Pathol* 2007, **36**, 1-13.
2. **Barrow PA, Freitas Neto OC.** Pullorum disease and fowl typhoid—new thoughts on old diseases: a review. *Avian Pathol* 2011, **40**, 1-13.
3. **Basnet HB, Kwon HJ, Cho SH, Kim SJ, Yoo HS, Park YH, Yoon SI, Shin NS, Youn HJ.** Reproduction of fowl typhoid by respiratory challenge with *Salmonella* Gallinarum. *Avian Dis* 2008, **52**, 156-159.
4. **Boruta A, Swierczewska E, Glebocka K, Nollet L.** Trace organic minerals as a replacement of inorganic sources for layers: effects on productivity and mineral excretion. In: *World Poultry Science Association, Proceedings of the 16th European Symposium on Poultry Nutrition*; 26-30 August 2007, Strasbourg, France.
5. **Bovee-Oudenhoven IM, Lettink-Wissink MLG, Van Doesburg W, Witteman BJM, Van Der Meer R.** Diarrhea caused by enterotoxigenic *Escherichia coli* infection of humans is inhibited by dietary calcium. *Gastroenterology* 2003, **125**, 469-476.
6. **Bovee-Oudenhoven IM, Termont DS, Weerkamp AH, Faassen-Peters MA, Van der Meer R.** Dietary calcium inhibits the intestinal colonization and translocation of *Salmonella* in rats. *Gastroenterology* 1997, **113**, 550-557.
7. **Chappell L, Kaiser P, Barrow P, Jones MA, Johnston C, Wigley P.** The immunobiology of avian systemic salmonellosis. *Vet Immunol Immunopathol* 2009, **128**, 53-59.
8. **Cromwell GL.** Why and how antibiotics are used in swine production. *Anim Biotechnol* 2002, **13**, 7-27.
9. **de Oliveira GH, Berchieri A Jr, Fernandes AC.** Experimental infection of laying hens with *Salmonella enterica* serovar Gallinarum. *Braz J Microbiol* 2005, **36**, 51-56.
10. **de Paiva JB, Penha Filho RAC, Argüello YMS, da Silva MD, Gardin Y, Resende F, Berchieri A Jr, Sesti L.** Efficacy of several *Salmonella* vaccination programs against experimental challenge with *Salmonella* Gallinarum in commercial brown layer and broiler breeder hens. *Rev Bras Cienc Avic* 2009, **11**, 65-72.
11. **Faitarone ABG, Pavan AC, Mori C, Batista LS, Oliveira RP, Garcia EA, Pizzolante CC, Mendes AA, Sherer MR.** Economic traits and performance of Italian quails reared at different cage stocking densities. *Rev Bras Cienc Avic* 2005, **7**, 19-22.
12. **Gyles CL.** Antimicrobial resistance in selected bacteria from poultry. *Anim Health Res Rev* 2008, **9**, 149-158.
13. **Hong SS, Jeong J, Lee J, Kim S, Min W, Myung H.** Therapeutic effects of bacteriophages against *Salmonella* Gallinarum infection in chickens. *J Microbiol Biotechnol* 2013, **23**, 1478-1483.
14. **Janssens GPJ, Millet S, Van Immerseel F, De Buck J, Hesta M.** The impact of prebiotics and salmonellosis on apparent nutrient digestibility and *Salmonella* Typhimurium var.

- Copenhagen excretion in adult pigeons (*Columba livia domestica*). *Poult Sci* 2004, **83**, 1884-1890.
15. **Jung BG, Ko JH, Lee BJ.** Dietary supplementation with a probiotic fermented four-herb combination enhances immune activity in broiler chicks and increases survivability against *Salmonella* Gallinarum in experimentally infected broiler chicks. *J Vet Med Sci* 2010, **72**, 1565-1573.
 16. **Kamphues J.** [Antibiotic growth promoters for the view of animal nutrition]. *Berl Munch Tierarztl Wochenschr* 1999, **112**, 370-379. German.
 17. **Kim MS, Yoon YS, Seo JG, Lee HG, Chung MJ, Yum DY.** A study on the prevention of *salmonella* infection by using the aggregation characteristics of lactic Acid bacteria. *Toxicol Res* 2013, **29**, 129-135.
 18. **Kortman GAM, Mulder MLM, Richters TJW, Shanmugam NKN, Trebicka E, Boekhorst J, Timmerman HM, Roelofs R, Wiegierinck ET, Laarakkers CM, Swinkels DW, Bolhuis A, Cherayil BJ, Tjalsma H.** Low dietary iron intake restrains the intestinal inflammatory response and pathology of enteric infection by food-borne bacterial pathogens. *Eur J Immunol* 2015, **45**, 2553-2567.
 19. **Maciel MP, Saraiva EP, de Fátima Aguiar É, Ribeiro PAP, Passos DP, Silva JB.** Effect of using organic microminerals on performance and external quality of eggs of commercial laying hens at the end of laying. *Rev Bras Zootec* 2010, **39**, 344-348.
 20. **Mastroeni P, Chabalgoity JA, Dunstan SJ, Maskell DJ, Dougan G.** Salmonella: immune responses and vaccines. *Vet J* 2001, **161**, 132-164.
 21. **Murray MJ, Murray AB, Murray MB, Murray CJ.** The adverse effect of iron repletion on the course of certain infections. *Br Med J* 1978, **2**, 1113-1115.
 22. **National Research Council.** Nutrient Requirements of Poultry: Ninth Revised Edition, 1994. pp. 3-18, National Academies Press, Washington, DC, 1994.
 23. **Nyakeriga AM, Troye-Blomberg M, Dorfman JR, Alexander ND, Bäck R, Kortok M, Chemtai AK, Marsh K, Williams TN.** Iron deficiency and malaria among children living on the coast of Kenya. *J Infect Dis* 2004, **190**, 439-447.
 24. **Patterson JA, Burkholder KM.** Application of prebiotics and probiotics in poultry production. *Poult Sci* 2003, **82**, 627-631.
 25. **Rasschaert G, Michiels J, Tagliabue M, Missotten J, De Smet S, Heyndrickx M.** Effect of organic acids on *Salmonella* shedding and colonization in pigs on a farm with high *Salmonella* prevalence. *J Food Prot* 2016, **79**, 51-58.
 26. **Sazawal S, Black RE, Ramsan M, Chwaya HM, Stoltzfus RJ, Dutta A, Dhingra U, Kabole I, Deb S, Othman MK, Kabole FM.** Effects of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomised, placebo-controlled trial. *Lancet* 2006, **367**, 133-143.
 27. **Schaible UE, Kaufmann SHE.** Iron and microbial infection. *Nat Rev Microbiol* 2004, **2**, 946-953.
 28. **Seo YM, Shin KS, Rhee AR, Chi YS, Han J, Paik IK.** Effects of dietary Fe-soy proteinate and MgO on egg production and quality of eggshell in laying hens. *Asian-Australas J Anim Sci* 2010, **23**, 1043-1048.
 29. **Shivaprasad HL.** Fowl typhoid and pullorum disease. *Rev Sci Tech* 2000, **19**, 405-424.
 30. **Smith IM, Licence ST, Hill R.** Haematological, serological and pathological effects in chicks of one or more intravenous injections of *Salmonella* Gallinarum endotoxin. *Res Vet Sci* 1978, **24**, 154-160.
 31. **St Louis ME, Morse DL, Potter ME, DeMelfi TM, Guzewish JJ, Tauxe RV, Blake PA; the *Salmonella enteritidis* Working Group.** The emergence of grade A eggs as a major source of *Salmonella enteritidis* infections. New implications for the control of salmonellosis. *JAMA* 1988, **259**, 2103-2107.
 32. **Tompkins GR, O'Dell NL, Bryson IT, Pennington CB.** The effects of dietary ferric iron and iron deprivation on the bacterial composition of the mouse intestine. *Curr Microbiol* 2001, **43**, 38-42.
 33. **Vieira SL.** Chelated minerals for poultry. *Rev Bras Cienc Avic* 2008, **10**, 73-79.
 34. **Wang L, Johnson EE, Shi HN, Walker WA, Wessling-Resnick M, Cherayil BJ.** Attenuated inflammatory responses in hemochromatosis reveal a role for iron in the regulation of macrophage cytokine translation. *J Immunol* 2008, **181**, 2723-2731.
 35. **Weiss G.** Modification of iron regulation by the inflammatory response. *Best Pract Res Clin Haematol* 2005, **18**, 183-201.
 36. **Wigley P, Hulme S, Powers C, Beal R, Smith A, Barrow P.** Oral infection with the *Salmonella enterica* serovar Gallinarum 9R attenuated live vaccine as a model to characterise immunity to fowl typhoid in the chicken. *BMC Vet Res* 2005, **1**, 2.
 37. **Yenice E, Mizrak C, Gültekin M, Atik Z, Tunca M.** Effects of organic and inorganic forms of manganese, zinc, copper, and chromium on bioavailability of these minerals and calcium in late-phase laying hens. *Biol Trace Elem Res* 2015, **167**, 300-307.
 38. **Ziprin RL, Elissalde MH, Hinton A Jr, Beier RC, Spates GE, Corrier DE, Benoit TG, DeLoach JR.** Colonization control of lactose-fermenting *Salmonella* Typhimurium in young broiler chickens by use of dietary lactose. *Am J Vet Res* 1991, **52**, 833-837.