

Ischemia-Reperfusion Damage

İskemi Reperfüzyon Hasarı

Omer E. Yapca¹, Bunyamin Borekci², Halis Suleyman³

¹Department of Obstetrics and Gynecology, Sorgun State Hospital, Yozgat, Turkey

²Department of Obstetrics and Gynecology, Ataturk University Faculty of Medicine, Erzurum, Turkey

³Department of Pharmacology, Ataturk University Faculty of Medicine, Erzurum, Turkey

Abstract

Ischemia-reperfusion damage is a complex pathological process that begins with tissue anoxia and continues with the production of free oxygen radicals, expanding with the inflammatory response. The literature suggests the importance of antioxidant and anti-inflammatory treatment to treat ischemia-reperfusion-related tissue damage.

Key Words: Ischemia, reperfusion damage

Özet

İskemi-reperfüzyon hasarı, dokunun oksijensiz kalması ile başlayan, serbest oksijen radikallerinin üretimi ile devam eden ve inflamatuvar yanıtla genişleyen karmaşık patolojik bir süreçtir. Bu literatür bilgileri, iskemi-reperfüzyon doku hasarında antioksidan ve antiinflammatuar tedavinin önemini göstermektedir.

Anahtar Kelimeler: İskemi, reperfüzyon hasarı

Ischemic Damage

Ischemia is defined as an asphyxiated tissue state as a result of reduced or absent blood flow. In tissue ischemia, a series of chemical events that may extend to cell function disorders and cellular necrosis occurs. Healthy cellular functions are maintained through aerobic metabolism, whereas anaerobic metabolism is observed in ischemic cells [1]. Adenosine triphosphate (ATP) produced in healthy (aerobic) tissues is lysed into adenosine monophosphate (AMP), adenosine, inosine and hypoxanthine. The resulting hypoxanthine is metabolized into xanthine and uric acid. As nicotinamide adenine dinucleotide (NAD) is used in the metabolism of hypoxanthine by KDH in aerobic tissues, toxic oxygen radicals are not formed [2]. The metabolism of hypoxanthine in ischemic tissues is different from that in aerobic metabolism. Hypoxanthine is metabolized by xanthine oxidase (XO) in ischemic tissues, but not by KDH because KDH is transformed into XO during ischemia [3]. As molecular oxygen (O₂) is used in the metabolism of hypoxanthine by XO, toxic oxygen radicals are produced as intermediate products. However, because O₂ is not found in sufficient amounts in ischemic tissues, the accumulated hypoxanthine cannot be transformed into xanthine until reperfusion occurs, and toxic oxygen radicals cannot be produced as intermediate products [4]. However, cellular energy reservoirs are exhausted during long-term ischemia [5]. This exhaustion of energy reservoirs

results in the inhibition of low Na⁺/K⁺-ATP pumps found in the cell membrane. Low ATP pump activity prevents the transport of Na⁺ and Ca⁺² ions to the outside of the cell, and the intracellular concentrations of Na⁺ and Ca⁺² ions increase as a result [6]. The intracellular increase in Na⁺ increases water uptake by the cell by passive diffusion, causing cell swelling. This swelling decreases with the accumulation of anaerobic metabolites [7]. The intracellular increase in the Ca⁺² ion concentration leads to pathological events within the cells [8]. Therefore, the first intervention for ischemic tissue should be to induce blood flow (reperfusion) within the tissue.

Reperfusion Damage

Formed during ischemia with reoxygenation during reperfusion, XO leads to the formation of excessive amounts of free oxygen radicals while transforming the accumulated hypoxanthine into xanthine using O₂ [3, 9, 10]. These free oxygen radicals, known as reperfusion mediators, lead to the formation of toxic products such as aldehyde and malondialdehyde lipids by oxidizing cell membrane lipids [11]. Moreover, oxygen radicals induce DNA oxidative damage [12]. Following free radical reactions, alkaline changes in nucleic acids and chain breaks in DNA occur. If the acquired DNA alterations cannot be repaired, the DNA mutates. It is accepted that 8-hydroxyguanine (8-OHGua) is a mutagenic type of DNA [12, 13]. Parks DA and colleagues have shown

Received: March 19, 2013 / **Accepted:** May 13, 2013

Correspondence to: Halis Suleyman, Department of Pharmacology, Ataturk University Faculty of Medicine, 25240 Erzurum, Turkey
Phone: +90 442 231 15 47 Fax: +90 442 236 09 68 e-mail: halis.suleyman@gmail.com

doi:10.5152/eajm.2013.24



that reperfusion damage is more hazardous than ischemic damage alone [14]. Polymorphonuclear leukocytes (PMNLs) play an important role in the pathophysiology of reperfusion damage; PMNLs contain NADPH oxidase, which has oxidant effects, elastase, and myeloperoxidase in their azurophilic granules [15]. Respiratory burst occurs as a result of free oxygen radical release due to the increase of xanthine oxidase in active PMNLs. In the beginning of reperfusion following ischemia, approximately 70% of the oxygen provided to the tissue is oxidized to superoxide by xanthine oxidase. Superoxide is typically transformed into hydrogen peroxide by spontaneous dismutation. Hydrogen peroxide is reduced to hypochloric acid and chloride ions by myeloperoxidase. Hypochloric acid is a strong oxidant, and it can easily react with many biological molecules. Proteolytic enzymes, such as apolactoferrin, which is released from granules upon the activation of PMNLs, plasminogen activator, which activates complement, elastase, collagenase and gelatinase, damage the vessel endothelium [16]. Ischemia-reperfusion damage is also caused by the activation of phospholipase A2 with the increase of intracellular calcium during ischemia. Phospholipase A2 increases arachidonic acid, which is a membrane phospholipid. During tissue ischemia-reperfusion, cyclooxygenase-2 (COX-2) [15] is activated, leading to the formation of pro-inflammatory prostaglandins and free oxygen radicals from arachidonic acid [17-19]. Recent studies have demonstrated that histopathologic indications of inflammation and high COX-2 activity are observed in ovarian tissue following ischemia-reperfusion [20]. Reperfusion damage is known to be more severe than the damage caused by ischemia alone. Ischemia-reperfusion damage can be observed in almost all types of tissues. An increase in oxidant parameters and a decrease in antioxidant parameters have been reported following ischemia-reperfusion damage to the brain, myocardium, lung, kidney, ovary and stomach [21]. As a result, ischemia-reperfusion damage is a complicated pathological process beginning with tissue asphyxiation, continuing with the production of free oxygen radicals and expanding with the inflammatory response. The information reported in the literature underlines the importance of antioxidant and anti-inflammatory treatment to treat tissue with ischemia-reperfusion damage.

Conflict of interest statement: The authors declare that they have no conflict of interest to the publication of this article.

References

1. Semenza GL. Cellular and molecular dissection of reperfusion injury ROS within and without. *Circ Res* 2000; 86: 117-8. [\[CrossRef\]](#)
2. Grace PA. Ischaemia-reperfusion injury. *Br J Surg* 1994; 81: 637-47. [\[CrossRef\]](#)
3. Li C, Jackson RM. Reactive species mechanisms of cellular hypoxia-reoxygenation injury. *Am J Physiol Cell Physiol* 2002; 282: C227-41. [\[CrossRef\]](#)
4. Parks DA, Williams TK, Beckman JS. Conversion of xanthine dehydrogenase to oxidase in ischemic rat intestine: a reevaluation. *Am J Physiol* 1988; 254(5 Pt 1): G768-74.
5. Zimmerman BJ, Granger DN. Reperfusion injury. *Surg Clin North Am* 1992; 72: 65-83.
6. Green C, Gower J, Healing G, Cotterill L, Fuller B, Simpkin S. The importance of iron, calcium and free radicals in reperfusion injury: an overview of studies in ischaemic rabbit kidneys. *Free Radical Research* 1989; 7: 255-64. [\[CrossRef\]](#)
7. Siesjö B. Mechanisms of ischemic brain damage. *Critical care medicine* 1988; 16: 954-63. [\[CrossRef\]](#)
8. Orrenius S, Burkitt MJ, Kass GEN, Dypbukt JM, Nicotera P. Calcium ions and oxidative cell injury. *Ann Neurol* 1992; 32 Suppl: S33-42. [\[CrossRef\]](#)
9. Lindsay TF, Liauw S, Romaschin AD, Walker PM. The effect of ischemia/reperfusion on adenine nucleotide metabolism and xanthine oxidase production in skeletal muscle. *J Vasc Surg* 1990; 12: 8-15. [\[CrossRef\]](#)
10. Parks DA, Granger DN. Ischemia-induced vascular changes: role of xanthine oxidase and hydroxyl radicals. *Am J Physiol* 1983; 245: G285-9.
11. Del Maestro RF. An approach to free radicals in medicine and biology. *Acta Physiol Scand Suppl* 1980; 492: 153-68.
12. Marnett LJ. Oxyradicals and DNA damage. *Carcinogenesis* 2000; 21: 361-70. [\[CrossRef\]](#)
13. Huang HY, Helzlsouer KJ, Appel LJ. The effects of vitamin C and vitamin E on oxidative DNA damage: results from a randomized controlled trial. *Cancer Epidemiol Biomarkers Prev* 2000; 9: 647-52.
14. Parks DA, Granger DN. Contributions of ischemia and reperfusion to mucosal lesion formation. *Am J Physiol* 1986; 250 (6 Pt 1): G749-53.
15. Carden DL, Granger, DN. Pathophysiology of Ischemia-reperfusion injury. *J Pathol* 2000; 190: 255-66. [\[CrossRef\]](#)
16. Korthuis RJ, Granger DN. Reactive oxygen metabolites, neutrophils, and the pathogenesis of ischemic-tissue/reperfusion. *Clinical cardiology* 1993; 16: 19-26. [\[CrossRef\]](#)
17. Udassin R, Vromen A, Haskel Y. The time sequence of injury and recovery following transient reversible intestinal ischemia. *J Surg Res* 1994; 56: 221-5. [\[CrossRef\]](#)
18. Vilanova JM, Figueras-Aloy J, Roselló J, Gómez G, Gelpí E, Jiménez R. Arachidonic acid metabolites in CSF in hypoxic-ischaemic encephalopathy of newborn infants. *Acta Paediatrica* 1998; 87: 588-92. [\[CrossRef\]](#)
19. Fern R, Möller T. Rapid ischemic cell death in immature oligodendrocytes: a fatal glutamate release feedback loop. *J Neurosci* 2000; 20: 34-42.
20. Isaoglu U, Yilmaz M, Sener E, et al. The Impaired Balances of Oxidant/Antioxidant and COX-1/COX-2 in Ovarian Ischemia-Reperfusion Injury and Prevention by Nimesulide. *Lat Am J Pharm* 2012; 31: 1481-8.
21. Kisaoglu A, Borekci B, Yapca OE, Bilen H, Suleyman H. Tissue damage and oxidant/antioxidant balance. *Eurasian J Med* 2013; 45: 47-9. [\[CrossRef\]](#)