

## Vagus nerve stimulation: Invasive or noninvasive?

Autonomic imbalance with heightened sympathetic activity and withdrawal of vagal activity is critical for development and progression of chronic heart failure (1, 2). Primary characteristics of autonomic function comprise heart rate and heart rate variability. Abnormalities in both of these domains are associated with higher morbidity both in general population and in cardiovascular patients (3, 4). Both resting heart rate and a number of indices derived from heart rate variability are dominated by vagal vs. sympathetic tone, as the effects of muscarinic blockade are more significant than  $\beta$ -adrenergic blockade (5).

In this issue, Akdemir et al. (6) reviewed reports on the development of clinical vagus nerve stimulation (VNS) for cardiac applications. The article entitled "Vagus nerve stimulation: An evolving adjunctive treatment for cardiac disease" focuses on VNS for cardiac arrhythmias and heart failure, and discusses VNS device types and potential adverse effects. Although the review is highly relevant and sought after for discussion of invasive VNS techniques in cardiac diseases, it has to be emphasized that emerging noninvasive transcutaneous VNS (tVNS) approaches are not limited to treatments of migraine and headache.

Topical applications of tVNS are promising to exert specific and sometimes unique effects of tVNS on the cardiovascular system similar to those achieved through acupoints, which are reactive points on the surface of the body (7). For example, an increase of vagus nerve activity most likely is specifically involved in an increase of superior mesenteric artery blood flow volume induced by of manual acupuncture stimulation of ST36 (Zusanli) (8). Recent review by Murray et al. (9) entitled "The strange case of the ear and the heart: the auricular vagus nerve and its influence on cardiac control" provides fascinating discussion of the available evidence in support of modulating cardiac activity using the auricular nerve where tVNS can be delivered through electrical stimulation to the auricular branch of the vagus nerve (9).

Functional magnetic resonance imaging study demonstrates that the central projections of the auricular branch of the vagus nerve are consistent with the "classical" central vagal projections and can be accessed non-invasively via the external ear (10). Several human studies demonstrate beneficial cardiac effects following tVNS in cardiovascular patients with paroxysmal atrial fibrillation, angina pectoris, and severe heart failure (11–14). tVNS administration is associated with a significant modulation in systemic levels of tumor necrosis factor alpha, C-reactive protein (13), and heat shock proteins (14).

Interestingly, auricular stimulation may also influence the activity of the sympathetic nervous system in healthy individuals with no history of cardiovascular disease. Indeed, tVNS on the inner and outer surface of the tragus significantly improves heart rate variability (through a shift in cardiac autonomic activity toward relative parasympathetic/vagal dominance) and causes a significant decrease in muscle sympathetic nerve activity as recorded by microneurography in healthy human volunteers (15).

Considering potential specificity and better safety profile, tVNS appears to be a promising therapeutic alternative to invasive VNS. Further studies of mechanistic basis of tVNS and its clinical efficacy are required.

Yana Anfinogenova<sup>1,2,3</sup>

<sup>1</sup>Cardiology Research Institute, Tomsk National Research Medical Center; Tomsk-Russia

<sup>2</sup>National Research Tomsk Polytechnic University; Tomsk-Russia

<sup>3</sup>Siberian State Medical University; Tomsk-Russia

### References

1. Kishi T. Heart failure as an autonomic nervous system dysfunction. *J Cardiol* 2012; 59: 117-22.
2. Lachowska K, Gruchała M, Narkiewicz K, Hering D. Sympathetic activation in chronic heart failure: Potential benefits of interventional therapies. *Curr Hypertens Rep* 2016; 18: 51.
3. Wulsin LR, Horn PS, Perry JL, Massaro JM, D'Agostino RB. Autonomic imbalance as a predictor of metabolic risks, cardiovascular disease, diabetes, and mortality. *J Clin Endocrinol Metab* 2015; 100: 2443-8.
4. Zhang D, Shen X, Qi X. Resting heart rate and all-cause and cardiovascular mortality in the general population: a meta-analysis. *CMAJ* 2016; 188: E53-63.
5. Joyner MJ. Preclinical and clinical evaluation of autonomic function in humans. *J Physiol* 2016; 594: 4009-13.
6. Akdemir B, Benditt DG. Vagus nerve stimulation: An evolving adjunctive treatment for cardiac disease. *Anatol J Cardiol* 2016; 16: 804-10.
7. Giovanni M. *The Foundations of Chinese Medicine: A comprehensive text for acupuncturists and herbalists*, Churchill Livingstone, Edinburgh, UK, 1989.
8. Kaneko S, Watanabe M, Takayama S, Numata T, Seki T, Tanaka J, et al. Heart rate variability and hemodynamic change in the superior mesenteric artery by acupuncture stimulation of lower limb points: a randomized crossover trial. *Evid Based Complement Alternat Med* 2013; 2013: 315982.
9. Murray AR, Atkinson L, Mahadi MK, Deuchars SA, Deuchars J. The strange case of the ear and the heart: The auricular vagus nerve


**Address for correspondence:** Yana Anfinogenova, MD, PhD. Cardiology Research Institute, Tomsk NRMIC.  
111a Kievskaya Street, Tomsk 634012-Russia  
Phone: +7 (3822) 554-111. Fax: +7 3822 562 164 E-mail: anfiyj@gmail.com, anfy@tpu.ru

**Accepted Date:** 15.06.2016

©Copyright 2016 by Turkish Society of Cardiology - Available online at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)  
DOI:10.14744/AnatolJCardiol.2016.23113



- and its influence on cardiac control. *Auton Neurosci* 2016 Jun 28. Epub ahead of print.
10. Frangos E, Ellrich J, Komisaruk BR. Non-invasive access to the vagus nerve central projections via electrical stimulation of the external ear: fMRI Evidence in Humans. *Brain Stimul* 2015; 8: 624-36.
  11. Zamotrinsky AV, Kondratiev B, de Jong JW. Vagal neurostimulation in patients with coronary artery disease. *Auton Neurosci* 2001; 88: 109-16.
  12. Chen M, Yu L, Zhou X, Liu Q, Jiang H, Zhou S. Low-level vagus nerve stimulation: an important therapeutic option for atrial fibrillation treatment via modulating cardiac autonomic tone. *Int J Cardiol* 2015; 199: 437-8.
  13. Stavrakis S, Humphrey MB, Scherlag BJ, Hu Y, Jackman WM, Nakagawa H, et al. Low-level transcutaneous electrical vagus nerve stimulation suppresses atrial fibrillation. *J Am Coll Cardiol* 2015; 65: 867-75.
  14. Afanasiev SA, Pavliukova EN, Kuzmichkina MA, Rebrova TY, Anfinogenova Y, Likhomanov KS, et al. Nonpharmacological correction of hypersympatheticotonia in patients with chronic coronary insufficiency and severe left ventricular dysfunction. *Ann Noninvasive Electrocardiol* 2016 Mar 7. Epub ahead of print.
  15. Clancy JA, Mary DA, Witte KK, Greenwood JP, Deuchars SA, Deuchars J. Non-invasive vagus nerve stimulation in healthy humans reduces sympathetic nerve activity. *Brain Stimul* 2014; 7: 871-7.



1950-?, Zati Altay, MD, Germany