

19. Jaffe ES. The 2008 WHO classification of lymphomas: implications for clinical practice and translational research. *Hematology Am Soc Hematol Educ Program* 2009; 523–531.
20. Campo E, Swerdlow SH, Harris NL et al. The 2008 WHO classification of lymphoid neoplasms and beyond: evolving concepts and practical applications. *Blood* 2011; 117: 5019–5032.
21. Townsend W, Linch D. Hodgkin's lymphoma in adults. *Lancet* 2012; 380: 836–847.
22. Hawkes EA, Wotherspoon A, Cunningham D. The unique entity of nodular lymphocyte-predominant Hodgkin lymphoma: current approaches to diagnosis and management. *Leuk Lymphoma* 2012; 53: 354–361.
23. Amur S, Parekh A, Mummaneni P. Sex differences and genomics in autoimmune diseases. *J Autoimmun* 2011; 38: J254–J265.
24. Quintero OL, Amador-Pataroyo MJ, Montoya-Ortiz G et al. Autoimmune disease and gender: plausible mechanisms for the female predominance of autoimmunity. *J Autoimmun* 2012; 38: J109–J119.
25. Hemminki K, Li X, Sundquist K et al. Familial association of inflammatory bowel diseases with other autoimmune and related diseases. *Am J Gastroenterol* 2010; 105: 139–147.
26. Mercer LK, Davies R, Galloway JB et al. Risk of cancer in patients receiving non-biologic disease-modifying therapy for rheumatoid arthritis compared with the UK general population. *Rheumatology (Oxford)* 2013; 52: 91–98.
27. Ferlay J, Shin HR, Bray F et al. GLOBOCAN 2008 v2.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 [Internet]. Lyon, France: International Agency for Research on Cancer, 2010.
28. Kuppers R, Rajewsky K, Zhao M et al. Hodgkin disease: Hodgkin and Reed–Sternberg cells picked from histological sections show clonal immunoglobulin gene rearrangements and appear to be derived from B cells at various stages of development. *Proc Natl Acad Sci USA* 1994; 91: 10962–10966.
29. LeBien TW, Tedder TF. B lymphocytes: how they develop and function. *Blood* 2008; 112: 1570–1580.
30. Miller EB. Hodgkin lymphoma and autoimmunity: a two-way street. *Clin Adv Hematol Oncol* 2012; 10: 276–277.
31. Lossius A, Johansen JN, Torkildsen O et al. Epstein–Barr virus in systemic lupus erythematosus, rheumatoid arthritis and multiple sclerosis-association and causation. *Viruses* 2013; 4: 3701–3730.
32. Hjalgrim H. On the aetiology of Hodgkin lymphoma. *Dan Med J* 2012; 59: B4485.
33. Thorsby E, Lie BA. HLA associated genetic predisposition to autoimmune diseases: genes involved and possible mechanisms. *Transpl Immunol* 2005; 14: 175–182.

Annals of Oncology 25: 1404–1410, 2014
doi:10.1093/annonc/mdu166
Published online 5 May 2014

Antiviral treatment in patients with indolent B-cell lymphomas associated with HCV infection: a study of the Fondazione Italiana Linfomi

L. Arcaini^{1,2}, D. Vallisa³, S. Rattotti², V. V. Ferretti², A. J. M. Ferreri⁴, P. Bernuzzi³, M. Merli⁵, M. Varettoni², A. Chiappella⁶, A. Ambrosetti⁷, A. Tucci⁸, C. Rusconi⁹, C. Visco¹⁰, M. Spina¹¹, G. Cabras¹², S. Luminari¹³, M. Tucci¹⁴, P. Musto¹⁵, M. Ladetto¹⁶, F. Merli¹⁷, C. Stelitano¹⁸, A. d'Arco¹⁹, L. Rigacci²⁰, A. Levis²¹, D. Rossi²², P. Spedini²³, S. Mancuso²⁴, D. Marino²⁵, R. Bruno^{26,27}, L. Baldini²⁸ & A. Pulsoni²⁹

¹Department of Molecular Medicine, University of Pavia, Pavia; ²Department of Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia; ³Division of Hematology, Ospedale Giovanni da Saliceto, Piacenza; ⁴Unit of Lymphoid Malignancies, San Raffaele Scientific Institute, Milano; ⁵Division of Hematology, Department of Internal Medicine, Ospedale di Circolo, Fondazione Macchi, Varese; ⁶Division of Hematology 2, AO S Giovanni Battista, Torino; ⁷Department of Medicine, Section of Hematology, University of Verona, Verona; ⁸Division of Hematology, Spedali Civili, Brescia; ⁹Division of Hematology, Ospedale Niguarda Ca' Granda, Milano; ¹⁰Division of Hematology, Ospedale San Bortolo, Vicenza; ¹¹Division of Medical Oncology, National Cancer Center, Aviano; ¹²Division of Hematology, Osp A Businco, Cagliari; ¹³Department of Oncology and Hematology, University of Modena and Reggio Emilia, Modena; ¹⁴IMO, Department of Internal Medicine and Clinical Oncology, University of Bari, Aldo Moro, Bari; ¹⁵Scientific Direction, IRCCS, Centro di Riferimento Oncologico della Basilicata, Rionero in Vulture; ¹⁶Division of Hematology, Department of Experimental Medicine and Oncology, University of Torino, Torino; ¹⁷Hematology Unit, Department of Oncology, Arcispedale Santa Maria Nuova, Istituto di Ricovero e Cura a Carattere Scientifico, Reggio Emilia; ¹⁸Division of Hematology, Dipartimento di Ematologia, Ospedale M. Morelli, Reggio Calabria; ¹⁹Division of Oncohematology, Ospedale Umberto I, Nocera Inferiore; ²⁰Division of Hematology, Azienda Ospedaliero Universitaria Careggi, Firenze; ²¹Division of Hematology, Azienda Ospedaliera SS Arrigo e Biagio e Cesare Arrigo, Alessandria; ²²Division of Hematology, Department of Translational Medicine, Amedeo Avogadro University of Eastern Piedmont, Novara; ²³Division of Haematology, Hospital of Cremona, Cremona; ²⁴Division of Hematology, Policlinico, Palermo; ²⁵Department of Oncology, Istituto Oncologico Veneto, IRCCS, Padova; ²⁶Division of Infectious and Tropical Diseases, Fondazione IRCCS Policlinico San Matteo, Pavia; ²⁷Department of Clinical Surgical, Diagnostic and Pediatric Sciences, University of Pavia, Pavia; ²⁸Division of Hematology, Fondazione IRCCS Cà Granda OM Policlinico, Università degli Studi, Milano; ²⁹Division of Hematology, Sapienza University, Rome, Italy

Received 7 October 2013; revised 23 March 2014 and 7 April 2014; accepted 11 April 2014

Background: Tumor regression after antiviral therapy (AT) is in favor of an etiological role of hepatitis C virus (HCV) in non-Hodgkin's B-cell lymphomas (NHL).

*Correspondence to: Dr Luca Arcaini, Department of Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Viale Golgi 19, Pavia, Italy. Tel: +39-0382-501308; E-mail: luca.arcaini@unipv.it

Patients and methods: We carried out a cohort study of 704 consecutive HIV-negative, HCV-positive patients with indolent NHL diagnosed and treated from 1993 to 2009 in 39 centers of the Fondazione Italiana Linfomi; 134 patients were managed with AT for lymphoma control.

Results: For entire cohort, 5-year overall survival (OS) was 78% [95% confidence interval (CI): 74%–82%] and 5-year progression-free survival (PFS) was 48% (95% CI: 44%–53%). In multivariate analysis, the use of AT during the patients' life had positive impact on OS. Forty-four of the 100 patients treated with first-line AT achieved a complete remission (CR) and 33 a partial response (PR). HCV-RNA clearance was achieved in 80 patients and was related to lymphoma response. At a median follow-up of 3.6 years, 5-year PFS was 63% (95% CI: 50%–73%). CR + PR rate was 85% with AT as second-line treatment.

Conclusion: AT produces HCV-RNA clearance and consequent tumor regression in most patients with HCV-related indolent NHL. AT used at any time is associated with improved OS. Consequently, AT can be considered an option for patients with indolent lymphomas who do not need immediate cytoreductive treatment.

Key words: HCV, indolent lymphoma, antiviral treatment, outcome

introduction

In addition to liver involvement, hepatitis C virus (HCV) infection has been linked to lymphoproliferative disorders and the role of HCV infection in lymphomagenesis may be related to chronic antigenic stimulation of the virus [1].

Several epidemiological studies have been carried out since the mid-1990s to investigate the link between HCV and non-Hodgkin's lymphoma (NHL). Systematic concluded that HCV prevalence in patients with B-NHL is higher with respect to the general population [2]. In subtype-specific analyses, HCV prevalence was associated with marginal zone lymphoma (MZL) [odds ratio (OR) = 2.47], diffuse large B-cell lymphoma (DLBCL) (OR = 2.24), and lymphoplasmacytic lymphoma (LPL) (OR = 2.57) [3].

Interestingly, Kawamura et al. [4] demonstrated that sustained virologic response induced by interferon (IFN) therapy protects against the development of malignant lymphoma in HCV-infected patients.

The regression of HCV-associated lymphoma with antiviral treatment (AT) alone is the strongest argument in favor of an etiological link between lymphoma and infection [5, 6].

In relation to the relatively low number of patients treated with AT so far, the impact of this approach on the outcome of indolent lymphomas associated with HCV infection needs to be confirmed. To this task, a survey ('HCV-LNH outcome survey') was initiated in Italy under the sponsorship of the Fondazione Italiana Linfomi (FIL) in 2010 to better define the outcome and the efficacy of treatments of HCV-positive NHL. Moreover, we analyzed separately the 134 patients managed with AT as anti-lymphoma strategy.

methods

study design

In the nationwide database of FIL (before 2012 named Intergruppo Italiano Linfomi, IIL), we identified 704 HCV-positive (serology and/or HCV-RNA), HIV-negative patients affected by indolent NHL diagnosed between 1993 and 2009 in 39 centers members with available information about treatment. HCV infection is a substantial health problem in Italy and majority of patients with NHL are evaluated for HCV infection. HCV-positive cases were 15% of all registered cases with available HCV serology from participating centers; HCV serology was not available for 19.5% of all diagnosed cases

in the examined period. We included also cases of mantle-cell lymphoma with an indolent course.

Since 2000, FIL/IIL centers shared homogeneous procedures for lymphoma diagnosis, staging, response assessment, and follow-up. Before 2000, minimal requirements of homogeneity were adopted. A central pathology review was not formally carried out; however, participating centers are all characterized by high rate of lymphomas diagnosis and local pathologists systematically participate in the activities of updating the diagnostic procedures of the FIL. All lymphoma cases are consecutively included and followed by centers participating to the FIL.

The database of HCV-positive indolent NHL was analyzed to establish HCV infection features, distribution of lymphoma categories, clinical presentation, natural history, therapeutic management, outcome, and prognostic factors.

We also analyzed in detail virological and lymphoma response and outcome of 134 patients who received AT both as first-line ($n = 100$) and as second-line ($n = 34$) anti-lymphoma treatment: all these patients did not need immediate cytoreductive therapy at the start of AT.

Approval for this study, which was based on the use of archival data, was obtained from the Institutional Review Board. The report was prepared in accordance with the STROBE statement [7]. Data management and analysis were carried out in accordance with the ethical guidelines of the FIL and the tenets of the Declaration of Helsinki of 1964, as revised in 2000. All patients gave informed consent.

end points and variables

The primary outcome measure was overall survival (OS); secondary end points were progression-free survival (PFS), complete remission (CR), and partial remission (PR) rates. OS was taken from diagnosis until death from any cause; patients who had not died were censored at the date of their last follow-up visit. PFS was defined as the time from start of treatment until lymphoma progression or initiation of new treatment, or death [8]. Several variables were included in the analysis (Table 1; supplementary Table S1, available at *Annals of Oncology* online), including the use of AT at any time during a patient's life.

For patients analyzed for anti-lymphoma activity of AT, CR was defined by the complete disappearance of all detectable sites and symptoms; PR was defined as a more than 50% reduction. Responses different from CR/PR were defined as stable disease (SD); progressive disease was considered an increase in the size of more than 25% of previously documented disease or the appearance of new lesions. Specific examinations have been carried out for response assessment in peculiar clinical presentations (i.e. paraprotein level when present before AT, endoscopy for gastric MALT lymphoma). For these patients, OS and PFS were calculated from the start of therapy.

Table 1. Comparison of clinical features of patients who underwent antiviral therapy as anti-lymphoma treatment with respect to those who did not

	n	%	Antiviral therapy as anti-lymphoma treatment				Test and P-value
			Yes (134)		No (570)		
			n	%	n	%	
Male/female	289/415	41/59	57/77	43/57	232/338	41/59	$\chi^2 = 0.2, P = 0.698$
Age >60 years	483	69	70	52	413	73	$\chi^2 = 21.5, P < 0.001$
Histotypes							
Follicular lymphoma	134	19	12	9	122	21	$\chi^2 = 27.7, P < 0.001$
MALT MZL	157	22	29	22	128	22	
Primary nodal MZL	59	8	13	9	46	8	
Splenic MZL	137	20	36	27	101	18	
Small lymphocytic lymphoma	37	5	4	3	33	6	
Lymphoplasmacytic lymphoma	53	8	10	7	43	8	
Indolent mantle-cell lymphoma	30	4	1	1	29	5	
Indolent B-cell NHL NOS	97	14	29	22	68	12	
Ann Arbor stage III + IV	550	79	116	87	434	77	$\chi^2 = 6.3, P = 0.012$
Extranodal disease	271	42	80	60	191	38	$\chi^2 = 21.4, P < 0.001$
Liver involvement	53	8	16	12	37	7	$\chi^2 = 3.1, P = 0.080$
ECOG 2–3	65	10	6	4	59	11	$\chi^2 = 5.1, P = 0.025$
LDH > UNL	164	24	31	24	133	24	$\chi^2 = 0.0, P = 0.921$
Albumin < 3.5 mg/dl	100	16	16	13	84	17	$\chi^2 = 1.0, P = 0.308$

Counts (n) and percentage frequencies (%) with the Pearson's χ^2 test and the significance level (P-value) are reported.

MZL, marginal zone lymphoma; MALT, mucosa-associated lymphoid tissue; NOS, not otherwise specified; LDH, lactate dehydrogenase; UNL, upper normal limit.

statistical analysis

Association between categorical variables was tested by using the Pearson χ^2 test or the Fisher's exact test. Difference in the mean values of quantitative variables between two independent groups was tested by using the *t*-test for unpaired data. The Kaplan–Meier product-limit method was used to estimate OS and PFS.

The adjusted association of AT with OS and PFS was estimated by means of backward stepwise multivariate Cox's regression models, after checking for the applicability assumptions. The significance value for exclusion was set at 0.10, but both models were 'forced' to keep AT and diagnosis. AT was analyzed as a time-dependent covariate. Unadjusted and adjusted hazard ratios (HR) with 95% confidence interval (95% CI) are provided. The limit of significance for all analyses was defined as a P-value of ≤ 0.05 . All computations were carried out using Stata 12.1 (2007).

results

clinical and virological features

Clinical features of 704 patients with indolent lymphoma associated with HCV infection are summarized in Table 1. There was a prevalence of females and the median age was 66 years; MZL was the most common subtype. Virological features are summarized in supplementary Table S1, available at *Annals of Oncology* online.

outcome

Alkylators was the first-line treatment in 148 patients (21%), rituximab + alkylators in 49 (7%), CHOP in 106 (15%), R-CHOP in 77 (11%), other chemotherapy regimens in 56 (8%),

AT in 100 (14%), radiotherapy 21 (3%), surgery in 21 (3%), antibiotics in 13 (2%), and watch and wait in 113 (16%).

Three hundred and twenty-four patients experienced lymphoma relapse or progression at a median follow-up of 3.5 years (range 0.5–17). Five hundred and thirty-nine patients are alive (169 without evidence of disease), with a 5- and 10-year OS of 78% (95% CI: 74%–82%) and 62% (95% CI: 55%–67%), respectively (Figure 1); 5- and 10-year PFS of 48% (95% CI: 43%–53%) and 30% (95% CI: 25%–36%), respectively (Figure 2). Causes of death were: lymphoma 65%, cardiovascular events 12%, liver disease 9%, infection 6%, other neoplasia 6%, and other causes 2%.

univariate and multivariate analysis

In univariate analysis, AT showed a significant association both with OS and with PFS (supplementary Table S2, available at *Annals of Oncology* online). Furthermore, age >60 years, B symptoms, ECOG >1, albumin <3.5 g/dl, and increased LDH were also associated with OS and PFS, while Ann Arbor stage III–IV, absence of cryoglobulinemia and of symptomatic cryoglobulinemia, and the presence of cirrhosis were associated only with OS (supplementary Table S2, available at *Annals of Oncology* online).

Despite the multivariate analysis has shown a significant association of age >60 years, cirrhosis, cryoglobulinemia, and albumin <3.5 g/dl with OS, in multivariate analysis, AT retained a protective effect with OS (Table 2, Figure 3). On the contrary, the protective effect of AT on PFS was not confirmed after adjustment for confounders.




Figure 1. Overall survival (A) and progression-free survival (B) of entire series of 704 patients with hepatitis C virus-associated indolent lymphoma.




Figure 2. Overall survival according to the use of antiviral treatment at any time.

Table 2. Predictive model for OS and PFS derived from multivariate backward stepwise Cox analysis, considering antiviral treatment as a time-dependent covariate

	HR	95% CI	P-value
Overall survival			
Antiviral treatment	0.21	0.06–0.73	0.014
Age>60 years	3.89	1.71–8.85	0.001
B symptoms	1.99	0.88–4.51	0.099
Cirrhosis	2.81	1.24–6.35	0.013
Cryoglobulinemia	0.36	0.15–0.88	0.024
Albumin <3.5 g/dl	3.16	1.64–6.09	0.001
Progression-free survival			
Antiviral treatment	0.80	0.48–1.33	0.385
Age>60 years	1.43	0.95–2.17	0.089
ECOG>1	1.64	0.93–2.87	0.085
Albumin <3.5 g/dl	1.55	1.00–2.40	0.050

The mutually adjusted hazard ratios (HRs) with the 95% confidence interval (95% CI) and the significance level (P-value) are reported. HRs are also adjusted for diagnosis.

anti-lymphoma antiviral treatment

One hundred and thirty-four patients received AT for lymphoma control: 100 patients received AT as the first-line option, while 34 patients received AT as second-line treatment following a previous therapeutic failure. AT consisted of IFN in 47 patients (plus RBV in 36) and peg-IFN in 87 (plus RBV in 82). The median age was 60 years (range 23–80 years) with no difference between patients treated with AT as first or second line ($P = 0.07$). Histological, virological, and hematological features of the 134 patients are summarized in Table 3.

first-line antiviral treatment: tolerability and activity. Among 100 patients treated with AT as first line (33 with IFN and 67 with peg-IFN), 60 were affected by MZL. HCV genotype was 2 in 52 patients and 1 in 37. The median duration of first-line AT was 7 months (range 2–48). Eighty-seven patients completed the planned AT; 6 patients discontinued due to toxicity, while 7 patients interrupted AT early due to lymphoma progression and lack of virological response. HCV-RNA clearance was achieved in 80 patients (80%).

Forty-four (44%) of patients achieved CR and 33 (33%) PR, with an ORR of 77% (95% CI: 69%–85%); 14 (14%) had SD. The median response duration was 33 months.

Lymphoma response was related to achievement of HCV-RNA clearance ($P = 0.003$) (supplementary Table S3, available at *Annals of Oncology* online). Lymphoma response was not statistically different between patients with MZL and non-MZL (ORR: 82% versus 70%), whereas it was lower in splenic MZL with respect to other MZL cases (ORR: 65% versus 92%; $P = 0.02$). ORR was 83% in genotype two carriers and 70% in genotype one carriers ($P = 0.3$).

At a median follow-up of 3.6 years, 9 patients progressed and 13 experienced lymphoma relapse after initial response to AT, with a 5-year PFS of 63% (95% CI: 50%–73%) (Figure 3B). Five-year PFS for patients not treated with AT as first line was 45%




Figure 3. Overall survival (A) and progression-free survival (B) of 100 patients treated with AT as the first-line anti-lymphoma treatment.

(95% CI: 49%–50%) and was statistically shorter in comparison to treated patients (HR = 0.56, 95% CI: 0.39–0.80, $P = 0.001$).

Eighty-nine patients treated with first-line AT are alive, with a 5-year OS of 92% (95% CI: 83%–96%) (Figure 3A); only two patients died for lymphoma. The other causes of death were: hepatocellular carcinoma ($n = 3$), infections ($n = 2$), and cirrhosis, herpetic encephalitis, myelodysplastic syndrome, and suicide ($n = 1$ for each). Five-year OS for patients not treated with AT as first line was 75% (95% CI: 71%–80%) and was statistically shorter in comparison to treated patients (HR = 0.39, 95% CI: 0.20–0.73, $P = 0.004$). OS did not differ according to genotype, lymphoma category and serum levels of albumin, β_2 -microglobulin, and LDH. The impact of type of IFN (adjusted for levels of albumin, β_2 -microglobulin, and LDH) remained significant also in multivariate analysis (peg-IFN versus IFN, HR = 0.08, 95% CI: 0.01–0.81, $P = 0.03$).

second-line antiviral treatment. Nineteen (56%) of the 34 patients treated with second-line AT (14 with IFN and 20 with

Table 3. Histological, virological, and hematological features of 134 patients with hepatitis C virus-related indolent B-cell lymphoma treated with antiviral therapy as anti-lymphoma treatment

	First-line AT (100)		Second-line AT (34)	
	<i>n</i>	%	<i>n</i>	%
M/F	41/59	41/59	15/19	44/56
Marginal zone lymphoma	60	60	20	59
Splenic	23	23	12	35
Nodal	12	12	2	6
Extranodal of MALT	25	25	6	18
Follicular lymphoma	5	5	7	20
Lymphoplasmacytic lymphoma	7	7	2	6
Mantle-cell lymphoma	0	0	0	0
Small lymphocytic lymphoma	1	1	3	9
Indolent B-cell NHL NOS	27	27	2	6
Ann Arbor stage III–IV	90	90	28	82
Elevated β_2 -microglobulin	31	53	23	79
Albumin <3.5 g/dl	10	11	6	19
Serum MC	35	35	13	38
HCV genotype				
1	37	39	15	50
2	52	55	13	43
3	5	5	2	7
5	1	1	0	0
Cryoglobulinemia	34	34	10	29
Symptomatic cryoglobulinemia	19	56	7	70

Counts (*n*) and percentage frequencies (%) are reported.

peg-IFN) for relapse after a conventional first-line therapy achieved a CR and 10 (29%) a PR, with an ORR of 85% (95% CI: 73%–97%). HCV clearance was achieved in 22 patients (67%). The median response duration was 26 months. At a median follow-up of 4.1 years, four patients progressed and six experienced lymphoma relapse after initial response to AT, with a 5-year PFS of 63% (95% CI: 50%–73%).

discussion

To the best of our knowledge, this is the largest multicenter study focused on the anti-lymphoma efficacy of AT. Indeed, no reported study included more than 20 cases and related literature consists of around 100 patients treated with this strategy [9].

The present study shows that anti-lymphoma activity of AT is associated with viral load clearance and that AT is an active therapeutic option in these patients. The use of AT at any time during the life of these patients seems to be associated with improved outcome, which enforces the recommendation that AT must be considered as first-line approach for patients with indolent lymphomas who do not need immediate conventional treatment.

Major limitations of the present study are related to retrospective nature of the observation and the possible selection, loss-to-follow-up, referral bias of patients treated with AT; however, despite these aspects, reported cases are consecutively observed in centers and the decision of using AT is uniformly applied in all centers. Although PFS results might have been

biased by the retrospective nature of this study, the significant results observed for OS make clinically relevant the present findings. Considering the particular nature of this subset of lymphoma and the objective difficulty to perform prospective trials in significantly wide cohort of patients, our data may impact the standard of care.

It is possible that there are potential confounding factors that influence the difference in OS between the AT group and those who did not have AT, which we did not have data on (for the multivariate analysis in Table 2). However, the effect was so large (HR = 0.33), that it is unlikely other important confounders could explain this size of effect.

About 10 years ago, some reports attested the efficacy of AT in HCV-related SMZL [5, 10]. A previous study from our group extended this experience to a cohort of 13 patients with indolent lymphoma even different from SMZL [6]. Although that study included a relatively small number of patients, results showed no significant differences within different lymphomas categories and suggested that lymphoma response to AT was related to viral load decrease or disappearance.

ORR in the present group of patients who received upfront AT is similar to that reported in our first study (77% versus 74%) [11], with a slightly lower CR rate (44% versus 58%). Interestingly, this wide study is in line with previous studies [10–12], suggesting that HCV-RNA clearance is a *conditio sine qua non* to attain lymphoma response.

A relevant contribution of the present study regards the demonstration that tumor regression after AT can be obtained in diverse lymphoma categories. In fact, response rate was not different between MZL and non-MZL.

In the present study, few patients were lymphoma-unresponsive to AT, which could be related to HCV-independent phase of disease in which antigenic trigger is no longer necessary for lymphoproliferation, possibly because of additional genomic events. Although a relation between genotype and lymphoma response was not observed, genotype 1 carriers seemed to be less responsive than genotype 2 carriers. For the past decade, only 40%–45% of these patients achieved a sustained virological response when treated with peg-IFN/RBV [12]. Although the direct anti-lymphoma properties of IFN cannot be ruled out [11], it should be underlined the clear association between the lymphoma regression and the clearance of HCV. A possible evolution could be the addition of rituximab to the standard regimen as reported in HCV-associated cryoglobulinemia patients [13]. Since rituximab alone therapy has also been shown to induce lymphoma regression in patients with MZL, the safety and efficacy of combined AT and rituximab in patients with HCV-related MZL should be examined.

Finally, IFN-free regimens with direct antiviral agents only [14] could consent the access to AT also for HCV-positive NHL patients with contraindications to IFN use. In addition, considering the antiproliferative properties of IFN [11], lymphoma response could definitely demonstrated that lymphoma regression is strictly linked to the HCV eradication.

Overall, the present study could be considered as a good historical control group for future prospective trials.

Considering the entire series of 704 patients, AT had a positive prognostic influence on OS. Interestingly, a study has shown that HCV-infected patients who obtained a sustained virologic

response following AT had an HR of lymphoma risk significantly lower than patients who did not receive AT [4].

Thus, our data, together with Japanese findings [4], strongly support the efficacy of AT in indolent NHL associated with HCV infection. It remains to clarify the benefit of AT after (immuno)-chemotherapy if AT is not feasible (for instance, in the case of symptoms and/or high tumor burden).

In conclusion, the present study shows anti-lymphoma activity of AT in a large series of HCV-infected patients with indolent NHL. Consequently, this strategy could be considered the first-line option for patients with indolent lymphomas who do not need for immediate conventional treatment. The observation that AT used at any time during the patients' life is associated with improved OS enforces such a therapeutic opportunity and suggests a relevant advantage of AT in this setting beyond the objective detection of lymphoma regression.

acknowledgements

We gratefully acknowledge following investigators who contributed to the study through data collection: M. Gotti, L. Morello, M.L. Guerrero, V. Fiaccadori (Department of Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy); M. Goldaniga (Division of Hematology, Fondazione IRCCS Cà Granda OM Policlinico, Università degli studi, Milano, Italy); M. Bruno Ventre (Unit of Lymphoid Malignancies, San Raffaele Scientific Institute, Milano, Italy); A. Ferrario (Division of Hematology, Department of Internal Medicine, Ospedale di Circolo, Fondazione Macchi, Varese, Italy); S. Ferrero (Division of Hematology, Department of Experimental Medicine and Oncology, University of Torino, Torino, Italy); C. Ingenito (Oncohematology, Ospedale Umberto I, Nocera Inferiore, Italy) DM (Department of Oncology, Istituto Oncologico Veneto, IRCCS, Padova, Italy); A. Andriani (Department of Hematology, Nuovo Regina Margherita Hospital, Roma, Italy); A. Aresta, M. Ferrarini (Division of Medical Oncology C, Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy); G. Gini (Division of Hematology, Università Politecnica delle Marche, Ancona, Italy); C. Mazzaro (Department of Internal Medicine, Pordenone General Hospital, Pordenone, Italy); E. Pennese (Hematology and Stem Cell Transplantation Unit, 'Vito Fazzi' Hospital, Lecce, Italy); L. Maiocchi, P. Sacchi (Division of Infectious and Tropical Diseases, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy).

funding

This work was supported by Associazione Italiana Ricerca contro il Cancro (AIRC), My first AIRC grant 2011 #11415 (to LA)

disclosure

The authors have declared no conflicts of interest.

references

1. Peveling-Oberhag J, Crisman G, Schmidt A et al. Dysregulation of global microRNA expression in splenic marginal zone lymphoma and influence of chronic hepatitis C virus infection. *Leukemia* 2012; 26: 1654–1662.

2. Dal Maso L, Franceschi S. Hepatitis C virus and risk of lymphoma and other lymphoid neoplasms: a meta-analysis of epidemiologic studies. *Cancer Epidemiol Biomarkers Prev* 2006; 15: 2078–2085.
3. de Sanjose S, Benavente Y, Vajdic CM et al. Hepatitis C and non-Hodgkin lymphoma among 4784 cases and 6269 controls from the International Lymphoma Epidemiology Consortium. *Clin Gastroenterol Hepatol* 2008; 6: 451–458.
4. Kawamura Y, Ikeda K, Arase Y et al. Viral elimination reduces incidence of malignant lymphoma in patients with hepatitis C. *Am J Med* 2007; 120: 1034–1041.
5. Hermine O, Lefrere F, Bronowicki JP et al. Regression of splenic lymphoma with villous lymphocytes after treatment of hepatitis C virus infection. *N Engl J Med* 2002; 347: 89–94.
6. Vallisa D, Bernuzzi P, Arcaini L et al. Role of anti-hepatitis C virus (HCV) treatment in HCV-related, low-grade, B-cell, non-Hodgkin's lymphoma: a multicenter Italian experience. *J Clin Oncol* 2005; 23: 468–473.
7. von Elm E, Altman DG, Egger M et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008; 61: 344–349.
8. Cheson BD, Pfistner B, Juweid ME et al. Revised response criteria for malignant lymphoma. *J Clin Oncol* 2007; 25: 579–586.
9. Arcaini L, Bruno R. Hepatitis C virus infection and antiviral treatment in marginal zone lymphomas. *Curr Clin Pharmacol* 2010; 5: 74–81.
10. Saadoun D, Suarez F, Lefrere F et al. Splenic lymphoma with villous lymphocytes, associated with type II cryoglobulinemia and HCV infection: a new entity? *Blood* 2005; 105: 74–76.
11. Kimby E, Jurlander J, Geisler C et al. Long-term molecular remissions in patients with indolent lymphoma treated with rituximab as a single agent or in combination with interferon alpha-2a: a randomized phase II study from the Nordic Lymphoma Group. *Leuk Lymphoma* 2008; 49: 102–112.
12. Manns MP, McHutchison JG, Gordon SC et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001; 358: 958–965.
13. Saadoun D, Resche Rigon M, Sene D et al. Rituximab plus Peg-interferon-alpha/ribavirin compared with Peg-interferon-alpha/ribavirin in hepatitis C-related mixed cryoglobulinemia. *Blood* 2010; 116: 326–334; quiz 504–505.
14. Sulkowski MS, Gardiner DF, Rodriguez-Torres M et al. Daclatasvir plus sofosbuvir for previously treated or untreated chronic HCV infection. *N Engl J Med* 2014; 370: 211–221.

Annals of Oncology 25: 1410–1416, 2014
doi:10.1093/annonc/mdu167
Published online 5 May 2014

Prognostic significance of human papillomavirus in recurrent or metastatic head and neck cancer: an analysis of Eastern Cooperative Oncology Group trials

A. Argiris^{1*}, S. Li², M. Ghebremichael³, A. M. Egloff⁴, L. Wang⁵, A. A. Forastiere⁶, B. Burtness⁷ & R. Mehra⁷

¹Division of Hematology/Oncology Cancer Therapy and Research Center, Department of Medicine, University of Texas Health Science Center at San Antonio, San Antonio; ²Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute, Boston; ³Ragon Institute of Harvard, MIT and MGH and Harvard Medical School, Boston; Departments of ⁴Otolaryngology; ⁵Pathology, University of Pittsburgh, Pittsburgh; ⁶Department of Medical Oncology, Johns Hopkins University, Baltimore; ⁷Department of Medical Oncology, Fox Chase Cancer Center, Philadelphia, USA

Received 23 October 2013; revised 27 March 2014; accepted 16 April 2014

Background: The purpose of this article was to study the association of human papillomavirus (HPV) with clinical outcomes in patients with recurrent or metastatic squamous cell carcinoma of the head and neck (SCCHN).

Patients and methods: Archival baseline tumor specimens were obtained from patients treated on two clinical trials in recurrent or metastatic SCCHN: E1395, a phase III trial of cisplatin and paclitaxel versus cisplatin and 5-fluorouracil, and E3301, a phase II trial of irinotecan and docetaxel. HPV DNA was detected by *in situ* hybridization (ISH) with a wide-spectrum probe. p16 status was evaluated by immunohistochemistry. Clinical outcomes of interest were objective response, progression-free survival (PFS) and overall survival (OS).

Results: We analyzed 64 patients for HPV ISH and 65 for p16. Eleven tumors (17%) were HPV+, 12 (18%) were p16+, whereas 52 (80%) were both HPV– and p16–. The objective response rate was 55% for HPV-positive versus 19% for HPV-negative ($P = 0.022$), and 50% for p16-positive versus 19% for p16-negative ($P = 0.057$). The median survival was 12.9 versus 6.7 months for HPV-positive versus HPV-negative patients ($P = 0.014$), and 11.9 versus 6.7 months for p16-

*Correspondence to: Dr. Athanasios Argiris, Division of Hematology/Oncology, UT Health Science Center at San Antonio, Cancer Therapy & Research Center, 7979 Wurzbach Road MC8232, Zeller Building – 4th Floor, Room Z418, San Antonio, TX 78229, USA. Tel: +1-210-450-3838; Fax: +1-210-692-7502; E-mail: athanassios.argiris@gmail.com