

Physiological Research Pre-Press Article

Protective Effect of Ginsenoside against Acute Renal Failure via Reduction of Renal Oxidative Stress and Enhanced Expression of ChAT in the Proximal Convoluted Tubule and ERK 1/2 in the Paraventricular Nuclei

Zhou Jun^{1,2*}, Zhang Heng-ai^{1,3*}, Lin Yuan⁴, Liu Hongmei⁵, Cui Yumeng⁵, Xu Yin⁵, Zhao Nan⁵, Ma Jianmei⁶, Fan Kai⁶, Jiang Chunling¹

¹Department of Physiology, Dalian Medical University, Dalian, China, ² Department of

Dermatology, Dalian Economic & Technological Developmental Area Hospital, Dalian, China,

³Division of Cardiovascular Research, State Key Laboratory of Biomembrane and Membrane

Biotechnology, Institute of Zoology, Chinese Academy of Sciences, Beijing, China, ⁴Department

of Pharmacology, ⁵Department of Clinical Medicine, ⁶Department of Anatomy, Dalian Medical

University, Dalian, China

Summary

Generation of reactive oxygen species significantly contributes to the pathogenesis of acute renal failure (ARF) induced by myoglobin release. Ginsenosides (GS), the principal active ingredients of ginseng, is considered as an extremely good antioxidative composition of Chinese traditional and herbal drugs. The purpose of the present study was to investigate the protective effect of ginsenoside in rats with ARF on the changes of cholinergic nervous system in the kidney as well as on the involvement of mitogen-activated protein kinases (MAPK) in the hypothalamic paraventricular nuclei (PVN). In our assay, glycerol-induced acute renal failure in rats was employed to study the protective effects of ginsenoside. Our results indicated that the treatment of

ARF rats with ginsenosides for 48 h significantly reduced lipid peroxidation, restored the superoxide dismutase (SOD) level. Meanwhile, the obvious increase of choline acetyltransferase -immunoreactivity (ChAT-IR) in the proximal convoluted tubular cells (PCT) was observed by immunohistochemistry in ARF+GS group. The same effect was also observed in the changes of p-ERK1/2-IR in the hypothalamic paraventricular nuclei. Our results suggest that ginsenoside administered orally may have a strong renal protective effect against glycerol-induced ARF, reduce the renal oxidative stress, and ginsenoside can also activate the cholinergic system in PCT, simultaneously MAPK signal pathway in the PVN was also activated.

Key words: Ginsenoside, Acute renal failure, Proximal convoluted tubular cells, Choline acetyltransferase, Paraventricular nuclei

*The first two authors contributed equally to this work.

Corresponding author

Chunling Jiang, Department of Physiology, Dalian Medical University, Dalian, Liaoning, China.

E-mail: chunling_jiang2006@163.com

Introduction

Almost 10%~15% of all cases of acute renal failure (ARF) may account for traumatic and non-traumatic clinical conditions manifested with muscle fiber disintegration or rhabdomyolysis (Vlahović *et al.* 2007). There is increasing evidence indicating that oxidative stress plays an important role in the pathogenesis of rhabdomyolysis-induced myoglobinuric acute renal failure (Ustundag *et al.* 2009). ARF is usually reversible, the kidney being able to recover from almost

complete loss of function. Nevertheless, ARF is associated with major inpatient morbidity and mortality, reflecting the severity of the casual illnesses and the high frequency of complications (Yoon *et al.* 2008). Thus, early and effective renoprotective treatments are of utmost importance. In recent years, great effort has been focused on traditional and herbal medicine without toxic effects to provide a novel therapeutic agent for ARF.

Ginsenosides (GS), the unique constituents and secondary metabolites of the Panax species, have been known to be the pharmacologically active ingredients of ginseng and has a wide range of pharmacological and physiological actions due to its antioxidant and free radicals scavenging properties (Attele *et al.* 1999). It has been reported that Ginsenosides Rh4 significantly reduced the cisplatin-induced nephrotoxicity in cultured renal proximal tubular epithelial cells in a dose-dependent manner (Baek *et al.* 2006). Similarly, ginsenoside-Rd ameliorated the renal injury induced by cephaloridine through up-regulating the activities of the antioxidation enzymes superoxide dismutase and catalase (Yokozawa *et al.* 1999). However, the effect of GS administered orally on ARF has been seldom reported yet and little is known about its cellular and molecular mechanism.

The primary role of the kidney is to maintain the volume and electrolyte composition of body fluids and, in this way, to regulate the blood pressure. In proximal tubular cells, cholinergic agonists have been shown to rapidly increase basolateral Na-HCO₃ cotransport (NBC) activity. And the classic mitogen-activated protein kinase (MAPK) pathway activation is coupled to increase NBC activity (Robey *et al.* 2001). It suggested that renal cholinergic system participated in regulation of epithelial ion transport and there was an association between MAPK pathway and cholinergic system. Recent studies suggested that acetylcholine prevented the hypoxia-induced

apoptosis of mouse embryonic stem (ES) cells by inhibiting the reactive oxygen species (ROS) production. Meanwhile, acetylcholine also prevented angiotensin II-induced oxidative stress and apoptosis in H9c2 cells (Liu *et al.* 2011). These findings suggested that cholinergic system play a positive role in oxidative stress . However, whether renal cholinergic system activity is changed in ARF rats involved in oxidative stress is unknown.

The hypothalamic paraventricular nuclei (PVN) play a prominent role in the central regulation of water intake, urinary volume and sodium excretion, maintaining the homeostasis and regulate organismal function (de Arruda Camargo *et al.* 2010). MAPKs are a family of Ser/Thr protein kinases widely conserved among eukaryotes and are involved in many cellular programs such as cell proliferation, differentiation, survival movement, and cell death. The extracellular signal-regulated kinases (ERK) represent one of the four subfamilies of MAP kinases. It is reported that survival factors inhibit cell death by activating specific signaling pathways, including stimulation of protein-tyrosine phosphorylation and activation of ERK-1/2 , which then lead to the inhibition of the apoptotic signaling cascade (Baldanzi *et al.* 2002). This finding implied activation of MAPK pathways may participate in cytoprotection. Increasing evidences have shown that ACh plays a key role in the regulation of cell morphogenesis, proliferation and differentiation in many species; the growth-regulatory action of ACh is associated with signal transduction pathways involving MAPK signaling pathways (Ma *et al.* 2000). The paraventricular nuclei plays an important role in the regulation of fear, anxiety and stress (Herring *et al.* 2004). The neural cell adhesion molecule (NCAM) mimetic peptide C3d promoted ChAT activity in septal neurons through the stimulation of fibroblast growth factor receptor (FGFR) and MAPK cascade, in this way, played a protective role (Burgess *et al.* 2009). Therefore we hypothesized

that MAPK might be involved in the protective role of cholinergic system. Our previous studies showed MAPK signaling pathways was activated in locus ceruleus of ARF rats (Zhang *et al.* 2010). However, whether hypothalamic MAPK signal pathway was activated in glycerol-induced ARF rats remains unclear.

Based on these findings, it could be speculated that ginsenosides might have protective effects and there might be some interactions between cholinergic system in the kidney and MAPK signal pathway in the hypothalamic paraventricular nuclei in glycerol-induced ARF rats. Therefore, we examined 1) the changes of malondialdehyde (MDA) and superoxide dismutase (SOD) in renal cortex homogenate of ARF rats treated with ginsenoside for 48 h, 2) the changes of choline acetyltransferase -immunoreactivity (ChAT-IR) in the PCT of glycerol-induced ARF rats treated with ginsenoside for 48 h, 3) the changes of phospho-ERK1/2-immunoreactivity (p-ERK1/2-IR) in the hypothalamic paraventricular nuclei of glycerol-induced ARF rats treated with ginsenoside for 48 h.

Methods

Animals

Healthy male Sprague-Dawley rats (Dalian Medical University Animal Center, China) weighing 180~220 g were kept on a 12 h/12 h light/dark cycle and received food and water ad libitum at room temperature. During this time, rats were handled daily to avoid stress-induced expression on the day of the experiment. Special care was taken to minimize animal suffering and to reduce the number of animals used to the minimum necessary.

Animals' Treatment

Sixty four male rats were used in this study. After several days of adaptation, they were deprived of water for 16 h from 5 p.m. to 9 a.m.

Forty rats for the experiment *in vivo* were divided randomly into four groups(n=10 per group): ARF + physiological saline (NS) group, ARF + GS group, NS+NS group and NS+GS group. After deprived of water for 16 h, ARF+NS group and ARF+GS group were given intramuscularly with 10 ml/ kg body weight of 50% (vol/vol) glycerol solution distributed equally in both hind limbs. As soon as the model was established, ARF+GS group was given 25 mg GS in 2 ml NS using a stomach tube, at the same time, ARF+NS group was given 2 ml NS. The treatment of the other two groups were similar to them. They were treated with GS or NS for two consecutive days (once per 6 h, thrice per day).

Another 24 rats for immunohistochemistry were also divided into 4 groups and treated as described in the experiment *in vivo*.

Experiment in vivo

After treatment with ginsenoside or physiological saline for 48 h, kidney was removed, then renal cortex was homogenized with ice-cold physiological saline. MDA content, a measure of lipid peroxidation, was assayed in the form of thiobarbituric acid reacting substances (TBARS) using commercial reagents (Nanjing Jiancheng Bioengineering Institute, P.R. China). The SOD was measured by the method of xanthine oxidase using commercial reagents (Nanjing Jiancheng Bioengineering Institute, P.R. China). It was observed by measuring the absorbance at 550 nm.

Immunohistochemistry analysis

8 μ M thick renal sections were cut on Cryostat at -12 °C and placed onto low-iron clear glass slides. The sections above were incubated in 30 % H₂O₂ in methanol for 10 min and incubated with 0.4% pepsin for 1h at 37°C. The sections were washed three times in PBS 10 min and incubated in the primary antibody (ChAT-Ab, 1:200 Boster Company, P.R. China) 1 h at room temperature. The sections were then rinsed three times in PBS for 5 min and incubated in the biotinylated-second antibody (Boster Company, P.R. China) at room temperature for 1 h. Then the sections were again rinsed three times in PBS for 10 min and incubated with the avidin-biotin complex ABC (Boster Company, P.R. China) at room temperature for 1 h. Diaminobenzidine (DAB, Sigma Co, USA) and hematoxylin were used for signal detection. The control sections were incubated with PBS instead of primary antibody. Image Pro Plus image analysis system was used to analyze ChAT-IR positive granules. Kidney sections were observed in a microscope at 20 \times magnification. Optical density of ChAT-IR positive granules was calculated per area and per group.

50 μ M thick coronal brain sections were sliced on a vibratome. The identification of paraventricular nucleus (bregma – 1.80 mm to – 2.12 mm) was based on the atlas by Paxinos and Watson. The sections above were rinsed three times in PBS 10 min and then incubated with 0.2% Triton for 5 min. The sections were rinsed three times in PBS 10 min and incubated with bovine serum albumin (2% BSA) (Sigma Co., USA) for 1 h. Thereafter the sections were incubated in the primary antibody (p-ERK1/2-Ab, 1:100, Boster Company, P.R. China) overnight at 4 °C. The sections were then rinsed three times in PBS for 10 min and incubated with 2 % BSA for 20 min. Subsequently, the sections were rinsed three times in PBS for 10 min, and further incubated in the

biotinylated-second antibody (Boster Company, P.R. China) at room temperature for 2 h. Finally, the sections were again rinsed three times in PBS for 10 min and incubated with the avidin-biotin complex ABC (Boster Company, P.R. China) at room temperature for 2 h. Diaminobenzidine (DAB, Sigma Co, USA) was used for signal detection. The control sections were incubated with PBS instead of primary antibody. The HPIAS series colorful pathology photographic system was used to analyze p-ERK1/2-IR positive neurons. The brain sections were observed in a 20× magnification. The number and optical density of p-ERK1/2-IR positive neurons were calculated per area and per group.

Statistical Analysis

All data were expressed as mean \pm S.E.M. Statistical evaluation was done using ANOVA with post hoc test of LSD in Equal Variances Assumed. In all comparisons, statistical significance was set at $P<0.05$.

Results

1. Effect of GS on renal MDA and SOD level

Glycerol-treated rats (ARF+NS group) showed a significant lipid peroxidation as indicated by a marked increase of renal MDA level and decreased SOD level as compared to those in NS+NS group. However, treatment with GS for 48 h (ARF+GS group) significantly reduced the MDA level and restored the SOD level as compared to ARF+NS group (Fig. 1, $P<0.05$).




Fig.1 Changes of MDA and SOD in renal cortex homogenate in the ARF rats treated with GS for 48 h. A. MDA in renal cortex; B. SOD in renal cortex. Data are mean \pm S.E.M. n=10
(* P<0.05 ARF+NS group vs. NS+NS group, # P<0.05 ARF+GS group vs. ARF+NS group).

2. Effect of GS on the changes of ChAT-IR in the PCT in ARF rats induced by glycerol

In the NS+NS group, ChAT-IR positive granules were detected very weakly in proximal convoluted tubular cells (Fig. 2C). Optical density of ChAT-IR positive granules in the ARF+NS group was significantly increased in PCT compared with that in the NS+NS group (Fig. 2A, P<0.05). There was a further increase of optical density of ChAT-IR positive granules in PCT of ARF+GS group (Fig. 2B, P<0.05) when compared with that in ARF+NS group. No significant difference was found in optical density of ChAT-IR positive granules in PCT, when the NS +GS group (Fig.2D, P<0.05)was compared with the NS+NS group. The data on ChAT-IR in the PCT were summarized in Figure 2.




Fig. 2 Changes of ChAT-IR in PCT treated with GS for 48 h in ARF rats. A, B, C and D were stained by DAB. A:ARF+NS group; B:ARF+GS group; C:NS+NS group; D:NS+GS group. Bar indicates 60.6 μ M, arrow points to ChAT-IR positive granules. E: Quantitative analysis of ChAT-IR positive granules in PCT. *P < 0.05 vs. NS+NS group; #P < 0.05 vs. ARF+NS group.

3. Effect of GS on the changes of p-ERK1/2-IR in the PVN in ARF rats induced by glycerol

Immunohistochemistry also showed an obvious increase of p-ERK 1/2 -IR in the PVN of ARF+NS group (Fig. 3A, P<0.05); but p-ERK-IR 1/2 was further enhanced in ARF+GS group compared with ARF+NS group (Fig. 3B, P<0.05)




Fig. 3 Changes of p-ERK1/2-IR in PVN in the ARF rats treated with GS for 48 h. A, B, C and D

were stained by DAB. A:ARF+NS group; B:ARF+GS group; C:NS+NS group; D:NS+GS group. Bar indicates 60.6 μ M, arrow points to p-ERK1/2-IR positive neurons. E and F: quantitative analysis of positive neurons and optical mean density of p-ERK1/2-IR in PVN. *P < 0.05 vs. NS+NS group; #P < 0.05 vs. ARF+NS group.

Discussion

Intramuscular administration of hypertonic glycerol is the most common used animal model of experimental ARF. The mechanisms of this model is rhabdomyolysis induced by hypertonic glycerol (Ustundag *et al.* 2009). Hypertonic glycerol produced a large amount of myoglobin to leak into circulation. Myoglobin, along with uric acid, can damage the kidney directly by forming

casts within the renal tubules leading to their obstruction. This blockage of renal tubules induced necrosis of renal tubular epithelial cell and produced a large amount of free radical. Myoglobin also can lead to intrarenal vasoconstriction and decreased glomerular filtration rate. Finally, all of these factors cause ARF (Aydogdu *et al.* 2006, Boutaud *et al.* 2011). Our previous paper reported that glycerol administration resulted in a significant increase in BUN and Cre levels as compared to those in NS+NS group, and the kidneys of rats treated with glycerol showed marked histological changes in cortex (Zhang *et al.* 2010). Therefore, the animal model of experimental ARF was successful.

It was reported that the iron released from the heme group of the myoglobin. Iron induces the formation of high-activity oxygen free radicals that increase oxidative stress and provoke lipid peroxidation and cellular death (Johnson *et al.* 2010). It was reported that glycerol caused the renal oxidative stress. Reactive oxygen metabolites were proved to be the key mediators of tissue injury in glycerol-induced ARF model (Ustundag *et al.* 2009).

SOD used to scavenge free radicals was decreased when free radicals was increased (Yoon *et al.* 2008); SOD is a significant index in the research about oxidative damage. In the present study, we observed that MDA level in renal cortex homogenate from ARF rats was significantly increased and the level of SOD in renal cortex homogenate of ARF rats was markedly decreased, indicating that reactive oxidative species were generated in this model in which oxidative damage was enhanced. In addition, we also demonstrated that the level of MDA was significantly decreased and the level of SOD was obviously increased after the treatment with ginsenosides for 48 h. Similarly, ginsenosides exerted a protective effects on the progression of renal damage via inhibiting oxidative stress in type 2 diabetic rats (Kang *et al.* 2010). Ginsenoside Rb2 through

transcription factor AP2 binding sites and its induction greatly activated the SOD gene, upregulated expression of SOD and enhanced antioxidant activity (Kim *et al.* 1996). Taken together, we supposed ginsenosides, as inhibitors of lipid peroxidation resulting from oxidative stress and as an oxidative free radical scavenger through the increase of SOD, played a crucial role in the protection against the damage associated with rhabdomyolysis.

Previous study have indicated in the dog renal cortex the existence of high-affinity choline uptake and the presence of ChAT, the enzyme that converts choline to acetylcholine and acetylcholine grains were also found in dog renal cortex (Pirola *et al.* 1991). It was reported that hypoxia increased the intracellular level of reactive oxygen species and induced cell apoptosis in mouse ES cells; but this effect can be inhibited by ACh administration (Kim *et al.* 2008). It suggested that acetylcholine prevented the hypoxia-induced apoptosis of mouse ES cells by inhibiting the reactive oxygen species production. These evidences implied renal cholinergic system played a important role during oxidative stress. In the present study, we observed that the ChAT-IR was markedly increased in PCT cell of ARF rats, it may be a compensatory mechanism of cholinergic system aganist the increase of oxyradical and renal ischemia, hypoxia. In addition, we also observed a further increase of the ChAT-IR in the PCT of ARF+GS group, compared with that in ARF+NS groups. Our results suggested that ginsenoside further up-regulated renal cholinergic system activity.

Immunostaining for ERK was intense in various brain regions such as neocortex, hippocampus, hypothalamus, brainstem (Flood *et al.* 1998). It was reported that several physiologically stressful stimuli, including seizure induction, ischemic insult and electroconvulsive shock, could activate MAPK signal in various brain regions (Imbe *et al.* 2004).

Meanwhile, the MAPK pathway was activated in the process of Ang II-induced ROS production (Palomeque *et al.* 2009). NA-induced p38 MAPK activation, through alpha-adrenoceptor, had a protective role in cardiomyocytes to antagonize NA-induced cell death (Tsang *et al.* 2009). It prompted that MAPK signaling pathways activation had a cytoprotective action. Other investigators have reported that the ERK pathway activation was an important step in the regulation of oxidative stress-response genes by arecoline (Thangjam *et al.* 2009). The nature of the signal transduction suggested that activation of ERK1/2 may serve as a convergence site for mAChR activation in COS-7 cells (Rosenblum *et al.* 2000). It implied that the MAPK signaling pathways activation may influence cholinergic nervous system. A parasite-derived neurotrophic factor stimulated mRNA and protein expression of ChAT in PC12 cell, and activating MAPK signaling pathways is necessary for this effect (Akpan *et al.* 2008). The neural cell adhesion molecule mimetic peptide C3d promoted ChAT activity in septal neurons through the stimulation of fibroblast growth factor receptor and the activation of the MAPK signaling pathway (Burgess *et al.* 2009). Therefore, we can speculate that the MAPK signaling pathway was involved in the regulation of ChAT activity.

The results of the present study also showed that of the p-ERK1/2-IR in PVN of ARF rats was significantly increased. It indicates that MAPK signal pathway in the PVN was excited after intramuscular injection of glycerol and might be involved in the regulation of cholinergic system in proximal convoluted tubule. In addition, we also observed that glycerol-induced ARF rats treated with ginsenosides for 48 h enhanced the p-ERK1/2-IR as well as the ChAT-IR in PCT.

On the basis of the above results, we propose that glycerol-induced ARF causes the changes not only in the kidney such as acute tubular necrosis, the increase of oxidative stress and the

upregulation of the ChAT-IR in the PCT, but also in central nervous system such as the upregulation of the p-ERK1/2-IR in the hypothalamic PVN. We suggest that both changes in the PCT and PVN should be one of the compensatory mechanisms both in the kidney and in central nervous system in ARF rats. Our study also shows that oral administration of ginsenosides not only further increase renal antioxidative activity by some humoral mechanism, such as activating cholinergic system activity in the PCT, but also activate hypothalamic PVN MAPK signal pathway through some unknown pathways. The possible relationship between the activation of cholinergic system in the PCT and the activation of hypothalamic PVN MAPK signal pathway, and also its correlation with renal protective effect of ginsenosides against ARF will be a focal point in our lab for the future.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

This work was supported by grant No. 9910500711 from the Natural Science Foundation of Liaoning Province, China. The present address of Hong-mei Liu: Tangshan Gongren Hospital, Tangshan , Hebei, China.

References

AKPAN N, CARADONNA K, CHUENKOVA MV, PEREIRAPERRIN M: Chagas' disease parasite-derived neurotrophic factor activates cholinergic gene expression in neuronal PC12

cells. Brain Res 1217:195-202, 2008.

ATTELE AS,WU JA,YUAN CS: Ginseng pharmacology: multiple constituents and multiple actions. Biochem Pharmacol 58(11): 1685-1693, 1999 .

AYDODGU N,ATMACA G,YALCIN O,TASKIRAN R,TASTEKIN E,KAYMAK K:Protective effects of L-carnitine on myoglobinuric acute renal failure in rats. Clin Exp Pharmacol Physiol 33(1-2) : 119-124, 2006.

BEAK SH,PIAO XL,LEE UJ,KIM HY,PARK JH: Reduction of Cisplatin-induced nephrotoxicity by ginsenosides isolated from processed ginseng in cultured renal tubular cells. Biol Pharm Bull 29(10):2051-2055, 2006.

BALDANZI G,FILIGHEDDU N,CUTRUPI S,CATAPANO F,BONISSONI S,FUBINI A,MALAN D,BAJ G,GRANATA R,BROGLIO F,PAPOTTI M,SORICO N,BUSSOLINO F,ISGAARD J,DEGHENGHI R,SINIGAGLIA F,PRAT M,MUCCIOLI G,GHIGO E,GRAZIANI A: Ghrelin and des-acyl ghrelin inhibit cell death in cardiomyocytes and endothelial cells through ERK1/2 and PI 3-kinase/AKT. J Cell Biol 159(6):1029-1037, 2002.

BOUTAUD O, ROBERTS LJ ANG : Mechanism-based therapeutic approaches to rhabdomyolysis-induced renal failure. Free Radic Biol Med 51(5):1062-1067, 2011.

BURGESS A,SAINI S,WENG YQ,AUBERT I: Stimulation of choline acetyltransferase by C3d, a neural cell adhesion molecule ligand. J Neurosci Res 87(3):609-616, 2009.

DE ARRUDA CAMARGO GM, DE ARRUDA CAMARGO LA, SAAD WA: On a possible dual role for the lateral septal area 5-HT(1A) receptor system in the regulation of water intake and urinary excretion. Behav Brain Res 215(1):122-128, 2010.

FLOODDG,FINNJP,WALTONKM,DIONNE CA,CONTRERASPC,MILLER MS,BHAT RV:

Immunolocalization of the mitogen-activated protein kinases p42MAPK and JNK1, and their regulatory kinases MEK1 and MEK4, in adult rat central nervous system. *J Comp Neurol* 398(3):373-392, 1998 .

HERRING BE, MAYFIELD RD, CAMP MC AND ALCANTARA AA: Ethanol-induced Fos

immunoreactivity in the extended amygdala and hypothalamus of the rat brain: focus on cholinergic interneurons of the nucleus accumbens. *Alcohol Clin Exp Res* 28(4):587-597, 2004.

IMBE H, MURAKAMI S, OKAMOTO K, IWAI-LIAO Y, SENBA E: The effects of acute and

chronic restraint stress on activation of ERK in the rostral ventromedial medulla and locus coeruleus. *Pain* 112(3): 361-371, 2004 .

JOHNSON AC, BECKER K, ZAGER RA. Parenteral iron formulations differentially affect

MCP-1, HO-1, and NGAL gene expression and renal responses to injury. *Am J Physiol Renal Physiol* 299(2):F426-435, 2010.

KANG KS, YAMABE N, KIM HY, PARK JH, YOKOZAWA T: Effects of heat-processed

ginseng and its active component ginsenoside 20(S)-Rg3 on the progression of renal damage and dysfunction in type 2 diabetic Otsuka Long-Evans Tokushima Fatty rat. *Biol Pharm Bull* 33(6):1077-1081, 2010.

KIM MH, KIM MO, HEO JS, KIM JS, HAN HJ: Acetylcholine inhibits long-term

hypoxia-induced apoptosis by suppressing the oxidative stress-mediated MAPKs activation as well as regulation of Bcl-2, c-IAPs, and caspase-3 in mouse embryonic stem cells.

Apoptosis 13(2):295-304, 2008.

KIM YH, PARK KH, RHO HM: Transcriptional activation of the Cu, Zn-superoxide dismutase gene through the AP2 site by ginsenoside Rb2 extracted from a medicinal plant, Panax ginseng. *J Biol Chem* 271(40):24539-24543, 1996.

LIU JJ, LI DL, ZHOU J, SUN L, ZHAO M, KONG SS, WANG YH, YU XJ, ZHOU J, ZANG WJ: Acetylcholine prevents angiotensin II-induced oxidative stress and apoptosis in H9c2 cells. *Apoptosis* 16(1):94-103, 2011.

MA W, MARIC D, LI BS, HU Q, ANDREADIS JD, GRANT GM, LIU QY, SHAFFER KM, CHANG YH, ZHANG L, PANCRAZIO JJ, PANT HC, STENGER DA, BARKER JL: Acetylcholine stimulates cortical precursor cell proliferation in vitro via muscarinic receptor activation and MAP kinase phosphorylation. *Eur J Neurosci* 12(4):1227-1240, 2000.

PALOMEQUE J, RUEDA OV, SAPIA L, VALVERDE CA, SALAS M, PETROFF MV, MATTIAZZI A: Angiotensin II-induced oxidative stress resets the Ca²⁺ dependence of Ca²⁺-calmodulin protein kinase II and promotes a death pathway conserved across different species. *Circ Res* 105(12):1204-1212, 2009.

PIROLA CJ, ALVAREZ AL, FINKIELMAN S, NAHMOD VE: Release of acetylcholine from isolated canine renal tissue. *Am J Physiol* 260(2 Pt 2):F198-203. 1991.

ROBEY RB, RUIZ OS, BANIQUED J, MAHMUD D, ESPIRITU DJ, BERNARDO AA, ARRUDA JA: SFKs, Ras, and the classic MAPK pathway couple muscarinic receptor activation to increased Na-HCO₃ cotransport activity in renal epithelial cells. *Am J Physiol Renal Physiol* 280(5):F844-850, 2001.

ROSENBLUM K, FUTTER M, JONES M, HULME EC, BLISS TV: ERK1/II regulation by the muscarinic acetylcholine receptors in neurons. *J Neurosci* 20(3):977-985, 2000.

THANGJAM GS, KONDAIAH P: Regulation of oxidative-stress responsive genes by arecoline in human keratinocytes. *J Periodontal Res* 44(5): 673-682, 2009.

TSANG MY, RABKIN SW: p38 mitogen-activated protein kinase (MAPK) is activated by noradrenaline and serves a cardioprotective role, whereas adrenaline induces p38 MAPK dephosphorylation. *Clin Exp Pharmacol Physiol* 36(8): e12-19, 2009.

USTUNDAG S, SEN S, YALCIN O, CIFTCI S, DEMIRKAN B, TURE M: L-Carnitine ameliorates glycerol-induced myoglobinuric acute renal failure in rats. *Ren Fail* 31(2):124-133, 2009.

VLAHOVIC P, CVETKOVIC T, SAVIC V, STEFANOVIC V: Dietary curcumin does not protect kidney in glycerol-induced acute renal failure. *Food Chem Toxicol* 45:1777-1782, 2007.

YOKOZAWA T, OWADA S: Effect of ginsenoside-Rd in cephaloridine-induced renal disorder. *Nephron* 81(2):200-207, 1999.

YOON HJ, MOON ME, PARK HS, KIM HW, IM SY, LEE JH, KIM YH: Effects of chitosan oligosaccharide (COS) on the glycerol-induced acute renal failure in vitro and in vivo. *Food Chem Toxicol* 46(2):710-716, 2008.

ZHANG HA, WANG M, ZHOU J, YAO QY, MA JM, JIANG CL: Protective effect of ginsenoside against acute renal failure and expression of tyrosine hydroxylase in the locus coeruleus. *Physiol Res* 59(1):61-70, 2010.