

## ***Ascophyllum nodosum* extract improves phenolic compound content and antioxidant activity of medicinal and functional food plant *Achillea millefolium* L.**

Ana Cláudia Pacheco\*<sup>1</sup>, Leonardo Araujo Sobral<sup>1</sup>, Pedro Henrique Gorni<sup>1</sup>, Marcia Eugenia Amaral Carvalho<sup>2</sup>

<sup>1</sup>Universidade do Oeste Paulista, Presidente Prudente – SP, Brazil

<sup>2</sup>Escola Superior de Agricultura “Luiz de Queiroz”, Piracicaba – SP, Brazil

\*Corresponding author: [anaclau@unoeste.br](mailto:anaclau@unoeste.br)

### Abstract

Genetic, biochemical and physiological parameters can be changed by applying seaweed extract-based products. However, there is scarce information about the influence of seaweed extract on yarrow performance (*Achillea millefolium* L.), which is widely used in the folk medicine. Therefore, this study aimed to evaluate the effects of *Ascophyllum nodosum* extract on plant development (leaf and root biomass, and leaf area), physiological indexes (leaf weight ratio, and root: leaf ratio), secondary metabolite (phenolic compounds) content and antioxidant activity of yarrow. The experiment was carried out in a completely randomized design with 4 treatments (seaweed extract concentrations 0, 3, 6 and 9 mL L<sup>-1</sup>) and 10 replications. The higher concentration of seaweed extract caused higher total dry weight of plants (from 17.8 to 19%), especially due to increases in the root biomass (up to 28.5%). Only plants that received the highest concentration of seaweed-based product presented increments in the number of leaves when compared to the control plants (18.3 %). Furthermore, the use of *A. nodosum* extract 9 mL L<sup>-1</sup> provided increases in the antioxidant activity and content of phenolic compounds in leaves (up to 30.44%). In conclusion, application of *A. nodosum* is a potential tool strategy to improve the quality of raw material from yarrow plants, since it increased the phenolic compound content and antioxidant activity in leaves, which are the plant organs commonly used in folk medicine.

**Keywords:** biostimulant, elicitor, medicinal plant, leaf pigments, plant defense, seaweed extract, secondary compounds.

**Abbreviations:** ANE\_ *Ascophyllum nodosum* extract, LDM\_leaf dry mass, LWR\_leaf weight ratio, LDLs\_low-density lipoproteins, NL\_number of leaves, ROS\_reactive oxygen species, RDM\_root dry mass, TDM\_total dry mass, LA\_total leaf area, LA/NL\_individual leaf area.

### Introduction

For millenniums, medicinal plants have been employed by humans in diverse areas that include medicine, nutrition, flavouring, beverages, dyeing, repellents, fragrances and cosmetics (Najafi and Deokule, 2010). *Achillea millefolium* L. (yarrow) belongs to Asteraceae family that is represented by about 85 species, which are mostly found in Europe and Asia (Turner and Wasson, 1997). Antioxidant properties of *A. millefolium* have previously been reported in hydroalcoholic, methanolic and aqueous extracts, as also in its essential oils (Trumbeckaite et al., 2011; Vitalini et al., 2011). This property is related to presence of phenolic compounds in leaves, especially flavonoids (hyperoside as the major class) and phenolic acids (chlorogenic acid and p-coumaric acid are the most abundant compounds) (Georgieva et al., 2015).

In order to supply the pharmaceuticals, food additives, flavours and herbal industries with high-quality raw materials, and add value to these products, strategies such as treating with elicitors have been used in both plant cell culture and in intact plants (Zhao et al., 2005). Elicitors refer to chemicals or natural products that can trigger physiological and morphological responses and

accumulation of secondary metabolites, such as polyphenols in plants (Dong et al., 2010). For instance, the use of salicylic acid or utilization of a product based on a mix of hormones (cytokinins, auxins and gibberellic acid) elicited the production of phenolic compounds in marigold, yarrow, and fennel leaves, so increasing the quality of their raw material (Machado et al., 2014; Gorni and Pacheco, 2016; Gorni et al., 2017).

Medicinal plant farming should be carried out under organic system (Mógor et al., 2008). In this context, the use of seaweed extract is an environmentally friendly alternative to replace some fertilizers and biostimulants (Craigie, 2011). *Ascophyllum nodosum* (L.) Le Jolis extract contains several hormones, and also other compounds (such as amino acids and polysaccharides) that stimulate plant growth and yield, also improving tolerance to biotic and abiotic stresses through elicitor mechanisms (Rayorath et al., 2008; Craigie, 2011; Vera et al., 2011; Carvalho et al., 2013, 2014; Carvalho and Castro, 2014; Battacharyya et al., 2015; Castro et al., 2017). One of these mechanisms is related to the presence of polysaccharides and its derivated-oligosaccharides that trigger oxidative stress in plants, which is able to activated

defense pathways, hence increasing gene expression and synthesis of secondary metabolites (Vera et al., 2011). Previously, an increased production of phenolic compounds and flavonoids were reported in cauliflower inflorescence (*Brassica oleracea* cv. Caraflex) after seaweed-based product application (Lola-Luz et al., 2013). Taking account this information, the use of seaweed extracts may potentially enhance both synthesis of bioactive compounds and biomass production of medicinal and functional food plants. However, there are scarce information about the influence of seaweed-based products on yarrow development or on quality of its leaves, which are used as raw material in phytomedicine and beverage industries. Therefore, this work aimed to evaluate the effects of *A. nodosum* extract on plant growth, as well as on phenolic compounds and antioxidant activity in leaves of *A. millefolium*.

## Results and Discussion

### Yarrow development

The use of seaweed extract in agriculture have increased in the last decades due to its positive effects on plant development and quality of edible portions (Craigie, 2011; Lola-Luz et al., 2013; Carvalho and Castro, 2014; Battacharyya et al., 2015). In this study, root dry weight increased linearly to the increases of seaweed extract concentration (up to 28.5% when compared to control, Fig 1). Application of *A. nodosum* extract also improved the root architecture of wheat cv. Pusa Gold, which presented an increased lateral root formation and, consequently, a higher root volume than the control plants (Kumar and Sahoo, 2011). This response can be related to the fact that *A. nodosum* extract modulates concentration and localization of auxin, which is the main hormone that regulates root development (Rayorath et al., 2008; Taiz and Zeiger, 2010). Therefore, seaweed extract potentially enhances nutrient uptake by plants through stimulus of root development, as reported by Mancuso et al. (2006).

Santos et al. (2013) also shown an increased root dry weight in maize (15.59%), in addition to increments in the leaf dry weight (up to 33.9%) after application of seaweed-based product. However, there were no statistical differences among treated and non-treated plants for both parameters (Santos et al., 2013). Application of seaweed extract on *A. millefolium* did not change the leaf dry weight, when compared to the control (Fig 1). Therefore, use of seaweed extract increased the total dry mass (from 17.8 to 19%) (Fig 1) and root: shoot ratio (Fig 2) because root biomass represented the highest proportion of plant weight. For the same reason, a decreased leaf weight ratio (Table 1) was observed in seaweed-treated plants when compared to control ones. On the other hand, wheat cv. IAC 364 exhibited an improved shoot development after application of *A. nodosum* extract as either seed treatment or soil irrigation (Carvalho et al., 2014), reinforcing reports that seaweed extract effects depend on several factors, such as dose, application mode and frequency, plant species and even cultivars (Carvalho and Castro, 2014).

In this study, yarrow plants presented an increased number of leaves (18.3%) after seaweed extract application (Fig 2),

corroborating Silva et al. (2012) who reported an improved yield in cabbage (*Brassica oleracea* due to the increments in both number and biomass of leaves after seaweed application. However, use of *A. nodosum* extract trends to reduce the leaf area per yarrow plant (from 33.85 to 40.8%), due to decreases in the individual leaf area after the use of seaweed-based product (Table 2). These responses can be related to the alterations in cytokinin production by plants, a hormone that drives the leaf development (Taiz and Zeiger, 2010), since *A. nodosum* extract can modifies its endogenous synthesis (Zhang and Ervin, 2004). Taking together these results, the effects of *A. nodosum* on yarrow plants is probably due to the changes in hormonal synthesis that disturbs photoassimilate production and partitioning, inducing biomass accumulation in roots rather than in shoots (Fig 1).

### Chlorophyll, carotenoid and anthocyanin contents

Seaweed extract application did not affect anthocyanin, carotenoid, total chlorophyll and chlorophyll *b* contents, but decreased chlorophyll *a* content when applied at 6 mL L<sup>-1</sup> (Table 2). The biosynthesis of chlorophylls *a* and *b* in 'IAC 364' wheat was not changed after use of *A. nodosum* extract either on seeds or as soil application (Carvalho et al., 2014). However, Blunden et al. (1996) reported that application *A. nodosum* extract as foliar spray resulted in higher chlorophyll concentration in tomato, wheat, barley, and maize leaves. Moreover, these authors associated the increased chlorophyll production in treated plants to the presence of betaines in seaweed extract. However, according to the literature, there is a close relation between chlorophyll synthesis and the applied dose of seaweed extract; where lower doses would be the most effective in promoting increases in chlorophyll content (Jothinayagi and Anbazhagan, 2009). The method of application is also referred as crucial factor to trigger increases in the chlorophyll content (Matysiak et al., 2011).

### Seaweed extract effects on plant defense


The total phenolic content increased linearly to the increases in the seaweed extract concentration (Fig 3). Phenolic compounds are the most widely distributed secondary metabolites, ubiquitously present in the plant kingdom, that fulfill a very broad range of physiological roles in plants related to the plant growth and survival (Cheynier et al., 2013). The expression "plant phenolics" encompasses a highly diverse group with an extremely large structural diversity: tens of thousands of diverse structures have been identified, with the number continually increasing (Quideau et al., 2011). Lola-Luz et al. (2013) also reported an increased (1.3 fold) production of phenolic compounds in cauliflower's inflorescence that was treated with two different seaweed-based products. Usually, these products contain polysaccharides and its derivated-oligosaccharides, which are able to induce synthesis of secondary metabolites (such as terpenes, terpenoids, phenols and/or alkaloids) through change in the gene expression (Vera et al., 2011).

The most remarkable feature of polyphenols is their capability to scavenge reactive oxygen species (ROS), which

**Table 1.** Total leaf area (LA, cm<sup>2</sup> plant<sup>-1</sup>), superficial area per leaf unit (LA/NL, cm<sup>2</sup> leaf<sup>-1</sup>), and leaf weight ratio (LWR) of *Achillea millefolium* that was treated with different concentrations (mL L<sup>-1</sup>) of *Ascophyllum nodosum* extract (ANE).

ANE	LA <sup>ns</sup>	LA/NL*	LWR**
0	1315.15 ± 106.80 a	32.51 ± 3.49 a	0.22 ± 0.011 a
3	1080.03 ± 90.31 a	25.84 ± 2.96 ab	0.16 ± 0.010 b
6	1015.91 ± 48.06 a	29.51 ± 1.41 ab	0.17 ± 0.009 b
9	1091.72 ± 101.66 a	24.65 ± 5.91 b	0.16 ± 0.009 b

NL = number of leaves. Means followed by distinct letters differ by Tukey's test. Bars represent the standard deviation of the mean (\*\* significant at 1%, \* significant at 5% and ns not significant).




**Fig 1.** Leaf (LDM), root (RDM) and total dry mass (TDM) of *Achillea millefolium* that was treated with different concentrations of *Ascophyllum nodosum* extract. Distinct letters differ by Tukey's test at  $p \leq 0.05$ . Bars represent the standard deviation of the mean. Regression equation significant at  $p \leq 0.05$  (\*).


**Table 2.** Chlorophyll a (Chl a), chlorophyll b (Chl b) total chlorophyll (Chl t), anthocyanins (Ant), and carotenoid (Car) contents of *Achillea millefolium* L. that was treated with different concentrations (mL L<sup>-1</sup>) of *Ascophyllum nodosum* extract (ANE).

ANE	Chl a*	Chl b*	Chl t*	Ant <sup>ns</sup>	Car <sup>ns</sup>
	( $\mu\text{g ml}^{-1}$ )				
0	3.2619 ± 0.3377 a	2.2637 ± 0.1726 ab	5.2843 ± 0.4967 ab	3.2138 ± 0.1632 a	0.6005 ± 0.0783 a
3	3.1212 ± 0.4060 ab	3.1898 ± 0.5823 a	6.3111 ± 0.9284 a	4.4297 ± 1.1222 a	0.7089 ± 0.1080 a
6	1.8515 ± 0.1207 b	1.6554 ± 0.1331 b	3.5069 ± 0.2520 b	2.5050 ± 0.2202 a	0.4694 ± 0.0257 a
9	2.5465 ± 0.3056 ab	2.1895 ± 0.1624 ab	4.7361 ± 0.4655 ab	2.8044 ± 0.1140 a	0.6664 ± 0.0915 a

Means followed by distinct letters differ by Tukey's test. Bars represent the standard deviation of the mean (\* significant at 5% and ns not significant).



**Fig 2.** Number of leaves and root: shoot ratio of *Achillea millefolium* that was treated with different concentrations of *Ascophyllum nodosum* extract. Distinct letters differ by Tukey's test at  $p \leq 0.05$ . Bars represent the standard deviation of the mean. Regression equation significant at  $p \leq 0.05$  (\*) and  $p \leq 0.01$  (\*\*).



**Fig 3.** Content of total phenols and antioxidant activity of *Achillea millefolium* that was treated with different concentrations of *Ascophyllum nodosum* extract. Distinct letters differ by Tukey's test at  $p \leq 0.05$ . Bars represent the standard deviation of the mean. Regression equation significant at  $p \leq 0.01$ (\*\*).

include radical and nonradical oxygen species such as  $O_2^{\cdot-}$ ,  $HO\cdot$ ,  $NO\cdot$ ,  $H_2O_2$ ,  $^1O_2$ , and  $HOCl$ , as well as oxidatively generated free radicals  $RO\cdot$  and  $ROO\cdot$  derived from biomolecules like low-density lipoproteins (LDLs) (Neudorffer, 2006) proteins, and oligonucleic acids (Shi et al., 2000). All these species can have deleterious effects on human health (Ferguson, 2001). The antioxidant activity in ethanolic leaf extract of seaweed-treated yarrow exhibited lower  $IC_{50}$  values, which indicate a higher antioxidant activity than control plants. It can be noted that *A. nodosum* extract  $9\text{ mL L}^{-1}$  significantly decreased  $IC_{50}$  (70%), when compared to the non-treated plants (Fig 3). Significant increases in the antioxidant activities were associated with the increased phenolic and flavonoid content in plants treated with *A. nodosum* extract (Elansary et al., 2016; Lola-Luz et al., 2014). Although phenolic compounds are the major antioxidant class in plants, other compounds (such as betacyanins,  $\alpha$ -tocopherol, ascorbic acid and  $\beta$ -carotene) may act as free radical scavengers, hence contributing to the oxidative stress stabilization (Janda et al., 2014; Brandão et al., 2014). Several studies have indicated that the regular intake of plant-origin products [such as fruits, vegetables, spices and medicinal herbs (as tea, for example)] prevents and /or reduces the risk of chronic and degenerative diseases that are triggered by cellular oxidative stress, particularly due to presence of substances with antioxidant activity like as phenolic compounds (Rumbaoa et al., 2009; Cardoso Silva et al., 2010; Engel et al., 2016). Despite their structural diversity, phenolic compounds are categorized into several classes. Among them, phenolic acids, flavonoids and tannins are regarded as the main dietary phenolic compounds

(Balasundram et al., 2006). In this context, water extracts of *A. millefolium* (obtained from infusion or decoction methods) can be intake in the everyday life, since this extract maintain total polyphenolic content and antioxidant capacity (Georgieva et al., 2015).

## Materials and Methods

### Plant materials and growth conditions

The experiment was carried out in greenhouse with controlled temperature ( $26\text{ }^\circ\text{C}$ ) and humidity (70%), in Presidente Prudente, São Paulo state, Brazil ( $22^\circ 7' 39''\text{ S}$ ,  $51^\circ 23' 8''\text{ W}$ , 471 m.a.s.l.). The seedlings were collected from yarrow plant matrices that were grown at Medicinal Plant Garden of Universidade do Oeste do Estado de São Paulo. Specimens were deposited in the Universidade Federal de Uberlândia's Herbarium (voucher # 74428 HUFU). Seedlings with three completely expanded leaves were transplanted to  $20\text{-dm}^3$  vessels that were filled with 18 kg of soil (Table 1), and plants were managed as recommended by *Bulletim 100 – IAC* for perennial herbaceous species. Foliar sprays with solutions containing 0 (only water); 3; 6 and  $9\text{ mL L}^{-1}$  of seaweed extract were applied three times - 20, 30 and 100 days after seedlings transplanting (DAT). As surfactant agent, nonyl phenoxy poly (ethyleneoxy) ethanol (Agral<sup>®</sup>)  $50\text{ }\mu\text{L L}^{-1}$  was added to the solutions that were sprayed through a hand sprayer. Physical and chemical properties of the soil that was used in the experiment are described as following: pH ( $\text{CaCl}_2$ ) 4.3, Ca  $5.5\text{ (mmol}_c\text{ dm}^{-3})$ , Mg  $4.2\text{ (mmol}_c\text{ dm}^{-3})$ , K 3.6, P 3.0, S 24.5, Mn 0.4, Fe 3.8, Cu 1.0, Zn 0.5 and B  $0.14\text{ (mg dm}^{-3})$ .

### Growth analyses and physiological indexes

At 120 DAT, leaves and roots (rhizomes + radicles) were collected. Next, the number of leaves (NL) and leaf area (LA – cm<sup>2</sup>) per plant were evaluated. All organs were placed into an oven with air circulation at 40 °C, in order to obtain their dry mass (g). With these data, (i) leaves, roots and total dry mass per plant (LDM, RDM and TDM, respectively), (ii) root: leaf ratio (RDM/LDM), (iii) leaf weight ratio (LDM/TDM), and (iv) individual leaf area (LA/NL) were calculated.

### Quantification of the leaf pigments

Chlorophyll (*a*, *b* and total), carotenoid and anthocyanin contents were spectrophotometrically determined following the extraction on TRIS-acetone buffered solution (hydroximetil-aminomethan), according to the method of Sims and Gamon (2002).

### Determination of the total phenol content and antioxidant capacity

Total phenol content in leaves was measured by Folin-Ciocalteu method, using gallic acid as standard in sodium carbonate solution (Stagos et al., 2012). In order to evaluate the antioxidant activity of yarrow, different volumes of leaf extract, which were required to decrease initial concentration of 2,2-diphenyl-1-picrylhydrazyl (DPPH) by 50% - inhibitory concentration (IC<sub>50</sub>), were used, as described by Brand-Williams et al. (1995). For assessments of total phenol content and antioxidant activity, ethanolic extracts from dried leaves were used.

### Statistical analyses

The experiment was carried out in a completely randomized design with 4 treatments (seaweed extract doses) with 10 and 5 replications (*i.e.* one plant per experimental unit) for biometric and biochemical analyses, respectively. The obtained data were submitted to analysis of variance ( $p \leq 0.05$ ), and then to both regression analysis and Tukey's test (all them considering  $p \leq 0.05$ ).

### Conclusion

Application of *A. nodosum* is a potential tool to improve the quality of raw material from yarrow, since it increased the phenolic compound content and antioxidant activity in leaves. Rises in the content of phenolic compounds may be related to both seaweed extract-induced production of new young leaves and presence of elicitor substance in *A. nodosum* extract, which are able to change secondary metabolism of plants. Data also revealed that this seaweed extract affects photoassimilate partitioning by inducing biomass allocations to roots without changing leaf dry weight.

### Acknowledgements

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 for granted to PHG and MEAC.

### References

Balasundram N, Sundram K, Samman S (2006) Phenolic compounds in plants and agri-industrial by-products:

antioxidant activity, occurrence, and potential uses. *Food Chem.* 99(1): 191-203.

Battacharyya D, Babgohari MZ, Rathor P, Prithiviraj B (2015) Seaweed extracts as biostimulants in horticulture. *Sci Hortic.* 196 (30): 39-48.

Blunden G, Jenkins T, Liu YW (1996) Enhanced leaf chlorophyll levels in plants treated with seaweed extract. *J Appl Phycol.* 8(6): 535-543.

Brandão I, Kleinowski AM, Einhardt AM, Lima MC, Amarante L, Peters JA, Braga EJB (2014) Salicylic acid on antioxidant activity and betacyanin production from leaves of *Alternanthera tenella*. *Cienc Rural.* 44(10): 1893-1898.

Brand-Williams W, Cuvelier ME, Berset C (1995) Use of a free radical method to evaluate antioxidant activity, *Lebensmittel-Wissenschaft und -Technologie.* *Food Sci Technol Int.* 28(1): 25-30.

Cardoso Silva ML, Costa RS, Santana AS, Koblitz MGB (2010) Compostos fenólicos, carotenóides e atividade antioxidante em produtos vegetais. *Semina Cienc Agrar.* 31(3): 669-682.

Carvalho MEA, Castro PRC (2014) Extratos de algas e suas aplicações na agricultura. *ESALQ/Divisão de Biblioteca, Piracicaba, Brazil.* 58 p.

Carvalho MEA, Castro PRC, Gallo LA, Ferraz Junior MVC (2014) Seaweed extract provides development and production of wheat. *Agrarian.* 7(23): 166-170.

Carvalho MEA, Castro PRC, Novembre ADC, Chamma HMCP (2013) Seaweed extract improves the vigor and provides the rapid emergence of dry bean seeds. *Am Eurasian J Agric Environ Sci.* 13(8): 1104-1107.

Castro PRC, Carvalho MEAC, Mendes ACCM, Angelini BG (2017) Manual de estimulantes vegetais: nutrientes, biorreguladores, bioestimulantes, bioativadores, fosfitos e biofertilizantes na agricultura tropical. *Agronômica Ceres, Ouro Fino, Brazil.* 453 p.

Cheyrier V, Comte G, Davies KM, Lattanzio V, Martens S (2013) Plant phenolics: recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiol Biochem.* 72: 1-20.

Craigie JS (2011) Seaweed extract stimuli in plant science and agriculture. *J Appl Phycol.* 23(3): 371-393.

Dong J, Wan G, Liang Z (2010) Accumulation of salicylic acid-induced phenolic compounds and raised activities of secondary metabolic and antioxidative enzymes in *Salvia miltiorrhiza* cell culture. *J Biotechnol.* 148(2-3): 99-104.

Elansary HO, Skalicka-Woźniak K, King IW (2016) Enhancing stress growth traits as well as phytochemical and antioxidant contents of *Spiraea* and *Pittosporum* under seaweed extract treatments. *Plant Physiol Biochem.* 4: 1-4.

Engel R, Szabó K, Abrankó L, Rendes K, Füzü A, Takács T (2016) Effect of arbuscular mycorrhizal fungi on the growth and polyphenol profile of marjoram, lemon balm, and marigold. *J Agr Food Chem.* 64 (19): 3733-3742.

Ferguson LR (2001) Role of plant polyphenols in genomic stability. *Mutat Res.* 475(1): 89-111.

Georgieva L, Gadjalova A, Mihaylova D, Pavlov A (2015) *Achillea millefolium* L. – phytochemical profile and *in vitro* antioxidant activity. *Int Food Res J.* 22(4): 1347-1352.

Gorni PH, Brozulato MO, Lourencao RS, Konrad ECG (2017) Increased biomass and salicylic acid elicitor activity in fennel (*Foeniculum vulgare* Miller). *Braz J Food Technol.* 20: e2016172.

- Gorni PH, Pacheco AC (2016) Growth promotion and elicitor activity of salicylic acid on *Achillea millefolium* L. *Afri J Biotechnol.* 15(16): 657-665.
- Janda T, Gondor OK, Yordanova R, Szalai G, Pal M (2014) Salicylic acid and photosynthesis: signaling and effects. *Acta Physiol Plant.* 36(10): 2537-2546.
- Jothinayagi N, Anbazhagan C (2009) Effect of seaweed liquid fertilizer of *Saragassum wightii* on the growth and biochemical characteristics of *Abelmoschus esculentus* (L.) Medikus. *Rec Res Sci Technol.* 1(4): 155-158.
- Kumar G, Sahoo D (2011) Effect of seaweed liquid extract on growth and yield of *Triticum aestivum* var. Pusa Gold. *J Appl Phycol.* 23(2): 251-255.
- Lola-Luz T, Hennequart F, Gaffney M (2013) Enhancement of phenolic and flavonoid compounds in cabbage (*Brassica oleraceae*) following application of commercial seaweed extracts of the brown seaweed (*Ascophyllum nodosum*). *Agr Food Sci.* 22(2): 288-295.
- Lola-Luz T, Hennequart F, Gaffney M (2014) Effect on health promoting phytochemicals following seaweed application, in potato and onion crops grown under a low input agricultural system. *Sci Hortic.* 170: 224-227.
- Machado VPO, Pacheco AC, Carvalho MEA (2014) Effect of biostimulant application on production and flavonoid content of marigold (*Calendula officinalis* L.). *Rev Ceres.* 61(6): 983-988.
- Mancuso S, Barlow PW, Volkmann D, Baluska F (2006) Actin turnover-mediated gravity response in maize root apices: gravitropism of decapped roots implicates gravisensing outside of the root cap. *Plant Signal Behav.* 1(2): 52-58.
- Matysiak K, Kaczmarek S, Krawczyk R (2011) Influence of seaweed extracts and mixture of humic and fulvic acids on germination and growth of *Zea mays* L. *Acta Sci Pol Agr.* 10(1): 33-45.
- Mógor AF, Ono EO, Rodrigues JD, Mógor G (2008) Aplicação foliar de extrato de alga, ácido 1-glutâmico e cálcio em feijoeiro. *Sci Agrar.* 9(4): 431-437.
- Najafi S, Deokule SS (2010) Pharmacognostic study of *Tylophora dalzellii* Hook. f. *J Med Plants Res.* 4(5): 403-406.
- Neudörffer A, Desvergne JP, Bonnefont-Rousselot D, Legrand A, Fleury MB, Largeton M (2006) Protective effects of 4-hydroxycinnamic ethyl ester derivatives and related dehydrodimers against oxidation of LDL: Radical scavengers or metal chelators?. *J Agr Food Chem.* 54(5): 1898-1905.
- Quideau S, Deffieux D, Douat-Casassus C, Pouysegou L (2011) Plant polyphenols: chemical properties, biological activities, and synthesis. *Angew Chem Int Ed Engl.* 50(3): 586-621.
- Rayorath P, Jithesh MN, Farid A, Khan W, Palanisamy R, Hankins SD, Critchley AT, Prithiviraj B (2008) Rapid bioassays to evaluate the plant growth promoting activity of *Ascophyllum nodosum* (L.) Le Jol. using a model plant, *Arabidopsis thaliana* (L.) Heynh. *J Appl Phycol.* 20(4): 423-429.
- Rumbaoa RGO, Cornago DF, Geronimo IM (2009) Phenolic content and antioxidant capacity of Philippine sweet potato (*Ipomoea batatas*) varieties. *Food Chem.* 113(4): 1133-1138.
- Santos VM, Melo AV, Cardoso DP, Gonçalves AH, Varanda MAF, Taubinger M (2013) Uso de bioestimulantes no crescimento de plantas de *Zea mays* L. *Rev Bras Milho Sorgo* 12(3): 307-318.
- Shi X, Ye J, Leonard S, Ding M, Vallyathan V, Castranova V, Dong Z (2000) Antioxidant properties of (-)-epicatechin-3-gallate and its inhibition of Cr (VI)-induced DNA damage and Cr (IV)-or TPA-stimulated NF-κB activation. *Mol Cell Biochem.* 206(1-2): 125-132.
- Silva CP, Garcia KGV, Silva RM, Oliveira LAA, Tosta MS (2012) Desenvolvimento inicial de mudas de couve-folha em função do uso de extrato de alga (*Ascophyllum nodosum*). *Rev Verde.* 6(1): 7-11.
- Sims DA, Gamon JA (2002) Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages. *Remote Sens Environ.* 81(2-3): 337-354.
- Stagos D, Portesis N, Spanou C, Mossialos D, Aligiannis N, Chaita E, Panagoulis C, Reri E, Skaltsounis L, Tsatsakis AM, Kouretas D (2012) Correlation of total polyphenolic content with antioxidant and antibacterial activity of 24 extracts from Greek domestic Lamiaceae species. *Food Chem Toxicol.* 50(11): 4115-4124.
- Taiz L, Zeiger E (2010) *Plant Physiology.* 5th edn. Sinauer Associates, Sunderland, Netherlands. 781 p.
- Trumbeckaite S, Benetis R, Bumblauskienė L, Burdulis D, Janulis V, Toleikis A, Viškelis P, Jakštas V (2011) *Achillea millefolium* L. herb extract: antioxidant activity and effect on the rat heart mitochondrial functions. *Food Chem.* 127(4): 1540-1548.
- Turner RJ, Wasson E (1997) *Botanica: the illustrated A-Z of over 10,000 garden plants and how to cultivate them.* Random House Australia, Sidney, Australia. 1020 p.
- Vera J, Castro J, Gonzalez A, Moenne A (2011) Seaweed polysaccharides and derived oligosaccharides stimulate defense responses and protection against pathogens in plants. *Mar Drugs.* 9(12): 2514-2525.
- Vitalini S, Beretta G, Iriti M, Orsenigo S, Basilico N, Dall'acqua S, Iorizzi M, Fico G (2011) Phenolic compounds from *Achillea millefolium* L. and their bioactivity. *Acta Biochim Pol.* 58(2): 203-209.
- Zhang X, Ervin EH (2004) Cytokinin-containing seaweed and humic acid extracts associated with creeping bentgrass leaf cytokinins and drought resistance. *Crop Sci.* 44(5): 1737-1745.
- Zhao J, Davis LC, Verpoorte R (2005) Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol Adv.* 23(4): 283-333.