

Identification of Developmental-Behavioral Problems in Primary Care: A Systematic Review

abstract

CONTEXT: Recent mandates and recommendations for formal screening programs are based on the claim that pediatric care providers underidentify children with developmental-behavioral disorders, yet the research to support this claim has not been systematically reviewed.

OBJECTIVE: To review research literature for studies regarding pediatric primary care providers' identification of developmental-behavioral problems in children.

METHODS: On the basis of a Medline search conducted on September 22, 2010, using relevant key words, we identified 539 articles for review. We included studies that (1) were conducted in the United States, (2) were published in peer-reviewed journals, (3) included data that addressed pediatric care providers' identification of developmental-behavioral problems in individual patients, (4) included an independent assessment of patients' developmental-behavioral problems, such as diagnostic interviews or validated screening instruments, and (5) reported data sufficient to calculate sensitivity and specificity. Studies were not limited by sample size. Eleven articles met these criteria. We used Quality Assessment of Diagnostic Accuracy Studies (QUADAS) criteria to evaluate study quality. Although the studies were similar in many ways, heterogeneous methodology precluded a meta-analysis.

RESULTS: Sensitivities for pediatric care providers ranged from 14% to 54%, and specificities ranged from 69% to 100%. The authors of 1 outlier study reported a sensitivity of 85% and a specificity of 61%.

CONCLUSIONS: Pediatricians are often the first point of entry into developmental and mental health systems. Knowing their accuracy in identifying children with developmental-behavioral disabilities is essential for implementing optimal evaluation programs and achieving timely identification. Moreover, these statistics are important to consider when planning large-scale screening programs. *Pediatrics* 2011; 128:356–363

AUTHORS: R. Christopher Sheldrick, PhD, Shela Merchant, MA, and Ellen C. Perrin, MD

Department of Pediatrics, Floating Hospital, Tufts Medical Center, Boston, Massachusetts

KEY WORDS

pediatric, screening, behavior disorders/problems, developmental disabilities, systematic reviews

ABBREVIATIONS

PPV—positive predictive value

NPV—negative predictive value

QUADAS—Quality Assessment of Diagnostic Accuracy Studies

All the authors made substantive contribution to the study and/or manuscript and reviewed the final paper before its submission.

www.pediatrics.org/cgi/doi/10.1542/peds.2010-3261

doi:10.1542/peds.2010-3261

Accepted for publication Apr 20, 2011

Address correspondence to R. Christopher Sheldrick, PhD, Department of Pediatrics, Floating Hospital, Tufts Medical Center, 800 Washington St, Box 854, Boston, MA 02111. E-mail: rsheldrick@tuftsmedicalcenter.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2011 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: *The authors have indicated they have no financial relationships relevant to this article to disclose.*

Estimates indicate that at least 1 in 5 children has a developmental and/or behavioral disability.^{1,2} Several recent recommendations demonstrate the growing consensus that early identification is essential for providing adequate treatment to children with such disabilities.^{3,4} To facilitate early identification, the American Academy of Pediatrics Council on Children With Disabilities has recommended that pediatricians and other child health care providers perform ongoing developmental surveillance during all routine health supervision visits, supplemented with standardized screening instruments at specified ages.³ More recently, several states have instituted programs that encourage or even require all child health providers to administer screening instruments at well-child visits.^{5,6}

Such mandates and recommendations are based on the claim that pediatric care providers underidentify children with developmental-behavioral disorders. A systematic review of the evidence for this claim is critical for several reasons. First, to evaluate the utility of any new screening program or recommendation, its effectiveness must be compared with the clinical accuracy of standard pediatric practice that does not include validated screening instruments. Second, most claims about pediatric providers' accuracy focus on their sensitivity and largely ignore their specificity. Hence there is a focus on correctly identifying children who have disorders while ignoring whether the providers correctly identify children without disorders. For a full understanding of clinicians' accuracy in identifying developmental and behavioral problems, both sensitivity and specificity are essential. Third, for many families, especially those with young children, pediatric care providers serve as gatekeepers to mental health and developmental services.⁷ If

providers are, in fact, underidentifying children with developmental-behavioral disorders, finding feasible methods for improving identification is essential for effective treatment.

We have systematically reviewed the research literature for evidence regarding pediatric primary care providers' identification of developmental and behavioral problems in children. We make a distinction between developmental and behavioral disorders because most professional recommendations,³ screening instruments, and diagnostic tests follow this dichotomy and focus on 1 or the other but not both. We recognize that developmental and behavioral disorders (1) cause significant impairment in children and deserve meticulous attention in the pediatric care setting, (2) are not distinct categories (some disorders, such as attention-deficit/hyperactivity disorder, are alternately classified in either category),^{8–10} and (3) are often present in the same child (research has suggested that 75%–80% of children with an autism spectrum disorder also have a comorbid behavioral disorder).^{11,12} Nevertheless, most research studies focus on either behavioral or developmental problems, so we review each as an independent category.

METHOD

To identify articles for this literature review, we conducted a search of the Medline database on September 22, 2010. There was no preexisting protocol for this type of search, so we devised our own method and included search terms particular to the topic at hand.

Criteria for Inclusion

We included studies that (1) were conducted in the United States, (2) were published in a peer-reviewed journal, (3) included data regarding whether pediatric care providers identified a

developmental or behavioral problem in individual patients, (4) included an independent assessment of patients' developmental-behavioral problems, such as diagnostic interviews or validated screening instruments, and (5) reported data for a sufficient number of cases to determine sensitivity and specificity. Studies were not limited by sample size.

Search Strategy

We began by accessing all the articles in which an assessment of childhood developmental or behavioral disorders in pediatric settings was described. Specifically, we searched for the intersection (ie, using the "and" term) of articles in the following groups:

- articles with subject headings that included "developmental disabilities," "delay," "mental health," "mental disorders," "child behavior disorders," "language development disorders," "depression," "anxiety," or "autistic disorders" (note that these terms were combined with an "or" term);
- articles with the subject heading "pediatrics" or "pediatric providers"; and
- articles with subject headings "diagnosis," "identification," "screening," or "surveillance" (note that these terms were combined with an "or" term).

Because research on this topic is sparse and began only in the 1980s, we did not set a date limit on the studies we searched. The final yield was 539 articles.

We included 2 additional search strategies. First, we reviewed the reference section for additional citations in any article that focused on screening or surveillance. We included this additional step because some studies that met inclusion criteria were conducted

for a purpose other than to assess the accuracy of pediatric care providers (eg, to validate a screening instrument). Second, for all articles that met inclusion criteria, we used the “cited by” function in Medline to identify articles that refer to studies that met our inclusion criteria. These 2 additional search strategies yielded 174 new articles for review.

Two of us (Dr Sheldrick and Ms Merchant) conducted the literature review. Studies with titles or abstracts that referred to pediatric care providers’ recognition of developmental or behavioral problems were reviewed in greater detail. Examples of excluded studies are those that focused on pediatric issues not relating to child behavior and development, such as screening for parent psychopathology; neurologic bases of psychiatric disorders; treatment of psychiatric disorders; studies not conducted in the United States; and general summaries of common developmental and behavioral problems seen in the pediatric population. For the remaining articles, we read the abstracts, introductions, and results sections to determine if articles (1) met the inclusion criteria described earlier or (2) included references to other articles that might meet inclusion criteria.

Dr Sheldrick and Ms Merchant reviewed the final set of studies for data from which sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) could be calculated. In the case of multiple studies that reported on the same data set, we included only the earliest study in our analyses. In the case of studies that seemed to meet inclusion criteria (ie, the methods section described a procedure for eliciting pediatricians’ assessment of developmental or behavioral problems as well as a diagnostic interview) but from which such statistics were not reported, the au-

thor was contacted to determine if any such analyses were conducted. Two authors were contacted. In 1 case,¹³ the author reported that analyses relevant to our study had not been performed; thus the study was not included in our analyses. In the second case,¹⁴ the author referred us to another study that used the same data set as a study that was already included in our sample.

Assessment of Clinical Accuracy Studies’ Methodology

To determine if the studies in our final sample were comparable, we assessed all final studies that met inclusion criteria by using the Quality Assessment of Diagnostic Accuracy Studies (QUADAS) criteria. QUADAS provides a framework for evaluating the quality of each study’s methodology and conclusions. It helps to determine if the quality of each study is high enough to have reliable and valid conclusions and whether information across different studies can be integrated.

Clinicians’ Accuracy

Sensitivity, specificity, PPV, NPV, and a diagnostic odds ratio were calculated for each study along with their 95% confidence intervals. For studies with weighted data, true-positive, false-negative, true-negative, and false-positive results were estimated on the basis of the size of the final sample that completed the gold-standard reference test, the sensitivity and specificity reported, and the weighted prevalence and identification rate. For unweighted data, these numbers were recorded verbatim from the original article or estimated directly from reported results. In addition to conducting assessments with parents, some authors also reported data from assessments directly with children. Because only some articles reported such information, we abstracted data only from par-

ent interviews and screening instruments for all studies.

RESULTS

Of the 713 articles identified by the electronic search strategy, 445 were eliminated because of duplications and/or because they did not focus on pediatricians’ recognition of developmental-behavioral problems. Detailed review of the remaining 268 articles yielded 11 articles that met the criteria outlined earlier such that we were able to derive indices of clinicians’ accuracy (see Fig 1).

There were many consistent findings with regard to study quality. For all studies, the time period between the pediatric assessment and the criterion (ie, gold-standard) test was reasonable; all participants in each study received the same criterion test; criterion tests were conducted regardless of the results of the pediatric examination; and pediatric examinations and criterion tests were conducted independently. Withdrawals were generally well explained with 1 exception, in which the 56% enrollment rate was not explained.¹⁵

In other ways, studies varied greatly in quality and design with respect to the purpose of our review. No accepted summary of study quality is available,¹⁶ so relevant study details and QUADAS criteria are instead listed individually in Table 1. For example, 5 studies included only screening instruments as their criterion tests. Because 4 of these 5 studies had large sample sizes, and all 5 of them administered the criterion test to all participants, we deemed their inclusion worthwhile. The remaining 6 studies relied on diagnostic interviews, but the specific interviews used varied. For 10 studies, data regarding whether pediatric providers identified a developmental-behavioral problem derived from brief, provider-completed question-




FIGURE 1
Flow diagram of the literature review.

naires; 1 study relied on an extensive record review. Although most study samples were representative of pediatric populations, 2 were drawn from ongoing studies with high-risk samples. Studies also varied greatly in the proportion of the study population that were successfully enrolled in the study (range: 36%–91%), which indicates different levels of external validity. Some studies focused only on young children, others focused on adolescents, and yet others focused on more comprehensive age ranges.

Nine studies assessed behavioral problems. Only 2 studies^{17,18} assessed developmental problems, 1 of which relied on a screening instrument as a criterion test. No study we identified included diagnostic tests for autism or pervasive developmental disorder.

Because of the high degree of heterogeneity in study methodology and quality, we determined that meta-analytic statistics were inappropriate. Therefore, we list descriptive statistics for each study in Table 1 and show each

study's sensitivity and specificity (with 95% confidence intervals) in Fig 2. One study seems to be an outlier. Brown and Wissow¹⁹ reported that physicians identified 48.6% of their patients as having a behavior problem, which yielded a sensitivity of 0.85 and a specificity of 0.61 when compared with a screening instrument that identified 21.5% of the sample as being at risk. For the remaining studies, sensitivity ranged from 14% to 54% and specificity ranged from 69% to 100%. PPVs ranged from 24% to 66%, and NPVs ranged from 61% to 94%. However, because PPV rises and NPV falls as prevalence increases (all else being equal), these statistics should be interpreted in light of the prevalence rates according to the gold-standard diagnostic or screening test.

DISCUSSION

The American Academy of Pediatrics recommends that developmental screening instruments maintain sensitivity and specificity rates higher

than 70%.⁴ Among the studies examined, pediatric care providers who worked without screening instruments achieved specificity (ie, the proportion of children correctly identified as not having a problem) that was consistently near or higher than 70%. The sensitivity of pediatric care providers was also consistent among the studies but in the opposite direction: in all cases except 1 it was lower than 54%. Thus, the number of children correctly identified as having a developmental or behavioral problem was quite low.

The lack of data on identification of developmental disorders was notable. Nine studies in our sample examined the identification of behavioral problems only, 1 focused on language delays, and 1 focused on a range of developmental problems. This distinction is important, because pediatricians' accuracy might vary according to the type of problem in question. We were unable to identify any study that directly compared pediatric care providers to a clinically determined diagnosis of a developmental disorder. Thus, although diagnoses of autism or pervasive developmental disorder are often reported to be delayed,²⁰ we have no direct evidence regarding the role of pediatric care providers in causing this delay. Moreover, no study assessed both developmental and behavioral disorders in the same children, which makes it impossible to determine if identification of 1 type of problem influences detection of the other.

We note several limitations to this study. The major limitation to improved understanding of pediatricians' accuracy in identifying developmental and behavioral disorders stems from the paucity of research that focuses on this topic (especially identification of developmental delays). In many cases, the primary purpose of studies we included was not to determine if pediatricians accurately

TABLE 1 Study Characteristics, Quality, and Results

Study	Construct Assessed	Criterion Test		Pediatric Care Provider		Sample		Clinical Accuracy					
		Assessment	Prevalence, %	ID Rate, %	Enrollment Rate, %	n	Age Range, y	QUADAS Concern Noted	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	DOR (95% CI)
Studies with diagnostic interviews as criterion tests													
Costello et al ²⁵ (1988)	Behavioral problems	DISC	11.7	5.7	64.6	300 ^a	7–11	None	0.17 (0.08–0.33)	0.96 (0.93–0.98)	0.36 (0.18–0.59)	0.90 (0.86–0.93)	4.9 (1.7–14.1)
Lavigne et al ¹⁵ (1993)	Behavioral problems	Agreement between 2 diagnostic interviews	12.9	8.7	56.0	495 ^a	2–5	Whole or random sample did not receive criterion test ^b	0.21 (0.12–0.32)	0.93 (0.90–0.95)	0.31 (0.19–0.45)	0.89 (0.86–0.91)	3.5 (1.7–7.1)
Merritt et al ²⁶ (1993)	Behavioral problems	CAS	41.5	24.4	59.8	41	6–12	None	0.29 (0.13–0.53)	0.79 (0.60–0.91)	0.50 (0.24–0.76)	0.61 (0.42–0.78)	1.6 (0.4–6.6)
Chang et al ¹⁷ (1988)	Behavioral problems	SADS	43.5	36.5	38.6	85	6–17	Sample not representative of pediatric primary care patients ^c	0.43 (0.29–0.59)	0.69 (0.55–0.80)	0.52 (0.35–0.68)	0.61 (0.48–0.73)	1.7 (0.7–4.1)
Studies with screening instruments as criterion tests													
Costello et al ²⁸ (1985)	Behavioral problems	DISC	70.0	10.0	91.0	40	7–11	Inclusion criteria unclear	0.14 (0.06–0.31)	1.00 (0.76–1.00)	NA	NA	NA
King et al ¹⁸ (2005)	Language delay	Preschool Language Scale-3rd edition	10.1	19.5	36.0	513	3	Sample not representative of pediatric primary care patients ^e	0.33 (0.22–0.46)	0.82 (0.78–0.85)	0.17 (0.11–0.26)	0.92 (0.88–0.94)	2.2 (1.2–4.1)
Kelleher et al ²³ (1997)	Behavioral problems	PSC-35	12.2	17.7	82.0	9766	4–15	None	0.54 (0.51–0.56)	0.87 (0.86–0.88)	0.37 (0.35–0.39)	0.93 (0.92–0.94)	7.9 (6.9–9.0)
Leaf et al ¹⁶ (2004)	Behavioral problems	CBCL	13.6	27.9	78.0	1629	5–9	None	0.50 (0.44–0.57)	0.76 (0.73–0.78)	0.24 (0.21–0.29)	0.91 (0.89–0.92)	3.1 (2.3–4.2)
Murphy et al ³¹ (1992)	Behavioral problems	PSC-35	21.1	18.4	52.8	38	6–16	None	0.50 (0.22–0.79)	0.90 (0.74–0.97)	0.57 (0.25–0.84)	0.87 (0.71–0.95)	9.0 (1.5–56.1)
Brown and Wissow ¹⁹ (2010)	Behavioral problems	SDQ	21.5	48.6	88.1	767	5–16	None	0.85 (0.79–0.90)	0.61 (0.57–0.65)	0.38 (0.33–0.43)	0.94 (0.91–0.96)	8.9 (5.6–14.0)
Hix-Smith et al ¹⁷ (2007)	Developmental problems	ASQ	21.6	10.6	48.9	699	1–2	None	0.32 (0.26–0.40)	0.95 (0.93–0.97)	0.66 (0.55–0.76)	0.84 (0.81–0.86)	10.1 (5.9–17.0)

ID indicates identification; DOR, diagnostic odds ratio; CI, confidence interval; DISC, Diagnostic Interview Schedule for Children; CAS, Child Assessment Schedule; SADS, Schedule for Affective Disorders and Schizophrenia for School-Aged Children; NA, not applicable/not available; PSC-35, Pediatric Symptom Checklist (raw score ≥ 28), 35-item version; CBCL, Child Behavior Checklist (7 score ≥ 64), SDQ, Strengths & Difficulties Questionnaire (high symptoms + impairment); ASQ, Ages & Stages Questionnaire (12- and 24-month forms).

^a Sample that received criterion test; total sample size was larger.

^b Oversampled children with high CBCL scores; children with low CBCL scores were matched, not randomly sampled.

^c Sampled from study of depressed mothers and matched controls (Yale-National Institute of Mental Health Collaborative Family Study).

^d For any mention of concern regarding developmental delay.

^e High-risk sample.




FIGURE 2
Sensitivity, specificity, and 95% confidence intervals.

identified children's developmental and behavioral disorders. In fact, 1 study that collected data that could have been used to address this question did not analyze it (S. M. Horwitz, PhD, Department of Pediatrics, Stanford University, written correspondence, June 5, 2009). Six studies used a structured diagnostic interview as the criterion, and 5 used validated parent-report screening instruments. Given that the reliability and validity of these criterion instruments vary, this information could affect the estimate of pediatricians'

accuracy as well. In addition, studies included in this review predominantly relied on parent report for criterion assessments, but there might be an advantage to including standardized examinations of the child (eg, the Bayley scales) to assess development. Moreover, cutoff scores for various screening instruments also differ; some are set with the aim of maximizing accuracy with respect to diagnostic interviews, whereas others are set to conform to state-level policies. Results of studies that compared pediatric care providers to

screening instruments should, therefore, be interpreted with caution. For reasons such as these, we were not able to identify enough articles with similar design, quality, and sample characteristics to support a meta-analysis; thus, interpretation must occur at the level of each individual study. Finally, our search strategy might have missed some studies that would otherwise have met our inclusion criteria. Because relevant articles may be published under diverse subject headings, identification of studies for inclusion was difficult.

Pediatric care providers have increasingly been encouraged to use standardized screening instruments to identify children with behavioral and developmental problems. The utility of screening instruments depends on how much they are able to improve on pediatricians' standard care, which was the focus of our study. Many screening instruments have been reported to display higher sensitivity than the pediatricians in the studies sampled for the review. However, few screening instruments claim to improve on pediatricians' specificity, and many fall far short.^{21,22}

Given what we know about pediatric care providers' accuracy in identifying developmental and behavioral problems, it is important to consider what

the downstream effects of implementing a screening program might be. Assuming that the screening instrument is truly effective, the proportion of true cases identified could be expected to rise. However, this rise would not occur without cost. Along with the rise in the number of true cases would come an increase in the number of "false-positives," or patients who are incorrectly identified as having a condition. False-positive results require additional assessment, and if they occur in increasing numbers, it would have an effect on pediatricians' time and the capacity of referral sources. These changes might also affect patient satisfaction. A positive screening result suggests the presence of disorder even if that suggestion is later reversed. For parents who have real con-

cerns about their children despite sub-clinical symptoms, such attention might be welcome. However, other parents might still experience anxiety or stigma associated with false-positive results,^{23,24} as has been noted in other areas of medicine.

CONCLUSIONS

Before instituting a screening program, it is essential to consider how that program will change both sensitivity and specificity as well as the downstream effects of these changes. In this article we have summarized what is known about the accuracy of pediatricians' judgments of children's developmental and behavioral status in the absence of formal screening programs.

REFERENCES

- Boyle CA, Decouffé P, Yeargin-Allsopp M. Prevalence and health impact of developmental disabilities in US children. *Pediatrics*. 1994;93(3):399–403
- Merikangas KR, He JP, Burstein M, et al. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication—Adolescent Supplement. *J Am Acad Child Adolesc Psychiatry*. 2010;49(10):980–989
- American Academy of Pediatrics, Council on Children With Disabilities; Section on Developmental Behavioral Pediatrics; Bright Futures Steering Committee; Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening [published correction appears in *Pediatrics*. 2006;118(4):1808–1809]. *Pediatrics*. 2006;118(1):405–420
- Developmental surveillance and screening of infants and young children. *Pediatrics*. 2001;108(1):192–196
- Early and Periodic Screening, Diagnosis, and Treatment, 130 MA ADC 450.140 (2008)
- National Academy for State Health Policy, May J, Kaye N. State strategies to support practice changes that improve identification of children at risk for or with developmental delays: findings from the ABCD Screening Academy. Available at: http://nashp.org/sites/default/files/State_Strategies.pdf?q=files/State_Strategies.pdf. Accessed May 31, 2011
- Dulcan MK, Costello EJ, Costello AJ, Edelbrock C, Brent D, Janiszewski S. The pediatrician as gatekeeper to mental health care for children: do parents' concerns open the gate? *J Am Acad Child Adolesc Psychiatry*. 1990;29(3):453–458
- Panzer A, Viljoen M. Supportive neurodevelopmental evidence for ADHD as a developmental disorder. *Med Hypotheses*. 2005;64(4):755–758
- US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health. *Mental Health: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services; 1999
- Merck Sharp & Dohme Corp. Attention-deficit/hyperactivity disorder (ADHD, ADD). Available at: www.merck.com/mmpe/sec19/ch299/ch299b.html. Accessed October 28, 2010
- De Bruin L, Ferdinand RF, Meester S, de Nijs PF, Verheij F. High rates of psychiatric comorbidity in PDD-NOS. *J Autism Dev Disord*. 2007;37(5):877–886
- Sheldrick RC, Perrin EC. Medical home services for children with behavioral health conditions. *J Dev Behav Pediatr*. 2010;31(2):92–99
- Briggs-Gowan MJ, Horwitz SM, Schwab-Stone ME, Leventhal JM, Leaf PJ. Mental health in pediatric settings: distribution of disorders and factors related to service use. *J Am Acad Child Adolesc Psychiatry*. 2000;39(7):841–849
- Gardner W, Pajer KA, Kelleher KJ, Scholle SH, Wasserman RC. Child sex differences in primary care clinicians' mental health care of children and adolescents. *Arch Pediatr Adolesc Med*. 2002;156(5):454–459
- Lavigne JV, Binns HJ, Christoffel KK, et al. Behavioral and emotional problems among preschool children in pediatric primary care: prevalence and pediatricians' recognition. *Pediatrics*. 1993;91(3):649–655
- Whiting P, Harbord R, Kleijnen J. No role for quality scores in systematic reviews of diagnostic accuracy studies. *BMC Med Res Methodol*. 2005;5:19
- Hix-Small H, Marks K, Squires J, Nickel R. Impact of implementing developmental screening at 12 and 24 months in a pediatric practice. *Pediatrics*. 2007;120(2):381–389
- King TM, Rosenberg LA, Fuddy L, McFarlane E, Sia C, Duggan AK. Prevalence and early identification of language delays among at-risk three year olds. *J Dev Behav Pediatr*. 2005;26(4):293–303
- Brown JD, Wissow LS. Screening to identify

- mental health problems in pediatric primary care: considerations for practice. *Int J Psychiatry Med*. 2010;40(1):1–19
20. Volkmar F, Chawarska K, Klin A. Autism in infancy and early childhood. *Annu Rev Psychol*. 2005;56:315–336
 21. Camp BW. Evaluating bias in validity studies of developmental/behavioral screening tests. *J Dev Behav Pediatr*. 2007;28(3):234–240
 22. Rydz D, Srour M, Oskoui M, et al. Screening for developmental delay in the setting of a community pediatric clinic: a prospective assessment of parent report questionnaires. *Pediatrics*. 2006;118(4). Available at: www.pediatrics.org/cgi/content/full/118/4/e1178
 23. Poulakis Z, Barker M, Wake M. Six month impact of false positives in an Australian infant hearing screening programme. *Arch Dis Child*. 2003;88(1):20–24
 24. Bergman AB, Stamm SJ. The morbidity of cardiac nondisease in schoolchildren. *N Engl J Med*. 1967;276(18):1008–1013
 25. Costello EJ, Edelbrock C, Costello AJ, Dulcan MK, Burns BJ, Brent D. Psychopathology in pediatric primary care: the new hidden morbidity. *Pediatrics*. 1988;82(3 pt 2):415–424
 26. Merritt KA, Thompson RJ, Keith BR, Johndrow DA, Murphy LB. Screening for behavioral and emotional problems in primary care pediatrics. *J Dev Behav Pediatr*. 1993;14(5):340–343
 27. Chang G, Warner V, Weissman M. Physicians' recognition of psychiatric disorders in children and adolescents. *Am J Dis Child*. 1988;142(7):736–739
 28. Costello EJ, Edelbrock GS. Detection of psychiatric disorders in pediatric primary care: a preliminary report. *J Am Acad Child Psychiatry*. 1985;24(6):771–774
 29. Kelleher KJ, Childs GE, Wasserman RC, McInerney TK, Nutting PA, Gardner WP. Insurance status and recognition of psychosocial problems: a report from the Pediatric Research in Office Settings and the Ambulatory Sentinel Practice Networks. *Arch Pediatr Adolesc Med*. 1997;151(11):1109–1115
 30. Leaf PJ, Owens PL, Leventhal JM, et al. Pediatricians' training and identification and management of psychosocial problems. *Clin Pediatr (Phila)*. 2004;43(4):355–365
 31. Murphy MJ, Arnett HL, Bishop SJ, Jellinek M, Reede JY. Screening for psychosocial dysfunction in pediatric practice: a naturalistic study of the Pediatric Symptom Checklist. *Clin Pediatr (Phila)*. 1992;31:660–667

THE VINES HAVE IT: *Each spring and fall, I clear the grape vines from the trees that edge our property. If I don't, the vines proliferate wildly and overwhelm the tree. Eventually the weight of the vines pulls down branches leading to ill health and even the death of some trees. The problem of robust vine growth does not seem confined to northern Vermont. In the Central and South American rain forests, woody vines appear to be winning their clash with the trees. As reported in The New York Times (Science: May 23, 2011), woody vines, such as liana, are increasing in tropical forests. Lianas are long-stemmed woody vines rooted in the soil that use trees to support their growth toward the sunlight in the forest canopy. Once the vine reaches the top, it produces a cornucopia of leaves. In several areas that have been monitored for the past few decades, the tops of three-fourths of all trees with trunks greater than eight inches in diameter are infested with lianas. While the vines may form bridges between trees that make it easier for arboreal animals to move about the forest, they compete very successfully with the trees for water, soil nutrients, and sunlight. Why the vines are so successful is not known but it may be that they are better able than trees to utilize atmospheric carbon dioxide and also to extract water from the soil, particularly in the dry season. Trees infested with liana have stunted growth and higher mortality rates. The sheer mass of the vines can also lead to mechanical failure and death of the tree. The long term effects of liana growth are worrisome. Trees are essential for storing terrestrial carbon. Because lianas store less carbon than trees a reduction in tree mass will result in a net reduction in carbon storage. Carbon storage is essential for regulating atmospheric carbon dioxide. Some experts suggest that the liana proliferation has led to a 10 percent net decrease in carbon storage and could have implications for global warming. As for me, while my tree line is not like a tropical forest, removing the vines is hard work. Still, I like my trees and their role in the environment so I will continue to battle the vines twice a year.*

Noted by WVR, MD