

Preferential Expression of Biotransformation Enzymes in the Olfactory Organs of *Drosophila melanogaster*, the Antennae*

(Received for publication, October 30, 1998, and in revised form, January 22, 1999)

Qin Wang[‡], Gaiti Hasan[§], and Claudio W. Pikielny^{‡¶}

From the [‡]Department of Neuroscience and Cell Biology, Robert Wood Johnson Medical School/University of Medicine and Dentistry of New Jersey, Piscataway, New Jersey 08854 and [§]National Centre for Biological Sciences, TIFR Centre, IISc Campus, P. O. Box 1234, Bangalore 560012, India

Biotransformation enzymes have been found in the olfactory epithelium of vertebrates. We now show that in *Drosophila melanogaster*, a UDP-glycosyltransferase (UGT), as well as a short chain dehydrogenase/reductase and a cytochrome P450 are expressed specifically or preferentially in the olfactory organs, the antennae. The evolutionarily conserved expression of biotransformation enzymes in olfactory organs suggests that they play an important role in olfaction. In addition, we describe five *Drosophila* UGTs belonging to two families. All five UGTs contain a putative transmembrane domain at their C terminus as is the case for vertebrate UGTs where it is required for enzymatic activity. The primary sequence of the C terminus, including part of the transmembrane domain, differs between the two families but is highly conserved not only within each *Drosophila* family, but also between the members of one of the *Drosophila* families and vertebrate UGTs. The partial overlap of the conserved primary sequence with the transmembrane domain suggests that this part of the protein is involved in specific interactions occurring at the membrane surface. The presence of different C termini in the two *Drosophila* families suggests that they interact with different targets, one of which is conserved between *Drosophila* and vertebrates.

All organisms live in environments that contain potentially harmful chemicals, both natural and man-made. Extensive studies of detoxification in the vertebrate liver provide a framework to the study of detoxification mechanisms in other systems (1–3). Detoxification often occurs in two phases; in phase I, the initial compound is transformed into a more reactive species. A variety of different chemical transformations are involved, including redox reactions catalyzed by enzymes of the cytochrome P450 superfamily (4, 5) and members of the short chain dehydrogenase/reductase (SDR)¹ family (6). Phase II re-

actions consist in the addition, either to a product of a phase I reaction or directly to many toxic chemicals, of a highly polar group such as UDP-glucuronosyl (catalyzed by UDP-glucuronosyltransferases) (2) or glutathione (catalyzed by glutathione S-transferases) (7). Products of phase II reactions are hydrophilic; they can no longer cross membranes and are eliminated by secretion. In addition to the elimination of environmental toxins, phase I and II biotransformation enzymes participate in the removal of toxic side products of normal metabolism (e.g. bile acids), participate in drug clearance, and play an important role in the synthesis of hormones such as prostaglandins and some steroids (3). Finally, the involvement of these enzymes in the production of carcinogens (8), drug clearance, and some hereditary diseases (9) makes an understanding of their function important for human health.

Biotransformation enzymes related to those found in vertebrates have also been found in insects and are likely to play equally important roles. Cytochrome P450s and glutathione S-transferases in particular have been implicated in insect resistance to pesticides (10). UDP-glucuronosyltransferases are part of a superfamily of UDP-glycosyltransferases (UGTs) present in plants, animals, and bacteria (11). These enzymes transfer the sugar moiety of UDP-glucose, UDP-glucuronic acid, UDP-galactose, or UDP-xylose to a variety of hydrophobic substrates (11). Insects contain UGT activities that can use UDP-glucose but not UDP-glucuronic acid as a glycosyl donor (12–14). Although no molecular information on any insect UGT was available until this work, baculoviruses infecting several species of moths have been shown to encode ecdysteroid UDP-glycosyltransferases (15, 16). These viral enzymes specifically inactivate ecdysteroids, the molting hormones of the infected hosts, and thus prolong the larval stage permissive to viral replication. ecdysteroid UDP-glycosyltransferases lack a C-terminal transmembrane domain and are secreted in the hemolymph where ecdysone is present (15).

Here we report that, in *Drosophila*, several phase I and II biotransformation enzymes are expressed preferentially in the olfactory organs, the antennae. This observation is reminiscent of the preferential or exclusive expression of a cytochrome P450, UGT, and glutathione S-transferase in the vertebrate olfactory epithelium (17–20). The presence of these enzymes in the olfactory organs of such evolutionarily distant organisms supports the notion that they play an important role in olfaction. In addition, the availability of the first sequences of UGTs from insects sheds light on the structure and function of the C-terminal domain of vertebrate UGTs.

EXPERIMENTAL PROCEDURES

Generation of an Appendage cDNA Library, Cloning, and Sequencing—Partial cDNA clones for *AntP450*, *AntDH*, and *DmeUgt35b* were initially found through random sequencing of clones in an antennae-minus-heads subtracted cDNA library that was described previously (21). All cDNA sequences discussed in this paper were obtained from

* This research was supported in part by a Shannon award from the NICHD, National Institutes of Health and a small grant from the Foundation for the University of Medicine and Dentistry of New Jersey (to C. P.) and a Biotechnology Career Fellowship from the Rockefeller Foundation (to G. H.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

The nucleotide sequence(s) reported in this paper has been submitted to the GenBank™/EBI Data Bank with accession number(s) AF116553, AF116555, AF116554.

¶ To whom correspondence should be addressed: Dept. of Neuroscience and Cell Biology, Robert Wood Johnson Medical School/UMDNJ, 675 Hoes Lane, Piscataway, NJ 08854. Tel.: 732-235-5658; Fax: 732-235-4029; E-mail: pikielc@umdnj.edu.

¹ The abbreviations used are: SDR, short chain dehydrogenase/reductase; UGT, UDP-glycosyltransferase; ORF, open reading frame; EST, expressed sequence tag.

full-length cDNA clones isolated by using the partial clones as probes to screen an appendage cDNA library in Lambda-ZAP (Stratagene). Appendage RNA was generated from poly(A)⁺ RNA isolated from an appendage fraction (see below and Ref. 22).

Analysis of Gene Expression—Total RNA for Northern blots was isolated either from hand-dissected antennae or legs or from mass-produced body fractions generated as follows (22). Frozen flies are vortexed, and the resulting body parts are then sieved to yield three fractions: appendages (antennae, legs, and wings), heads (without antennae), and bodies (abdomen and thorax, decapitated and without legs or wings). Because all the proteins under study belong to multigene families, the probes used were first tested on Southern blots under identical conditions to ensure that there was no detectable cross-reactivity to related genes (not shown). The probes used are indicated in the legend to Fig. 1.

Sequence Analysis—Sequences were assembled and analyzed using Wisconsin Package Version 9.1, Genetics Computer Group, Madison, WI. Data base searches were performed using BLAST (23) both on the Berkeley *Drosophila* Genome Project and National Center for Biotechnology Information www servers (<http://www.fruitfly.org/blast/> and <http://www.ncbi.nlm.nih.gov/cgi-bin/BLAST/nph-newblast?Jform=0>, respectively).

The central portion of the *DmeUgt35a* cDNA is represented by nucleotides number 1–324 of P1 clone DS07339 that has been mapped by the BDGP to 86D5–10. Similarly, the 3' end of *DmeUgt35b* is partially represented in preliminary sequence (nucleotides 1–326) of DS08785 mapped to 86D1–D2. *AntP450* is a partial cDNA sequence identical to EST number GH06928 and with significant sequence similarity to the C terminus of the *cyp6* family of cytochrome P450s (24).

The three members of the *DmeUgt37* family result from conceptual translation of sequences from the *Drosophila* Genome Project. The sequences encoding the three open reading frames (ORFs) are present at nucleotides number 68193 to 66570 (reverse strand) of P1 clone DS51087 for *DmeUgt37a1*; positions 22203–22207 of DS00108 for *DmeUgt37b1* and nucleotides 111614 to 113188 of DS07321 for *DmeUgt37c1*. To generate the *DmeUgt37a1* ORF, we removed a likely intron at positions 67516–67464. In the case of *DmeUgt37b1**, deletion of a single T in a stretch of 5 Ts at positions 22203–22207 results in the creation of an ORF with high similarity to the other *Drosophila* UGT37 ORFs (see text). The frameshift in the database sequence could be caused by a sequencing error or a recent mutation resulting in a pseudogene. In either case, conceptual translation of the “corrected” *DmeUgt37b1** sequence represents a UGT with high similarity to the other UGT37 proteins throughout its open reading frame and therefore likely represents a real UGT, even if it no longer exists in present day laboratory canton S strains. For the purpose of this publication we will keep the asterisk to denote the ambiguity. Note that the inclusion of the *DmeUgt37b1** ORF is not necessary to reach the conclusions about the domain structure of *Drosophila* UGTs that are discussed in the text. The sequences of the three novel cDNAs discussed here have been deposited in the GenBank™ data base and their accession numbers are as follows: *AntDH*, AF116553; *DmeUgt35a*, AF116555; and *DmeUgt35b*, AF116554. Non-*Drosophila* UGTs are designated according to the names given by the UGT Nomenclature Committee and accession numbers are given in the figures.

RESULTS

Several Biodegradation Enzymes Are Expressed Preferentially in the Antennae of *Drosophila*—We have previously described a subtracted cDNA library (antennae-minus-heads) enriched in cDNAs expressed specifically or preferentially in the antennae of *Drosophila melanogaster*. Analysis of a number of those cDNAs led to the discovery of several putative odorant-binding proteins with distinct expression patterns on the surface of the antennae, suggesting a role for odorant-binding proteins in olfactory discrimination (21). Here we report that, in addition to odorant-binding proteins, sampling of our library has led to the discovery of cDNAs coding for a cytochrome P450 (*AntP450*, see “Experimental Procedures”), a UGT (*DmeUgt35b*, see below for explanation of the nomenclature), and a short chain dehydrogenase/reductase (*AntDH*). The proteins encoded by these three cDNAs are related to vertebrate enzymes involved in detoxification. In addition to *DmeUgt35b* isolated from our subtractive library, a second UGT-encoding cDNA (*DmeUgt35a*) was isolated from an appendage cDNA




FIG. 1. Several novel biodegradation enzymes are preferentially expressed in *Drosophila* antennae. Northern blots were performed with RNA extracted from different parts of the fly (see “Experimental Procedures”) as indicated above each lane. *Appendages*: legs, third antennal segments, and wings; *bodies*: decapitated bodies without legs or wings; *heads*: heads without third antennal segments. RNA from third antennal segments and legs was obtained after manual dissection. To ensure that each signal corresponds to expression from a single gene, ³²P probes were generated from relatively nonconserved regions of each gene that give rise to a single band on genomic Southern blots under identical hybridization conditions (data not shown). Expression of the ubiquitously expressed *rp49* gene (42) was monitored in all fractions as a loading control. 1 μg of total RNA was loaded in each lane. Probes used for the two UGTs were 5' cDNA fragments of 560 base pairs (*EcoRI-NheI*) and 610 base pairs (*EcoRI-NruI*) for *DmeUgt35a* and *DmeUgt35b*, respectively. The probe used for *AntDH* was the full-length cDNA clone and that for *AntP450* was the partial cDNA clone obtained in the subtracted library.

library by cross-hybridization to *DmeUgt35b* (see below).

Analysis of expression patterns was performed by Northern blots using probes specific for *DmeUgt35a*, *DmeUgt35b*, *AntDH*, and *AntP450*. In every case the probes were generated from sequences that show little similarity with other genes of the same family and the lack of cross-hybridization was verified on Southern blots performed under identical hybridization conditions (data not shown). The Northern blot shown in Fig. 1 analyzes RNAs prepared from two types of samples as indicated above the lanes. First, fly parts were separated into three fractions (see “Experimental Procedures”): appendages (third antennal segments, legs, and wings), heads (without third antennal segments), and bodies (decapitated and without legs). Second, to differentiate between different appendages, we separately hand-collected third antennal segments and legs from approximately 200 flies. Expression of *AntDH* is restricted to appendages, and within appendages it is much higher in third antennal segments than in legs. *AntP450* and *DmeUgt35b* are more ubiquitous because they are detected in heads and bodies albeit at slightly lower levels than in appendages. Nevertheless, both genes are expressed at highest levels in the third antennal segment. In contrast to these three genes, *DmeUgt35a* shows lower expression in the third antennal segment than in legs. Although most sensilla involved in taste can be found on the legs, head, and wings of *Drosophila*, the third antennal segment contains the vast majority of olfactory sensilla (25). The preferential expression of *AntDH*, *DmeUgt35b*, and *AntP450* in the third antennal segment is therefore suggestive of a role in olfaction.

To further delineate expression patterns we used *in situ* hybridization on cryosections of heads and antennae (Fig. 2). Of the genes discussed here, only *AntDH* expression was detected




FIG. 2. *AntDH* is specifically expressed in the third antennal segment. Horizontal cryosections of heads were hybridized with digoxigenin-labeled DNA probes that were visualized using standard experimental procedures with anti-digoxigenin antibodies conjugated to alkaline phosphatase (21). In the presence of a chromogenic substrate, a blue/purple precipitate is formed. The section shown is at the level of the third antennal segment and is typical of many others. In no case was signal observed in other parts of the head or in the second antennal segment (data not shown).

by this method, most likely because of its higher expression levels (data not shown). Consistent with the Northern blot analysis, *in situ* hybridization to *AntDH* mRNA is restricted to the third antennal segment; no expression is detected in the head (Fig. 2) or in the second antennal segment (not shown). This observation further supports a role for *AntDH* in olfaction, because the second antennal segment does not contain chemosensory hairs (25). Within the third antennal segment however, *AntDH* expression appears uniformly distributed (Fig. 2 and data not shown), in contrast with several odorant-binding proteins, each of which is restricted to a single morphological type of sensillum with a nonuniform distribution on the antennal surface (21, 26, 27).²

AntDH Is a Short Chain Dehydrogenase/Reductase Specifically Expressed in Third Antennal Segments—The initial *AntDH* cDNA clone was used as a probe to isolate a full-length cDNA clone from an appendage cDNA library (see “Experimental Procedures”). A single ORF has sequence similarity to the members of a large family of SDRs (Fig. 3) (28) found in organisms ranging from prokaryotes and plants to humans (28). The overall degree of sequence identity is relatively low as is typical in this family of proteins (the closest sequence we have found is aldehyde reductase from the bacterium *Streptomyces clavuligerus*, Cla9_Sc in Fig. 3, which has 36% identity with *AntDH*). However, *AntDH* has all the residues that have been demonstrated to be important for the function of SDRs (Fig. 3). In particular, the GlyXXXGlyXGly motif close to the amino terminus corresponds to a coenzyme binding pocket for either NAD or NADP. In addition, the TyrXXXLys motif necessary for catalysis can be found at positions 164 through 168 and Ser¹⁴⁴ is the likely homologue of the essential Ser¹³⁹ of alcohol dehydrogenase (29). Although most SDRs are cytoplasmic, some

members of this family are microsomal or even extracellular (30). Contrary to the case of the membrane-associated mouse corticosteroid 11- β -dehydrogenase (dhi1_mouse in Fig. 3) there is no apparent amino-terminal signal sequence in the *AntDH* ORF. Because the short sequence preceding the apparent translational start in our *AntDH* clone does not contain any stop codon, we cannot entirely rule out the possibility that we are missing some 5' sequences that code for a signal peptide. However, the presence of an AUG at the almost identical position as it is found in many cytoplasmic SDRs (Fig. 3) suggests that we have identified the correct amino terminus and that *AntDH* is a cytoplasmic protein.

Identification of Nine Putative Drosophila UGTs—When probing the appendage library with our partial *Ugt* clone we found two classes of clones that hybridize at different intensities. Southern blotting and sequence analysis shows that these phages correspond to two different cDNAs each encoded by a separate gene, which we will call *DmeUgt35a* and *-b* (see the last paragraph under “Results” for a justification of this nomenclature). We have mapped both sequences by *in situ* hybridization to cytogenetic locations 86C-D in the *Drosophila* genome, suggesting that these two genes have their origin in a relatively recent duplication. More recently, the Berkeley and European *Drosophila* genome projects have sequenced parts of both *DmeUgt35* genes, refining the mapping to 86D5-10 and 86D1-2 for the *a* and *b* genes, respectively.

We have also found that several other likely *Ugt* sequences are present in the *Drosophila* genome project data. These include three genomic DNA sequences containing full-length ORFs that define three members of a second family of *Drosophila* UGTs, UGT37: *DmeUgt37a1*, *-b1* and *-c1* (see below for an explanation of the nomenclature). Five other likely *Ugt* genes are represented by partial cDNA sequences or ESTs (EST numbers GH06505, GH09393, GM04645, LD25345, and LD15335). The first four are 5' sequences coding for NH₂ termini (Fig. 4A), whereas the last one is a 3' sequence coding for a C terminus (Fig. 5). In all, we describe nine or ten putative *Drosophila* UGTs (because each of the above ESTs has only been sequenced from one end, LD15335 may be identical to one of the other clones). The two *DmeUgt35* cDNAs as well as the three genomic *DmeUgt37* sequences appear to represent complete ORFs because they begin with ATG codons and end with stop codons at positions that match closely those expected for this family of genes (see “Experimental Procedures” for further discussion of DNA sequence analysis). In contrast to the use of alternative splicing for the generation of diversity, as is the case of the human *UGT1A1* gene (2), we have found no evidence of alternative mRNA splicing, and an intron is present in only one of the three genomic sequences (see “Experimental Procedures”).

Some of the highest similarity between the five complete ORFs is found near a sequence present in all known UGTs and defined by the string: [FVA]-[LIVMF]-[TS]-[HQ]-[SGAC]-G-X(2)-[STG]-X(2)-[DE]-X(6)-P-[LIVMFA]-[LIVMFA]-X(2)-P-[LMVFIQ]-X(2)-[DE]-Q, in which all amino acids that can occur at a given position are listed inside brackets (11). The presence of this sequence strongly supports the identification of these five proteins as UGTs (Fig. 4B). In addition, the five complete ORFs contain C-terminal hydrophobic domains followed by several basic residues (see below and Fig. 5). In the case of vertebrate UGTs, similar sequences have been identified as a transmembrane domain and a positively charged “stop-transfer” sequence that in combination are responsible for the anchoring of the protein to the endoplasmic reticulum membrane (1) and are necessary for enzymatic activity (31). Baculovirus ecysteroid UDP-glucosyltransferases, which are solu-

² S.-K. Park, S. Shanbhag, A. Dubin, G. Hasan, Q. Wang, P. Yu, G. Harris, A. Steinbrecht, and C. W. Pikielny, manuscript in preparation.




FIG. 3. **AntDH is a member of the short chain dehydrogenase family of proteins.** The sequence of AntDH was aligned with those of representative members of the large family of SDRs using Pileup. Residues identical between at least three of the five sequences are boxed in black and similar residues in gray. The NAD/NADP-binding domain and residues required for activity in other SDRs are indicated by asterisks under the sequences. PksB_Dd, PksB gene product from *Dictyostelium discoideum* (accession number AF019986); Cla9_Sc, Cla9 clavulanate-9-aldehyde reductase from *Streptomyces clavuligerus* (accession number AJ000671); dhi1_mouse, corticosteroid 11- β -dehydrogenase from mouse (accession number P50172); adh_drome, alcohol dehydrogenase from *Drosophila melanogaster* (accession number P00334).

ble and secreted in the hemolymph of the hosts lack such C-terminal transmembrane domains (15).

Finally, at least nine of the ten putative *Drosophila* UGTs display a region of high similarity at their very amino terminus, immediately following the signal peptides (32, 33) (Fig. 4A). In the case of the tenth putative UGT: LD15335 only the C-terminal sequence is presently known. In vertebrate UGTs this region of the molecule is involved in the formation of dimers (34), which may be the active form of the protein. In at least one case a heterodimer has enzymatic activities that differ from those of either homodimer (35), suggesting that the combinatorial association of different subunits into heterodimers may provide added functional diversity.

UGT35a and -b Contain C-terminal Transmembrane Domains Similar to Those of Vertebrate UGTs but Different from Those of the Drosophila UGT37 Protein Family—In consultation with the UGT nomenclature committee (11), we have assigned the two cDNAs we have cloned to a single family, *Ugt35*. The other three full-length sequences found through the genome projects fall into a second family, *DmeUgt37*. Both families fit the commonly accepted criteria for protein families (more than 45% overall identity within a family and less than 45% between different families, data not shown). Finally, although the C-terminal sequence of LD15335 suggests that it is a member of the *DmeUgt37* family (Fig. 5), the definitive assignment of the five UGTs presently only known as ESTs (Fig. 4A) to one of these two families, or yet new ones, will require their complete sequences.

Amino acid residues present in all nine *Drosophila* proteins occur in the first sixty residues (Fig. 4A) as well as in the C-terminal half of the protein, particularly around the signature sequence (Fig. 4B). After the first sixty amino acids, the

amino-terminal halves of the proteins are highly divergent (not shown), as is the case for vertebrate UGTs, perhaps corresponding to different substrate specificities.

Strikingly, although the C-terminal halves of all the *Drosophila* proteins are closely related, there is a strong discontinuity of this similarity at their very C termini. After a highly conserved segment, the sequences of the two families diverge abruptly, each encoding a different C-terminal domain containing putative transmembrane stretches and stop-transfer sequences (Fig. 5). Within each family, however, there is a high degree of sequence conservation. Interestingly, the region of sequence similarity overlaps with the likely transmembrane helix for members of the UGT35 (36) as well as UGT37 families (Fig. 5). Five of the first six amino acids in the putative transmembrane domain are identical between *Drosophila* UGT35b and human UGT1A1, and conservation of similarly located residues is apparent for the UGT37 family. This pattern of conservation suggests specific and different roles for the C-terminal domains of the two *Drosophila* families.

DISCUSSION

The Evolutionarily Conserved Presence of Biodegradation Enzymes Argues for an Important Function in Olfaction—The results presented here suggest that, as in the olfactory epithelium of vertebrates (17–20), several biodegradation enzymes are expressed specifically or at higher levels in the antennae of *Drosophila*. This conserved expression of biodegradation enzymes in olfactory organs argues for an important function in olfaction. Such a role is also consistent with the presence in the antennae of an enzyme involved in cytochrome P450 activation, NADPH P450 oxidoreductase (37).

In a highly specialized case of olfactory behavior, male moths




FIG. 4. Nine putative *Drosophila* UGTs. A, nine *Drosophila* UGTs have putative NH₂-terminal dimerization sequences. The amino-terminal portion of the nine *Drosophila* putative UGTs were included in a multiple sequence alignment with representative UGTs from vertebrates and baculovirus. The organism or group of organisms from which each sequence was obtained is abbreviated on the left. *Dm*, *Drosophila melanogaster*; *Bac*, baculoviruses; *Vert*, vertebrates. An asterisk indicates the location of a leucine residue required for dimerization of the UGT2B1 gene (34). Although the nomenclature for full-length members of the UGT35 and UGT37 protein families is discussed under "Experimental Procedures," four additional *Drosophila* putative UGTs are presently only known by an EST and are indicated according to the name of the cDNA clone within the Berkeley *Drosophila* Genome Project EST database. UGTs from vertebrates and baculoviruses are named as suggested by the UGT Nomenclature Committee (11). UGT31A2 (accession number Q88168), UGT31D3 (accession number P18569), and UGT32 (accession number Q98166) are encoded by different baculoviruses. UGT1A1 (accession number M84125) and UGT2B1 (accession number P09875) are two UDP-glucuronosyltransferases from different vertebrate families; UGT8, UDP-galactosyltransferase from human brain (accession number Q09426). B, five *Drosophila* UGTs contain the "UGT signature sequence." The UGT signature sequence (11) is indicated below a multiple sequence alignment of UGTs from a variety of organisms. The five full-length *Drosophila* UGTs are indicated by arrowheads. UGTs 18E1, 16B1, and 17A1 are from *Caenorhabditis elegans* and 71A1, 77A1, and 78C1 are from plants (see Ref. 11 for accession numbers).

are able to find females located many miles away by rapidly alternating between two types of behavior, upwind flight when inside a pheromone plume and casting from side to side as soon as the pheromone is no longer detected (38). The ability to monitor concentration changes without a lag requires that the half-life of odorants inside the olfactory organs be short relative to the time course of the outside fluctuations. Based on these considerations, researchers have looked for and found enzymes that can specifically metabolize pheromones in the antennae of several species of moths (39, 40). In vertebrate olfaction, a similar role has been attributed to biotransformation enzymes that are better known for their role in detoxification in the liver. The same sequence of events that eliminates toxic chemicals may be involved in odorant degradation, thereby preventing continuing stimulation of olfactory receptors. Consistent with this hypothesis, an olfactory-specific UGT, UGT_{olf}, modifies odorants more efficiently than liver UGTs (18).

A second function for these enzymes might be in the protection of olfactory organs from environmental toxins to which they are, by necessity, preferentially exposed. Although odorant turnover and toxin degradation are not mutually exclusive functions and any given protein may be involved in both, the expression pattern of each gene may suggest the relative con-

tribution to either function. Because detoxification occurs in many organs, proteins whose expression is highly specific to the antennae, such as AntDH and UGT35b, are likely to be involved in odorant turnover. On the other hand, proteins that have more ubiquitous expression patterns, such as UGT35a, may participate primarily in detoxification.

Parallel Conservation of Different Primary Sequences Suggests Different Functions for the C-terminal Domains of the Two Drosophila UGT Families—We report here that the *Drosophila* genome encodes at least nine different UGTs. The five genes for which complete coding sequences are available contain the signature motif characteristic of this superfamily and thus represent the first reported UGTs from any insect other than ecdysteroid UDP-glucosyltransferases from baculoviruses (11). In addition to the overall similarity to UGTs in a variety of organisms, both *Drosophila* and vertebrate proteins have at their C terminus a transmembrane domain followed by a stop-transfer sequence composed of several positively charged amino acids (1, 36). Although this domain is absent from viral-encoded ecdysteroid UDP-glucose transferases and some plant UGTs (11), mutations in either the transmembrane domain or the stop-transfer sequence of vertebrate UGT2B1 eliminate and reduce its activity, respectively (31).




FIG. 5. *Drosophila* UGT35a and -b have C-terminal domains similar to those of vertebrate UGTs but different from those of *Drosophila* UGT37a1, -b1 and -c1. Vertebrate UGTs (UGT1A1, UGT2A1, and UGT8) can be aligned with all five *Drosophila* UGTs (indicated by arrowheads) up to a point where the sequences of the two families diverge dramatically. The three alignments were generated independently using Pileup, but the sequences shown are contiguous and their order within the alignments has been preserved and does not correspond to similarity scores. Boxed residues indicate identities in five of eight, three of five, and two of three for the left, top, and bottom alignments, respectively. The transmembrane domain indicated for members of the UGT35 family and its vertebrate relatives is the one proposed for vertebrate UGT2B1 (36). Hydrophathy calculations for members of both *Drosophila* families are consistent with the prediction that the transmembrane domains start immediately after the LD sequence (data not shown). Hydrophathy was calculated by the Kyte and Doolittle method using the peptides structure program of Genetics Computer Group. See the legend to Fig. 4 for UGT nomenclature and accession numbers.

More surprising, however, is the high degree of primary sequence identity between the *Drosophila* members of the UGT35 family and vertebrate sequences in a stretch immediately NH₂-terminal to and partially overlapping with the putative transmembrane domain (61 and 70% identity to the human UGT1A1 gene over a 31 amino acid stretch for the *a* and *b* genes, respectively). In addition, although the sequences of members of the UGT37 family are very different from those of the UGT35 family in this region, they are also highly conserved within this second family (Fig. 5). This parallel conservation of primary sequences suggests that the C-terminal domains of UGTs are involved in specific interactions at the membrane surface that differ between UGT35a, -b and the vertebrate enzymes on one hand and the members of the UGT37 family on the other. Despite these differences, the domains of the two classes of proteins have some shared features. In all cases except for UGT37b1*, the two amino acids at positions 15 and 16 after the start of sequence divergence are LD, which are immediately followed by a series of hydrophobic residues likely to be part of the transmembrane domain (Ref. 36 and Fig. 5). In the case of UGT37b1*, a three amino acid insertion moves LD to positions 18 and 19 and the putative transmembrane domain starts at position 20. These similarities suggest that despite the divergent sequences the two different types of C-terminal domains have similar secondary structures and may therefore interact with related proteins.

What is the function of these alternative C-terminal domains? The primary sequence conservation within each family suggests it may be involved in an interaction with another protein that occurs at least in part within the membrane. The enzymatic reactions catalyzed by UGTs take place in the lumen of the endoplasmic reticulum and are therefore dependent on specific transporters that allow entry of nucleotides into this subcellular compartment (41). One intriguing possibility is that the C-terminal domain of UGTs participates in interactions

with specific transporters, perhaps corresponding to the specificities of these enzymes for different glycosyl donors. However, because permeabilization of membranes with detergent does not restore activity to proteins with mutations in the C-terminal domain (31), substrate transport cannot be its only function.

Although scans of the existing data bases have not revealed any UGT from any organism with a C-terminal domain similar to that of the UGT37 family, the ongoing sequence of the human genome may yet uncover such genes. Alternatively, if this domain arose after the divergence of the ancestors of insects and vertebrates, it may constitute an insect-specific domain and therefore a possible target for rational pesticide design.

The availability of the genes coding for all these enzymes in *Drosophila* should allow the test of their involvement in olfaction as well as a dissection of the function of the UGT C-terminal domains using both biochemical and reverse genetic approaches.

Acknowledgments—We thank Yashi Ahmed and Ren-Ping Zhou for helpful comments on the manuscript and express gratitude also to Michael Rosbash in whose laboratory this work was initiated.

REFERENCES

1. Tephly, T. R., and Burchell, B. (1990) *Trends Pharmacol. Sci.* **11**, 276–279
2. Brierley, C. H., and Burchell, B. (1993) *Bioessays* **15**, 749–754
3. Meyer, U. A. (1996) *J. Pharmacokinet. Biopharm.* **24**, 449–459
4. Nebert, D. W., and Gonzalez, F. J. (1987) *Annu. Rev. Biochem.* **56**, 945–993
5. Guengerich, F. P. (1993) *Sci. Am.* **81**, 440–447
6. Maser, E., and Opperman, U. C. T. (1997) *Eur. J. Biochem.* **249**, 365–369
7. Shen, H., Kauvar, L., and Tew, K. D. (1997) *Oncol. Res.* **9**, 295–302
8. Nebert, D. W., McKinnon, R. A., and Puga, A. (1996) *DNA Cell Biol.* **15**, 273–280
9. Owens, I. S., Ritter, J. K., Yeatman, M. T., and Chen, F. (1996) *J. Pharmacokinet. Biopharm.* **24**, 491–508
10. Feyereisen, R. (1995) *Toxicol. Lett. (Shannon)* **82**, 83–90
11. Mackenzie, P. I., Owens, I. S., Burchell, B., Bock, K. W., Bairoch, A., Belanger, A., Fournet-Gigleux, S., Green, M., Hum, D. W., Iyanagi, T., and Lancet, D. (1997) *Pharmacogenetics* **7**, 255–269
12. Dutton, G. J. (1962) *Comp. Biochem. Physiol.* **7**, 39–46
13. Ahmad, S. A., and Hopkins, T. L. (1992) *Arch. Insect Biochem. Physiol.* **21**,

- 207–224
14. Rausel, C., Llorca, J., and Real, M. D. (1997) *Arch. Insect Biochem. Physiol.* **34**, 347–358
15. O'Reilly, D. R., and Miller, L. K. (1989) *Science* **245**, 1110–1112
16. Hu, Z. H., Broer, R., Westerlaken, J., Martens, J. W. M., Jin, F., Jehle, J. A., Wang, L. M., and Vlak, J. M. (1997) *Virus Res.* **47**, 91–97
17. Lazard, D., Tal, N., Rubinstein, M., Khen, M., Lancet, D., and Zupko, K. (1990) *Biochemistry* **29**, 7433–7440
18. Lazard, D., Zupko, K., Poria, Y., Nef, P., Lazarovits, J., Horn, S., Khen, M., and Lancet, D. (1991) *Nature* **349**, 790–793
19. Dear, T. N., Campbell, K., and Rabbitts, T. H. (1991) *Biochemistry* **30**, 10376–10382
20. Ben-Arie, N., Khen, M., and Lancet, D. (1993) *Biochem. J.* **292**, 379–384
21. Pikielny, C. W., Hasan, G., Rouyer, F., and Rosbash, M. (1994) *Neuron* **12**, 39–45
22. Oliver, D. V., and Philips, J. P. (1970) *Drosophila Information Service* **45**, 58
23. Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. (1997) *Nucleic Acids Res.* **25**, 3389–3402
24. Cohen, M. B., Koener, J. F., and Feyereisen, R. (1994) *Gene* **146**, 267–272
25. Stocker, R. F. (1994) *Cell Tissue Res.* **275**, 3–26
26. McKenna, M. P., Hekmat-Scafe, D. S., Gaines, P., and Carlson, J. R. (1994) *J. Biol. Chem.* **269**, 16340–16347
27. Hekmat-Scafe, D., Steinbrecht, R. A., and Carlson, J. (1997) *J. Neurosci.* **17**, 1616–1624
28. Jornvall, H., Persson, B., Krook, M., Atrian, S., Gonzalez-Duarte, R., Jeffery, J., and Ghosh, D. (1995) *Biochemistry* **34**, 6003–6013
29. Cols, N., Atrian, S., Benach, J., Ladenstein, R., and Gonzalez-Duarte, R. (1997) *FEBS Lett.* **413**, 191–193
30. Lee, B.-K., Lee, K., Mendez, J., and Shimkets, L. J. (1995) *Genes Dev.* **9**, 2964–2973
31. Meech, R., Yogalingam, G., and Mackenzie, P. I. (1996) *DNA Cell Biol.* **15**, 489–494
32. von Heijne, G. (1982) *J. Mol. Biol.* **159**, 537–541
33. von Heijne, G. (1985) *J. Mol. Biol.* **184**, 99–105
34. Meech, R., and MacKenzie, P. I. (1997) *J. Biol. Chem.* **272**, 26913–26917
35. Ikushiro, S., Emi, Y., and Iyanagi, T. (1997) *Biochemistry* **36**, 7154–7161
36. Mackenzie, P. I. (1986) *J. Biol. Chem.* **261**, 6119–6125
37. Hovemann, B. T., Sehlmeier, F., and Malz, J. (1997) *Gene* **189**, 213–219
38. Baker, T. C., and Vogt, R. G. (1988) *J. Exp. Biol.* **137**, 29–38
39. Vogt, R. G., Riddiford, L. M., and Prestwich, G. D. (1985) *Proc. Natl. Acad. Sci. U. S. A.* **82**, 8827–8831
40. Rybczynski, R., Vogt, R. G., and Lerner, M. R. (1990) *J. Biol. Chem.* **265**, 19712–19715
41. Bossuyt, X., and Blanckaert, N. (1997) *Biochem. J.* **323**, 645–648
42. O'Connell, O., and Rosbash, M. (1984) *Nucleic Acids Res.* **12**, 5495–5513

**Preferential Expression of Biotransformation Enzymes in the Olfactory Organs of
Drosophila melanogaster, the Antennae**
Qin Wang, Gaiti Hasan and Claudio W. Pikielny

J. Biol. Chem. 1999, 274:10309-10315.

doi: 10.1074/jbc.274.15.10309

Access the most updated version of this article at <http://www.jbc.org/content/274/15/10309>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 42 references, 13 of which can be accessed free at
<http://www.jbc.org/content/274/15/10309.full.html#ref-list-1>