

Ribosome Recycling Factor and Release Factor 3 Action Promotes TnaC-Peptidyl-tRNA Dropoff and Relieves Ribosome Stalling during Tryptophan Induction of *tna* Operon Expression in *Escherichia coli*

Ming Gong, Luis R. Cruz-Vera and Charles Yanofsky
J. Bacteriol. 2007, 189(8):3147. DOI: 10.1128/JB.01868-06.
Published Ahead of Print 9 February 2007.

Updated information and services can be found at:
<http://jb.asm.org/content/189/8/3147>

REFERENCES

These include:

This article cites 32 articles, 19 of which can be accessed free
at: <http://jb.asm.org/content/189/8/3147#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Ribosome Recycling Factor and Release Factor 3 Action Promotes TnaC-Peptidyl-tRNA Dropoff and Relieves Ribosome Stalling during Tryptophan Induction of *tna* Operon Expression in *Escherichia coli*[∇]

Ming Gong,[†] Luis R. Cruz-Vera, and Charles Yanofsky*

Department of Biological Sciences, Stanford University, Stanford, California

Received 12 December 2006/Accepted 31 January 2007

Upon tryptophan induction of *tna* operon expression in *Escherichia coli*, the leader peptidyl-tRNA, TnaC-tRNA₂^{Pro}, resists cleavage, resulting in ribosome stalling at the *tnaC* stop codon. This stalled ribosome blocks Rho factor binding and action, preventing transcription termination in the *tna* operon's leader region. Plasmid-mediated overexpression of *tnaC* was previously shown to inhibit cell growth by reducing uncharged tRNA₂^{Pro} availability. Which factors relieve ribosome stalling, facilitate TnaC-tRNA₂^{Pro} cleavage, and relieve growth inhibition were addressed in the current study. In strains containing the chromosomal *tna* operon and lacking a *tnaC* plasmid, the overproduction of ribosome recycling factor (RRF) and release factor 3 (RF3) reduced *tna* operon expression. Their overproduction in vivo also increased the rate of cleavage of TnaC-tRNA₂^{Pro}, relieving the growth inhibition associated with plasmid-mediated *tnaC* overexpression. The overproduction of elongation factor G or initiation factor 3 did not have comparable effects, and tmRNA was incapable of attacking TnaC-tRNA₂^{Pro} in stalled ribosome complexes. The stability of TnaC-tRNA₂^{Pro} was increased appreciably in strains deficient in RRF and RF3 or deficient in peptidyl-tRNA hydrolase. These findings reveal the existence of a natural mechanism whereby an amino acid, tryptophan, binds to ribosomes that have just completed the synthesis of TnaC-tRNA₂^{Pro}. Bound tryptophan inhibits RF2-mediated cleavage of TnaC-tRNA₂^{Pro}, resulting in the stalling of the ribosome translating *tnaC* mRNA. This stalling results in increased transcription of the structural genes of the *tna* operon. RRF and RF3 then bind to this stalled ribosome complex and slowly release TnaC-tRNA₂^{Pro}. This release allows ribosome recycling and permits the cleavage of TnaC-tRNA₂^{Pro} by peptidyl-tRNA hydrolase.

Expression of the tryptophanase (*tna*) operon of *Escherichia coli* is regulated by catabolite repression and tryptophan-induced transcription antitermination. Induction by tryptophan requires the translation of a 24-residue peptide coding region, *tnaC*, located in the 319-bp transcribed leader region. This region precedes *tnaA*, the structural gene for tryptophanase. Both Trp12 and Pro24 of TnaC have been shown to be essential for induction (5, 8). The key feature of this antitermination mechanism is tryptophan's ability to bind to ribosomes containing TnaC-peptidyl-tRNA; the bound tryptophan inhibits the cleavage of this peptidyl-tRNA. This inhibition results in the stalling of the ribosome translating *tnaC* mRNA (6). The stalled ribosome blocks the *rut* binding site in *tna* leader RNA, preventing Rho factor from binding and terminating transcription (7). Plasmid-mediated overexpression of *tnaC* results in growth inhibition (5). This inhibition was shown to be primarily a consequence of the reduced availability of tRNA₂^{Pro} for general protein synthesis (9). This deficiency was caused by the accumulation of uncleaved TnaC-tRNA₂^{Pro} within stalled, translating ribosomes (9). Growth inhibition was relieved by overproducing tRNA₂^{Pro} (9).

During the biosynthesis of polypeptides in bacteria, amino acid assembly is terminated when the mRNA stop codon of a coding region is encountered in the A site of the 70S ribosomal

complex. The nascent polypeptide chain is then cleaved from its associated tRNA in a factor-dependent manner. The translating ribosome with the deacylated tRNA initially remains bound to mRNA and is subsequently separated by the action of accessory translation factors (13, 15). Six different factors are known to participate in the translation termination process: (i) release factors RF1 and RF2 (these factors are responsible for the recognition of the three stop codons and for activating the cleavage of each peptidyl-tRNA) (26), (ii) release factor RF3 (this factor increases the rate of dissociation of RF1 and RF2 from the ribosome following polypeptide release) (4, 32, 33), (iii) ribosome recycling factor (RRF) (this factor is responsible for ribosome separation from a transcript, and it releases the mRNA and tRNA from the ribosomal complex) (14), (iv) elongation factor G (EF-G) (this factor and RRF are believed to be jointly responsible for recycling bacterial ribosomes following the termination of polypeptide synthesis) (13), and (v) translation initiation factor IF3 (this factor prevents the reassociation of the dissociated ribosomal subunits by binding to the transiently formed 30S subunit during or following the action of RRF and EF-G) (13, 16, 27).

In addition to the normal mechanism of translation termination, there are two auxiliary mechanisms that bacteria use to hydrolyze a peptidyl-tRNA: the action of peptidyl-tRNA hydrolase (Pth) (19) and the participation of tmRNA (17). An uncleaved peptidyl-tRNA may be released from a stalled ribosome by a phenomenon called peptidyl-tRNA dropoff (21). The peptidyl-tRNA liberated is hydrolyzed by Pth, separating the peptide from its tRNA (19). It has also been reported that

* Corresponding author. Mailing address: Department of Biological Sciences, Stanford University, Stanford, CA 94305. Phone: (650) 725-1835. Fax: (650) 725-8221. E-mail: yanofsky@cmgm.stanford.edu.

[†] Present address: Department of Biology and Biochemistry, UCLA Medical School, Los Angeles, CA 90095.

[∇] Published ahead of print on 9 February 2007.

TABLE 1. Bacterial strains and plasmids

Strain or plasmid	Relevant genotype or description	Reference or source
Strains		
SVS1144	W3110 <i>bglR551</i> $\Delta(lac-argF)U169$ (λ _{tna_p} <i>tnaA-lacZ</i>)	30
SVS1144(<i>prfC</i> Δ 2: <i>kan</i>)	<i>prfC</i> gene deleted	30
SVS1144[<i>frr</i> (Ts)]	<i>frr</i> (Ts)	Lab collection
SVS1144[<i>frr</i> (Ts) <i>prfC</i> Δ 2: <i>kan</i>]	<i>frr</i> (Ts); <i>prfC</i> gene deleted	This study
C600	Wild type	3
C600[<i>pth</i> (Ts)]	<i>pth</i> (Ts)	3
Plasmids		
pUC18	ColE1 vector; Ap ^r	Lab collection
pACYC184	p15E replicon vector; Tet ^r Cm ^r	Lab collection
PTnaC	pUC18 derivative; <i>tna_p</i> <i>tnaC-rpoBC</i> Ap ^r	9
PTnaC-RRF	Derived from pTnaC, contains the <i>frr</i> gene with its own promoter and terminator; Ap ^r	This study
pUC-IF3	Contains the <i>infC</i> gene with its promoter and initiation codon AUU; Ap ^r	27
pKW1	Derived from pACYC184; Cm ^r gene deleted, Tet ^r	25
pKW24	Derived from pKW1, contains the tmRNA-His ₆ mutant gene; Tet ^r	25
PHSG299	Derived from pUC18; Ap ^r gene deleted, Km ^r	14
pRR1	Derived from PHSG299; contains the <i>frr</i> gene with its promoter and terminator; Km ^r	14
pIQ-RF3	Derived from pACYC 184, contains <i>lacI</i> ^q and <i>plac-prfC</i> ; Cm ^r	30
pEF-G	Derived from pACYC184, contains the <i>fusA</i> gene; Cm ^r	K. Ito and Y. Nakamura
PACD-IF3	Derived from pACYC184, contains the <i>infC</i> gene with its promoter and initiation codon AUU; Tet ^r	U. Varshney

nascent peptide chains encoded by truncated mRNAs containing contiguous rare codons, or containing an inefficient stop codon, are targeted and tagged by the action of tmRNA (17, 24, 25). When *trans*-translation is completed, the unfinished peptide or polypeptide will have an 11-amino-acid-residue tag (AANDENYALAA) added at its carboxy-terminal end. This tag and its associated peptide or polypeptide are subjected to rapid degradation by various cellular proteases (10, 17). This process allows a translating ribosome to be recycled so that it may participate in a new round of protein synthesis (23).

In this study, various plasmids were used to overproduce the different translation factors, and *in vivo* levels of TnaC-tRNA₂^{Pro} and free tRNA₂^{Pro} were continuously monitored. The assay procedure used to measure the levels of TnaC-tRNA₂^{Pro} was developed previously (9). Our data indicate that when *tnaC* is overexpressed in the presence of tryptophan, (i) RF2 action is prevented, (ii) RRF and RF3 cause TnaC-tRNA₂^{Pro} dropoff, (iii) the released TnaC-tRNA₂^{Pro} is cleaved by Pth, (iv) RRF and RF3 promote ribosome recycling, and (v) RRF, RF3, and Pth action relieves the growth inhibition caused by tRNA₂^{Pro} depletion. tmRNA did not act on tryptophan-induced stalled ribosome complexes containing TnaC-tRNA₂^{Pro}. Under normal induction conditions, in strains lacking a *tnaC* plasmid, the extent of TnaC-tRNA₂^{Pro} accumulation is insufficient to cause growth inhibition. However, RRF and RF3 appear to play the same role, promoting TnaC-tRNA₂^{Pro} dropoff and ribosome recycling. This allows the released tRNA₂^{Pro} and freed ribosome to participate in new rounds of protein synthesis.

MATERIALS AND METHODS

Bacterial strains and plasmids. All the strains and plasmids used in this study are listed in Table 1. The parental *E. coli* strain SVS1144, bearing a *tnaA'*-*lacZ* translational fusion (30), and strains C600 and a C600 derivative with a temperature-sensitive Pth, C600[*pth*(Ts)], have been described previously (3). Strain SVS1144[*frr*(Ts) *prfC* Δ 2:*kan*] was prepared by P1 transduction.

Plasmid pTnaC-RRF was constructed by cloning an EcoRI-HindIII DNA fragment containing the *frr* gene with its own promoter, obtained from plasmid pRR1, into plasmid pTnaC. Plasmid pRR1 was kindly provided by Akira Kaji (14). Its *frr*-containing fragment was inserted immediately following the *rpoBC* terminator sequence (see Fig. 2A).

Growth conditions and viability curves. Cultures were grown aerobically with shaking at the temperatures indicated. Strains C600 and C600[*pth*(Ts)] were

grown in Luria-Bertani (LB) medium. Strain SVS1144 and its derivatives were grown in Vogel-Bonner minimal medium (29) supplemented with 0.2% glycerol and 0.05% acid-hydrolyzed casein with or without the addition of 100 μ g/ml L-tryptophan. The following antibiotics were used: ampicillin (100 μ g/ml), tetracycline (25 μ g/ml), kanamycin (50 μ g/ml), and chloramphenicol (30 μ g/ml). Growth rates were determined for cultures grown with or without the appropriate antibiotics at 37°C. Cell growth was monitored by measuring the optical density at 600 nm (OD₆₀₀).

β -Gal assays. β -Galactosidase (β -Gal) assays were performed on cultures grown with shaking at 37°C in minimal medium with or without 100 μ g/ml tryptophan and with or without 10 mM isopropyl- β -D-thiogalactopyranoside (IPTG). β -Gal assays were performed as described previously (22); β -Gal activity is reported in Miller units.

Measurement of levels of TnaC-tRNA₂^{Pro} *in vivo*. Various *E. coli* strains, several bearing the plasmids listed in Table 1, were grown in appropriate media at 30°C, 37°C, or 42°C. Cells were harvested at an OD₆₀₀ of 0.8. The levels of TnaC-tRNA₂^{Pro} in samples taken from each culture were determined by preparing sonic extracts, centrifugation, and analysis of equal amounts of protein in the supernatants by electrophoresis and Northern blotting. The probe used was unique to tRNA₂^{Pro}. The procedures used have been described previously (9). TnaC-tRNA₂^{Pro} and tRNA₂^{Pro} were detected as bands at 26 kDa and 14 kDa, respectively, under the conditions used (9). For quantification, each gel was exposed to films for different lengths of time, the developed films were scanned, and values were compared to ensure the accuracy of the final values (densitometry imager; Bio-Rad). The band densities (densitometry units) were determined using imaging software (Molecular Analysis; Bio-Rad).

RESULTS

Overproduction of both RRF and RF3 *in vivo* decreases *tna* operon expression (TnaC-LacZ production). In *E. coli*, when a translating ribosome encounters one of the three stop codons, release factor RF1 or RF2 mediates peptidyl-tRNA cleavage. RRF and EF-G as well as other factors, e.g., RF3 and IF3, then participate in the ribosome release/recycling process to provide ribosomal subunits for additional protein synthesis (13). Interestingly, the inactivation of *prfC*, the structural gene for RF3, has been reported to increase basal-level expression of the *tna* operon two- to threefold (31). Overexpression of *prfC* had the opposite effect: it reduced both the basal and induced levels of *tna* operon expression (31). On the basis of this knowledge, experiments were performed to determine whether the overproduction of any of the translation factors, alone or in combination, would affect *tna* operon expression. We initially determined β -Gal levels produced by a *lacZ* deletion strain,

TABLE 2. TnaA-LacZ expression by strain SVS1144 containing different combinations of plasmids expressing mutant and nonmutant translational factors^a

Strain	Gene(s) (protein[s]) overexpressed	β-Gal (TnaA-LacZ) activity (Miller units) ^c		+Trp/-Trp ratio
		-Trp	+Trp	
SVS1144		499	8,310	17
SVS1144/pACYC184/pHSG299		430	7,430	17
SVS1144/pACYC184/pUC18		483	7,830	16
SVS1144/pACYC184/pRR1	<i>frr</i> (RRF)	<u>272</u>	<u>5,220</u>	19
SVS1144/pIQ-RF3/pHSG299	<i>prfC</i> (RF3)	457	7,670	17
		224 (IPTG) ^b	4,480 (IPTG)	20
SVS1144/pACYC184/pUC-IF3	<i>infC</i> (IF3)	436	8,170	19
SVS1144/pEF-G/pHSG299	<i>fusA</i> (EF-G)	456	7,900	17
SVS1144/pIQ-RF3/pUC-IF3	<i>prfC</i> and <i>infC</i> (RF3 and IF3)	466	7,860	17
		238 (IPTG)	4,600 (IPTG)	19
SVS1144/pEF-G/pRR1	<i>fusA</i> and <i>frr</i> (EF-G and RRF)	<u>202</u>	<u>5,300</u>	26
SVS1144/pEF-G/pUC-IF3	<i>fusA</i> and <i>infC</i> (EF-G and IF3)	402	7,300	18
SVS1144/pACD-IF3/pRR1	<i>infC</i> and <i>frr</i> (IF3 and RRF)	<u>302</u>	<u>5,090</u>	17
SVS1144/pIQ-RF3/pRR1	<i>prfC</i> and <i>frr</i> (RF3 and RRF)	<u>184</u>	<u>5,430</u>	30
		74 (IPTG)	840 (IPTG)	11
SVS1144/pKW1		525	7,900	15
SVS1144/pKW24	<i>ssrA</i> (tmRNA)	560	8,180	14

^a Cultures were grown in minimal medium plus 0.2% glycerol and 0.05% acid-hydrolyzed casein with or without additions. Added compounds were 100 μg/ml tryptophan, 30 μg/ml chloramphenicol (pIQ-RF3 and pEF-G), 25 μg/ml tetracycline (pACD-IF3, pKW1, and pKW24), 100 μg/ml ampicillin (pUC-IF3), or 50 μg/ml kanamycin (pHSG299 and pRR1).

^b A total of 10 mM IPTG was added for the production of RF3 from pIQ-RF3.

^c β-Gal assays and the averaged results are presented. Repeat experiments gave values that varied approximately 10%. Values reflecting significant changes are underlined.

SVS1144, containing a *tnaA'*-*lacZ* translational fusion (Table 1). This strain was transformed with combinations of two plasmids that either do or do not overproduce individual translation factors (Table 1). Whenever two plasmids were introduced into the same strain, they were compatible with one another. Initially, we observed that the overproduction of either the RRF or the RF3 protein consistently reduced both the basal and induced levels of β-Gal observed by about 25%, compared to levels of control cultures (Table 2). However, when either IF3 or EF-G was overproduced, no detectable effect on the β-Gal level was observed (Table 2). When RRF and RF3 were overproduced in the same cell, the β-Gal levels were reduced even further (Table 2). When RRF was overproduced together with EFG or IF3 or when RF3 was overproduced with IF3, the reduction in the β-Gal level observed was no greater than the reduction observed upon overproducing either RRF or RF alone (Table 2). These results indicate that these other factors, when tested in combination with RRF or RF3, do not enhance their action. Also, the overproduction of EFG and IF3 in the same cell did not affect the β-Gal levels (Table 2). Thus, the overproduction of both RRF and RF3 has the greatest negative effect on *tna* operon expression in vivo, reducing both the basal and induced levels. The overproduction of tmRNA had no observable effect on the β-Gal level. Strains containing the empty vector or the plasmid overexpressing tmRNA had the same β-Gal levels (Table 2).

Growth inhibition caused by plasmid-mediated overexpression of *tnaC* is relieved by overproduction of both RRF and RF3. On the basis of the decrease in TnaA-LacZ synthesis observed upon the overproduction of RRF and/or RF3 in a strain with chromosomal *tnaC* (Table 2), we next determined whether the overproduction of both of these factors could rescue ribosomes stalled as a consequence of *tnaC* overex-

pression. It was shown previously that when *tnaC* was overexpressed from a multicopy plasmid, pTnaC, the availability of free tRNA₂^{Pro} for continued protein synthesis was reduced (9). This was due to the accumulation of uncleaved TnaC-tRNA₂^{Pro} within the ribosomes translating *tnaC* mRNA (9). This inhibition was relieved upon tRNA₂^{Pro} overproduction (9). If the overproduction of RRF and RF3 could promote TnaC-tRNA₂^{Pro} cleavage, their overproduction would be expected to relieve the growth inhibition resulting from TnaC-tRNA₂^{Pro} accumulation (9). To examine this possibility, it was necessary to overproduce TnaC, RRF, and RF3 in the same bacterium. To achieve this objective, the *frr* gene (encoding RRF), driven by its own promoter, *pfrr*, was inserted into plasmid pTnaC immediately following the *rhoBC* terminator sequence (Fig. 1A). After confirming the location and sequence of the DNA insert, the overproduction of RRF in vivo was verified by Western blotting using anti-RRF antibodies. The amount of RRF produced was estimated to be three- to fourfold greater than that produced by the control strain (data not shown). Reverse transcription-PCR analyses also indicated that there was a 2.5-fold increase in *tnaC* mRNA in strains with the *tnaC* plasmid and that the insertion of the *frr* gene and its promoter had no effect on the level of *tnaC* mRNA (data not shown). SVS1144 strains transformed with plasmids (i) pTnaC, (ii) pTnaC and pIQ-RF3, (iii) pTnaC-RRF, and (iv) pTnaC-RRF and pIQ-RF3 were grown in supplemented minimal medium with or without 100 μg/ml tryptophan and with or without 10 mM IPTG (Fig. 1B). The production of RF3 by plasmid pIQ-RF3 is dependent upon the presence of the inducer, IPTG (the RF3 level has been reported to increase 10-fold when 10 mM IPTG is present in the growth medium) (20). Our results indicate that the growth inhibition caused by the presence of plasmid pTnaC in strains grown in the presence of tryptophan




FIG. 1. Overproduction of RRF and RF3 relieves the growth inhibition caused by *tnaC* overexpression. (A) Schematic representation of the organization of plasmids pTnaC and pTnaC-RRF. Both plasmids are pUC18 derivatives and contain the *tna* operon promoter, the *tnaC* open reading frame, the noncoding region located between *tnaC* and *tnaA* genes, and an added *rpoBC* terminator. In preparing pTnaC-RRF, a fragment containing the RRF coding region and its own promoter from plasmid pRR1 was subcloned into plasmid pTnaC immediately following the *rpoBC* terminator. (B) Overproduction of both RRF and RF3 relieves the growth inhibition caused by *tnaC* overexpression by pTnaC in the presence of tryptophan. *E. coli* cells bearing the different plasmids listed in the figure (Table 1) were diluted from cultures grown overnight and were grown with shaking at 37°C in minimal medium supplemented with 0.05% acid-hydrolyzed casein and 0.2% glycerol with (+) or without (-) 100 μ g/ml tryptophan (Trp). Cell growth was monitored at an OD₆₀₀. A total of 10 mM IPTG (+) was added at time zero for RF3 overproduction. Four independent growth experiments were performed. The growth curves for the different strains grown with or without added tryptophan were all similar to those shown in this figure.

was relieved only when both RRF and RF3 were overproduced in transformants containing both plasmids pTnaC-RRF and pIQ-RF3 grown with 10 mM IPTG (Fig. 1B). Transformants overproducing RRF or RF3 alone did not exhibit relief from this growth inhibition (Fig. 1B). These findings demonstrate that an excess of both factors RRF and RF3 is necessary in order to reverse the growth inhibition caused by plasmid-mediated *tnaC* overexpression in strains grown in the presence of tryptophan.

In the parental strain and a strain overexpressing *tnaC*, the half-life of TnaC-tRNA₂^{Pro} is 8 min under induction conditions in vivo. Cultures of strain SVS1144 with or without pTnaC (Table 1), a plasmid overexpressing *tnaC*, were grown to an OD₆₀₀ of 0.6 to 0.8 in supplemented minimal medium in the presence or absence of 100 μ g/ml tryptophan. Glucose was then added to each culture (final level, 1%) to inhibit the initiation of transcription of the *tna* operon, and portions of each culture were harvested at the times indicated (Fig. 2). The levels of TnaC-tRNA₂^{Pro} were then measured, and its half-life was determined (Fig. 2). The parental SVS1144 culture, expressing *tnaC* chromosomally, when grown in the absence of tryptophan, had no detectable TnaC-tRNA₂^{Pro} (Fig. 2A). When grown in the presence of tryptophan, its TnaC-tRNA₂^{Pro} level in the sample at time zero was low, and its half-life was approximately 8 min (Fig. 2A and C). In previous in vitro-coupled transcription-translation assays performed with RF2-deficient cell extracts, it was shown that the transcription of *tnaC* and translation of *tnaC* mRNA were not affected by the ab-

sence of tryptophan (8). Thus, the absence of detectable TnaC-tRNA₂^{Pro} in cultures grown without tryptophan is probably due to its rapid cleavage; its half-life under these conditions is presumably <8 min. When strain SVS1144 containing plasmid pTnaC was grown in the presence of tryptophan, the TnaC-tRNA₂^{Pro} half-life was also approximately 8 min (Fig. 2B and C). In the absence of tryptophan, no TnaC-tRNA₂^{Pro} was detected; therefore, its half-life is presumably <8 min. These findings demonstrate that under induction conditions, with or without *tnaC* overexpression, TnaC-tRNA₂^{Pro} accumulates. However, it is eventually hydrolyzed, yielding TnaC and tRNA₂^{Pro}. The freed tRNA₂^{Pro} and ribosomes thus would become available for additional protein synthesis.

Overproduction of RRF and RF3 in a strain overexpressing *tnaC* reduces TnaC-tRNA₂^{Pro} accumulation and shortens its half-life. Next, experiments to determine if the stability of TnaC-tRNA₂^{Pro} was affected in strains containing plasmid pTnaC when RF3 and/or RRF was limiting or overproduced were performed (Fig. 3). Cultures of wild-type and mutant strains with plasmid pTnaC were grown at 37°C in supplemented minimal medium in the presence of 100 μ g/ml tryptophan to an OD₆₀₀ of 0.6 to 0.8. At this temperature, it was observed by Western blot analysis that the RRF level is reduced in strains containing *frr*(Ts) (data not shown). Glucose was then added to each culture (final level, 1%) to shut off *tna* operon transcription initiation. Portions of each culture were harvested at the times indicated, and the level and half-life of TnaC-tRNA₂^{Pro} were measured (Fig. 3). In the control culture




FIG. 2. *tnaC* overexpression does not affect the half-life of TnaC-tRNA₂^{Pro} in vivo. Strain SVS1144 (A) or SVS1144 harboring plasmid pTnaC (B) was grown at 37°C in minimal medium supplemented with 0.05% acid-hydrolyzed casein and 0.2% glycerol with (+) or without (-) the addition of 100 µg/ml tryptophan (Trp). The OD₆₀₀ at the time of glucose addition (final concentration of 1.0%) to each culture was 0.6 to 0.8. Samples were taken at the indicated times, harvested by centrifugation, and disrupted by sonication, and equivalent amounts of total protein were electrophoresed on a 10% Tricine-sodium dodecyl sulfate acrylamide gel. Northern blotting was then performed to quantify the level of TnaC-tRNA₂^{Pro} (TC-tRNA₂^{Pro}). The TnaC-tRNA₂^{Pro} band was detected at 26 kDa. The tRNA₂^{Pro} (tRNA₂^{Pro}) molecule was also detected at 14 kDa; however, measurements of tRNA₂^{Pro} levels were not accurate quantitatively (10). The measured half-life (t1/2) of TC-tRNA₂^{Pro}, calculated using the curves in C, is given to the right of each set of data. Little or no TnaC-tRNA₂^{Pro} was detected in the samples from uninduced cultures. The percentage of TnaC-tRNA₂^{Pro} relative to the level at 0 min was calculated by dividing the densitometry units obtained from the TnaC-tRNA₂^{Pro} band in each lane by the units obtained for the TnaC-tRNA₂^{Pro} band in the 0-min lane. Note that there is less TnaC-tRNA₂^{Pro} in the 0-min sample from chromosomal *tnaC* expression (compare +Trp TC-tRNA₂^{Pro} values in A with those in B). (C) Plot of TnaC-tRNA₂^{Pro} decay curves based on the data shown in A and B. Plots are not shown for the cultures grown without tryptophan. At least three independent experiments were performed, and the TnaC-tRNA₂^{Pro} values obtained for any strain and condition varied by less than 20%.

containing plasmid pTnaC (SVS1144 plus pTnaC), the TnaC-tRNA₂^{Pro} half-life was approximately 8 min. When either RF3 (pIQ-RF3) or RRF (pTnaC-RRF) was overproduced alone in strains with pTnaC, the TnaC-tRNA₂^{Pro} half-life was reduced from 8 min to 7 min and from 8 min to 4 min, respectively (Fig. 3A and B). When both RRF and RF3 were overproduced in the same cell, the half-life of TnaC-tRNA₂^{Pro} was reduced further, to 2 min (Fig. 3A and B). In pTnaC strains producing a temperature-sensitive RRF protein [*frr*(Ts)] incubated at 37°C for 1 h after shifting from 30°C, the half-life of TnaC-tRNA₂^{Pro} was increased only slightly (Fig. 3A and B). A similar result was observed with a strain lacking the RF3 protein (*prfCΔ2::kan*) (Fig. 3A and B). However, when both RRF and RF3 were limiting {strain SVS1144[*frr*(Ts) *prfCΔ2::kan*] plus pTnaC}, the half-life of TnaC-tRNA₂^{Pro} was increased appreciably, from 8 min to 20 min (Fig. 3A and B). These findings demonstrate that changing the levels of these two factors, RRF and RF3, has the most profound effect on the half-life of TnaC-tRNA₂^{Pro}. Consistent with this conclusion and the calculated half-lives are the levels of TnaC-tRNA₂^{Pro} detected in the 0-min samples that were analyzed (Fig. 3A, second column). These 0-min levels reflect the sensitivity of TnaC-tRNA₂^{Pro} to cleavage in the respective strains. The half-life of TnaC-

tRNA₂^{Pro} was also examined in strains overproducing other translational factors such as EF-G and IF3; these had no observable effects (data not shown).

Limiting Pth availability affects the half-life of TnaC-tRNA₂^{Pro}. When the synthesis of a peptidyl-tRNA is completed, either it is hydrolyzed in response to the action of one of the release factors or, if peptidyl-tRNA dropoff occurs, it is cleaved by Pth (21). Since the presence of tryptophan inhibits RF2-initiated hydrolysis of TnaC-tRNA₂^{Pro}, we wished to determine if the slow cleavage of TnaC-tRNA₂^{Pro} observed under tryptophan induction conditions was due to peptidyl-tRNA dropoff and Pth cleavage. The stability and half-life of TnaC-tRNA₂^{Pro} were determined in strains C600 and C600[*pth*(Ts)] as well as in strains C600 pTnaC and C600[*pth*(Ts)] pTnaC grown at 42°C (Fig. 4). These strains were grown under conditions chosen to maximally reduce the level of the Pth enzyme. Cultures were initially grown at 30°C in LB broth, a medium rich in tryptophan, to an OD₆₀₀ of 0.8 (mid-log phase). The cultures were then shifted to 42°C for an hour of additional growth, with the objective of reducing the cellular level of the Pth(Ts) protein (3). Glucose was then added, and cell samples were harvested at the times indicated. The TnaC-tRNA₂^{Pro} level was then determined. In wild-type strain C600




FIG. 3. Overproduction of RRF and RF3 reduces the half-life of TnaC-tRNA₂^{Pro} in vivo. (A) Strains bearing the indicated plasmid or plasmids were grown at 30°C or 37°C in minimal medium supplemented with 0.05% acid-hydrolyzed casein and 0.2% glycerol with the addition of 100 μg/ml tryptophan (Trp). In strain SVS1144(*prfCA2::kan*), the *prfC* gene was deleted. Strain SVS1144[*frr*(Ts)] produces an RRF protein that is temperature sensitive. The *frr*(Ts) allele was also present in SVS1144[*frr*(Ts) *prfCA2::kan*]. Strains with plasmid pIQ-RF3 were grown with 10 mM IPTG, which induces RF3 production. Plasmid pTnaC-RRF was used to overexpress *tnaC* and overproduce RRF. Strain SVS1144[*frr*(Ts)] pTnaC was grown initially at 30°C and then shifted to 37°C for 1 h. Presumably, most of its RRF protein would be inactivated during growth at 37°C. The OD₆₀₀ at the time of glucose addition (final concentration of 1.0%) to each culture was 0.6 to 0.8. Samples were taken and sonicated, and Northern blot assays performed to detect TnaC-tRNA₂^{Pro} (TC-tRNA₂^{Pro}) and tRNA₂^{Pro} (tRNA₂^{Pro}), as indicated in the legend of Fig. 2. The half-life (t_{1/2}) of TnaC-tRNA₂^{Pro}, calculated using the curve in C, is shown to the right of each set of data. The relative TnaC-tRNA₂^{Pro} level at 0 min is shown to the left of each panel. These values were calculated by the densitometry units obtained for the TnaC-tRNA₂^{Pro} band in each 0-min lane by the densitometry units obtained for the TnaC-tRNA₂^{Pro} band in the 0-min lane for the control SVS1144(pTnaC) culture. (B) The TnaC-tRNA₂^{Pro} decay curves are based on the percentage of TnaC-tRNA₂^{Pro} detected relative to the 0-min value, as indicated in Fig. 2. At least three independent experiments were performed, and the TnaC-tRNA₂^{Pro} values obtained for any strain and condition varied by less than 20%.

grown at 42°C, the half-life of TnaC-tRNA₂^{Pro} was 11 min. A similar value was obtained for strain C600 pTnaC; the half-life of its TnaC-tRNA₂^{Pro} was 9 min. (Fig. 4, compare A and B; also see Fig. 4C). When the TnaC-tRNA₂^{Pro} half-life was determined in mutant strains C600[*pth*(Ts)] and C600[*pth*(Ts)] pTnaC grown at 42°C, half-life values greater than 20 min were calculated for both strain (Fig. 4, compare A and B; also see C). These findings indicate that TnaC-tRNA₂^{Pro} was essentially uncleaved in these strains during the 20-min test period. This important result indicates that peptidyl-tRNA dropoff fol-

lowed by Pth cleavage is primarily responsible for the processing of this peptidyl-tRNA under these conditions.

DISCUSSION

Previous studies on the mechanism of tryptophan induction of *tna* operon expression have established that ribosome recognition of features of the newly synthesized TnaC peptidyl-tRNA, TnaC-tRNA₂^{Pro}, creates a tryptophan binding site in the ribosome at which bound tryptophan inhibits TnaC-




FIG. 4. Pth (peptidyl-tRNA hydrolase) is responsible for the cleavage of TnaC-tRNA₂^{Pro} in vivo. Cultures of strains C600 (wild type) and C600[*pth*(Ts)] (temperature-sensitive Pth) either without (A) or with (B) plasmid pTnaC (+pTnaC) were grown in LB medium {C600[*pth*(Ts)]} at 30°C to an OD₆₀₀ of 0.8. They were then shifted to 42°C and incubated for 1 h. Glucose was then added to 1%, and samples were harvested at the indicated times. Cultures of wild-type (wt) strain C600 grown at 42°C were used as controls. Northern blot assays were performed to detect TnaC-tRNA₂^{Pro} (TC-tRNAP) and tRNA₂^{Pro} (tRNAP), and the percentage of TnaC-tRNA₂^{Pro} detected relative to the 0-min value was calculated as described in the legend of Fig. 2. The half-life (t_{1/2}) of TnaC-tRNA₂^{Pro}, calculated using the curve in C, is shown to the right of each set of data in A and B. (C) TnaC-tRNA₂^{Pro} decay curves based on the data in A and B. At least two independent experiments were performed, and the TnaC-tRNA₂^{Pro} values obtained for any strain and condition varied by less than 15%.

tRNA₂^{Pro} cleavage (1, 2, 6). This uncleaved TnaC-tRNA₂^{Pro} is temporarily retained within the translating ribosome, stalling its movement (7, 9). This stalling at the *tnaC* stop codon blocks Rho factor binding to *tna* operon leader RNA, thereby permitting a paused RNA polymerase to continue transcription into the structural genes of the *tna* operon (7). It has also been shown that tryptophan induction in strains containing a plasmid overexpressing *tnaC* results in the retention of sufficient uncleaved TnaC-tRNA₂^{Pro} within the translating ribosome to create a tRNA₂^{Pro} deficiency (9). This deficiency reduces the cell growth rate. Providing additional uncharged tRNA₂^{Pro} was shown to restore normal growth (9).

If tryptophan induction of *tna* operon expression led to permanent ribosome stalling, induced cells would suffer from the accumulation of inactive ribosomes and a deficiency of tRNA₂^{Pro}. In previous studies, the half-life of TnaC-tRNA₂^{Pro} in cultures grown under inducing conditions was observed to be 10 to 15 min, implying that *E. coli* employs some natural mechanism to cleave the accumulated TnaC-tRNA₂^{Pro} (9). In the present study, the mechanisms used by *E. coli* to release and cleave ribosome-bound TnaC-tRNA₂^{Pro} were investigated. It was found that when *tnaC* was overexpressed in vivo, increased production of RRF plus RF3 reduced the half-life of TnaC-tRNA₂^{Pro} from 8 min to 2 min (Fig. 3). It was also observed that the inactivation of RRF in a temperature-sensitive *frr* mutant and the deletion of *prfC*, the structural gene for RF3, stabilized TnaC-tRNA₂^{Pro} appreciably; its half-life in-

creased from 8 min to >20 min (Fig. 3). It was further established that the principal means of cleavage of TnaC-tRNA₂^{Pro} was by the action of the enzyme Pth following TnaC-tRNA₂^{Pro} dropoff from the translating ribosome. In a mutant producing a temperature-sensitive Pth grown at an elevated, nonpermissive temperature, the TnaC-tRNA₂^{Pro} produced was essentially stable (Fig. 4). This stability was observed regardless of whether *tnaC* was expressed chromosomally or was overexpressed from a plasmid (Fig. 4). These findings suggest that the peptidyl-tRNA dropoff mediated by RRF and RF3 action is the normal mechanism employed by *E. coli* to relieve tryptophan-induced ribosome stalling at the *tnaC* stop codon. Dropoff presumably makes the TnaC-tRNA₂^{Pro} available for Pth cleavage, and this cleavage provides additional tRNA₂^{Pro} for protein synthesis. RF2 does not appear to play a major role in activating (or inhibiting) TnaC-tRNA₂^{Pro} cleavage under tryptophan induction conditions (data not shown). Rather, RRF and RF3 appear to bind to TnaC-tRNA₂^{Pro}-stalled ribosomes and release the TnaC-tRNA₂^{Pro}. This would free the previously stalled ribosomes and permit Pth to cleave the TnaC-tRNA₂^{Pro}.

E. coli is known to use at least three procedures to promote the release of a ribosome stalled at a sense or stop codon: (i) the action of the release factor RF1 or RF2 (26), (ii) RRF/RF3-mediated peptidyl-tRNA dropoff (28), and (iii) trans-translation by tmRNA (17). The in vivo and in vitro experiments described previously suggest that RF2 is ineffective in releasing ribosomes stalled at the *tnaC* UGA stop codon fol-

lowing the synthesis of TnaC-tRNA₂^{Pro}. This is due to bound tryptophan preventing RF2 action (6). Rather, as shown in this study, RRF and RF3 appear to promote TnaC-tRNA₂^{Pro} dropoff from the ribosome, and this peptidyl-tRNA is then cleaved by Pth.

The possible action of tmRNA on ribosome-bound TnaC-tRNA₂^{Pro} was also studied in vivo and in vitro (data not shown). It was found that tmRNA was incapable of attacking ribosomes containing TnaC-tRNA₂^{Pro} in vivo (overexpression of tmRNA had no observable effect on the β-Gal [TnaA-LacZ] level in induced cells) (Table 2). tmRNA action was also examined in vitro using two truncated inducible and noninducible (Trp12 replaced by Arg12) *tnaC* mRNAs (data not shown). In both transcripts, the UGA stop codon of *tnaC* was replaced by UU, with these nucleotides at the 3' end of the transcript. In reactions performed in the absence of added tryptophan, tmRNA was able to attack the TnaC peptidyl-tRNA produced from either transcript. However, when tryptophan was added, it blocked tmRNA action on the inducible transcript but not on the noninducible transcript. These data provide additional support for the conclusion that Trp12 of TnaC is essential for tryptophan binding and inhibition of TnaC-tRNA₂^{Pro} cleavage; they also indicate that tmRNA is not involved in TnaC-tRNA₂^{Pro} metabolism.

In other unpublished studies, Feng Gong, then in our laboratory, and Akira Kaji showed that RRF plays a role in determining the level of expression of a noninducible *tna* operon. They found that the expression of a noninducible *tnaA'*-*lacZ* fusion increased when RRF was limiting. They performed these studies using a plasmid with a mutant *tnaC* gene with the Trp12-to-Arg12 codon replacement. This result suggests that in the absence of RRF, a ribosome translating the *tnaC* Arg12 sequence is not liberated normally from the *tnaC* stop codon. It was also shown previously that the overproduction of RF3 influences the expression of *tnaA* of *E. coli*; however, the explanation for this observation was not known at that time (31). Previous studies also showed that RRF and RF3 were involved in peptidyl-tRNA dropoff from a translating ribosome (12, 28). In vitro analyses have also suggested that a disruption of *prfC*, or a down-regulation of RRF expression, results in a decreased rate of peptidyl-tRNA dropoff (12). It was also shown that RF3 can substitute for EF-G in RRF-dependent ribosome recycling reactions in vitro, during the dropoff of a short peptidyl-tRNA (11). The cryoelectron microscopy structure of the translating ribosome suggests that RF3 may induce a conformational change in the ribosome that facilitates the binding of RRF (18). In our studies, other factors such as EF-G or IF3 had no observable effects on either basal or induced *tna* operon expression (Table 2).

On the basis of studies performed previously by others and the results described in this paper, we propose the following explanation for the events that occur during tryptophan induction of *tna* operon expression. During induction, features of the TnaC-leader peptidyl-tRNA, most notably, Trp12 and Pro24 (1, 8), create a tryptophan binding site in the A site of the translating ribosome (1). Tryptophan binds at this site, interfering with RF2 action and preventing the cleavage of the newly synthesized TnaC-tRNA₂^{Pro}. RF3 and RRF then slowly bind to the ribosome and promote TnaC-tRNA₂^{Pro} dropoff from the ribosome. Pth then hydrolyzes the released TnaC-

tRNA₂^{Pro}, freeing tRNA₂^{Pro} for additional rounds of protein synthesis. The release of TnaC-tRNA₂^{Pro} also allows ribosome dissociation from the transcript. Our results raise several additional questions. How is tmRNA prevented from acting on TnaC-tRNA₂^{Pro}? What roles do RF3 and RRF play during the liberation of TnaC-tRNA₂^{Pro}? Additional studies will be required to answer these questions and identify the exact site(s) and mechanism(s) of action of the various translation termination factors.

ACKNOWLEDGMENTS

We are greatly indebted to Koichi Ito, Yoshikazu Nakamura, Akira Kaji, and Umesh Varshney for providing materials needed for these investigations. We thank Robert Sauer for generous gifts of tmRNA plasmids. We are also grateful to Catherine Squires, Yoshikazu Nakamura, and Akira Kaji for their helpful advice and for their comments on the manuscript. We thank Feng Gong and Akira Kaji for permission to cite their unpublished results.

These studies were supported by National Science Foundation grants MCB-0093023 and MCB-0615390 to C.Y.

REFERENCES

1. Cruz-Vera, L. R., M. Gong, and C. Yanofsky. 2006. Changes produced by bound tryptophan in the ribosome peptidyl transferase center in response to TnaC, a nascent leader peptide. *Proc. Natl. Acad. Sci. USA* **103**:3598–3603.
2. Cruz-Vera, L. R., S. Rajagopal, C. Squires, and C. Yanofsky. 2005. Features of ribosome-peptidyl-tRNA interactions essential for tryptophan induction of *tna* operon expression. *Mol. Cell* **19**:333–343.
3. Cruz-Vera, L. R., I. Toledo, J. Hernandez-Sanchez, and G. Guarneros. 2000. Molecular basis for the temperature sensitivity of *Escherichia coli pth* (Ts). *J. Bacteriol.* **182**:1523–1528.
4. Freistroffer, D. V., M. Y. Pavlov, J. MacDougall, R. H. Buckingham, and M. Ehrenberg. 1997. Release factor RF3 in *E. coli* accelerates the dissociation of release factors RF1 and RF2 from the ribosome in a GTP-dependent manner. *EMBO J.* **16**:4126–4133.
5. Gish, K., and C. Yanofsky. 1993. Inhibition of expression of the tryptophanase operon in *Escherichia coli* by extrachromosomal copies of the *tna* leader region. *J. Bacteriol.* **175**:3380–3387.
6. Gong, F., K. Ito, Y. Nakamura, and C. Yanofsky. 2001. The mechanism of tryptophan induction of tryptophanase operon expression: tryptophan inhibits its release factor-mediated cleavage of TnaC-peptidyl-tRNA(Pro). *Proc. Natl. Acad. Sci. USA* **98**:8997–9001.
7. Gong, F., and C. Yanofsky. 2002. Analysis of tryptophanase operon expression in vitro: accumulation of TnaC-peptidyl-tRNA in a release factor 2-depleted S-30 extract prevents Rho factor action, simulating induction. *J. Biol. Chem.* **277**:17095–17100.
8. Gong, F., and C. Yanofsky. 2002. Instruction of translating ribosome by nascent peptide. *Science* **297**:1864–1867.
9. Gong, M., F. Gong, and C. Yanofsky. 2006. Overexpression of *tnaC* of *Escherichia coli* inhibits growth by depleting tRNA₂^{Pro} availability. *J. Bacteriol.* **188**:1892–1898.
10. Gottesman, S., E. Roche, Y. Zhou, and R. T. Sauer. 1998. The ClpXP and ClpAP proteases degrade proteins with carboxy-terminal peptide tails added by the SsrA-tagging system. *Genes Dev.* **12**:1338–1347.
11. Grentzmann, G., P. J. Kelly, S. Laalami, M. Shuda, M. A. Firpo, Y. Cenatiempo, and A. Kaji. 1998. Release factor RF-3 GTPase activity acts in disassembly of the ribosome termination complex. *RNA* **4**:973–983.
12. Heurgue-Hamard, V., R. Karimi, L. Mora, J. MacDougall, C. Leboeuf, G. Grentzmann, M. Ehrenberg, and R. H. Buckingham. 1998. Ribosome release factor RF4 and termination factor RF3 are involved in dissociation of peptidyl-tRNA from the ribosome. *EMBO J.* **17**:808–816.
13. Hirokawa, G., R. M. Nijman, V. S. Raj, H. Kaji, K. Igarashi, and A. Kaji. 2005. The role of ribosome recycling factor in dissociation of 70S ribosomes into subunits. *RNA* **11**:1317–1328.
14. Ichikawa, S., and A. Kaji. 1989. Molecular cloning and expression of ribosome releasing factor. *J. Biol. Chem.* **264**:20054–20059.
15. Kaji, A., M. C. Kiel, G. Hirokawa, A. R. Muto, Y. Inokuchi, and H. Kaji. 2001. The fourth step of protein synthesis: disassembly of the posttermination complex is catalyzed by elongation factor G and ribosome recycling factor, a near-perfect mimic of tRNA. *Cold Spring Harb. Symp. Quant. Biol.* **66**:515–529.
16. Karimi, R., M. Y. Pavlov, R. H. Buckingham, and M. Ehrenberg. 1999. Novel roles for classical factors at the interface between translation termination and initiation. *Mol. Cell* **3**:601–609.
17. Keiler, K. C., P. R. Waller, and R. T. Sauer. 1996. Role of a peptide tagging system in degradation of proteins synthesized from damaged messenger RNA. *Science* **271**:990–993.

18. **Klaholz, B. P., A. G. Myasnikov, and M. van Heel.** 2004. Visualization of release factor 3 on the ribosome during termination of protein synthesis. *Nature* **427**:862–865.
19. **Kossel, H.** 1970. Purification and properties of peptidyl-tRNA hydrolase from *Escherichia coli*. *Biochim. Biophys. Acta* **204**:191–202.
20. **Matsumura, K., K. Ito, Y. Kawazu, O. Mikuni, and Y. Nakamura.** 1996. Suppression of temperature-sensitive defects of polypeptide release factors RF-1 and RF-2 by mutations or by an excess of RF-3 in *Escherichia coli*. *J. Mol. Biol.* **258**:588–599.
21. **Messenger, J. R.** 1976. Peptidyl transfer RNA dissociates during protein synthesis from ribosomes of *Escherichia coli*. *J. Biol. Chem.* **251**:3392–3398.
22. **Miller, J. H.** 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
23. **Moore, S. D., and R. T. Sauer.** 2005. Ribosome rescue: tmRNA tagging activity and capacity in *Escherichia coli*. *Mol. Microbiol.* **58**:456–466.
24. **Roche, E. D., and R. T. Sauer.** 2001. Identification of endogenous SsrA-tagged proteins reveals tagging at positions corresponding to stop codons. *J. Biol. Chem.* **276**:28509–28515.
25. **Roche, E. D., and R. T. Sauer.** 1999. SsrA-mediated peptide tagging caused by rare codons and tRNA scarcity. *EMBO J.* **18**:4579–4589.
26. **Scolnick, E., R. Tompkins, T. Caskey, and M. Nirenberg.** 1968. Release factors differing in specificity for terminator codons. *Proc. Natl. Acad. Sci. USA* **61**:768–774.
27. **Seshadri, A., and U. Varshney.** 2006. Mechanism of recycling of post-termination ribosomal complexes in eubacteria: a new role of initiation factor 3. *J. Biosci.* **31**:281–289.
28. **Singh, N. S., G. Das, A. Seshadri, R. Sangeetha, and U. Varshney.** 2005. Evidence for a role of initiation factor 3 in recycling of ribosomal complexes stalled on mRNAs in *Escherichia coli*. *Nucleic Acids Res.* **33**:5591–5601.
29. **Vogel, H. J., and D. M. Bonner.** 1956. Acetylornithinase of *Escherichia coli*: partial purification and some properties. *J. Biol. Chem.* **218**:97–106.
30. **Yanofsky, C., V. Horn, and P. Gollnick.** 1991. Physiological studies of tryptophan transport and tryptophanase operon induction in *Escherichia coli*. *J. Bacteriol.* **173**:6009–6017.
31. **Yanofsky, C., V. Horn, and Y. Nakamura.** 1996. Loss of overproduction of polypeptide release factor 3 influences expression of the tryptophanase operon of *Escherichia coli*. *J. Bacteriol.* **178**:3755–3762.
32. **Zavialov, A. V., R. H. Buckingham, and M. Ehrenberg.** 2001. A posttermination ribosomal complex is the guanine nucleotide exchange factor for peptide release factor RF3. *Cell* **107**:115–124.
33. **Zavialov, A. V., L. Mora, R. H. Buckingham, and M. Ehrenberg.** 2002. Release of peptide promoted by the GGQ motif of class 1 release factors regulates the GTPase activity of RF3. *Mol. Cell* **10**:789–798.