

---

# Human *Mycobacterium bovis* Infection and Bovine Tuberculosis Outbreak, Michigan, 1994–2007

Melinda J. Wilkins,\* Joshua Meyerson,†  
Paul C. Bartlett,‡ Susan L. Spieldenner,\*  
Dale E. Berry,\* Laura B. Mosher,\*  
John B. Kaneene,‡ Barbara Robinson-Dunn,§  
Mary Grace Stobierski,\* Matthew L. Boulton¶

*Mycobacterium bovis* is endemic in Michigan's white-tailed deer and has been circulating since 1994. The strain circulating in deer has remained genotypically consistent and was recently detected in 2 humans. We summarize the investigation of these cases and confirm that recreational exposure to deer is a risk for infection in humans.

Historically, *Mycobacterium bovis* infection in humans was associated with consumption of unpasteurized milk and dairy products (1,2) and this is still the most important route of exposure in developing countries. US populations are exposed to unpasteurized dairy products imported from countries where *M. bovis* is prevalent (3,4). *M. bovis* infection in humans is of concern to health officials in Michigan because of its endemicity in the state's wild white-tailed deer population and its discovery in several cattle herds. *M. bovis* in deer represents possible occupational and recreational routes of exposure to humans, especially for hunters, trappers, taxidermists, venison processors, and venison consumers (5).

Although *M. bovis* is a zoonotic agent, surveillance indicates no increase in its incidence in Michigan residents since an outbreak began in 1994. Since 1995, the incidence rate of *M. bovis* infection in Michigan residents has remained very low, with  $\approx 1$  new case per year for a total of 13. No genetic or epidemiologic link to the deer/cattle outbreak strain has been identified among 11 of these human *M. bovis* cases, based on restriction fragment-length

polymorphism analysis, spoligotyping, or mycobacterial interspersed repeat units (MIRU) typing (M. Wilkins, unpub. data, Michigan Department of Community Health, March 2007). Table 1 shows the spoligotyping and MIRU typing results from 9 available human specimens that were unrelated to the deer/cattle outbreak strain. All genotyping of isolates mentioned in this report was performed at the Michigan Department of Community Health, Bureau of Laboratories, Lansing, MI, USA, by using currently recommended guidelines (Centers for Disease Control and Prevention, Atlanta, GA, USA) (8–10). The remaining 2 human cases of *M. bovis* occurred in US-born, Michigan residents; the cases had epidemiologic and molecular links to the genotypically consistent deer/cattle outbreak strain circulating in Michigan.

## The Cases

### Patient 1, 2002

In January 2002, a 74-year-old man sought medical care, reporting malaise, anorexia, and fever. Past medical history included ischemic bowel disease, vascular disease, partial gastrectomy for peptic ulcers, and left upper lobectomy for squamous cell carcinoma (December 1999). On February 1, he was hospitalized with persistent fever and nonproductive cough; results of a chest radiograph were consistent with necrotizing pneumonia. A tuberculosis (TB) skin test (TST) result was negative, and a sputum smear was negative for acid-fast bacilli (AFB). After 5 days, the patient had not improved clinically; chest radiograph showed increasing infiltrate on the left side. Diagnostic bronchoscopy was performed, which yielded an AFB-positive smear. The condition of the patient deteriorated clinically over the next 10 days; he died on day 16 of his hospitalization.

Laboratory confirmation for TB, speciation, and antimicrobial drug susceptibility testing were pending at the time of his death. Genotyping analysis showed that the *M. bovis* isolated from this patient matched the circulating deer/cattle strain (Table 2).

In his youth, patient 1 lived on a farm geographically distant from the current bovine TB-endemic area. His first wife had a reported diagnosis of TB after their divorce >40 years before, and his second wife reports he drank unpasteurized milk as a youth. He moved to the edge of Deer Management Unit (DMU) 452 in 1994, which is the focal area for the bovine TB outbreak in deer. There, he ran a business with a buck pole where hunters displayed killed deer. Additional potential exposures included hunting white-tailed deer and consuming venison (>10 years before his death), handling a deer carcass from the DMU 452 vicinity in 2000, and recreational feeding of deer.

This patient was in poor health at the time of death, having acute and chronic illness. His poor health would

---

\*Michigan Department of Community Health, Lansing, Michigan, USA; †Health Department of Northwest Michigan, Charlevoix, Michigan, USA; ‡Michigan State University, East Lansing, Michigan, USA; §Beaumont Hospital, Royal Oak, Michigan, USA; and ¶University of Michigan, Ann Arbor, Michigan, USA



He continued to receive antimicrobial drug therapy for 9 months without further complications

As an experienced hunter, patient 2 recognized the tan nodules in the deer's chest cavity as *M. bovis* and promptly buried the carcass. In December, he led Michigan Department of Natural Resources staff back to the carcass, which was retrieved; the chest cavity was filled with lesions (Figure). Although the carcass was buried for >9 weeks, chest cavity samples were submitted for culture. After numerous attempts with alternative decontamination techniques, a viable culture was obtained. Genotyping results of the carcass isolate were identical to that recovered from patient 2 and the circulating deer/cattle strain (Table 2). The investigation of the infection in patient 2 provided strong evidence of transmission of *M. bovis* infection from deer to human through percutaneous injection with a contaminated hunting knife. The patient's history of hunting exposure was essential to diagnosis and treatment of this rare form of TB.

## Conclusions

Although epidemiologic evidence presented for patient 1 is not irrefutable, we conclude that both cases are part of a cluster that is epidemiologically and genotypically confirmed (11). The initial TST result was negative in both of these cases, likely due to cutaneous anergy (patient 1) and administration too soon after exposure (patient 2). Initial negative skin test results made diagnosis problematic for healthcare providers.

The confirmation of a hunter's acquiring cutaneous *M. bovis* from an infected deer supports the need for public health precautions. First, hunters should wear heavy latex or rubber gloves while field dressing deer. Second, hunter education was important in the second case



Figure. Chest cavity of a deer shot by patient 2; the deer was retrieved after being buried for 9 weeks. The photo shows the classic nodular lesions of *Mycobacterium bovis* infection. Photo: J.S. Fierke, D.J. O'Brien, S.M. Schmitt, Wildlife Disease Laboratory, Michigan Department of Natural Resources.

because the hunter recognized the deer as infected and specifically mentioned his exposure each time he sought medical treatment. Third, efforts to raise the index of suspicion of the medical community regarding cutaneous and other occupational or recreational exposures to TB continues to be important, so that appropriate diagnoses can be made. Finally, in both cases, the initially negative TST result complicated the diagnostic efforts. It is an ongoing challenge to ensure that providers appropriately apply and interpret the TST.

## Acknowledgments

We thank Sue Lovelace and Bridget Patrick for patient 2 follow-up, Steven V. Church for his laboratory contributions, and Dan J. O'Brien and Steven M. Schmitt for Michigan Department of Natural Resources assistance.

Dr Wilkins is the director of the Division of Communicable Disease, Michigan Department of Community Health. Her research interests include the epidemiology and control of zoonotic diseases, specifically *M. bovis*, and surveillance system development and evaluation. She is also a doctoral candidate at Michigan State University, College of Veterinary Medicine.

## References

1. Enarson DA, Rieder HL, Meslin FX, Cosivi O. The importance of *Mycobacterium bovis* to the tuberculosis epidemic in humans. In: Thoen CO, Steel JH, editors. *Mycobacterium bovis: infection in animals and humans*. Ames (IA): Iowa State University Press; 2006. p. xxii–xxv.
2. Wigle WD, Ashley MJ, Killough EM, Cosens M. Bovine tuberculosis in humans in Ontario: the epidemiologic features of 31 active cases occurring between 1964 and 1970. *Am Rev Respir Dis*. 1972;106:528–34.
3. Dankner WM, Waecker NJ, Essey MA, Moser K, Thompson M, Davis CE. *Mycobacterium bovis* infections in San Diego: a clinicoepidemiologic study of 73 patients and a historical review of a forgotten pathogen. *Medicine (Baltimore)*. 1993;72:11–37.
4. Centers for Disease Control and Prevention. Human tuberculosis caused by *Mycobacterium bovis*—New York City, 2001–2004. *MMWR Morb Mortal Wkly Rep*. 2005;54:605–8.
5. Wilkins MJ, Bartlett PC, Frawley B, O'Brien DJ, Miller CE, Boulton ML. *Mycobacterium bovis* (bovine TB) exposure as a recreational risk for hunters: results of a Michigan Hunter Survey, 2001. *Int J Tuberc Lung Dis*. 2003;7:1001–9.
6. Cowan LS, Mosher L, Diem L, Massey JP, Crawford JT. Variable-number tandem repeat typing of *Mycobacterium tuberculosis* isolates with low copy numbers of IS6110 by using mycobacterial interspersed repetitive units. *J Clin Microbiol*. 2002;40:1592–602.
7. *Mycobacterium bovis* international database [cited 2007 Mar 14]. Available from <http://www.mbovis.org>
8. Yang Z, Barnes PF, Chaves F, Eisenach KD, Weis SE, Bates JH, et al. Diversity of DNA fingerprints of *Mycobacterium tuberculosis* isolates in the United States. *J Clin Microbiol*. 1998;36:1003–7.
9. Cowan LS, Diem L, Brake MC, Crawford JT. Transfer of a *Mycobacterium tuberculosis* genotyping method, spoligotyping, from a reverse line-blot hybridization, membrane-based assay to the Luminex multianalyte profiling system. *J Clin Microbiol*. 2004;42:474–7.

10. Kwara A, Schiro R, Cowan LS, Hyslop NE, Wiser MF, Harrison SR, et al. Evaluation of the epidemiologic utility of secondary typing methods for differentiation of *Mycobacterium tuberculosis* isolates. *J Clin Microbiol.* 2003;41:2683–5.
11. National TB. Controllers Association/CDC Advisory Group on Tuberculosis Genotyping. Guide to the application of genotyping to tuberculosis prevention and control. Atlanta: Centers for Disease Control and Prevention; June 2004 [cited 2007 Mar 14]. Available from <http://www.cdc.gov/nchstp/tb/genotyping/toc.htm>

---

Address for correspondence: Melinda J. Wilkins, Michigan Department of Community Health, 201 Townsend, PO Box 30195, Lansing, MI, 48909; email: [wilkinsm@michigan.gov](mailto:wilkinsm@michigan.gov)

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.