

## Antiprotozoal activities of *Vincetoxicum stocksii* and *Carum copticum*

Arsala Mansoor<sup>1</sup>, Mohammad Ali Ibrahim<sup>1</sup>, Mudassir Asrar Zaidi<sup>2</sup> and Moinuddin Ahmed<sup>2</sup>

<sup>1</sup>Department of Pharmacognosy, School of Pharmacy, University of Mississippi, MS38677, USA; <sup>2</sup>Department of Botany, University of Balochistan, Quetta, Pakistan.

### Article Info

Received:	23 August 2011
Accepted:	26 August 2011
Available Online:	10 September 2011
Keywords:	
Antileishmanial	
Antimalarial	
Carum copticum	
Vincetoxicum stocksii	
Number of Tables:	3
Number of Refs:	9
Correspondence: MAZ e-mail: mudassirisrar@hotmail.com	

### Abstract

The antileishmanial and antimalarial activities of different fractions of *Vincetoxicum stocksii*, *Carum copticum*, *Gentiana olivieri* and *Zygophyllum fabago* were determined. Amongst these, *V. stocksii* and *C. copticum* showed moderate to good antimalarial activity. The ethyl acetate fractions of *C. copticum* and hexane fraction of *V. stocksii* showed 94 % and 52% inhibition of *Plasmodium falciparum* (D6 Clone) respectively. The hexane ethyl acetate fraction of *C. copticum* *V. stocksii* exhibited antileishmanial activity against, *Leishmania donovani* promastigotes with  $IC_{50}$  of 30  $\mu$ g/mL and 51  $\mu$ g/mL, respectively.

### Introduction

The parasitic diseases such as malaria and leishmaniasis have a high death rate in developing countries. Every year more than 500 million people, mostly those living in the world's poorest countries are at risk of malaria (Mansoor, 2003). Approximately 350 million people living in tropical and sub-tropical countries, suffer from this disease (Bharate, 2007). Therefore, there is a need to develop effective antimalarial and antileishmanial drugs for the treatment of these diseases.

The natives of Balochistan from centuries have been using more than 400 species of medicinal plants. The medicinally important plants *Carum copticum* Linn. Syn (Umbelliferae), *Vincetoxicum stocksii* (Asclepiadaceae) Ali & Khatoon, *Gentiana olivieri* Griseb (Gentianaceae) and *Zygophyllum fabago* (Zygophyllaceae) are widely distributed in Pakistan. *C. copticum* is used for curing diarrhea, amebiasis, dyspepsia and killing worms in stomach and other stomach disorders. It is also used as an antihelminthic, antispasmodic and antiseptic agent (Sahaf, 2007). It is traditionally used, as antileishmanial

herb *Vincetoxicum stocksii* is used by the natives as poultice for the treatment of wounds, injuries and external cancers (Staerk et al., 2000). *G. olivieri* is used in traditional use folk medicine for the treatment of skin diseases, abscesses, ulcer, blood pressure (Mansoor et al., 1998) and fever caused by malaria. *Z. fabago* is used to treat skin diseases, septic, injuries and external wounds (Zaidi, 2006) and Leishmania. In this paper the antimalarial and antileishmanial effects of these four medicinal plants were determined.

### Materials and Methods

**Plant material:** The whole plant of *C. copticum*, *V. stocksii*, *G. olivieri* and *Z. fabago* (2 kg each) were collected in June 2006 from the mountains of Quetta. These were identified and a voucher specimen of all the plants were deposited in the herbarium of Botanical Garden, University of Balochistan Quetta. Plants were cleaned, dried in shade, ground, packed and then sent to USA for analysis. These were soaked in ethanol and extracted repeatedly 3-4 times at room temperature.



This work is licensed under a Creative Commons Attribution 3.0 License. You are free to copy, distribute and perform the work. You must attribute the work in the manner specified by the author or licensor.

The extract was filtered and concentrated under reduced pressure in a rotary evaporator at 40°C, to obtain thick crude extract, which yielded *C. copticum* (23.4 g), *V. stocksii* (20.7 g), *G. olivieri* (27.5 g) and *Z. fabago* (22.8 g). The crude ethanolic extract of these plants was evaluated for antimarial and antileishmanial activity.

**Preparation of fractions:** Only biologically active plant extracts were subjected to further fractionation. The crude ethanol extract of *C. copticum* was fractionated by vacuum liquid chromatography. Elution was initiated by hexane (Fraction ACCH-1 4.6 g), followed by hexane/ethyl acetate (Fraction ACCHE-2, 2.8 g), ethyl acetate (Fraction ACCE-3, 1.9 g), ethyl acetate/acetone (Fraction ACCEA-4, 2.5 g), acetone (Fraction ACCA-5, 2.8 g), acetone/methanol (Fraction ACCAM-6, 2.7 g), methanol (Fraction AccM-7, 3.5 g) and finally with water (ACCW-8, 2.1 g).

Ethanol extract of *V. stocksii* was dissolved in water and partitioned first with hexane (Fraction AVH, 3.9 g) and then with ethyl acetate (Fraction AVE, 5.3 g) and finally with butanol (Fraction AVB, 9.8 g). The activity of each fraction was determined for antimarial and antileishmanial assay.

**Antileishmanial activity:** Antileishmanial activity of the extracts of the plants was tested in vitro against a culture of *Leishmania donovani* promastigotes grown at 26°C, in RPMI 1640 medium supplemented with 10% fetal calf serum (Gibco Chem. Co.). A 3-day-old culture was diluted to  $5 \times 10^5$  promastigotes/mL. Extract dilutions were prepared in cell suspension in 96-well plates, which were incubated at 26 °C for 48 hours. Almar blue assay was used to determine the growth of *L. donovani* promastigotes (Mikus and Steverding, 2000). Fluostar Galaxy plate reader (BMG Lab Technologies) was used to measure standard fluorescence at excitation wavelength of 544 nm and emission wavelength of 590 nm. Amphotericin B and pentamidine were used as the standard antileishmanial agents. IC<sub>50</sub> and IC<sub>90</sub> values were computed from dose-response curves generated by plotting percent growth versus drug concentration.

**Assay for in vitro antimarial activity:** The plasmodial LDH Activity is the bases for the determination of antimarial activity. A red blood cells suspension was infected with D6 or W2 strains of *P. falciparum* D6 Clone (200 µL, with 2%hematocrit and 2%parasitemia, in RPMI 1640 medium, with 60 µg/mL amikacin and 10% human serum) was added to the wells of a 96 well plate containing 10 µg/mL of test samples diluted in medium at various concentrations. The plate was kept in a modular incubation chamber (Billups-Rothenberg, CA) flushed with 90%N<sub>2</sub>, 5%O<sub>2</sub>, and 5%CO<sub>2</sub> gas mixture, and incubated at 37°C, for 72 hours. Malstat™ reagent was used to determine parasitic LDH activity (Flow Inc., Portland, OR) as described by elsewhere (Makler

and Hinrichs, 1993). Incubation mixture (20 µL) was mixed with 100 µL of the Malstat™ reagent at room temperature and incubated for 30 min. Then twenty microliters of NBT/PES (Sigma, St. Louis, MO) 1:1 mixture was added. The plate was further incubated for 1 hour in the dark. Hundred microliter of a 5%acetic acid solution was then added to stop the reaction. EL-340 Biokinetics Reader (Bio-Tek Instruments, Vermont) was used to read the plate at 650 nm. IC<sub>50</sub> values were computed from the dose-response curves. Chloroquine and artemisinin were used as the drug controls in each assay. DMSO (0.25%) was used as vehicle for control.

## Results and Discussion

Ethanol extracts of all the four plants were evaluated for in vitro antileishmanial and antimarial, activities. Only the extracts of *C. copticum* and *V. stocksii* showed moderate to mild antileishmanial activities with IC<sub>90</sub> values of 60 µg/mL and 70 µg/mL respectively. IC<sub>50</sub> of 50 µg/mL was detected for *V. stocksii* (Table I). No such

Table I: Antiprotozoal activity of the ethanol extracts of *Gentiana olivieri*, *Carum copticum*, *Zygophyllum fabago* and *Vincetoxicum stocksii*

Sample Name	P. falciparum	L. donovani	
	% Inhibition	IC <sub>50</sub> (µg/mL)	IC <sub>90</sub> (µg/mL)
<i>Gentiana olivieri</i>	28	Inactive	Inactive
<i>Carum copticum</i>	60	22	60
<i>Zygophyllum fabago</i>	Inactive	Inactive	Inactive
<i>Vincetoxicum stocksii</i>	40	50	70
Pentamidine	Not tested	1.2	6
Amphotericin B	Not tested	0.28	0.74
Chloroquine	99.2	Not tested	Not tested
Artemisinin	98.8	Not tested	Not tested

IC<sub>50</sub> and IC<sub>90</sub> are the sample concentrations that kill 50% and 90% Leishmanial cells compared to the solvent controls; Test = 3 concentrations: 100, 20, 4 µg/mL

activity was shown by *G. olivieri* and *Z. fabago*. *C. copticum* showed good antimarial activity, whereas, moderate activity was shown by *V. stocksii*. Poor activity was shown by *G. olivieri* and no such activity was observed by *Z. fabago*. Therefore, further fractionation of the extracts of the two active medicinal plants i.e., *C. copticum* and *V. stocksii* were carried out.

The most potent antimalarial activity (94%) was exhibited by fraction ACCE-3 and ACCHE-2 (90%) of *C. copticum* (Table II). The significant antimarial activity of ACCE-2 and ACCHE-3 indicates that the active compounds must be present in the less-polar fractions

Table II: In vitro antimalarial and antileishmanial activities of fractions of *Carum copticum*

Samples Name	P. falciparum	L. donovani	
	% Inhibition	IC <sub>50</sub> (µg/mL)	IC <sub>90</sub> (µg/mL)
Carum copticum (ACCH-1)	0	90	Inactive
Carum copticum (ACCHE-2)	90	30	90
Carum copticum (ACCE-3)	94	50	90
Carum copticum (ACCEA-4)	82	50	98
Pentamidine	Not tested	2.1	6.2
Amphotericin B	Not tested	0.09	0.2
Chloroquine	99	Not tested	Not tested
Artemisinin	99.4	Not tested	Not tested

IC<sub>50</sub> and IC<sub>90</sub>, are the concentrations that affords 50% and 90% inhibition of leishmanial growth compared to the solvent controls; Test = 3 concentrations; Carum copticum concentration: 100, 20, 4 µg/mL

Table III: In vitro antimalarial and antileishmanial activity of the fractions of *Vincetoxicum stocksii*

Samples Name	P. falciparum	L. donovani	
	% Inhibition	IC <sub>50</sub> (µg/mL)	IC <sub>90</sub> (µg/mL)
Vincetoxicum stocksii AVH	52	51	90
Vincetoxicum stocksii AVE	21	Inactive*	Inactive*
Vincetoxicum stocksii AVB	15	Inactive*	Inactive*
Pentamidine	Not tested	1.2	5.2
Amphotericin B	Not tested	0.45	0.9
Chloroquine	99.6	Not tested	Not tested
Artemisinin	99	Not tested	Not tested

IC<sub>50</sub> and IC<sub>90</sub>, are the concentrations that affords 50% and 90% inhibition of leishmanial growth compared to the solvent controls; \*at 40 µg/mL

of hexane/ethyl acetate (Fraction ACCHE-2) and ethyl acetate (ACCE-3). The chemical composition of the essential oil from dry seeds of *C. copticum* was found to have thymol (41.3%), α-terpinolene (17.5%) and β-cymene (11.8%) as the major constituents of the oil and have shown to exhibit strong insecticidal activity. The mortalities of the insect species reached 100% at

concentrations higher than 185.2 µL/L and 12-hours exposure time (Sahaf, 2007). The findings indicate that *C. copticum* oil had potential antiprotozoal activities.

Since good antiprotozoal activity was shown by *Vincetoxicum stocksii*, so its ethanol extract was dissolved in water and partitioned first with hexane (Fraction AVH) and then with ethyl acetate (Fraction AVE) and finally with butanol (Fraction AVB) (Table III). Moderate activity was shown against *P. falciparum* and *L. donovani* by the hexane fraction of *V. stocksii* whereas poor or no activity was shown by the other two fractions.

*Vincetoxicum* is known for its poisonous as well as its medicinal properties. It has also been found to be allelopathic and active against a number of microorganisms (Zaidi and Crow, 2005). Pronounced cytotoxicity has also been reported against *Candida albicans* by Staerk et al. (2000). This effect is probably due to the presence of phenanthroindolizidine alkaloids reported from other species of *Vincetoxicum*.

In conclusion, this study demonstrates good antimalarial and moderate antileishmanial activity of the extracts of *C. copticum* and *V. stocksii*, which parallels the traditional use of these herbs as an antiprotozoal medicine.

#### Acknowledgment

This research was supported by postdoctoral research program of Higher Education Commission of Pakistan. Special thanks to Dr. M.T. Hamman and Shabana I. Khan for providing facilities and guidance during this study.

#### References

- Bharate SB, Khan SI, Yunus NA, Chauthe SK, Jacob MR, Tekwani BL, Khan IA, Singh IP. Antiprotozoal and antimicrobial activities of O-alkylated and formylated acylphloroglucinols. Bioorg Med Chem. 2007; 15: 87-96.
- Makler MT, Hinrichs DJ. Measurement of the lactate dehydrogenase activity of *Plasmodium falciparum* as an assessment of parasitemia. Am J Trop Med Hyg. 1993; 48: 205-09.
- Mansoor A, Samad A, Zaidi MI. Hypotensive effect of *G. olivieri* and its alkaloid gentianine in rats. Pharm Pharmacol Comm. 1998; 4: 229-30.
- Mansoor A. Toxicological evaluation of the extracts and pure compounds of *G. olivieri*. Pak J Biol Sci. 2003; 6: 1949-50.
- Mikus J, Steverding D. A simple colorimetric method to screen drug cytotoxicity against *Leishmania* using the dye alamar blue. Parasitol Int. 2000; 48, 265-69.
- Sahaf BZ, Moharrampour S, Meshkalsadat MH. Chemical constituents and fumigant toxicity of essential oil from *Carum copticum* against two stored product beetles. Insect

- Sci. 2007; 14: 213-18.
- Staerk D, Christensen J, Lemmich E, Duus JØ, Olsen CE, Janoezewsk JW. Cytotoxic activity of some phenanthroindolizidine N-oxide alkaloids from *Cynanchum vincetoxicum*. *J Nat Prod*. 2000; 63: 1584-86.
- Zaidi MI. Phytotoxicity and insecticidal activity of *Vincetoxicum stocksii*. In: Plant ecology and biodiversity research. Barak SK (ed). New Delhi, Regency Publishers, 2006, pp 193-97.
- Zaidi MA, Crow SA Jr. Biologically active traditional medicinal herbs from Balochistan, Pakistan. *J Ethnopharmacol*. 2005; 96: 331-34.
-