

## **Anti-CD20 Monoclonal Antibody (Rituximab) Treatment for Cryoglobulinemic Vasculitis: Where do We Stand?**

Patrice CACOUB, M.D., Professor (1),  
Aurélien DELLUC, M.D., (1),  
David SAADOUN, M.D. (1),  
Dan Avi LANDAU, M.D (1)  
Damien SENE, M.D., (1)

<sup>1</sup>Université Pierre et Marie Curie-Paris6, CNRS, UMR 7087, Paris, F-75013 France ;  
AP-HP, Hôpital Pitié-Salpêtrière, Service de Médecine Interne, Paris, F-75013 France ;

**Running title:** Rituximab Treatment for Cryoglobulinemia Vasculitis

**Key words:** cryoglobulinemia vasculitis, anti-CD20, rituximab, hepatitis C virus, vasculitis, interferon alpha.

**This paper includes:** abstract (230 words), 2107 words, 8 pages, 25 references, 3 tables.

**Correspondence should be addressed to:** Professor Patrice Cacoub, M.D., AP-HP, Hôpital Pitié-Salpêtrière, Service de Médecine Interne, Paris, F-75013 France.  
e-mail: [patrice.cacoub@psl.aphp.fr](mailto:patrice.cacoub@psl.aphp.fr)

**Licence statement**

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in ARD and any other BMJPG products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<http://ARD.bmj.com/ifora/licence.pdf>)

**Abstract**

**Background:** Mixed cryoglobulinemia (MC) vasculitis represents a complication of the B cell response to a variety of chronic inflammatory diseases. Recent reports describe the use of monoclonal antibodies directed to CD20 antigen (Rituximab), a transmembrane protein expressed on pre-B lymphocytes and mature lymphocytes. The goal of this article is to review published data in order to better analyze the efficacy and tolerance of Rituximab treatment in patients with MC vasculitis.

**Methods and results:** After systematic review of the literature and exclusion of review papers, 13 references reported on a total number of 57 cases of MC secondary to hepatitis C virus (HCV) infection (75.4%) or essential mixed cryoglobulinemia (24.6%). Previous treatments failed to control the main signs of vasculitis; these were either HCV (n=37) or immunomodulating treatments. Most patients (48/57) received four weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of Rituximab. The duration of follow-up after Rituximab therapy was 9.7 months. Rituximab infusions had great efficacy on the main vasculitis signs, with a clinical response in 80-93% patients. A relapse of MC was noted in 14/36 (39%) patients. A relatively small number of side effects were reported.

**Conclusion:** Rituximab therapy for patients with mixed cryoglobulinemia vasculitis, HCV-induced or essential, shows a great efficacy on the main vasculitis signs in the majority of reported patients. A relapse of cryoglobulinemia vasculitis was frequently noted. Randomised controlled trials with long term study are needed to conclude definitively on the benefit/risk ratio of Rituximab therapy in such patients.

Mixed cryoglobulinemia (MC) is a systemic vasculitis characterized by the proliferation of B-cell clones producing pathogenic IgM with rheumatoid factor activity. MC leads to clinical manifestations ranging from MC syndrome (purpura, arthralgia, asthenia) to more serious lesions with neurological and renal involvement [1, 2]. Since 1990 it has been established that Hepatitis C virus (HCV) infection is associated with most cases of MC. On one hand, 60 to 90% of patients with MC are HCV-infected. In the other hand, 36 to 55% of HCV-infected patients have a positive mixed cryoglobulin. Up to 15-20% of HCV-MC patients will present a MC-associated systemic vasculitis that may be severe enough to cause death [3, 4].

Limited data are available regarding the treatment of patients with MC systemic vasculitis. In patients with HCV infection, interferon (IFN)- $\alpha$  monotherapy is associated with a relatively poor response and a high relapse rate, especially in severe cases [5, 6]. The combination of peg-IFN  $\alpha$  with ribavirin, which is the current standard for the treatment of patients with chronic HCV infection, has shown better efficacy, with up to 77% of patients showing complete remission [7]. The poor tolerability (in many cases) of conventional anti-HCV therapy, including flu-like symptoms, lends further importance to the development of additional therapeutic approaches. The use of corticosteroids, cyclophosphamide, and plasmapheresis, both in HCV-MC patients with no response or intolerance to HCV treatment and in patients with non-HCV MC, may lead to life-threatening complications and is difficult to manage in the long term.

After initial single case reports [8-10], two Italian groups have reported the efficacy of anti-CD20 monoclonal antibody (Rituximab) treatment in patients with HCV-MC vasculitis resistant or intolerant to IFN- $\alpha$  monotherapy [11, 12]. Such approach involves the use of monoclonal antibodies directed to CD20 antigen, a transmembrane protein expressed on pre-B lymphocytes and mature B lymphocytes. Rituximab proved effective on skin vasculitis manifestations, subjective symptoms of peripheral neuropathy, arthralgia and low grade B-cell lymphoma. Most clinical responders also had decrease in serum cryoglobulin levels and increased in C4 serum levels, though not to undetectable or normal levels. However, one potential concern regarding the use of Rituximab is its propensity to worsen HCV viremia [12], which may lead patients to develop more severe HCV-induced liver lesions and/or cryoglobulinemic relapses in ensuing years [13]. These studies did not allow conclusions to be drawn concerning the efficacy of anti-CD20 monoclonal antibody on peripheral neuropathy and nephropathy.

The goal of this article is to review published data in order to better analyze the efficacy and tolerance of Rituximab treatment in patients with cryoglobulinemia vasculitis.

## Methods

A systematic Pubmed search with the following key words "Cryoglobulinemia, Rituximab and anti-CD20" resulted in 54 references. Review papers were excluded. Fifteen references which reported on at least one case of cryoglobulinemia vasculitis treated with rituximab were included in the present analysis [8-22]. Two references were excluded because they included initial reports of patients which had already been reported on in other publications [8, 9].

The response to treatment was analysed by comparing clinical and immunologic parameters before Rituximab infusion and at the end of follow-up. Clinical response was defined by analyzing the evolution of the following main clinical signs: skin involvement (absence of purpura), peripheral neuropathy (clinical and/or electrophysiologic improvement), renal involvement (normalization of serum creatinine level and

disappearance of proteinuria), and the absence of arthralgia. Response to Rituximab treatment was defined as complete (disappearance of baseline manifestations) or partial (improvement of baseline clinical manifestations); all other patients were classified as non responders. Relapse was defined as the reappearance of at least one clinical sign of vasculitis. A complete immunologic response was defined by the absence of serum cryoglobulin, and a partial immunologic response was defined by a decrease of more than 50% from the baseline cryoglobulin level.

## Results

Overall, 13 references reported on a total number of 57 cases which were analyzed in detail for the present study [10-23]. There were two large uncontrolled series of 20 [12] and 15 [11] patients, and two smaller series of 6 [13] and 5 [21] patients. All other publications reported either single or two case reports.

The baseline characteristics of patients with cryoglobulinemia vasculitis who received anti-CD20 antibody (Rituximab) treatment are summarized in **Table 1**. Fifty-seven patients, mainly of female gender (79%) with a mean age of 59 years, had cryoglobulinemia vasculitis secondary to chronic active HCV infection (75.4%) or essential mixed cryoglobulinemia (24.6%). Previous treatments failed to control the main signs of vasculitis; these were either HCV treatment (n=37) or immunomodulating treatments (corticosteroids, immunosuppressive drugs, plasma exchanges). The main clinical manifestations of cryoglobulinemia vasculitis were skin involvement (84.2%), arthralgia (61.4%), peripheral neuropathy (54.4%) and glomerulonephritis (31.6%).

The main indication for Rituximab therapy was non-responsiveness to other previous treatments (n=50), intolerance to previous treatments (n=3), associated lymphoma (n=2), or first-line therapy for cryoglobulinemia vasculitis (n=2).

Most patients (48/57) received four weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of Rituximab. In other cases (9/57), however, the Rituximab treatment protocol was different. In the Rocatello's series (13), 5 out of 6 patients received four weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of Rituximab followed by two more infusions at months 1 and 2. In the patient reported by Ghobrial et al. [15], who also had Waldenström's disease, eight weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of rituximab were given. In the case reported by Koukoulaki et al. [19], the patient received four weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of Rituximab, followed by two 1 gram infusions at months 4 and 4.5. In the case reported by Lamprecht et al. [14], 500 mg of Rituximab was given every three weeks for a total of six infusions. The patient of Ghijssels et al. [16] received four weekly consecutive IV infusions of 375 mg/m<sup>2</sup> of Rituximab, followed by a second series of four courses of Rituximab at the same dosage two months later.

The mean duration of follow-up after Rituximab therapy was 9.7 months (0.3 to 24). The evolution of the cryoglobulinemia vasculitis features after Rituximab infusions are given in **Table 2**. Rituximab infusions had great efficacy on the main vasculitis signs, with a clinical response (partial + complete/total) in 32/40 (80%) patients for skin involvement, 27/34 (79.4%) for arthralgia, 27/29 (93.1%) for neuropathy, and 15/18 (83.3%) for glomerulonephritis. However, a relapse of cryoglobulinemia vasculitis was noted in 14/36 (39%) patients (13 HCV-infected, 1 HCV negative) within a few days to nineteen months (mean 6.7 months) after the last Rituximab infusion. Eight out of 14 relapsers had complete remission after a second course of Rituximab. Baseline plus follow up HCV viral load was available in only five reported patients, whereas in the large study of Sansonno the mean HCV viral load increased at month 6 and month 12

[16]. There was no significant difference in the efficacy of Rituximab therapy when patients presented with HCV-induced or essential cryoglobulinemia vasculitis (*Table 3*).

A relatively small number of side effects were reported. During the short term follow-up they included: bradycardia (3), hypotension (2), infection (in three renal transplant patients), mild ALT elevation (3), retinal arterial thrombosis (1), panniculitis of elbows and knees (1), and serum sickness (1). Two deaths were reported; one occurred twelve months after Rituximab infusion in an HCV-infected patient with renal insufficiency, and the second occurred two months after Rituximab infusion in an HCV-negative renal transplant patient due to *Cryptococcus neoformans* meningoencephalitis [18]. During the long term (>12 months), there were two cases of lymphoma and one of breast cancer [11].

## Discussion

Although this literature review does not allow a definite conclusion to be drawn regarding the role of Rituximab in the therapeutic options of patients presenting with cryoglobulinemia vasculitis, it gives important information concerning its efficacy and short term tolerance. When we analyzed the efficacy, a complete response was reported in one-third to two-thirds of patients, depending on the main targeted organ (skin, nerve, kidney, joint). Complete plus partial responses were noted in 80% to 93% of reported patients. Such results are encouraging considering that most patients presented with a very severe form of systemic cryoglobulinemia vasculitis which was resistant or intolerant to other previous treatments. Such a disease remains life threatening, with a mortality rate ranging from 15% to 20% in HCV-MC patients and up to 50% in non HCV-MC patients [12]. The main predictors of poor outcome are older age and renal insufficiency (present in one-third of the cases treated with Rituximab).

Most reported cases (75%) were secondary to HCV infection. There has been some debate about the best therapeutic management of HCV MC patients, i.e. anti-HCV, immunomodulating drugs or both [6]. It is noteworthy that in one of the two large open studies using Rituximab [12], HCV MC patients were described as “non responders or intolerant to interferon alpha monotherapy”. Such treatment, however, is known to have a poor response on the main symptoms of cryoglobulinemia vasculitis, i.e. less than 10% except for purpuric skin lesions [5, 6]. This poor efficacy is related to the low percentage (less than 15%) of long term sustained viral clearance. More recent data with the best available HCV treatment (peg interferon alpha plus ribavirin), which results in a sustained viral clearance in up to 55% of patients, showed much better responses, with a complete clinical response of cryoglobulinemia vasculitis in up to 77% of patients [7]. However, the poor tolerability of HCV therapy often leads to decrease the doses or duration of treatment with lower clinical and virological responses. Using Rituximab, a clinical response was noted in two-thirds of patients, but 39% of them developed a relapse after a mean delay of 6.7 months. In this setting, Rituximab cannot be seen as a curative treatment as long as the viral antigenic trigger of the vasculitis remains. A sequential treatment strategy may therefore give better results. This would begin with Rituximab to block the B-cell activation and the TH1 cytokine-chemokine cascade involved in the inflammatory part of vasculitis [23], and be followed by anti-HCV treatment with pegylated interferon alpha plus ribavirin to clear the antigenic trigger and block the causative agent [12]. Such a strategy may also allow major side effects due to conventional treatments such as corticosteroids, immunosuppressive drugs and plasma exchanges to be obviated.

Due to the very small number of patients with non-HCV systemic cryoglobulinemia vasculitis, it is difficult to come to a clear conclusion in this setting,

although Rituximab therapy seems to have similar efficacy on the main involved organs (**Table 3**). This form of the disease is close to non-Hodgkin's lymphoma and may also be sensitive to CD20 blockade. Furthermore, limitations due to HCV disease (viremia) will be absent, suggesting that autoimmune-disease-associated MC would be the perfect situation to use Rituximab.

The safety of Rituximab monotherapy has been well established in patients with non-Hodgkin's lymphoma, with the main adverse events being mild to moderate infusion-related reactions occurring after the first infusion. The database of patients having received Rituximab in clinical trials and/or clinical practice now includes almost 750,000 patients, making Rituximab the only antibody in this setting to have such an extensive safety analysis [24, 25]. Therefore, we were not surprised at the small number of side effects reported in patients with cryoglobulinemia vasculitis. However, caution is still warranted in this latter indication for the following reasons: no control trial was available, all data came from only small series or case reports, most cases reported on short-term follow-up, and in a large number of cases there was insufficient data to analyse, particularly regarding the course of HCV viral load and liver enzymes. Further assessment of the long-term effects of prolonged B-cell depletion is still required.

We acknowledge some limitations to our analysis. First, there is a potential role that negative publication bias may play in the currently favourable view of Rituximab therapy for MC in the literature. Most studies employing Rituximab do not indicate what types of glucocorticoid regimens were used concomitantly with the Rituximab. This gives the potentially misleading impression that Rituximab alone might have been responsible for the beneficial effects reported. Outside of the 2 series, the remaining reports are small, and different with regards to patient population, Rituximab regimen, outcome definitions, and use of concurrent therapies. Most published cases reported on short or mid term follow up, and we need to know on the long-term effects of prolonged B-cell depletion.

**In conclusion**, Rituximab therapy for patients with mixed cryoglobulinemia vasculitis, HCV-induced or essential, shows a great efficacy on the main vasculitis signs in the majority of reported patients. A relapse of cryoglobulinemia vasculitis was frequently noted. Randomised controlled trials with long term study are necessary to conclude definitively on the benefit/risk ratio of Rituximab therapy in such patients.

**Table 1: Main baseline characteristics of patients with Cryoglobulinemia Vasculitis who received anti-CD20 antibody (Rituximab) treatment.**

	Number of patients with available data	Number of positive patients	Percentage of positive patients
Age years, mean (range),	57	-	59 (31-79)
Sex (F) %	57	45	79 %
<b>Vasculitis</b>			
Duration months, mean (range)	57	-	60.1 (6-240)
Skin involvement	57	48	84.2 %
Arthralgia	57	35	61.4 %
Neuropathy	57	31	54.4 %
Glomerulonephritis	57	18	31.6 %
<b>Immunology</b>			
Cryoglobulin positive	57	57	100 %
Type I		2	3.5 %
Type II		41	71.9 %
Type III		10	17.6 %
Type unknown		4	7.0 %
Rheumatoid factor positive	57	30	52.6 %
C4 serum level (mg/dl), mean	57	-	7.1
<b>HCV status</b>	57		
HCV RNA negative or unknown		14	24.6 %
HCV RNA positive		43	75.4 %
genotype 1-4		24	55.8 %
genotype 2-3		18	41.9 %
genotype not available		1	2.3 %
viral load > 2 million IU/mL	8	6	75.0 %
ALT (IU/l), mean	31	-	54.3
<b>Previous treatment</b>			
<b>HCV infection</b>	37		
Interferon alpha		27	72.8 %
Pegylated Interferon alpha plus Ribavirin		4	14.8 %
None		12	32.4 %
<b>Vasculitis</b>			
Corticosteroids	36*	31	86.1 %
Immunosuppressive drug	56	18	32.1 %
Plasma exchange	56	12	21.4 %

\* This number is underestimated because patients reported in the series of Sansonno et al [12] were not included due to the lack of details (“most patients were previously given low to moderate doses of corticosteroids”)



**Table 2: Main course of Cryoglobulinemia Vasculitis features after anti-CD20 antibody (Rituximab) infusion.**

	Number of patients positive at baseline	Number of patients with available data at follow up	Percentage of patients with available data at follow up
<b>Vasculitis</b>			
<b><i>Skin involvement</i></b>	48	40	
Complete response (CR)	-	27	67.5 %
Partial response (PR)	-	5	12.5 %
Non response (NR)	-	8	20.0 %
<b><i>Arthralgia</i></b>	35	34	
CR	-	18	52.9 %
PR	-	9	26.5 %
NR	-	7	20.6 %
<b><i>Neuropathy</i></b>	31	29	
CR	-	9	31.0 %
PR	-	18	62.1 %
NR	-	2	6.9 %
<b><i>Glomerulonephritis</i></b>	18	18	
CR	-	12	66.6 %
PR	-	3	16.7 %
NR	-	3	16.7 %
<b>Cryoglobulin</b>	57	22*	
CR	-	16	72.7 %
PR	-	2	9.1 %
NR	-	4	18.2 %
<b>Follow up after Rituximab therapy</b>			
Duration months, mean (range)	57	56	9.7 (0.3-24)
Relapses	-	14/36	39 %

\*The serum cryoglobulin status at the end of follow-up was available in 22 patients.

**Table 3: Main course of Cryoglobulinemia Vasculitis features after anti-CD20 antibody (Rituximab) infusion in patients with or without HCV infection.**

	HCV+ patients		HCV- patients	
	N = 43		N = 14	
	Number*	%	Number*	%
<b>Skin involvement</b>	33		7	
Complete response (CR)	24	73	3	42
Partial response (PR)	3	9	2	29
Non response (NR)	6	18	2	29
<b>Arthralgia</b>	30		4	
CR	16	53	2	50
PR	8	27	1	25
NR	6	20	1	25
<b>Neuropathy</b>	25		4	
CR	9	36	0	0
PR	14	56	4	100
NR	2	8	0	0
<b>Glomerulonephritis</b>	13		5	
CR	9	70	4	80
PR	2	15	0	0
NR	2	15	1	20
<b>Cryoglobulin</b>	15		7	
CR	11	73.4	5	71.4
PR	2	13.3	0	0
NR	2	13.3	2	28.6

\* The reported numbers correspond to patients whose data were available at the end of follow-up.

## REFERENCES

1. Gorevic PD, Kassab HJ, Levo Y, Kohn R, Meltzer M, Prose P, *et al.* Mixed cryoglobulinemia: clinical aspects and long-term follow-up of 40 patients. *Am J Med* 1980;**69**:287-308.
2. Brouet JC, Clauvel JP, Danon F, Klein M, and Seligmann M. Biologic and clinical significance of cryoglobulins. A report of 86 cases. *Am J Med* 1974;**57**:775-88.
3. Cacoub P, Poynard T, Ghillani P, Charlotte F, Olivi M, Piette JC, *et al.* Extrahepatic manifestations of chronic hepatitis C. MULTIVIRC Group. Multidepartment Virus C. *Arthritis Rheum* 1999;**42**:2204-12.
4. Sene D, Limal N, and Cacoub P. Hepatitis C virus-associated extrahepatic manifestations: a review. *Metab Brain Dis* 2004;**19**:357-81.
5. Misiani R, Bellavita P, Fenili D, Vicari O, Marchesi D, Sironi PL, *et al.* Interferon alfa-2a therapy in cryoglobulinemia associated with hepatitis C virus. *N Engl J Med* 1994;**330**:751-6.
6. Cacoub P, Saadoun D, Sene D, Limal N, and Piette JC. Treatment of hepatitis C virus-related systemic vasculitis. *J Rheumatol* 2005;**32**:2078-82.
7. Cacoub P, Saadoun D, Limal N, Sene D, Lidove O, and Piette JC. PEGylated interferon alfa-2b and ribavirin treatment in patients with hepatitis C virus-related systemic vasculitis. *Arthritis Rheum* 2005;**52**:911-5.
8. Zaja F, Russo D, Fuga G, Patriarca F, Ermacora A, and Baccarani M. Rituximab for the treatment of type II mixed cryoglobulinemia. *Haematologica* 1999;**84**:1157-8.
9. Zaja F, De Vita S, Russo D, Michelutti A, Fanin R, Ferraccioli G, *et al.* Rituximab for the treatment of type II mixed cryoglobulinemia. *Arthritis Rheum* 2002;**46**:2252-4; author reply 2254-5.
10. Arzoo K, Sadeghi S, and Liebman HA. Treatment of refractory antibody mediated autoimmune disorders with an anti-CD20 monoclonal antibody (rituximab). *Ann Rheum Dis* 2002;**61**:922-4.
11. Zaja F, De Vita S, Mazzaro C, Sacco S, Damiani D, De Marchi G, *et al.* Efficacy and safety of rituximab in type II mixed cryoglobulinemia. *Blood* 2003;**101**:3827-34.
12. Sansonno D, De Re V, Lauletta G, Tucci FA, Boiocchi M, and Dammacco F. Monoclonal antibody treatment of mixed cryoglobulinemia resistant to interferon alpha with an anti-CD20. *Blood* 2003;**101**:3818-26.
13. Roccatello D, Baldovino S, Rossi D, Mansouri M, Naretto C, Gennaro M, *et al.* Long-term effects of anti-CD20 monoclonal antibody treatment of cryoglobulinaemic glomerulonephritis. *Nephrol Dial Transplant* 2004;**19**:3054-61.

14. Lamprecht P, Lerin-Lozano C, Merz H, Dennin RH, Gause A, Voswinkel J, *et al*. Rituximab induces remission in refractory HCV associated cryoglobulinaemic vasculitis. *Ann Rheum Dis* 2003;**62**:1230-3.
15. Ghobrial IM, Uslan DZ, Call TG, Witzig TE, and Gertz MA. Initial increase in the cryoglobulin level after rituximab therapy for type II cryoglobulinemia secondary to Waldenstrom macroglobulinemia does not indicate failure of response. *Am J Hematol* 2004;**77**:329-30.
16. Ghijssels E, Lerut E, Vanrenterghem Y, and Kuypers D. Anti-CD20 monoclonal antibody (rituximab) treatment for hepatitis C-negative therapy-resistant essential mixed cryoglobulinemia with renal and cardiac failure. *Am J Kidney Dis* 2004;**43**:e34-8.
17. Catuogno M, Rezai S, Priori R, Magrini L, and Valesini G. Serum sickness associated with rituximab in a patient with hepatitis C virus-related mixed cryoglobulinaemia. *Rheumatology (Oxford)* 2005;**44**:406.
18. Basse G, Ribes D, Kamar N, Mehrenberger M, Esposito L, Guitard J, *et al*. Rituximab therapy for de novo mixed cryoglobulinemia in renal transplant patients. *Transplantation* 2005;**80**:1560-4.
19. Koukoulaki M, Abeygunasekara SC, Smith KG, and Jayne DR. Remission of refractory hepatitis C-negative cryoglobulinaemic vasculitis after rituximab and infliximab. *Nephrol Dial Transplant* 2005;**20**:213-6.
20. Nehme-Schuster H, Korganow AS, Pasquali JL, and Martin T. Rituximab inefficiency during type I cryoglobulinaemia. *Rheumatology (Oxford)* 2005;**44**:410-1.
21. Quartuccio L, Soardo G, Romano G, Zaja F, Scott CA, De Marchi G, *et al*. Rituximab treatment for glomerulonephritis in HCV-associated mixed cryoglobulinaemia: efficacy and safety in the absence of steroids. *Rheumatology (Oxford)* 2006;**45**:842-6.
22. Cai FZ, Ahern M, and Smith M. Treatment of cryoglobulinemia associated peripheral neuropathy with rituximab. *J Rheumatol* 2006;**33**:1197-8.
23. Saadoun D, Bieche I, Maisonobe T, Asselah T, Laurendeau I, Piette JC, *et al*. Involvement of chemokines and type 1 cytokines in the pathogenesis of hepatitis C virus-associated mixed cryoglobulinemia vasculitis neuropathy. *Arthritis Rheum* 2005;**52**:2917-25.
24. Ghielmini M, Rufibach K, Salles G, Leoncini-Francini L, Leger-Falandry C, Cogliatti S, *et al*. Single agent rituximab in patients with follicular or mantle cell lymphoma: clinical and biological factors that are predictive of response and event-free survival as well as the effect of rituximab on the immune system: a study of the Swiss Group for Clinical Cancer Research (SAKK). *Ann Oncol* 2005;**16**:1675-82.

25. Solal-Celigny P. Safety of rituximab maintenance therapy in follicular lymphomas. *Leuk Res* 2006;**30 Suppl 1**:S16-21.



## Anti-CD20 monoclonal antibody (rituximab) treatment for cryoglobulinemia vasculitis: where do we stand?

Patrice Cacoub, Aurélien Delluc, David Saadoun, Dan Avi Landau and Damien Sene

*Ann Rheum Dis* published online July 20, 2007

---

Updated information and services can be found at:  
<http://ard.bmj.com/content/early/2007/07/20/ard.2006.065565>

---

*These include:*

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (637)

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>