

Accepted Elasticity in Local Arithmetic Congruence Monoids

Lorin Crawford · Vadim Ponomarenko ·
Jason Steinberg · Marla Williams

Abstract For certain $a, b \in \mathbb{N}$, the *Arithmetic Congruence Monoid* $M(a, b)$ is a multiplicatively closed subset of \mathbb{N} given by $\{x \in \mathbb{N} : x \equiv a \pmod{b}\} \cup \{1\}$. An *irreducible* in this monoid is any element that cannot be factored into two elements, each greater than 1. Each monoid element (apart from 1) may be factored into irreducibles in at least one way. The *elasticity of a monoid element* (apart from 1) is the length of the longest factorization into irreducibles, divided by the length of the shortest factorization into irreducibles. The *elasticity of the monoid* is the supremum of the elasticities of the monoid elements. A monoid has *accepted elasticity* if there is some monoid element that has the same elasticity as the monoid. An Arithmetic Congruence Monoid is *local* if $\gcd(a, b)$ is a prime power (apart from 1). It has already been determined whether Arithmetic Congruence Monoids have accepted elasticity in the non-local case; we make significant progress in the local case, i.e. for many values of a, b .

Keywords non-unique factorization · arithmetical congruence monoid · accepted elasticity · elasticity of factorization

Mathematics Subject Classification (2000) 20M14 · 20D60 · 13F05

Lorin Crawford
Clark Atlanta University

Vadim Ponomarenko
San Diego State University, San Diego, CA 92182-7720
E-mail: vponomarenko@mail.sdsu.edu

Jason Steinberg
Princeton University

Marla Williams
Willamette University

1 Introduction

Factorization theory studies the arithmetic properties of domains or commutative, cancellative monoids where unique factorization fails to hold. For a reference see any of the recent works [1, 2, 9, 12]. The present work determines a standard arithmetic invariant for a particular type of monoid. Previous work in this direction left a significant gap, and we close much of this gap. Unfortunately, the problem becomes complex so it appears to be quite difficult to close the gap completely.

Let \mathbb{N} denote the set of positive integers, and \mathbb{N}_0 denote the set of non-negative integers. Let $a, b \in \mathbb{N}$ with $a < b$ and $a^2 \equiv a \pmod{b}$. Set $M(a, b) = \{x \in \mathbb{N} : x \equiv a \pmod{b}\} \cup \{1\}$. This set is a monoid under multiplication. Such sets are called *arithmetic congruence monoids*, and their arithmetic has received considerable attention recently [3–8, 11, 13, 15]. If $\gcd(a, b) = 1$, then the ACM is a Krull monoid, whose arithmetic is well-studied (see [10]). The case where $\gcd(a, b)$ is neither 1 nor a prime power was resolved in [5] with regards to the issue of accepted elasticity. Consequently, we restrict our attention to the remaining case wherein $\gcd(a, b)$ is a prime power, in which case $M(a, b)$ is called a *local* (singular) arithmetic congruence monoid. Specifically, we consider the local arithmetic congruence monoid, henceforth ACM, given as $M = M(p^\alpha \xi, p^\alpha n)$, for some $\xi, n, p, \alpha \in \mathbb{N}$ with p prime and $\gcd(\xi, n) = \gcd(p, n) = 1$.

For monoid M , we say nonunit $x \in M$ is *irreducible* if there are no factorizations $x = y \cdot z$ where y, z are nonunits from M . ACM's are examples of C-monoids (for a reference see the monograph [12]); consequently each nonunit $x \in M = M(p^\alpha \xi, p^\alpha n)$ has at least one factorization into irreducibles. Set $\mathcal{L}(x) = \{n \mid x = x_1 x_2 \cdots x_n, \text{ with each } x_i \text{ irreducible in } M\}$. We define the *elasticity* of x , denoted $\rho(x)$, as $\frac{\max \mathcal{L}(x)}{\min \mathcal{L}(x)}$. We define the elasticity of M as the supremum of $\rho(x)$ over all nonunits $x \in M$. If the supremum is actually a maximum, i.e. if there is some $x \in M$ where $\rho(x) = \rho(M)$, we say that M has *accepted elasticity*. Our goal is to extend the work in [6] in determining which ACM's have accepted elasticity. We will show that the answer depends on the (multiplicative) group structure of \mathbb{Z}_n^\times , and on the cyclic subgroup generated by the element $[p] \in \mathbb{Z}_n^\times$. Broadly, if this subgroup has “large” index, elasticity will be accepted for all or almost all α . Otherwise, the answer is more complicated.

We now recall some standard notation from nonunique factorization theory. Let G be a finite abelian group. Although in our context we write G multiplicatively, our definitions will be compatible with the traditional ones in which groups are written additively. We use $\mathcal{F}(G)$ to denote the set of all finite length (unordered) sequences with terms from G , refer to the elements of $\mathcal{F}(G)$ as sequences, and write all sequences multiplicatively, so that a sequence $S \in \mathcal{F}(G)$ is written in the form

$$S = g_1 \cdot g_2 \cdots g_l = \prod_{g \in G} g^{\nu_g(S)}, \text{ with } \nu_g(S) \in \mathbb{N}_0 \text{ for all } g \in G.$$

We call $\nu_g(S)$ the *multiplicity* of g in S . For $d \in \mathbb{N}$, we call

$$S^d = \prod_{g \in G} g^{d\nu_g(S)} \in \mathcal{F}(G) \quad \text{the } d\text{-fold product of } S.$$

The notation $S_1|S$ indicates that S_1 is a subsequence of S , that is $\nu_g(S_1) \leq \nu_g(S)$ for all $g \in G$. For S_1, S_2, \dots, S_m , each a subsequence of S , if

$$\sum_{i=1}^m \nu_g(S_i) = \nu_g(S) \quad \text{for all } g \in G,$$

we write $S_1 S_2 \cdots S_m = S$ and call this a *partition* of S . If instead

$$\sum_{i=1}^m \nu_g(S_i) \leq \nu_g(S) \quad \text{for all } g \in G,$$

we write $S_1 S_2 \cdots S_m|S$ and call this a *subpartition* of S .

For a sequence $S = g_1 \cdot g_2 \cdots g_l = \prod_{g \in G} g^{\nu_g(S)} \in \mathcal{F}(G)$, we call

$$|S| = l = \sum_{g \in G} \nu_g(S) \in \mathbb{N}_0 \quad \text{the length of } S,$$

$$\sigma(S) = \prod_{i=1}^l g_i = \prod_{g \in G} g^{\nu_g(S)} \in G \quad \text{the sum of } S,$$

$$\Sigma(S) = \left\{ \prod_{i \in I} g_i : I \subseteq [1, l], 0 \neq |I| \right\} \subseteq G \quad \text{the set of subsequence sums of } S,$$

$$\text{and } \Sigma'(S) = \left\{ \prod_{i \in I} g_i : I \subseteq [1, l], 0 \neq |I| \neq l \right\} \subseteq G$$

the *set of proper subsequence sums* of S .

For fixed n , let $x \in \mathbb{Z}$ satisfy $\gcd(x, n) = 1$. We denote by $[x]$ the equivalence class in \mathbb{Z}_n^\times containing x . We define the valuation $\nu_p(x)$ as the unique integer d such that $p^d|x$ and $p^{d+1} \nmid x$; more generally we will use the same valuation for $p \in G$ and $x \in \mathcal{F}(G)$. The following are elementary results about ACM's that are either found in, or are easy to derive from, the previous ACM papers.

Lemma 1 *Let $M = M(p^\alpha \xi, p^\alpha n)$ be an ACM. Then*

1. For any $u \in \mathbb{N}$, $u \in M \setminus \{1\}$ if and only if $[u] = 1$ and $\nu_p(u) \geq \alpha$.
2. If $u \in M$ is irreducible, then $\alpha \leq \nu_p(u) \leq \alpha + \beta - 1$.
3. $\rho(M) = \frac{\alpha + \beta - 1}{\alpha}$.
4. For any $u \in M$, there are some $a, l \in \mathbb{N}_0$ such that $a \geq \alpha$ and $u = p^\alpha q_1 q_2 \cdots q_l$, where each q_i satisfies $\gcd(q_i, pn) = 1$.
5. We may determine ξ as the unique integer in $[1, n-1]$ satisfying $[\xi] = [p]^{-\alpha}$.
6. Let β be the unique minimal integer satisfying $\beta \geq \alpha$ and $[p]^\beta = [1]$. Then $p^\beta \in M$ and $p^s \notin M$ for all $s \in [1, \beta)$.

Consequently, an ACM $M(p^\alpha \xi, p^\alpha n)$ may be determined by p, α, n alone, and we will write $M(p, \alpha, n)$ for convenience, with ξ and β defined implicitly whenever needed.

2 Configurations

Our primary tool in determining whether an ACM has accepted elasticity will be the study of configurations, as defined below.

Let G be a finite abelian group, and let $g \in G$. We denote the order of g in G by $|g|_G$, or $|g|$ when unambiguous.

Definition 1 Let G be a finite abelian group. Let $g \in G$. Let $\delta, \gamma \in \mathbb{N}$ satisfy $\delta \geq |g| > \gamma \geq 0$. Suppose that there is some sequence $S \in \mathcal{F}(G)$ and some $c, d \in \mathbb{N}$ with $\frac{c}{d} \geq 1 + \frac{\delta-1}{\delta-\gamma}$ satisfying

1. There is some partition $S_1 S_2 \cdots S_d = S$ such that for each $i \in [1, d]$,
 - (a) $\sigma(S_i) = g^{\gamma+1}$, and
 - (b) $\Sigma(S_i) \cap \{g, g^2, \dots, g^\gamma\} = \emptyset$; and also
2. There is some subpartition $T_1 T_2 \cdots T_c | S$, satisfying $\sigma(T_i) = g^\gamma$ for each $i \in [1, c]$.

We call this sequence, partition, and subpartition a (G, g, δ, γ) -configuration.

Note that if (c, d) satisfy the conditions, then so do (kc, kd) for each $k \in \mathbb{N}$, by considering the subpartition $T_1^k T_2^k \cdots T_c^k | S^k = S_1^k S_2^k \cdots S_d^k$. Hence we will typically assume without loss of generality that $(\delta - \gamma) | d$.

The connection between (G, g, δ, γ) -configurations and accepted elasticity in ACMs, is given by the following.

Theorem 1 Let $M = M(p, \alpha, n)$ be an ACM. Then M has accepted elasticity if and only if there exists a $(\mathbb{Z}_n^\times, [p], \beta, \beta - \alpha)$ -configuration.

Proof Suppose first that M has accepted elasticity. Then there is some pair of factorizations into irreducibles $u_1 u_2 \cdots u_s = v_1 v_2 \cdots v_t$ with $\frac{s}{t} = \frac{\alpha + \beta - 1}{\alpha} = \rho(M)$. By Lemma 1, $s\alpha \leq \sum_{i=1}^s \nu_p(u_i) = \sum_{i=1}^t \nu_p(v_i) \leq t(\alpha + \beta - 1)$. All inequalities are therefore equalities, so $\nu_p(u_i) = \alpha, \nu_p(v_i) = \alpha + \beta - 1$ for all i .

Express each $v_i = p^{\alpha + \beta - 1} q_1^{(i)} q_2^{(i)} \cdots q_{l_i}^{(i)}$ as in Lemma 1. For each $i \in [1, s]$, we define a sequence from \mathbb{Z}_n^\times given by $S_i = [q_1^{(i)}][q_2^{(i)}] \cdots [q_{l_i}^{(i)}]$. We have $[1] = [v_i] = [p]^{\alpha + \beta - 1} \sigma(S_i)$, so $\sigma(S_i) = [p]^{\beta - \alpha + 1}$. Suppose there were a subsequence $T | S_i$ with $\sigma(T) = [p]^x$ for some $x \in [1, \beta - \alpha]$. Then we set $v'_i = p^{\beta - x} \prod q_j^{(i)}$, where the product is taken over all $[q_j^{(i)}] \in T$. We set $v''_i = \frac{v_i}{v'_i}$. We have $\nu_p(v'_i) \geq \alpha$ and $\nu_p(v''_i) = \alpha + x - 1 \geq \alpha$. Further $[v'_i] = [p]^{\beta - x} \sigma(T) = [p]^\beta = [1]$. Since $[1] = [v'_i v''_i] = [v'_i][v''_i]$, also $[v''_i] = 1$. Hence $v'_i, v''_i \in M$, which contradicts the irreducibility of v_i . Therefore, the S_i each satisfy the conditions of Definition 1.1. Set $S = S_1 S_2 \cdots S_t$.

Express each $u_i = p^\alpha r_1^{(i)} r_2^{(i)} \cdots r_{l_i}^{(i)}$ as in Lemma 1. For each $i \in [1, t]$, we define a sequence from \mathbb{Z}_n^\times given by $T_i = [r_1^{(i)}][r_2^{(i)}] \cdots [r_{l_i}^{(i)}]$. We have $[1] = [u_i] = [p]^\alpha \sigma(T_i)$, so $\sigma(T_i) = [p]^{-\alpha} = [p]^{\beta - \alpha}$. By unique factorization in \mathbb{N} , in fact $T_1 T_2 \cdots T_s = S$. Thus, $T_1 \cdots T_s$ is a partition (and hence a subpartition) of S . It remains to observe that $\frac{s}{t} = \frac{\alpha + \beta - 1}{\alpha} = 1 + \frac{\beta - 1}{\beta - (\beta - \alpha)}$.

Suppose now that there exists a $(\mathbb{Z}_n^\times, [p], \beta, \beta - \alpha)$ -configuration. Define $\phi : \mathbb{Z}_n^\times \rightarrow \mathbb{N}$ such that $\phi([x]) = q_x$ for some prime $q_x \neq p$ satisfying $[q_x] = [x]$. Such a ϕ exists by Dirichlet's theorem on primes. We now set $v_i = p^{\alpha+\beta-1} \prod_{x \in S_i} \phi([x])$ for $i \in [1, m]$. Note that $[v_i] = [p]^{\alpha+\beta-1} \sigma(S_i) = [p]^{\alpha+\beta-1} [p]^{\beta-\alpha+1} = [1]$, so $v_i \in M$. Suppose that v_i were reducible with factor v'_i . We have $v'_i = p^x \sigma(T)$ for some $x \geq \alpha$ and some $T | S_i$. We have $[1] = [v'_i] = [p]^x \sigma(T)$, so $\sigma(T) = [p]^{\beta-x}$, which is a contradiction. Hence each $v_i \in M$ is irreducible.

The second property gives us $\frac{c}{d} \geq 1 + \frac{\beta-1}{\beta-(\beta-\alpha)} = \frac{\alpha+\beta-1}{\alpha}$. We assume without loss that $\alpha | d$, and set $c' = \lfloor d(\frac{\alpha+\beta-1}{\alpha}) \rfloor = (\frac{d}{\alpha})(\alpha + \beta - 1)$. For $i \in [1, c' - 1] \subseteq [1, c]$, we take $u_i = p^\alpha \prod_{x \in T_i} \phi([x])$, and set $u_{c'} = \frac{S}{[u_1][u_2] \cdots [u_{c'-1}]}$. We have $[u_i] = [p]^\alpha \sigma(T_i) = [p]^\alpha [p]^{\beta-\alpha} = [1]$, so $u_i \in M$ for $i \in [1, c' - 1]$. Set $u = v_1 v_2 \cdots v_m = u_1 u_2 \cdots u_{c'}$. We have $[1] = [u] = [u_1][u_2] \cdots [u_{c'-1}][u_{c'}]$, so $[u_{c'}] = [1]$. Further, since $\alpha c' = d(\alpha + \beta - 1) = \nu_p(u) = (c' - 1)\alpha + \nu_p(u_{c'})$ we have $\nu_p(u_{c'}) = \alpha$. Hence $u_{c'} \in M$.

Finally, we have $\rho(u) \geq \frac{c}{d} \geq \frac{\alpha+\beta-1}{\alpha} = \rho(M)$, so M has accepted elasticity. \square

We now broadly outline the remainder of this paper. In the subsequent sections, we will find that if $G/\langle g \rangle$ is "large", then configurations will exist for all γ , provided that δ is sufficiently large. That is, if we fix p, n , then $M(p, \alpha, n)$ has accepted elasticity for all but finitely many α . However, if $G/\langle g \rangle$ is "small", then configurations will exist for "small" gamma and will not exist for "large" gamma (keeping in mind that $\gamma \in [1, |g| - 1]$). That is, if we again fix p, n , then $M(p, \alpha, n)$ has accepted elasticity for infinitely many α , and does not have accepted elasticity for infinitely many α , with the categorization occurring based on the congruence class α falls into modulo $\phi(n)$, the Euler totient.

3 Finding Configurations

We first present some results that produce (G, g, δ, γ) -configurations in certain special cases. Recall that by Theorem 1, we are only concerned with δ that are multiples of $|g|$. The following proposition, in the context of ACMs, states that $M(p, \alpha, n)$ has accepted elasticity, provided that $\alpha = \beta$. For other equivalent conditions, see Proposition 2.

Proposition 1 *Let G be any finite abelian group. Let $g \in G$, and let $\delta \in \mathbb{N}$ satisfy $\delta \geq |g|$. Then there is a $(G, g, \delta, 0)$ -configuration.*

Proof Set $d = 1$, and set $S = S_1 = (g)$. We have $\sigma(S_1) = g^{0+1}$, while $\{g, g^2, \dots, g^\gamma\} = \emptyset$. For the second condition, we take $c = \lceil 1 + \frac{\delta-1}{\delta} \rceil$ and set $T_1 = T_2 = \cdots = T_c = \emptyset$, which gives $\sigma(T_i) = 1 = g^0$. \square

Consequently, we will assume henceforth that $\gamma > 0$ and $\beta > \alpha$. By the following proposition, we equally assume that $\xi > 1$ and $\rho(M) \geq 2$.

Proposition 2 *Given ACM M , the following are equivalent: (1) $\xi = 1$; (2) $[p]^\alpha = [1]$; (3) $\beta = \alpha$; and (4) $\rho(M) < 2$.*

Proof If (1) holds, since $[\xi] = [p]^{-\alpha}$, in fact $[1] = [p]^\alpha$, so (2) holds. If (2) holds, since $\alpha \geq \alpha$ and $[p]^\alpha = [1]$, in fact $\beta = \alpha$, so (3) holds. If (3) holds, then $[\xi] = [p]^{-\alpha} = [p]^{-\beta} = [1]$. Because $1 \leq \xi \leq n-1$, in fact $\xi = 1$, so (1) holds. If (3) holds, then $\rho(M) = \frac{\beta+\alpha-1}{\alpha} = 2 - \frac{1}{\beta} < 2$, so (4) holds. Lastly, if (4) holds, then $\frac{\beta+\alpha-1}{\alpha} < 2$, so $\beta - 1 < \alpha \leq \beta$, so (3) holds. \square

The following proposition, in the context of ACMs, states that if $M(p, \alpha, n)$ has accepted elasticity, then so does $M(p, \alpha + t, n)$ for all $t \in \mathbb{N}$ satisfying $[p]^t = [1]$.

Proposition 3 *Suppose that there is a (G, g, δ, γ) -configuration with $\gamma \geq 1$. Let $\delta' \in \mathbb{N}$ with $\delta' > \delta$. Then there is a (G, g, δ', γ) -configuration.*

Proof We will show that the same configuration works. Because δ only appears in relation to c and d , we only need to check that inequality. Because $\gamma \geq 1$, we have $\frac{\delta-1}{\delta-\gamma} \geq \frac{\delta'-\delta}{\delta'-\delta}$. Their median is $\frac{\delta'-1}{\delta'-\gamma}$, which must be between these fractions and thus no more than $\frac{\delta-1}{\delta-\gamma}$. Consequently, $\frac{c}{d} \geq 1 + \frac{\delta-1}{\delta-\gamma} \geq 1 + \frac{\delta'-1}{\delta'-\gamma}$. \square

In the ACM context, the combination of the previous proposition with the following, states that if $M(p, 1, n)$ has accepted elasticity, then so does $M(p, \alpha, n)$ for all $\alpha \geq 1$.

Proposition 4 *Suppose that there is a $(G, g, |g|, |g| - 1)$ -configuration. Let $\gamma \in \mathbb{N}$ with $\gamma < |g| - 1$. Then there is a $(G, g, |g|, \gamma)$ -configuration.*

Proof Set $k = |g| - \gamma - 1$. We set $S'_i = S_i \cup \{(g^{-1})^k\}$ for $i \in [1, d]$. We have $S' = S'_1 S'_2 \cdots S'_d = SV$ for $V = (g^{-1})^{dk}$. We have $\sigma(S'_i) = g^{|g|-k} = g^{\gamma+1}$. Note that $\Sigma\{(g^{-1})^k\} = \{g^0, g^{-1}, \dots, g^{-k}\}$. Suppose that for some $s \in [1, \gamma]$ we had $g^s \in \Sigma(S'_i) = \Sigma(S_i) \cdot \Sigma\{(g^{-1})^k\}$. Hence $\Sigma(S_i)$ is not disjoint from $\{g^s, g^s, \dots, g^{s+k}\} \subseteq \{g, g^2, \dots, g^{|g|-1}\}$, a contradiction. Therefore $\Sigma(S'_i) \cap \{g, g^2, \dots, g^\gamma\} = \emptyset$.

Without loss, we may assume that $(k+1)|c$ and $(k+1)|d$. For each $i \in [1, \frac{c}{k+1}]$, we set $T'_i = T_{(i-1)(k+1)+1} T_{(i-1)(k+1)+2} \cdots T_{i(k+1)}$. We have $\sigma(T'_i) = [g]^{(k+1)(|g|-1)} = [g]^{-k-1} = [g]^\gamma$. For each $i \in [\frac{c}{k+1} + 1, \frac{c}{k+1} + \frac{kd}{k+1}]$, we set $T'_i = \{(g^{-1})^{k+1}\}$ and again $\sigma(T'_i) = [g]^\gamma$. By hypothesis $\frac{c}{d} \geq 1 + \frac{|g|-1}{|g|-(|g|-1)} = |g|$. Hence $\frac{c}{d} + k \geq |g| + k = (|g| - \gamma) + (|g| - 1) = (|g| - \gamma)(1 + \frac{|g|-1}{|g|-\gamma}) = (k+1)(1 + \frac{|g|-1}{|g|-\gamma})$. Consequently, $\frac{\frac{c}{k+1} + \frac{kd}{k+1}}{d} \geq 1 + \frac{|g|-1}{|g|-\gamma}$. \square

The following proposition, in the context of ACMs, makes clear that not n but \mathbb{Z}_n^\times is the important object.

Proposition 5 *Suppose that there is a (G, g, δ, γ) -configuration with $\gamma \geq 1$. Let $\phi : G \rightarrow H$ be a group homomorphism. Then there is an $(H, \phi(g), \delta, \gamma)$ -configuration.*

Proof For sequence $S = g_1 \cdots g_l \in \mathcal{F}(G)$, we define $\phi(S) = \phi(g_1) \cdots \phi(g_l) \in \mathcal{F}(H)$. We have $\phi(S_1) \cdots \phi(S_d) = \phi(S)$ is a partition that satisfies $\sigma(\phi(S_i)) = \phi(g)^{\gamma+1}$ and $\Sigma(\phi(S_i)) \cap \{\phi(g), \dots, \phi(g)^\gamma\} = \emptyset$. The same c, d as previously satisfy the necessary inequality, with subpartition $\phi(T_1) \cdots \phi(T_c) | \phi(S)$. Finally, we have $\sigma(\phi(T_i)) = \phi(g)^\gamma$. \square

The following proposition, in the context of ACMs, states that $M(p, \alpha, n)$ has accepted elasticity, provided that α is “large” and $|[p]|$ is composite. Specifically, if $|[p]| = rs$ in \mathbb{Z}_n^\times , then we need $\alpha \in (\beta - r, \beta)$. The remaining possibilities for α , namely $(\beta - rs, \beta - r]$, are not covered; however in some cases there are no configurations for these α , as will be shown in Proposition 7.

Proposition 6 *Let G be any finite abelian group. Let $g \in G$. Suppose that $|g| = rs$ with $r, s > 1$ and $rs > 4$. Let $\gamma \in \mathbb{N}$ satisfy $\gamma < r$. Then there is a (G, g, rs, γ) -configuration.*

Proof We first consider the special case $\{s = 2, \gamma = 1\}$; by hypothesis $r \geq 3$. We set $S_1 = (g^{-1})^{2r-2}, S_2 = (g^2)^{2r+1}, T = (g^{-1}) \cdot (g^2)$. We have $\sigma(S_1) = \sigma(S_2) = g^2 = g^{\gamma+1}$ and $\sigma(T) = g^\gamma$. Also, $\Sigma(S_1) = \langle g \rangle \setminus \{1, g\}$ and $\Sigma(S_2) = \langle g^2 \rangle$, which does not contain g since $|g|$ is even. We set $S = S_1 S_2$ and $d = 2$. We set $c = 2r - 2$ and see that $T^c | S$. Lastly we have $\frac{c}{d} = r - 1 \geq 2 = 1 + \frac{2r-1}{2r-1}$.

Henceforth we exclude $\{s = 2, \gamma = 1\}$. Set $S_1 = (g^{-1})^{rs-\gamma-1}$. We have $\sigma(S_1) = g^{\gamma+1-rs} = g^{\gamma+1}$, and $\Sigma(S_1) = \{g^{-1}, g^{-2}, \dots, g^{-rs+\gamma+1}\} = \{g^{\gamma+1}, g^{\gamma+2}, \dots, g^{rs-1}\}$, which has no intersection with $\{g^1, g^2, \dots, g^\gamma\}$. Set $S_2 = (g^r)^{2rs^2} (g^{\gamma+1})$. We have $\sigma(S_2) = g^{\gamma+1}$ and $\Sigma(S_2) = \langle g^r \rangle \cup g^{\gamma+1} \langle g^r \rangle$, which again has no intersection with $\{g^1, g^2, \dots, g^\gamma\}$. We set $d = rs - \gamma$ and $S = S_1^{d-1} S_2$.

We now set $c = s(rs - 2 + \gamma(s - 2)) + 1$. We set $T_0 = (g^{-1}) \cdot (g^{\gamma+1})$ and $T_i = (g^{-1})^{r-\gamma} \cdot (g^r)$ for $i \in [1, c - 1]$. Set $T = T_0 T_1 \cdots T_{c-1}$; we will prove that $T | S$. There are three group elements to consider. First, $(g^{\gamma+1})$ appears once in both T and S . Second, (g^r) appears $2rs^2$ times in S and $c - 1 \leq s(rs + rs) = 2rs^2$ times in T . Lastly, considering (g^{-1}) , we need $(rs - \gamma - 1)^2 \geq 1 + (c - 1)(r - \gamma)$. We chose c so that $(rs - \gamma - 1)^2 - (c - 1)(r - \gamma) = (\gamma(s - 1) - 1)^2$. This integer is zero only when $\gamma = 1$ and $s = 2$, a possibility which has been excluded.

We now prove that $\frac{c}{d} \geq 1 + \frac{rs-1}{rs-\gamma}$. This rearranges to $X \geq 0$, for $X = rs^2 + \gamma s^2 - 2\gamma s - 2s + 2 - 2rs + \gamma = (s - 1)^2 \gamma + s(r(s - 2) - 2) + 2$. If $s \geq 3$ we have $X \geq 4\gamma + 3(r - 2) + 2 \geq 0$; if $s = 2$ we have $X = \gamma - 2 \geq 0$ since $\gamma = 1$ has been excluded. \square

Let G be a nontrivial finite abelian group. Suppose that $g \in G$ generates G , i.e. $G = \langle g \rangle$. It is a well-known result from group theory that if $G \cong \mathbb{Z}_n^\times$ for some n , then $|G| = |g|$ is even. In this situation the following proposition states that the bound of Proposition 6 is tight (provided $|g| > 4$). It also shows that although (G, g, δ, γ) -configurations may be plentiful, they are not omnipresent – not all ACMs have accepted elasticity.

Proposition 7 *Let G be a finite abelian group. Let $g \in G$ satisfy $G = \langle g \rangle$. Suppose that $|g| = 2r \geq 4$. Let $\gamma, \delta \in \mathbb{N}$ satisfy $\delta \geq 2r > \gamma \geq r$. Then there is no (G, g, δ, γ) -configuration.*

Proof Set $A = \{\gamma + 1, \gamma + 2, \dots, 2r\}$ for convenience. We define $\phi : G \rightarrow \mathbb{N}$ via $\phi(g^{-i}) = i$, where we choose $i \in [0, 2r - 1]$. Note that $\phi(ab) \equiv \phi(a) + \phi(b) \pmod{2r}$. We extend ϕ to sequences in the natural way, via $\phi(a \cdot b) = \phi(a) + \phi(b)$. Set $k = 2r - \gamma - 1$, which satisfies $k \in [0, r - 1]$. Suppose there were a (G, g, δ, γ) -configuration. We will first prove that each S_i satisfies $\phi(S_i) = k$. We consider each subsequence $U|S_i$, and prove that $\phi(U) \leq k$ by induction on $|U|$. If $|U| = 1$, then $U = (g^s)$ for some $s \in A$. Hence $\phi(U) = \phi(g^s) \leq k$. Otherwise we write $U = U' \cdot (g^s)$ for some $s \in A$. We have $\sigma(U') = g^t$, where $t \in A$ and $\phi(U') \leq k$ by the inductive hypothesis. Now we have $\phi(U) = \phi(U') + \phi(g^s) \leq 2k$. Because $\phi(U) \equiv \phi(\sigma(U)) \pmod{2r}$, in fact $\phi(U) = \phi(\sigma(U)) = -s - t$. If $\phi(U) > k$ then $-s - t > k = 2r - \gamma - 1$ and $\sigma(U) \in \{g, g^2, \dots, g^\gamma\}$, a contradiction. Hence $\phi(U) \leq k$, and in particular $\phi(S_i) \leq k$. But $\phi(S_i) \geq \phi(\sigma(S_i)) = \phi(g^{\gamma+1}) = k$, so in fact $\phi(S_i) = k$.

Now, $\phi(T_i) \geq \phi(\sigma(T_i)) = \phi(g^\gamma) = k + 1$. Hence we have $dk = d\phi(S_i) = \phi(S) \geq \sum_{i=1}^c \phi(T_i) \geq c(k + 1)$. We rearrange to get $\frac{c}{d} \leq \frac{k}{k+1} < 1 + \frac{\delta-1}{\delta-\gamma}$, a contradiction. \square

We combine Propositions 6 and 7 into the following theorem, which was the main result of [6] (with different proof). It completely solves the special case where p is a primitive root modulo n . In particular, this requires \mathbb{Z}_n^\times to be cyclic, which in the ACM context occurs only when $n = 2, 4, q^k$, or $2q^k$ for some odd prime q .

Theorem 2 *Let G be a finite abelian group. Let $g \in G$ satisfy $G = \langle g \rangle$. Suppose that $|g|$ is even. Let $\delta, \gamma \in \mathbb{N}$ with $\delta \geq |g| > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration if and only if*

1. $|g| > 4$, and
2. $|g| \geq 2\gamma$.

Proof The only cases not covered by Propositions 6 and 7 are the following. $\{|g| = 4, \gamma = 1\}$: Because $\nu_g(S) = 0$, for all i we have $\nu_{g^3}(T_i) \geq 1$, while $\nu_{g^3}(S_i) \leq 2$. Hence we have $2d \geq \nu_{g^3}(S) \geq c$, but also $\frac{c}{d} \geq 1 + \frac{\delta-1}{\delta-1} = 2$. Hence all inequalities are equalities and $\nu_{g^3}(S_i) = 2$ for all i . Then $\nu_{g^2}(S_i) = 0$ for all i , and thus $\nu_{g^2}(S) = 0$. But now $\sigma(T_i) \neq g$, so in fact there is no configuration. $\{|g| = 2, \gamma = 1\}$: Because $\nu_g(S_i) = 0$, we have $\sigma(T_i) \neq g$. \square

4 $\langle g \rangle \oplus H$

With Theorem 2 we have resolved the case of $G = \langle g \rangle$, a cyclic group (provided $|G|$ is even, which holds for all nontrivial $G \cong \mathbb{Z}_n^\times$). Otherwise, $G/\langle g \rangle$ is nontrivial and in the remainder we explore its structure. In this section we consider nontrivial subgroups $H \leq G$ such that $\langle g \rangle \oplus H \leq G$. Such subgroups H need not exist, e.g. for $(G, g) \cong (\mathbb{Z}_{25}, 5)$. However they do exist in two important cases, given by Propositions 8 and 9. We recall first a lemma from the classical theory of finite abelian groups.

Lemma 2 *Let G be a finite abelian group with $|G| = y$. Let $x \in \mathbb{N}$ satisfy $x|y$. Then there is some subgroup $H \leq G$ with $|H| = x$.*

Proof See, e.g., [14, p. 77]. \square

The following proposition allows us to not only address noncyclic groups G , but also cyclic groups G provided that some prime divides $|G|$ but not $|g|$.

Proposition 8 *Let G be a finite abelian group with $g \in G$. Suppose that $|G| = xy$ and $\gcd(x, y) = \gcd(x, |g|) = 1$. Then there is some subgroup $H \leq G$ with $|H| = x$ and $\langle g \rangle \oplus H \leq G$.*

Proof By Lemma 2 there must be some $H \leq G$ with $|H| = x$. Let $z \in \langle g \rangle \cap H$. Then $|z|$ divides both $|g|$ and x , but then $|z| = 1$ so the conclusion follows. \square

Proposition 9 is an elementary result concerning finite abelian groups that seems like it should be well-known, but we have no reference. For noncyclic groups G , it provides a “large” subgroup H such that $\langle g \rangle \oplus H \leq G$. This directly generalizes the well-known result that if $|g| = \exp(G)$, then $\langle g \rangle \oplus (G/\langle g \rangle) \cong G$.

Proposition 9 *Let $G \cong \mathbb{Z}_{n_1} \oplus \mathbb{Z}_{n_2} \oplus \cdots \oplus \mathbb{Z}_{n_k}$ be a finite abelian group, with $n_1|n_2|\cdots|n_k$. Let $g \in G$. Then there is some $H \leq G$ such that $\langle g \rangle \oplus H \leq G$ and $H \cong \mathbb{Z}_{n_1} \oplus \mathbb{Z}_{n_2} \oplus \cdots \oplus \mathbb{Z}_{n_{k-1}}$.*

Proof We first assume that G is a p -group for some prime p , i.e. $G \cong \mathbb{Z}_{p^{a_1}} \oplus \mathbb{Z}_{p^{a_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{a_k}}$, for integers $a_k \geq a_{k-1} \geq \cdots \geq a_1 \geq 1$. We write G additively as k -tuples, and in particular $g = (g_1, g_2, \dots, g_k)$. For each $i \in [1, k]$, let m_i be the order of g_i in $\mathbb{Z}_{p^{a_i}}$. Let M be chosen so that m_M is maximal among $\{m_1, \dots, m_k\}$. By Lagrange’s theorem on finite groups, each m_i is a power of p for all $i \in [1, k]$, so in particular $m_i|m_M$. Hence m_M is the order of g , and therefore each nonzero element of $\langle g \rangle$ has a nonzero element in the M^{th} coordinate. We now set $H = \{(b_1, \dots, b_k) \in G : b_M = 0 \text{ and } p^{a_k - a_M}|b_k\}$, a subgroup of G . We have $\langle g \rangle \cap H = \{0\}$, so $\langle g \rangle \oplus H \leq G$. Further, by swapping the M^{th} and k^{th} coordinates, we see that $H \cong \mathbb{Z}_{n_1} \oplus \mathbb{Z}_{n_2} \oplus \cdots \oplus \mathbb{Z}_{n_{k-1}} \oplus \{0\} \cong \mathbb{Z}_{n_1} \oplus \mathbb{Z}_{n_2} \oplus \cdots \oplus \mathbb{Z}_{n_{k-1}}$.

Suppose now that there are distinct primes p_1, p_2, \dots, p_s and corresponding p -groups G_1, G_2, \dots, G_s , such that $G \cong G_1 \oplus G_2 \oplus \cdots \oplus G_s$. For each $i \in [1, s]$ we have $G_i \cong \mathbb{Z}_{p_i^{a(i,1)}} \oplus \cdots \oplus \mathbb{Z}_{p_i^{a(i,k_i)}}$, for integers $a(i, k_i) \geq \cdots \geq a(i, 1) \geq 1$. By the above, for each $i \in [1, s]$ we find $H_i \leq G_i$ such that $\langle g|_{G_i} \rangle \oplus H_i \leq G_i$ and $H_i \cong \mathbb{Z}_{p_i^{a(i,1)}} \oplus \cdots \oplus \mathbb{Z}_{p_i^{a(i,k_i-1)}}$. Let ϕ_i denote the natural embedding of each p -group G_i into G , and set $H = \phi_1(H_1) + \phi_2(H_2) + \cdots + \phi_s(H_s)$. Because the primes are distinct, in fact $\phi_1(H_1) \oplus \phi_2(H_2) \oplus \cdots \oplus \phi_s(H_s) \leq G$, and also $\langle g \rangle \oplus H \leq G$. We now have $H \cong \prod H_i$, and the result follows since $n_k = \prod_i p_i^{a(i,k_i)}$, $n_{k-1} = \prod_i p_i^{a(i,k_i-1)}$, \dots \square

Theorem 3 is the main result of this section, which requires the following definition.

Definition 2 Let $H \cong \mathbb{Z}_{m_1} \oplus \mathbb{Z}_{m_2} \oplus \cdots \oplus \mathbb{Z}_{m_k}$ be a finite abelian group, where $m_1 | m_2 | \cdots | m_k$. We define $d^*(H) = (m_1 + m_2 + \cdots + m_k) - k = \sum_{i=1}^k (m_i - 1)$.

Theorem 3 Let G be a finite abelian group and $g \in G$. Suppose that there is some $H \leq G$ with $\langle g \rangle \oplus H \leq G$. Let $\delta, \gamma \in \mathbb{N}$ that satisfy $\delta \geq |g| > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration, provided that the following inequality holds:

$$d^*(H) > \left(1 - \frac{1}{|g|}\right) \left(\frac{1}{|g| - \gamma} + \frac{\delta - 1}{\delta - \gamma}\right)$$

Proof We will construct the configuration explicitly. Let $\alpha \in \mathbb{N}$ be large. Let $h_1, \dots, h_k \in G$ with $\langle h_1 \rangle \oplus \cdots \oplus \langle h_k \rangle \oplus \langle g \rangle \leq G$, $|h_i| = m_i$ for $i \in [1, k]$, and $m_1 | m_2 | \cdots | m_k$. Set $S_1 = (g^{-1})^{|g| - \gamma - 1} \cdot \prod_{i=1}^k (h_i g^\gamma)^{m_i - 1} \cdot (h_i^{-1} g^{-\gamma})^{m_i - 1}$, $S_2 = (g^{-1})^{|g| - \gamma - 1} \cdot \prod_{i=1}^k (h_i^{-1})^{|g|^2 m_i^2 \alpha}$. We set $T_0 = (g^{-1})^{|g| - \gamma}$, and for $i \in [1, k]$ set $T_i = (h_i g^\gamma) \cdot (h_i^{-1})$, $T'_i = (h_i^{-1} g^{-\gamma})^{|g| - 1} \cdot (h_i^{-1})^{|g|(m_i - 1) + 1}$. Note that $\sigma(S_1) = \sigma(S_2) = g^{\gamma + 1}$ and for all $i \in [1, k]$, $\sigma(T_i) = \sigma(T'_i) = \sigma(T_0) = g^\gamma$. If $x \in \langle g \rangle \cap (\Sigma(S_1) \cup \Sigma(S_2))$ then in fact $x \in \Sigma((g^{-1})^{|g| - \gamma - 1})$ and consequently $x \notin \{g, g^2, \dots, g^\gamma\}$.

For convenience, set $a_1 = |g| - 1$, $a_\gamma = |g| - \gamma$. We set $d = a_1 a_\gamma \alpha + 1$ and $S = S_1^{a_1 a_\gamma \alpha} S_2$. We set $c = a_1(a_\gamma - 1)\alpha + d^*(H)|g|a_\gamma \alpha$ and $T = T_0^{a_1(a_\gamma - 1)\alpha} \prod_{i=1}^k T_i^{(m_i - 1)a_1 a_\gamma \alpha} T'_i^{(m_i - 1)a_\gamma \alpha}$. We now verify that $T|S$. For g^{-1} , we have $\nu_{g^{-1}}(T) = a_\gamma a_1(a_\gamma - 1)\alpha < (a_\gamma - 1)a_1 a_\gamma \alpha + (a_\gamma - 1) = \nu_{g^{-1}}(S)$. For any $i \in [1, k]$, we have $\nu_{h_i g^\gamma}(T) = (m_i - 1)a_1 a_\gamma \alpha = \nu_{h_i g^\gamma}(S)$. We also have $\nu_{h_i^{-1} g^{-\gamma}}(T) = (m_i - 1)a_1 a_\gamma \alpha = \nu_{h_i^{-1} g^{-\gamma}}(S)$. Lastly we have $\nu_{h_i^{-1}}(T) = (m_i - 1)a_1 a_\gamma \alpha + (m_i - 1)a_\gamma \alpha(|g|(m_i - 1) + 1) = (m_i - 1)m_i a_\gamma |g| \alpha \leq m_i^2 |g|^2 \alpha = \nu_{h_i^{-1}}(S)$. We now calculate $\frac{c}{d} = \frac{a_1(a_\gamma - 1)\alpha + d^*(H)|g|a_\gamma \alpha}{a_1 a_\gamma \alpha + 1} = 1 - \frac{1 + \frac{1}{a_1 \alpha}}{a_\gamma + \frac{1}{a_1 \alpha}} + d^*(H) \frac{|g|}{a_1 + \frac{1}{a_1 \alpha}} = 1 - \frac{1}{a_\gamma} + d^*(H) \frac{|g|}{a_1} + \epsilon$, where we choose α sufficiently large to ensure that $|\epsilon| < \frac{|g|}{a_1} d^*(H) - \left(\frac{1}{a_\gamma} + \frac{\delta - 1}{\delta - \gamma}\right)$, which we may do by hypothesis. Hence we have $\frac{c}{d} > 1 - \frac{1}{a_\gamma} + \frac{1}{a_\gamma} + \frac{\delta - 1}{\delta - \gamma}$, as desired. \square

Recall that in the ACM context we may assume that δ is a positive integer multiple of $|g|$. If we take $\delta = |g|$, the following corollary shows that it suffices to have $d^*(H) > \frac{|g| - 1}{|g| - \gamma}$. If $d^*(H) \geq |g|$ then this condition is met for all γ ; otherwise it is met only for $\gamma < |g| - \frac{|g| - 1}{d^*(H)}$.

Corollary 1 Let G be a finite abelian group and $g \in G$. Suppose that there is some $H \leq G$ with $\langle g \rangle \oplus H \leq G$. Let $\gamma \in \mathbb{N}$ such that $|g| > \gamma > 0$. Suppose that $d^*(H) > \frac{|g| - 1}{|g| - \gamma}$. Then there is a $(G, g, |g|, \gamma)$ -configuration.

Proof With $\delta = |g|$ we have $\left(1 - \frac{1}{|g|}\right) \left(\frac{1}{|g| - \gamma} + \frac{\delta - 1}{\delta - \gamma}\right) = \frac{|g| - 1}{|g| - \gamma}$. \square

Corollary 2 Let G be a finite abelian group, and let $\exp(G)$ denote the exponent of G . Suppose that $d^*(G) \geq 2 \exp(G) - 1$. Then there are (G, g, γ, δ) -configurations for all $g \in G$ and all $\gamma, \delta \in \mathbb{N}$ satisfying $\delta \geq |g| > \gamma > 0$.

Proof Let $g \in G$. Apply Proposition 9 to get $H \leq G$ with $\langle g \rangle \oplus H \leq G$. We have $d^*(H) + \exp(G) - 1 = d^*(G) \geq 2 \exp(G) - 1$, so $d^*(H) \geq \exp(G) \geq |g|$. We now apply Corollary 1. \square

If we exclude the smallest value of δ , namely $|g|$, we only need the weak condition that $d^*(H) \geq 3$ to get all possible γ .

Corollary 3 *Let G be a finite abelian group and $g \in G$. Suppose that there is some $H \leq G$ with $\langle g \rangle \oplus H \leq G$. Let $\delta, \gamma \in \mathbb{N}$ satisfy $\delta \geq 2|g|$ and $|g| > \gamma > 0$. Suppose that $d^*(H) \geq 3$. Then there is a (G, g, δ, γ) -configuration.*

Proof Since $\delta - \gamma > |g| > \gamma - 1$, we have $1 > \frac{\gamma-1}{\delta-\gamma}$. Therefore, $d^*(H) \geq 3 > 2 + \frac{\gamma-1}{\delta-\gamma} = 1 + \frac{\delta-1}{\delta-\gamma} > \left(1 - \frac{1}{|g|}\right) \left(\frac{1}{|g|-\gamma} + \frac{\delta-1}{\delta-\gamma}\right)$. \square

Corollary 3 gives configurations for all γ , provided that $d^*(H) \geq 3$ and δ is sufficiently large. If $d^*(H) = 2$ (i.e. $H \cong \mathbb{Z}_3$ or $\mathbb{Z}_2 \oplus \mathbb{Z}_2$), then Corollary 4 shows that again we get configurations for all γ provided that δ is sufficiently large. If $d^*(H) = 1$ (i.e. $H \cong \mathbb{Z}_2$), then we do not get configurations for all γ , no matter the size of δ , as will be shown later in Proposition 14.

Corollary 4 *Let G be a finite abelian group and $g \in G$. Suppose that there is some $H \leq G$ with $\langle g \rangle \oplus H \leq G$ and $d^*(H) = 2$. Let $\delta, \gamma \in \mathbb{N}$ satisfy $\delta > |g| \frac{|g|-1}{2}$ and $|g| > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration.*

Proof It suffices to prove that $2 > \left(1 - \frac{1}{|g|}\right) \left(\frac{1}{|g|-\gamma} + \frac{\delta-1}{\delta-\gamma}\right)$ for $\gamma = |g| - 1$. This is a rearrangement of $\delta > |g| \frac{|g|-1}{2}$. \square

In the special case of $H = \langle h \rangle$ with $|h| = |g|$, we have $d^*(H) = |g| - 1$. Here Theorem 3 does not apply for $\{\delta = |g|, \gamma = |g| - 1\}$. In fact there is a configuration for this case as well, and hence for all δ, γ by Proposition 4.

Proposition 10 *Let G be a finite abelian group. Let $g, h \in G$ with $\langle g \rangle \oplus \langle h \rangle \leq G$ and $|g| = |h|$. Then there is a $(G, g, |g|, |g| - 1)$ -configuration.*

Proof Set $k = |g|$ for convenience. Set $d = 2$, $S_1 = (hg^{-1})^{2k}$, $S_2 = (h^{-1})^{2k}$, and $S = S_1 S_2$. We have $\sigma(S_1) = (h^k)^2 (g^{-k})^2 = 1 = (h^{-k})^2 = \sigma(S_2)$. We have $\Sigma(S_1) = \{h^i g^{-i} : i \in [1, 2k]\}$. Suppose that for some $i, j \in \mathbb{N}$ we had $h^i g^{-i} = g^j$. Then we have $h^i = g^{j+i}$ so by hypothesis $h^i = 1$ and hence $k|i$ so $h^i g^{-i} = ((hg^{-1})^k)^{i/k} = 1$. We also have $\Sigma(S_2) = \langle h \rangle$ so $\Sigma(S_2) \cap \langle g \rangle = \{1\}$. We set $c = 2k$ and set $T = (hg^{-1}) \cdot (h^{-1})$. We have $\sigma(T) = g^{-1} = g^\gamma$, and $T^c = S$, in fact a partition of S . Lastly, we compute $\frac{c}{d} = |g| = 1 + \frac{\delta-1}{\delta-(\delta-1)} = 1 + \frac{\delta-1}{\delta-\gamma}$, as desired. \square

5 Minimal K-Sum Sequences

We now continue the study of $G/\langle g \rangle$, but drop the $\langle g \rangle \oplus H \leq G$ restriction which is too strong in some cases. For example, consider $(G, g) \cong (\mathbb{Z}_{49}, 7)$. In

this case, the previous section does not apply, but we will see in Theorem 4 that in fact there are configurations for almost every δ, γ .

To do this, we introduce another useful tool that may have some interest beyond this problem. Although we continue to write our groups multiplicatively, we will embrace the traditional additive terminology in the following definition.

Definition 3 Let G be a finite abelian group. Let K be a subgroup of G . Let $S \in \mathcal{F}(G)$. We call S a K -sum sequence if $\sigma(S) \in K$. We call S a *minimal K -sum sequence* if $\sigma(S) \in K$ and $\Sigma'(S) \cap K = \emptyset$.

Note that if $K = \{1\}$ then this definition coincides with that of zero-sum sequences, which are quite well-studied. For example, it is known that each such G has an associated number $D(G)$, called the Davenport constant, such that for every $t \in [1, D(G)]$, there is a minimal zero-sum sequence of length t . We can employ this fact to construct minimal K -sum sequences via the following.

Proposition 11 Let G be a finite abelian group with subgroup K . Let S be a minimal zero-sum sequence in G/K . Then there is a minimal K -sum sequence $T \in \mathcal{F}(G)$, such that $|T| = |S|$.

Proof Let $S = (g_1K) \cdot (g_2K) \cdot \dots \cdot (g_lK)$, for some $l \in \mathbb{N}_0$, and for some $g_1, \dots, g_l \in G$. Set $T = g_1 \cdot g_2 \cdot \dots \cdot g_l \in \mathcal{F}(G)$. Because $\sigma(S) = 1K$, we must have $\sigma(T) \in K$ and $|T| = |S|$. Suppose that there is some $I \subseteq [1, l]$ with $\prod_{i \in I} g_i \in K$. But then $\prod_{i \in I} (g_iK) \in KK = 1K$. Because S is minimal, either $|I| = 0$ or $|I| = l$. Hence $\Sigma'(T) \cap K = \emptyset$. \square

Once we have a minimal K -sum sequence, we can produce many more of the same length.

Proposition 12 Let G be a finite abelian group with subgroup K . Let $S = g_1 \cdot g_2 \cdot \dots \cdot g_l$ be a minimal K -sum sequence. Let $h_1, h_2, \dots, h_l \in K$. Then $T = (g_1h_1) \cdot (g_2h_2) \cdot \dots \cdot (g_lh_l)$ is a minimal K -sum sequence.

Proof We have $\sigma(T) = \prod_{i=1}^l g_i h_i = \sigma(S) \prod_{i=1}^l h_i \in K$. Further, suppose $I \subseteq [1, l]$ and $\prod_{i \in I} g_i h_i \in K$. But then $\prod_{i \in I} g_i = (\prod_{i \in I} g_i h_i) (\prod_{i \in I} h_i)^{-1} \in K$. Since S is a minimal K -sum sequence, either $|I| = 0$ or $|I| = l$. Hence $\Sigma'(T) \cap K = \emptyset$. \square

We now have the machinery to produce configurations, provided that $D(G/\langle g \rangle) \geq 6$ and subject to a mild restriction on γ .

Theorem 4 Let G be a finite abelian group. Let $g \in G$. Set $K = \langle g \rangle$. Suppose that there is a minimal K -sum sequence of length 6. Let $\delta, \gamma \in \mathbb{N}$ with $\delta \geq 2\gamma - 1$ and $|g| > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration.

Proof Let $R = g_1 \cdot g_2 \cdot g_3 \cdot g_4 \cdot g_5 \cdot g_6$ be a minimal K -sum sequence. Note that $R' = g_1^{-1} \cdot g_2^{-1} \cdot g_3^{-1} \cdot g_4^{-1} \cdot g_5^{-1} \cdot g_6^{-1}$ is also a minimal K -sum sequence. Because $\sigma(R) \in K$, there is some $s \in [1, |g|]$ such that $g^s = \sigma(R)$. Now, for each $i \in [1, |g|]$, we define two minimal K -sum sequences as follows. Set $S_i = (g_1 g^\gamma) \cdot (g_2 g^\gamma) \cdot g_3 \cdot g_4 \cdot (g_5 g^i) \cdot (g_6 g^{1-\gamma-s-i})$, and $S'_i = g_1^{-1} \cdot g_2^{-1} \cdot (g_3^{-1} g^\gamma) \cdot (g_4^{-1} g^\gamma) \cdot (g_5^{-1} g^{\gamma-i}) \cdot (g_6^{-1} g^{1-2\gamma+s+i})$. We set $S = S_1 S_2 \cdots S_{|g|} S'_1 S'_2 \cdots S'_{|g|}$. Note that for each $i \in [1, |g|]$, we have $\sigma(S_i) = \sigma(R) g^{1+\gamma-s} = g^{1+\gamma}$, and also $\sigma(S'_i) = \sigma(R') g^{1+\gamma+s} = g^{1+\gamma}$. Also for each $i \in [1, |g|]$, because they are minimal K -sum sequences, $\Sigma'(S_i) \cap \{g, g^2, \dots, g^{|g|-1}\} = \emptyset = \Sigma'(S'_i) \cap \{g, g^2, \dots, g^{|g|-1}\}$. Hence this partition of S satisfies the required definition with $d = 2|g|$.

We set $c = 6|g|$. For $i \in [1, 2]$, we define $T_i = (g_i g^\gamma) \cdot (g_i^{-1})$. For $i \in [3, 4]$, we define $T_i = (g_i^{-1} g^\gamma) \cdot (g_i)$. For $i \in [1, |g|]$, we define $U_i = (g_5 g^i) \cdot (g_5^{-1} g^{\gamma-i})$. Note that $\{g^{1-\gamma-s-i} : i \in [1, |g|]\} = K = \{g^{1-2\gamma+s+i} : i \in [1, |g|]\}$. Hence there is some permutation π of $\{1, 2, \dots, |g|\}$ such that $g^{1-\gamma-s-i} g^{1-2\gamma+s+\pi(i)} = g^\gamma$ for each $i \in [1, |g|]$. Now, for $i \in [1, |g|]$, we define $V_i = (g_6 g^{1-\gamma-s-i}) \cdot (g_6^{-1} g^{1-2\gamma+s+\pi(i)})$. We have $S = (T_1 T_2 T_3 T_4)^{|g|} U_1 \cdots U_{|g|} V_1 \cdots V_{|g|}$. By construction we see that $\sigma(T_i) = \sigma(U_i) = \sigma(V_i) = g^\gamma$, for all i in this subpartition (actually partition) of S .

Note that by assumption we have $\delta \geq 2\gamma - 1$, which is equivalent to $3 \geq 1 + \frac{\delta-1}{\delta-\gamma}$. The proof is now complete since $\frac{c}{d} = \frac{6|g|}{2|g|} = 3$. \square

There are only eight nonisomorphic nontrivial groups whose Davenport constant is less than 6, and the only cyclic ones are $\mathbb{Z}_2, \mathbb{Z}_3, \mathbb{Z}_4, \mathbb{Z}_5$. From the perspective that $G/\langle g \rangle$ is arbitrary, the restriction that $D(G/\langle g \rangle) \geq 6$ is quite mild. On the other hand, if we fix G of small rank and vary g , then this restriction appears stronger since a significant fraction of G may fail to meet it.

Recall that in the ACM context δ is always a multiple of $|g|$. Hence the condition in Theorem 4 that $\gamma \leq \delta/2$ always holds, unless $\delta = |g|$. That is, the condition that $\gamma \leq \delta/2$ is equivalent to $\alpha \geq \lfloor p \rfloor_{\mathbb{Z}_n^\times} / 2$.

We can also trade off the two restrictions in Theorem 4; if we increase the lengths of the S_i , we strengthen the Davenport constant restriction and weaken the already mild α restriction. However, no matter how big $D(G/\langle g \rangle)$ is, this approach cannot work for $\alpha = 1$. In the other direction, we can also weaken the Davenport constant restriction by one, but strengthen the α restriction, as given by the following.

Proposition 13 *Let G be a finite abelian group. Let $g \in G$. Set $K = \langle g \rangle$. Suppose that there is a minimal K -sum sequence of length 5. Let $\delta, \gamma \in \mathbb{N}$ with $\delta \geq 3\gamma - 2$ and $\gamma < |g|$. Then there is a (G, g, δ, γ) -configuration.*

Proof Let $R = g_1 \cdot g_2 \cdot g_3 \cdot g_4 \cdot g_5$ be a minimal K -sum sequence. Note that $R' = g_1^{-1} \cdot g_2^{-1} \cdot g_3^{-1} \cdot g_4^{-1} \cdot g_5^{-1}$ is also a minimal K -sum sequence. Because $\sigma(R) \in K$, there is some $s \in [1, |g|]$ such that $g^s = \sigma(R)$. Now, for each $i \in [1, |g|]$ and each $j \in [1, |g|]$, we define two minimal K -sum sequences as follows. Set $S_{i,j} = (g_1 g^\gamma) \cdot (g_2 g^\gamma) \cdot (g_3 g^\gamma) \cdot (g_4 g^i) \cdot (g_5 g^{1-2\gamma-s-i})$, and $S'_{i,j} =$

$(g_1^{-1}) \cdot (g_2^{-1}) \cdot (g_3^{-1}) \cdot (g_4^{-1}g^j) \cdot (g_5^{-1}g^{1+\gamma-s-j})$. We set $S = \prod_{i,j} S_{i,j}S'_{i,j}$, with $d = 2|g|^2$. Note that for each $i, j \in [1, |g|]$, we have $\sigma(S_{i,j}) = \sigma(S'_{i,j}) = g^{1+\gamma}$, so this partition satisfies the required definition.

We set $c = 5|g|^2$. For $i \in [1, 3]$, we define $T_i = (g_i g^\gamma) \cdot (g_i^{-1})$. Note that $\{g^{1-2\gamma-s-i} : i \in [1, |g|]\} = K = \{g^{1+\gamma-s-j} : j \in [1, |g|]\}$. For $i \in [1, |g|]$, we define $U_i = (g_4 g^i) \cdot (g_4^{-1} g^{\gamma-i})$. For $i \in [1, |g|]$, we define $V_i = (g_5 g^{1-2\gamma-s-i}) \cdot (g_5^{-1} g^{1+\gamma-s-(2-2\gamma-2s-i)}) = (g_5 g^{1-2\gamma-s-i}) \cdot (g_5^{-1} g^{-1+3\gamma+s+i})$. We have $S = (T_1 T_2 T_3)^{|g|^2} (U_1 \cdots U_{|g|})^{|g|} (V_1 \cdots V_{|g|})^{|g|}$. By construction we see that $\sigma(T_i) = \sigma(U_i) = \sigma(V_i) = g^\gamma$, for all i in this subpartition (actually partition) of S .

Note that by assumption we have $\delta \geq 3\gamma - 2$, which is equivalent to $\frac{5}{2} \geq 1 + \frac{\delta-1}{\delta-\gamma}$. The proof is now complete since $\frac{c}{d} = \frac{5|g|^2}{2|g|^2} = \frac{5}{2}$. \square

In the ACM context, the restriction that $\delta \geq 3\gamma - 2$ is equivalent to either $\alpha \in [\frac{2}{3}(x-1), x)$, or $\alpha \geq \frac{4x-2}{3}$, for $x = |p|_{\mathbb{Z}_n^\times}$. We cannot reduce the restriction on $D(G/\langle g \rangle)$ any further than 5 using this approach.

6 $\exp(G/\langle g \rangle)$

We now consider $G/\langle g \rangle$ in a third way, by considering its exponent. This is a particularly fruitful approach if G (and hence $G/\langle g \rangle$) is cyclic. The following result uses a construction similar to that in Theorem 3.

Theorem 5 *Let G be a finite abelian group and $g \in G$. Set $K = \langle g \rangle$, $m = \exp(G/K)$. Let $\delta, \gamma \in \mathbb{N}$ that satisfy $\delta \geq |g| > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration, provided that the following inequality holds:*

$$m \geq 1 + \frac{1}{|g| - \gamma} + \frac{\delta - 1}{\delta - \gamma}$$

Proof We will construct the configuration explicitly. Let $hK \in G/K$ satisfy $|hK| = \exp(G/K) = m$. Note that $h^x \notin K$ for $x \in [-(m-1), (m-1)] \setminus \{0\}$. For each $i \in [1, |g|]$, we set $S_i = (g^{-1})^{|g|-\gamma-1} \cdot (hg^i)^{m-1} \cdot (h^{-1}g^{-i})^{m-1}$. We set $T_0 = (g^{-1})^{|g|-\gamma}$, and for $i \in [1, |g|]$ set $T_i = (hg^{\gamma+i}) \cdot (h^{-1}g^{-i})$. Note that for all $i \in [1, |g|]$, we have $\sigma(S_i) = g^{\gamma+1}$ and $\sigma(T_i) = g^\gamma = \sigma(T_0)$. If $x \in K \cap \Sigma(S_i)$ then in fact $x \in \Sigma((g^{-1})^{|g|-\gamma-1})$ and consequently $x \notin \{g, g^2, \dots, g^\gamma\}$.

For convenience, set $a_\gamma = |g| - \gamma$. We set $d = |g|a_\gamma$ and $S = \prod_{i=1}^{|g|} S_i^{a_\gamma}$. We set $c = (a_\gamma - 1)|g| + (m-1)a_\gamma|g| = ma_\gamma|g| - |g|$ and $T = T_0^{(a_\gamma-1)|g|} \prod_{i=1}^{|g|} T_i^{(m-1)a_\gamma}$. We now verify that $T|S$ (in fact $T = S$). For g^{-1} , we have $\nu_{g^{-1}}(T) = a_\gamma(a_\gamma - 1)|g| = \nu_{g^{-1}}(S)$. For any $i \in [1, k]$, we have $\nu_{hg^i}(T) = (m-1)a_\gamma = \nu_{hg^i}(S)$ and equally $\nu_{h^{-1}g^i}(T) = (m-1)a_\gamma = \nu_{h^{-1}g^i}(S)$. Lastly, we calculate $\frac{c}{d} = \frac{ma_\gamma|g| - |g|}{|g|a_\gamma} = m - \frac{1}{a_\gamma} \geq 1 + \frac{\delta-1}{\delta-\gamma}$ by hypothesis. \square

As before, the theorem leads to several corollaries. Corollary 5 gives configurations for all but one γ , and all sufficiently large δ .

Corollary 5 *Let G be a finite abelian group. Let $g \in G$. Set $K = \langle g \rangle$. Suppose that $\exp(G/K) = 3$. Let $\delta, \gamma \in \mathbb{N}$ with $\delta \geq 3|g|$ and $|g| - 1 > \gamma > 0$. Then there is a (G, g, δ, γ) -configuration.*

Proof Suppose by way of contradiction that Theorem 5 fails to hold, i.e. $3 < 1 + \frac{1}{|g|-\gamma} + \frac{\delta-1}{\delta-\gamma} \leq 1 + \frac{1}{2} + \frac{3|g|-1}{2|g|+2}$, where we used the hypotheses regarding δ and γ . This rearranges to $3|g| + 3 < 3|g| + 1$, a contradiction. \square

Corollary 6 *Let G be a finite abelian group. Let $g \in G$. Set $K = \langle g \rangle$. Suppose that $\exp(G/K) = m$, for some $m \geq 4$. Let $\delta, \gamma \in \mathbb{N}$ with either*

1. $\delta \geq 2|g|$ and $|g| > \gamma > 0$; or
2. $\delta = |g|$ and $\frac{m-2}{m-1}|g| \geq \gamma > 0$.

Then there is a (G, g, δ, γ) -configuration.

Proof Suppose by way of contradiction that Theorem 5 fails to hold, i.e. $m < 1 + \frac{1}{|g|-\gamma} + \frac{\delta-1}{\delta-\gamma}$.

(1) Then $m < 1 + 1 + \frac{2|g|-1}{|g|+1}$, which rearranges to $(m-4)|g| < 1-m$, a contradiction.

(2) Then $m < 1 + \frac{|g|}{|g|-\gamma}$, which rearranges to $\gamma > \frac{m-2}{m-1}|g|$, a contradiction. \square

These corollaries show that there are configurations for all γ (for δ sufficiently large) if $\exp(G/K) \geq 4$, and all but one γ for $\exp(G/K) = 3$. The case of that missing γ is addressed in Proposition 15, while the case of $\exp(G/K) = 2$ is addressed in Proposition 14.

Proposition 14 *Let G be a finite abelian group. Let $g \in G$ with $|g| > 2$. Set $K = \langle g \rangle$. Suppose that $G/K \cong \mathbb{Z}_2$. Let $\delta \in \mathbb{N}$ with $\delta \geq |g|$. Then there is no $(G, g, \delta, |g| - 1)$ -configuration.*

Proof Suppose we had such a configuration. Set $\gamma = |g| - 1$ for convenience. Choose coset representative $h \in G \setminus K$. We have $G = K \cup (hK)$. For $X \in \mathcal{F}(G)$, we define X^+, X^- such that $X^+ \in \mathcal{F}(1K)$, $X^- \in \mathcal{F}(hK)$, and $X = X^+ \cdot X^-$. We define $Q = \{k \in G : |k| > 2\} \subseteq G$ and $\phi : \mathcal{F}(G) \rightarrow \mathbb{N}_0$ via $\phi(S) = \sum_{k \in Q} \nu_k(S)$. For each $i \in [1, c]$, we claim that $\phi(T_i) \geq 1$ because otherwise T_i would consist of elements of order at most 2, hence $\sigma(T_i)$ would be of order at most 2, but $\sigma(T_i) = g^{-1}$ which is of order $|g|$. We now claim that $\phi(S_i) \leq 2$ for each $i \in [1, d]$. Suppose to the contrary for some i we have $\phi(S_i) \geq 3$. We have $\phi(S_i^+) = 0$ so in fact $\phi(S_i^-) \geq 3$. Hence there are some $(hg^x), (hg^y), (hg^z) \in Q$ with $(hg^x) \cdot (hg^y) \cdot (hg^z) \in S_i^-$. Taking these pairwise, we get $h^2g^{x+z} = h^2g^{y+z} = 1$, since $\Sigma(S_i) \cap \{g, g^2, \dots, g^\gamma\} = \emptyset$. Modulo $|g|$, we have $x+z \equiv y+z \equiv 0$ and hence $x \equiv y$. But then $(hg^x)^2 = (hg^x)(hg^y) = 1$, so in fact $(hg^x) \notin Q$. Combining the above, we get $2d \geq \phi(S) \geq c$ hence $2 \geq \frac{c}{d} \geq 1 + \frac{\delta-1}{\delta-\gamma}$. This rearranges to $1 \geq \gamma = |g| - 1$, so $2 \geq |g|$, a contradiction. \square

Note that if $|g| = 2$ and $G/K \cong \mathbb{Z}_2$ then either $G \cong \mathbb{Z}_4$ and no configuration exists for $\gamma = |g| - 1 = 1$ (by a simple argument similar to the proof of Proposition 14), or $G \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ and configurations exist for all γ by Proposition 10.

Proposition 15 *Let G be a finite abelian group. Let $g \in G$. Set $K = \langle g \rangle$. Suppose that $G/K \cong \mathbb{Z}_3$ and $G \not\cong K \oplus \mathbb{Z}_3$. Let $\delta \in \mathbb{N}$ with $\delta \geq |g|$. Then there is no $(G, g, \delta, |g| - 1)$ -configuration.*

Proof Suppose we had such a configuration. Set $\gamma = |g| - 1$ for convenience. Choose coset representative $h \in G \setminus K$. We have $G = K \cup (hK) \cup (h^2K)$, with $h^3 \in K$. If there were some $s \in [1, |g| - 1]$ such that $h^3 = g^{3s}$, then we have $(hg^{-s})^3 = 1$ and hence $G \cong K \oplus \mathbb{Z}_3$, which violates the hypothesis. Similarly, there is no such s with $(h^2)^3 = g^{3s}$.

Let S_i be in our configuration; we claim that S_i contains at most 4 nonunit elements. First, S_i can contain no nonunit elements from K . Suppose that S_i contains four elements from hK , say $hg^{x_1}, hg^{x_2}, hg^{x_3}, hg^{x_4}$. Multiplying these three at a time, we get $h^3g^{x_1+x_2+x_3}, h^3g^{x_1+x_2+x_4} \in \Sigma S_i \cap K = \{1\}$. Hence $x_3 \equiv x_4 \pmod{|g|}$ and by symmetry $x_1 \equiv x_2 \equiv x_3 \equiv x_4 \pmod{|g|}$. But now $(hg^{x_1})^3 \in \Sigma S_i \cap K = \{1\}$, so $h^3 = (g^{-x_1})^3$, which contradicts our hypothesis. Hence S_i contains at most three nonunit elements from hK and by symmetry at most three nonunit elements from h^2K . Suppose now S_i contained at least 5 nonunit elements. At least three must be from the same coset, so without loss S_i contains $hg^{x_1}, hg^{x_2}, hg^{x_3}, h^2g^{x_4}$. But now $h^3g^{x_1+x_4} = h^3g^{x_2+x_4} = h^3g^{x_3+x_4} = 1$, so $x_1 \equiv x_2 \equiv x_3 \pmod{|g|}$ and again $(hg^{x_1})^3 \in S_i$, a contradiction.

Since $\nu_{g^{-1}}(S) = 0$ and $\sigma(T_i) = g^{-1}$, each T_i in our configuration must have at least two nonunit elements. Combining the above, we get $4d \geq 2c$, and hence $2 \geq \frac{c}{d} > 1 + \frac{\delta-1}{\delta-\gamma}$. This rearranges to $1 \geq \gamma = |g| - 1$, so $2 \geq |g|$. But then $G \cong \mathbb{Z}_6 \cong K \oplus \mathbb{Z}_3$, a contradiction. \square

The condition $G \not\cong K \oplus \mathbb{Z}_3$ in Proposition 15 is essential, since otherwise Corollary 4 gives us the opposite conclusion for large δ . Note that by Corollary 5, Proposition 15 is tight for $\delta \geq 3|g|$. That is, $\gamma = |g| - 1$ is the only value of γ that does not have a configuration.

7 Problems and Conjectures

Although we have made substantial progress on the existence question of (G, g, δ, γ) -configurations, there are still several gaps in our work.

First, there are some (G, g) pairs where there is no large enough H with $\langle g \rangle \oplus H \leq G$, and $G/\langle g \rangle$ has Davenport constant less than 5 and exponent less than 4. In these cases nothing at all is known. For example, if $G/\langle g \rangle \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$.

Second, in the case where $G/\langle g \rangle \cong \mathbb{Z}_2$, Proposition 14 gives the single value $\gamma = |g| - 1$ where no configuration exists, for any δ . However, we don't know

about the other values of γ . Preliminary work suggests that there is a cutoff $\tau \approx \sqrt{|g|}$, such that if $\gamma < \tau$ configurations exist for δ sufficiently large and if $\gamma > \tau$ configurations do not exist. This and other computational work leads us to the following conjecture, for general G, g .

Conjecture 1 Suppose that there is a (G, g, δ, γ) -configuration, and $\gamma > 0$. Then there is a $(G, g, \delta, \gamma - 1)$ -configuration.

Lastly, for all the cases where configurations are known to exist for all γ and all δ sufficiently large (i.e. all but finitely many α), there is still a small set of α where it is unknown whether configurations exist.

Acknowledgements This research was supported in part by NSF REU grant 1061366.

References

1. D. D. Anderson and Jonathan Preisser. Factorization in integral domains without identity. *Results Math.*, 55(3-4):249–264, 2009.
2. Paul Baginski, Scott T. Chapman, Christopher Crutchfield, K. Grace Kennedy, and Matthew Wright. Elastic properties and prime elements. *Results Math.*, 49(3-4):187–200, 2006.
3. Paul Baginski, Scott T. Chapman, and George J. Schaeffer. On the delta set of a singular arithmetical congruence monoid. *J. Théor. Nombres Bordeaux*, 20(1):45–59, 2008.
4. M. Banister, J. Chaika, S. T. Chapman, and W. Meyerson. On a result of James and Niven concerning unique factorization in congruence semigroups. *Elem. Math.*, 62(2):68–72, 2007.
5. M. Banister, J. Chaika, S. T. Chapman, and W. Meyerson. On the arithmetic of arithmetical congruence monoids. *Colloq. Math.*, 108(1):105–118, 2007.
6. M. Banister, J. Chaika, S. T. Chapman, and W. Meyerson. A theorem on accepted elasticity in certain local arithmetical congruence monoids. *Abh. Math. Semin. Univ. Hambg.*, 79(1):79–86, 2009.
7. M. Banister, J. Chaika, and W. Meyerson. Technical report, Trinity University REU, 2003.
8. S. T. Chapman and David Steinberg. On the elasticity of generalized arithmetical congruence monoids. *Results Math.*, 58(3-4):221–231, 2010.
9. Marco Fontana, Evan Houston, and Thomas Lucas. *Factoring ideals in integral domains*, volume 14 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Heidelberg, 2013.
10. Alfred Geroldinger. Additive group theory and non-unique factorizations. In *Combinatorial number theory and additive group theory*, Adv. Courses Math. CRM Barcelona, pages 1–86. Birkhäuser Verlag, Basel, 2009.
11. Alfred Geroldinger and Franz Halter-Koch. Congruence monoids. *Acta Arith.*, 112(3):263–296, 2004.
12. Alfred Geroldinger and Franz Halter-Koch. *Non-unique factorizations*, volume 278 of *Pure and Applied Mathematics (Boca Raton)*. Chapman & Hall/CRC, Boca Raton, FL, 2006. Algebraic, combinatorial and analytic theory.
13. Franz Halter-Koch. C-monoids and congruence monoids in Krull domains. In *Arithmetical properties of commutative rings and monoids*, volume 241 of *Lect. Notes Pure Appl. Math.*, pages 71–98. Chapman & Hall/CRC, Boca Raton, FL, 2005.
14. Thomas W. Hungerford. *Algebra*. Holt, Rinehart and Winston, Inc., New York, 1974.
15. M. Jenssen, D. Montealegre, and V. Ponomarenko. Irreducible factorization lengths and the elasticity problem within \mathbb{N} . *Amer. Math. Monthly*, 120(4):322–328, 2013.