

Griffith Research Online

<https://research-repository.griffith.edu.au>

Aberrant promoter hypermethylation and silencing of the critical 3p21 tumor suppressor gene, RASSF1A, in Chinese oesophageal squamous cell carcinoma

Author

Wong, Michelle, Tao, Qian, Fu, Li, Wong, Kai-Yau, Qiu, Guo-Hua, B. F. Law, Fian, Tin, Pui-Chi, Chenug, Wai-Ling, Lee, Pin-Yin, Tang, Jonny Cheuk-On, S. W. Tsao, George, Lam, Alfred, Law, Simon, Wong, John, Srivastava, Gopesh

Published

2006

Journal Title

International Journal of Oncology

Copyright Statement

Copyright 2006 Spandidos Publications. The attached file is reproduced here in accordance with the copyright policy of the publisher.

Downloaded from

<http://hdl.handle.net/10072/14121>

Link to published version

<http://www.spandidos.com/ijo/>

Aberrant promoter hypermethylation and silencing of the critical 3p21 tumour suppressor gene, *RASSF1A*, in Chinese oesophageal squamous cell carcinoma

MICHELLE LY WONG¹, QIAN TAO^{4,6}, LI FU⁴, KAI-YAU WONG¹, GUO-HUA QIU⁴, FIAN B.F. LAW¹, PUI-CHI TIN¹, WAI-LING CHEUNG¹, PING-YIN LEE², JOHNNY CHEUK-ON TANG^{1,7,8}, GEORGE S.W. TSAO³, KING-YIN LAM⁹, SIMON LAW², JOHN WONG² and GOPESH SRIVASTAVA¹

Departments of ¹Pathology, ²Surgery and ³Anatomy, The University of Hong Kong, Hong Kong, P.R. China; ⁴Johns Hopkins Singapore, Singapore; ⁵Sidney Kimmel Comprehensive Cancer Centre, Johns Hopkins School of Medicine, Baltimore, MD, USA; ⁶Cancer Epigenetics Laboratory, Cancer Centre, Department of Clinical Oncology, The Chinese University of Hong Kong; ⁷Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University; ⁸State Key Laboratory of Chinese Medicine and Molecular Pharmacology, Shenzhen, P.R. China; ⁹Department of Pathology, Griffith Medical School, Griffith University, Queensland, Australia

Received August 11, 2005; Accepted September 23, 2005

Abstract. 3p21 is an important locus harbouring critical tumour suppressor genes (TSG), which are implicated in the pathogenesis of multiple tumours, including oesophageal carcinoma. *RASSF1A* is a 3p21.3 candidate TSG frequently inactivated by promoter methylation in multiple tumours. We investigated *RASSF1A* promoter methylation and gene expression in Chinese oesophageal squamous cell carcinoma (ESCC) to compare it to data from Japanese patients. Methylation-specific PCR (MSP) showed that *RASSF1A* was partially methylated in 3/7 (43%) cell lines; 22/64 (34%) primary tumours and 3/64 (5%) corresponding non-tumour samples; and was not methylated in 2 immortalized normal oesophageal epithelial cell lines and 6 normal oesophageal epithelium samples. Bisulfite genome sequencing confirmed the MSP results. Promoter hypermethylation correlated well with *RASSF1A* mRNA down-regulation. Treatment of cell lines with 5-aza-2'-deoxycytidine activated *RASSF1A* mRNA expression along with promoter demethylation. *RASSF1A* hypermethylation in the Chinese cohort was much lower than

in a published report of Japanese ESCC patients (52%) and cell lines (74%). Our own analysis of Japanese ESCC cell lines for direct comparison also detected a high frequency of *RASSF1A* hypermethylation (8/10; 80%) and high levels of hypermethylation at each CpG site. No significant association between *RASSF1A* hypermethylation and histological differentiation ($p=0.953$), tumour staging ($p=0.117$), or survival ($p=0.7571$) was found in Chinese ESCC, unlike the results of Japanese patients. The incidence of oesophageal cancer shows marked variation by geographic area and ethnic group; it is almost three times higher in China than in Japan, indicating possible different pathogenetic mechanisms. Our results show that *RASSF1A* hypermethylation in ESCC has epidemiological/ethnic differences, and suggest that Chinese ESCC may result from different pathogenetic mechanisms.

Introduction

Oesophageal cancer is the sixth most common cancer in males in the world. Its age-standardized incidence rate is 11.5/10⁵ males (1). It is well known for its marked variation by geographic area and ethnic group. For example, it is the fourth most common cancer in China (27.4/10⁵), seventh in Japan (10/10⁵), ninth in United Kingdom (9.6/10⁵) and sixteenth in United States of America (5.9/10⁵) (1). No molecular mechanisms have yet been demonstrated to explain these variations. There are two major histological subtypes of oesophageal cancer: oesophageal squamous cell carcinoma (ESCC), which is the predominant subtype in Orientals; and oesophageal adenocarcinoma (EADC), which is more common in Caucasians. Despite advances in multimodality therapy, ESCC remains a significant problem with very low 5-year survival rates (2).

Gene amplification and/or overexpression are frequently involved in the pathogenesis of ESCC (2). Recent application of comparative DNA fingerprinting performed by our group

Correspondence to: Professor Gopesh Srivastava, Department of Pathology, The University of Hong Kong, Queen Mary Hospital, 102 Pokfulam Road, Hong Kong, P.R. China
E-mail: gopesh@pathology.hku.hk

Dr Qian Tao, Cancer Epigenetics Laboratory, Cancer Centre, Department of Clinical Oncology, The Chinese University of Hong Kong, Hong Kong, P.R. China
E-mail: qtao@clo.cuhk.edu.hk

Key words: oesophageal carcinoma, tumor suppressor gene, 3p21, *RASSF1A*, methylation

revealed that amplifications or deletions of specific chromosomal regions are common in both ESCC preneoplastic lesions and carcinomas (3). By cDNA arrays, our group and others have demonstrated that multiple cancer-related genes are differentially expressed in ESCC with several down-regulated tumour suppressor genes (TSGs) (4-8).

It is well established that epigenetic and genetic alterations are central to the pathogenesis of human cancers (9). Aberrant hypermethylation of CpG islands in TSG promoters has been found in a variety of cancers; with some promoters hypermethylated in most tumours and others being more cancer-type specific. In ESCC, frequent epigenetic inactivation by promoter hypermethylation has been documented in *CDKN2* (10); *FHIT* (11); *p14ARF*, *p16^{INK4a}* and *p15^{INK4b}* (12); *HLA class I* (13); *E-cadherin* (14); *O-6-methylguanine-DNA methyltransferase* (15); *VHL* (16); *RAR- β* (16,17); *ECRG4* (18) and *TSLC1* (19).

Allelic loss of 3p21 is frequent in many cancers (20), suggesting the presence of critical TSGs in this locus. *RASSF1*, a candidate TSG, is located within a critical 120-kb region at 3p21.3 that shows frequent homozygous deletions in lung tumour cell lines (21,22). *RASSF1* has a predicted Ras-association domain and homology to Ras-effector NORE1A (23). *RASSF1* encodes two major isoform transcripts, *RASSF1A* and *RASSF1C*, derived from alternative splicing and different promoter usage. The loss of *RASSF1A* has been suggested to play a key role in tumorigenesis. *RASSF1A* is frequently inactivated by promoter hypermethylation or loss of heterozygosity (LOH) in a variety of tumours, including small cell lung cancer (24,25), renal cell carcinoma (26), bladder cancer (27), thyroid carcinoma (28), nasopharyngeal carcinoma (29,30), melanoma (31), hepatocellular carcinoma (32), breast tumour (24), prostate cancer (33,34) and Hodgkin's disease (35). Mutation of *RASSF1A* is not commonly observed.

The most frequent chromosomal loss in Chinese ESCC in Hong Kong, detected by conventional comparative genomic hybridization analysis (CGH), is 3p (46.2%) (36). Through high-resolution array CGH (aCGH), we found that 3p21 is also heterozygously deleted in around half of the ESCC cell lines we studied, including both Chinese and Japanese cell lines, while homozygous deletion was not detected (Ying and Tao, unpublished data). A high frequency (72%) of LOH at 3p21.3 has also been described in primary ESCC from Hong Kong Chinese patients using microsatellite markers (37), suggesting that 3p21 TSGs could also be subjected to both genetic and epigenetic inactivation in ESCC. These studies highlight the 3p21 region as critical for ESCC pathogenesis.

As mentioned above, the incidence rate of ESCC is quite different between China and Japan, with China showing almost a three times higher rate. The possibility that different pathogenic mechanisms are operating in these two countries is appealing, but until now we have had little molecular evidence that this might be the case. *RASSF1A* hypermethylation has been recently reported in Japanese ESCC by Kuroki *et al* (16,38), but no study has been performed for Chinese ESCC. As part of our ongoing interest in the geographical distribution of disease, we investigated the epigenetic inactivation of *RASSF1A* by hypermethylation in a series of Chinese ESCC cell lines and primary tumour specimens, and we report here the rather surprising results.

Patients and methods

Tissue specimens. Primary ESCC specimens were collected prospectively from 64 Chinese patients that underwent resection for ESCC since 1996 in Queen Mary Hospital, Hong Kong, and were not pretreated with chemotherapy or radiotherapy. Representative tissue samples from tumours and matching morphologically normal oesophageal epithelium tissue (non-tumour) at least 6-10 cm away from the tumours were sampled for each patient. Sterile equipment was used for each sampling of tumour and non-tumour specimens. One half of each sample was fixed in 10% formalin for histological assessment and the other half was snap frozen in liquid nitrogen and stored at -85°C for DNA and RNA extraction. For tumour samples, non-tumour portions were trimmed off from the frozen tumour blocks and the selected areas had more than 80% tumour cells as shown by histological assessment. For every 20 cryostat sections cut from the frozen blocks for DNA and RNA extraction, an additional section with hematoxylin and eosin staining was prepared for histological assessment to confirm the presence or absence of tumour cells. Six morphologically normal oesophageal epithelium samples obtained from Chinese subjects who died of traffic accidents and showed no evidence of ESCC by histology were included in this study as controls.

Cell lines. Seven Chinese ESCC cell lines (HKESC-1, HKESC-2, HKESC-3, SLMT-1, EC1, EC18, EC109) (14,39-41) and two HPV16 E6E7-immortalized normal Chinese oesophageal epithelial cell lines (NE1 and NE3) (42) were included in the study. Subsequently, we also analyzed 10 Japanese ESCC cell lines (KYSE 30, KYSE 70, KYSE 140, KYSE 150, KYSE 180, KYSE 270, KYSE 410, KYSE 450, KYSE 510, KYSE 520; obtained from the German Collection of Microorganisms and Cell Cultures).

Treatment of cell lines with demethylation agent. The demethylation treatment of selected ESCC cell lines by 5-aza-2'-deoxycytidine (aza) (Sigma Chemical Co., St. Louis, MO) was carried out as described (35). The optimal concentration of aza was tested first by using 1-200 μ M of aza to treat the cells over 6 days while observing cell viability. It was found that aza concentrations over 50 μ M were toxic to the cells. Thus, cells were diluted to 3×10^5 and allowed to grow overnight and aza was added to the culture medium to the desired concentration (10 μ M). Cells were grown for 3 days or 6 days. Fresh medium containing 10 μ M aza was replaced every 24 h. After treatment, cells were pelleted and washed with phosphate-buffered saline (PBS).

DNA extraction, bisulfite treatment and methylation-specific PCR (MSP). Genomic DNA was extracted from the cell line pellets, frozen sections of paired tumour and non-tumour samples and normal oesophageal epithelium samples using proteinase K digestion and phenol/chloroform extraction. For each sample, 2-5 μ g DNA was used for DNA bisulfite modification using the CpGenome™ DNA modification kit (Intergen Co. New York). The methylation status of the *RASSF1A* promoter was analyzed using the MSP method recently developed by our group with methylated (m) and unmethylated (u)-specific primers and 50-100 ng of bisulfite-




Figure 1. Analysis of the *RASSF1A* methylation by MSP for 30 cycles (upper panel) and *RASSF1A* and *RASSF1C* mRNA by semi-quantitative RT-PCR for 30 cycles (lower panel) in (a) 7 Chinese ESCC cell lines and 2 immortalized Chinese normal oesophageal epithelial cell lines (NE1 and NE3), (b) representative Chinese ESCC tumour (T) and non-tumour (NT) samples in cases with methylated *RASSF1A* in tumours (T), (c) 6 Chinese normal oesophageal epithelium samples, and (d) 10 Japanese ESCC cell lines (KYSE series). Universal methylated DNA (UMD) was used as a positive control for MSP. *GAPDH* amplification served as a control for cDNA quality. H₂O was used as a negative control. (MW, molecular weight markers).

treated DNA (30). The MSP primers for the methylated *RASSF1A* promoter were RASSF1Am1 (5'-GGT TTT TTT TAG TTT TTT TTC GTC-3') and RASSF1Am6 (5'-CTA CCG TAT AAA ATT ACA CGC G-3'), and for the unmethylated promoter were RASSF1Au1 (5'-TGG TTT TTT TTA GTT TTT TTT TGT T-3') and RASSF1Au6 (5'-ACT ACC ATA TAA AAT TAC ACA CA-3'). This MSP for 30 cycles can detect as low as 1% of methylated DNA. Universal methylated DNA (Intergen Co. New York) was used as a positive control for MSP.

Bisulfite genomic sequencing. Detailed methylation analysis of each individual site in the *RASSF1A* CpG island by bisulfite genome sequencing was performed as described (35). Bisulfite-treated DNA samples (50-100 ng) were amplified with strand-specific primers, BGS4 (5'-CCC RCA ACT CAA TAA ACT CAA A-3') and BGS5 (5'-GTT AAG TGT GTT GTT TTA GTA AAT-3'), which amplify a 388-bp region (-338 to +50) spanning 34 CpG sites within the promoter and exon 1 of *RASSF1A* gene. PCR products were cloned into the pCR2.1-TA cloning vector (Invitrogen, Carlsbad, CA). For bisulfite genome sequencing, at least 8 bacterial colonies were analyzed for each DNA sample. Percentage methylation was determined as the percentage of methylated cytosines from 8-10 sequenced colonies.

RNA extraction and RT-PCR analysis. Total RNA was extracted from cell line pellets, frozen sections of tumours and non-tumour samples, and normal oesophageal epithelium samples using TRIzol (Life Technologies, Inc., Gaithersburg,

MD). RT-PCR for 30 cycles was performed using the GeneAmp RT-PCR kit (Perkin-Elmer-Cetus, Norwalk, CT) using 2 µg RNA. The primers for *RASSF1A* and *RASSF1C* isoform were as follows: RASSF1AC (for *RASSF1A*, in exon 1α), 5'-GAC CTC TGT GGC GAC TTC-3'; RDAA (for *RASSF1C*, in exon 2γ), 5'-GAG GCG CCT TCT TTC GAA-3'; and RDAB (reverse primer, for both isoforms in exon 4), 5'-CAA GGA GGG TGG CTT CTT-3' (35). RT-PCR of glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) transcript was performed with the use of forward primer, GAPDH-L (5'-GGC TCT CCA GAA CAT CAT CCC TGC-3') and reverse primer GAPDH-R (5'-GGG TGT CGC TGT TGA AGT CAG AGG-3') (35). *GAPDH* amplification served as a control for cDNA quality. H₂O was used as a negative control.

Clinicopathological correlation. Statistical analysis was performed using SPSS for Windows Release 8.0.0 (22 December 1997, SPSS Inc., Chicago, IL). For analyzing correlations between categorical variables, Pearson's χ^2 test was used. Kaplan-Meier was used to estimate survival distribution. A 2-tailed probability (p-value) <0.05 was considered as statistically significant.

Results

Analysis of *RASSF1A* methylation in Chinese ESCC cell lines. Analysis of *RASSF1A* promoter methylation in 7 ESCC cell lines and 2 immortalized normal oesophageal epithelial cell lines by MSP showed that *RASSF1A* was hypermethylated in 3/7 (43%) cell lines (HKESC-1, EC1, EC18) but not in the

Table I. Frequency of *RASSF1A* hypermethylation in ESCC cell lines and tumour samples.

Samples	Number of samples	Detection of hypermethylated <i>RASSF1A</i> (%)
Chinese ESCC		
ESCC cell lines	7	3 (43)
Immortalized normal oesophageal epithelial cell lines	2	0 (0)
ESCC primary tumours	64	22 (34)
ESCC paired non-tumour	64	3 (5)
Normal oesophageal epithelium	6	0 (0)
Japanese ESCC		
ESCC cell lines	10	8 (80)

2 immortalized normal cell lines (upper panel of Fig. 1a and Table I). The unmethylated *RASSF1A* promoter was detected in all of the cell lines except for EC1. RNA expression of the two isoforms of *RASSF1* (*RASSF1A* and *RASSF1C*) in these cell lines using isoform-specific semi-quantitative RT-PCR detected no *RASSF1A* mRNA in EC1 (lower panel of Fig. 1a). The 2 cell lines (HKESC-1 and EC18) which had stronger methylated and weaker unmethylated alleles, also had very low levels of *RASSF1A* expression. Higher levels of *RASSF1A* mRNA were detected in all cell lines which were unmethylated. As an internal control, *RASSF1C* expression was measured




Figure 3. Induction of *RASSF1A* mRNA by demethylating agent aza (10 μ M) in the Chinese EC1 cell line with the hypermethylated *RASSF1A* promoter for 3 days or 6 days. The Chinese EC109 cell line with the unmethylated *RASSF1A* promoter was also treated as a control.

and found in all cell lines. No methylation (upper panel of Fig. 1a) and high levels of *RASSF1A* mRNA were detected in the immortalized normal cell lines (lower panel of Fig. 1a).

Further bisulfite genome sequencing for *RASSF1A* in four ESCC cell lines (HKESC-3, EC1, EC109, SLMT-1) and one normal cell line (NE-1) showed densely methylated CpG sites in the *RASSF1A* promoter in EC1 (Fig. 2). Only scattered methylated CpG sites were detected in HKESC-3, EC109, SLMT-1, and NE-1. Bisulfite genome sequencing analysis confirmed the MSP results.

To determine whether hypermethylation of the *RASSF1A* promoter directly mediates its transcriptional repression, the ESCC cell line (EC1) with complete methylation was treated with aza. A gradual restoration of *RASSF1A* expression was




Figure 2. Detailed methylation analysis of *RASSF1A* by bisulfite genome sequencing of 4 representative Chinese ESCC cell lines, an immortalized Chinese normal oesophageal epithelial cell line (NE1), two hypermethylated Chinese ESCC tumour samples, and one hypermethylated Japanese ESCC cell line, KYSE 450. Top panel illustrates the CpG sites of *RASSF1A* and the amplified regions by MSP and bisulfite genome sequencing, covering its putative promoter and first exon, each vertical bar represents one of the CpG sites in the sequence. Percentage methylation was determined as percentage of methylated cytosines from 8-10 sequenced colonies. Each CpG site is shown in the row as an individual circle. M, methylated; U, unmethylated; (U), weakly unmethylated. On the right side are the MSP results and *RASSF1A* expression levels.

Table II. Correlation of *RASSF1A* methylation status with clinicopathological parameters in primary Chinese ESCC cases.^a

	<i>RASSF1A</i> methylation status		p-value
	Unmethylated	Hypermethylated	
Histological differentiation			
Well	9	5	0.953
Moderate	20	9	
Low	10	5	
T-stage			
1/2	8	1	0.117
3/4	31	19	
N-stage			
N0	19	8	0.525
N1	20	12	
M-stage			
M0	36	16	0.166
M1/1a/1b	3	4	
Stage of disease			
I	1	0	0.492
II	14	7	
III	21	9	
IV	3	4	

^aOnly 59/64 ESCC cases were statistically evaluated since complete clinical data could not be retrieved from 5 patients; p, Pearson's χ^2 test.

detected after 3 and 6 days of treatment as analyzed by semi-quantitative RT-PCR (lower panel of Fig. 3), concomitant with the demethylation of its promoter (upper panel of Fig. 3). Little or no change was detected in the expression of *RASSF1C* or the housekeeping gene, *GAPDH*. As a control, a *RASSF1A*-expressing cell line (EC109) was also treated with aza. After 3 and 6 days of treatment, no change in the expression of *RASSF1A*, *RASSF1C* and *GAPDH* was observed (lower panel of Fig. 3), and the *RASSF1A* methylation status was also unchanged, indicating that aza did not have an indirect effect on *RASSF1A* activation and demethylation (upper panel of Fig. 3). Thus, hypermethylation of its promoter directly repressed *RASSF1A* transcription in EC1.

Methylation of *RASSF1A* in clinical ESCC samples. MSP analysis of methylation in paired ESCC tumour and non-tumour samples showed *RASSF1A* hypermethylation in 22/64 (34%) tumour samples (upper panel of Fig. 1b and Table I). Unmethylated promoter alleles were detected in all ESCC tumour and non-tumour samples, indicating that both forms were detected and minimized the chance of false negative results. High-resolution methylation analysis by bisulfite genome sequencing of two tumours with hypermethylated *RASSF1A* detected by MSP confirmed that the promoter was indeed methylated in these tumours (Fig. 2), albeit only




Figure 4. Correlation of survival data with *RASSF1A* hypermethylation in Chinese ESCC cases. Only 59/64 ESCC cases were statistically evaluated since complete clinical data could not be retrieved from 5 patients.

partially. Only three cases (5%) showed hypermethylation in paired non-tumour tissue (e.g. no. 39). *RASSF1A* methylation was next analyzed in 6 normal oesophageal epithelium samples by MSP. No methylation was detected (Fig. 1c). RT-PCR analysis of paired samples showed that *RASSF1A* hypermethylation correlated well with mRNA down-regulation in primary tumours (lower panel of Fig. 1b). *RASSF1A* was expressed in all non-tumour oesophageal tissue samples.

No correlation of *RASSF1A* hypermethylation with clinicopathological parameters in Chinese ESCC patients. For 59/64 cases (complete clinical data could not be retrieved from 5 patients), the relationship between *RASSF1A* hypermethylation and clinicopathological parameters was analyzed, including TNM staging, histological differentiation, and survival (Table II). Statistical analysis showed no significant correlation of *RASSF1A* hypermethylation with histological differentiation ($p=0.953$), TNM staging ($p=0.117$), or survival ($p=0.7571$) (Table II and Fig. 4).

Comparative examination of *RASSF1A* methylation in Japanese ESCC cell lines. Analysis of 10 Japanese ESCC cell lines using MSP showed a significantly high percentage of *RASSF1A* hypermethylation (80%) in these cell lines (upper panel of Fig. 1d and Table I), compared to Chinese cell lines ($p<0.001$, Z-test). In addition, the degree of hypermethylation also appeared to be higher than in the Chinese (Fig. 2). *RASSF1A* hypermethylation correlated well with mRNA down-regulation in these cell lines (lower panel of Fig. 1d). Bisulfite genome sequencing of one of these cell lines (KYSE 450) confirmed that almost all *RASSF1A* alleles were hypermethylated (Fig. 2).

Discussion

Recent studies have shown that aberrant promoter hypermethylation is associated with loss of TSG functions, which promotes a cancer cell growth advantage or escape from apoptosis (43). *RASSF1A* is a novel candidate TSG at 3p21.3 methylated in tumours. We investigated *RASSF1A* methylation

in a series of Chinese ESCC cell lines and primary tumours, and determined its relationship to clinicopathological parameters. As shown in Table I, we found that *RASSF1A* is silenced by promoter methylation in 43% of Chinese ESCC cell lines, 34% of primary tumours and only 5% of corresponding non-tumour oesophageal tissue, but not in any immortalized normal oesophageal epithelial cell lines and normal oesophageal epithelium samples. Our data indicates that promoter hypermethylation contributes to the inactivation of *RASSF1A* in a fraction of Chinese ESCC tumours.

The unmethylated *RASSF1A* promoter was also detected in all ESCC cell lines except for EC1. The detection of both methylated and unmethylated *RASSF1A* alleles in 2/7 cell lines (HKESC-1, EC18) indicated a partial methylation in these cell lines, and both were found to have low levels of *RASSF1A* expression. All of the resected ESCC tumours and paired non-tumour samples contained unmethylated promoter alleles, which may be explained by contributions from tumour infiltrating stromal cells. While *RASSF1A* methylation was not detected in any normal epithelium samples, methylation was detected in 5% of the paired non-tumour oesophageal tissue in patients with ESCC. This finding may represent the appearance of premalignant lesions or, alternatively, it could be due to the presence of some small percentage of tumour cells disseminated into non-tumour tissue, as reported in other tumours.

Recently, promoter hypermethylation of *RASSF1A* in Japanese ESCC has been reported in 74% of Japanese ESCC cell lines and 52% of primary tumours (16,38). A significant correlation in Japanese ESCC was found between *RASSF1A* hypermethylation and tumour staging ($p=0.009$, Fisher's exact test), though none was found between expression and age and histological differentiation. The proportion of Chinese ESCC cell lines and primary tumours with *RASSF1A* hypermethylation is roughly half of that reported in Japanese ESCC and the difference is significant ($p<0.001$, Z-test). To investigate whether the differences observed reflected true differences between Chinese and Japanese ESCC rather than technical artefact, we subsequently analyzed 10 Japanese ESCC cell lines in a direct comparison, and detected a high percentage of *RASSF1A* hypermethylation (80%) in the Japanese ESCC cell lines. In addition, the percent of methylated sites, as determined by bisulfite genome sequencing, was also significantly different. Therefore, there are large differences between the extents of *RASSF1A* hypermethylation in Chinese and Japanese ESCC. Moreover, we did not find any significant correlation between *RASSF1A* hypermethylation and survival rates and other clinicopathological parameters, which is in contrast to that reported for the Japanese patients. The incidence rate of ESCC is quite different in China and Japan; and our results indicate for the first time that the molecular pathogenetic mechanisms of this highly prevalent tumour could also be different in these two populations.

In conclusion, our study demonstrated that hypermethylation-associated inactivation of *RASSF1A* can be detected in ESCC in the Chinese but not to the same extent as is found in the disease in Japan. Our interest in how ESCC manifests differently in these two countries has been piqued, and we plan additional studies to help identify whether there

is an environmental, cultural, or behavioural factor that can be modified to decrease the incidence of ESCC in China.

Acknowledgements

This work was supported by a CRCG research grant from the University of Hong Kong (to G.S.) and an A*STAR research grant to Johns Hopkins Singapore (to Q.T.).

References

1. Ferlay J, Bray F, Pisani P and Parkin DM: GLOBOCAN 2002: Cancer Incidence, Mortality and Prevalence Worldwide IARC CancerBase No. 5. Version 2.0. IARC Press, Lyon, 2004.
2. Lam KY: Molecular biology of oesophageal squamous cell carcinoma. *Crit Rev Oncol Hematol* 33: 71-90, 2000.
3. Tang JCO, Lam KY, Law S, Wong J and Srivastava G: Detection of genetic alterations in oesophageal squamous cell carcinomas and adjacent normal epithelia by comparative DNA fingerprinting using inter-simple sequence repeat PCR. *Clin Cancer Res* 7: 1539-1545, 2001.
4. Hu YC, Lam KY, Law S, Wong J and Srivastava G: Identification of differentially expressed genes in oesophageal squamous cell carcinoma (ESCC) by cDNA expression array: overexpression of *Fra-1*, *Neogenin*, *Id-1*, and *CDC25B* genes in ESCC. *Clin Cancer Res* 7: 2213-2221, 2001.
5. Hu YC, Lam KY, Law S, Wong J and Srivastava G: Profiling of differentially expressed cancer-related genes in oesophageal squamous cell carcinoma (ESCC) using human cancer cDNA arrays: overexpression of oncogene *MET* correlates with tumour differentiation in ESCC. *Clin Cancer Res* 7: 3519-3525, 2001.
6. Kan T, Shimada Y, Sato F, Maeda M, Kawabe A, Kaganoi J, Itami A, Yamasaki S and Imamura M: Gene expression profiling in human oesophageal cancers using cDNA microarray. *Biochem Biophys Res Commun* 286: 792-801, 2001.
7. Zhou J, Zhao LQ, Xiong MM, Wang XQ, Yang GR, Qiu ZL, Wu M and Liu ZH: Gene expression profiles at different stages of human oesophageal squamous cell carcinoma. *World J Gastroenterol* 9: 9-15, 2003.
8. Su H, Hu N, Shih J, Hu Y, Wang QH, Chuang EY, Roth MJ, Wang C, Goldstein AM, Ding T, Dawsey SM, Giffen C, Emmert-Buck MR and Taylor PR: Gene expression analysis of oesophageal squamous cell carcinoma reveals consistent molecular profiles related to a family history of upper gastrointestinal cancer. *Cancer Res* 63: 3872-3886, 2003.
9. Baylin SB and Herman JG: DNA hypermethylation in tumorigenesis: epigenetics joins genetics. *Trends Genet* 16: 168-174, 2000.
10. Maesawa C, Tamura G, Nishizuka S, Ogasawara S, Ishida K, Terashima M, Sakata K, Sato N, Saito K and Satodate R: Inactivation of the *CDKN2* gene by homozygous deletion and *de novo* methylation is associated with advanced stage oesophageal squamous cell carcinoma. *Cancer Res* 56: 3875-3878, 1996.
11. Tanaka H, Shimada Y, Harada H, Shinoda M, Hatooka S, Imamura M and Ishizaki K: Methylation of the 5' CpG island of the *FHIT* gene is closely associated with transcriptional inactivation in oesophageal squamous cell carcinomas. *Cancer Res* 58: 3429-3434, 1998.
12. Xing EP, Nie Y, Song Y, Yang GY, Cai YC, Wang LD and Yang CS: Mechanisms of inactivation of *p14ARF*, *p15INK4b*, and *p16INK4a* genes in human oesophageal squamous cell carcinoma. *Clin Cancer Res* 5: 2704-2713, 1999.
13. Nie Y, Yang G, Song Y, Zhao X, So C, Liao J, Wang LD and Yang CS: DNA hypermethylation is a mechanism for loss of expression of the *HLA* class I genes in human oesophageal squamous cell carcinomas. *Carcinogenesis* 22: 1615-1623, 2001.
14. Si HX, Tsao SW, Lam KY, Srivastava G, Liu Y, Wong YC, Shen ZY and Cheung AL: E-cadherin expression is commonly downregulated by CpG island hypermethylation in oesophageal carcinoma cells. *Cancer Lett* 173: 71-78, 2001.
15. Zhang L, Lu WF, Miao XP, Xing DY, Tan W and Lin DX: Inactivation of DNA repair gene *O-6-methylguanine-DNA methyltransferase* by promoter hypermethylation and its relation to p53 mutations in oesophageal squamous cell carcinoma. *Carcinogenesis* 24: 1039-1044, 2003.

16. Kuroki T, Trapasso F, Yendamuri S, Matsuyama A, Alder H, Mori M and Croce CM: Allele loss and promoter hypermethylation of *VHL*, *RAR-beta*, *RASSF1A*, and *FHIT* tumour suppressor genes on chromosome 3p in oesophageal squamous cell carcinoma. *Cancer Res* 63: 3724-3728, 2003.
17. Wang Y, Fang MZ, Liao J, Yang GY, Nie Y, Song Y, So C, Xu X, Wang LD and Yang CS: Hypermethylation-associated inactivation of *retinoic acid receptor beta* in human oesophageal squamous cell carcinoma. *Clin Cancer Res* 9: 5257-5263, 2003.
18. Yue CM, Deng DJ, Bi MX, Guo LP and Lu SH: Expression of *ECRG4*, a novel oesophageal cancer-related gene, downregulated by CpG island hypermethylation in human oesophageal squamous cell carcinoma. *World J Gastroenterol* 9: 1174-1178, 2003.
19. Ito T, Shimada Y, Hashimoto Y, Kaganoi J, Kan T, Watanabe G, Murakami Y and Imamura M: Involvement of *TSLC1* in progression of oesophageal squamous cell carcinoma. *Cancer Res* 63: 6320-6326, 2003.
20. Hogg RP, Honorio S, Martinez A, Agathangelou A, Dallol A, Fullwood P, Weichselbaum R, Kuo MJ, Maher ER and Latif F: Frequent 3p allele loss and epigenetic inactivation of the *RASSF1A* tumour suppressor gene from region 3p21.3 in head and neck squamous cell carcinoma. *Eur J Cancer* 38: 1585-1592, 2002.
21. Lerman MI and Minna JD: The 630-kb lung cancer homozygous deletion region on human chromosome 3p21.3: identification and evaluation of the resident candidate tumour suppressor genes. The International Lung Cancer Chromosome 3p21.3 Tumour Suppressor Gene Consortium. *Cancer Res* 60: 6116-6133, 2000.
22. Dammann R, LC, Yoon JH, Chin PL, Bates S and Pfeifer GP: Epigenetic inactivation of a RAS association domain family protein from the lung tumour suppressor locus 3p21.3. *Nat Genet* 25: 315-319, 2000.
23. Hesson L, Dallol A, Minna JD, Maher ER and Latif F: *NORE1A*, a homologue of *RASSF1A* tumour suppressor gene is inactivated in human cancers. *Oncogene* 22: 947-954, 2003.
24. Burbee DG, Forgacs E, Zochbauer-Muller S, Shivakumar L, Fong K, Gao B, Randle D, Kondo M, Virmani A, Bader S, Sekido Y, Latif F, Milchgrub S, Toyooka S, Gazdar AF, Lerman MI, Zbarovsky E, White M and Minna JD: Epigenetic inactivation of *RASSF1A* in lung and breast cancers and malignant phenotype suppression. *J Natl Cancer Inst* 93: 691-699, 2001.
25. Dammann R, Takahashi T and Pfeifer GP: The CpG island of the novel tumour suppressor gene *RASSF1A* is intensely methylated in primary small cell lung carcinomas. *Oncogene* 20: 3563-3567, 2001.
26. Dreijerink K, Braga E, Kuzmin I, Geil L, Duh FM, Angeloni D, Zbar B, Lerman MI, Stanbridge EJ, Minna JD, Protopopov A, Li J, Kashuba V, Klein G and Zbarovsky ER: The candidate tumour suppressor gene, *RASSF1A*, from human chromosome 3p21.3 is involved in kidney tumorigenesis. *Proc Natl Acad Sci USA* 98: 7504-7509, 2001.
27. Lee MG, Kim HY, Byun DS, Lee SJ, Lee CH, Kim JI, Chang SG and Chi SG: Frequent epigenetic inactivation of *RASSF1A* in human bladder carcinoma. *Cancer Res* 61: 6688-6692, 2001.
28. Schagdarsurengin U, Gimm O, Hoang-Vu C, Dralle H, Pfeifer GP and Dammann R: Frequent epigenetic silencing of the CpG island promoter of *RASSF1A* in thyroid carcinoma. *Cancer Res* 62: 3698-3701, 2002.
29. Kwong J, Lo KW, To KF, Teo PM, Johnson PJ and Huang DP: Promoter hypermethylation of multiple genes in nasopharyngeal carcinoma. *Clin Cancer Res* 8: 131-137, 2002.
30. Qiu GH, Tan LKS, Loh KS, Lim CY, Srivastava G, Tsai ST, Tsao SW and Tao Q: The candidate tumour suppressor gene *BLU*, located at the commonly deleted region 3p21, is an E2F-regulated, stress-responsive gene, and inactivated by epigenetic and genetic mechanisms in nasopharyngeal carcinoma. *Oncogene* 23: 4793-4806, 2004.
31. Spugnardi M, Tommasi S, Dammann R, Pfeifer GP and Hoon DS: Epigenetic inactivation of RAS association domain family protein 1 (*RASSF1A*) in malignant cutaneous melanoma. *Cancer Res* 63: 1639-1643, 2003.
32. Schagdarsurengin U, Wilkens L, Steinemann D, Flemming P, Kreipe HH, Pfeifer GP, Schlegelberger B and Dammann R: Frequent epigenetic inactivation of the *RASSF1A* gene in hepatocellular carcinoma. *Oncogene* 22: 1866-1871, 2003.
33. Szumin I, Gillespie JW, Protopopov A, Geil L, Dreijerink K, Yang Y, Vocke CD, Duh FM, Zbarovsky E, Minna JD, Rhim JS, Emmert-Buck MR, Linehan WM and Lerman MI: The *RASSF1A* tumour suppressor gene is inactivated in prostate tumours and suppresses growth of prostate carcinoma cells. *Cancer Res* 62: 3498-3502, 2002.
34. Liu L, Yoon JH, Dammann R and Pfeifer GP: Frequent hypermethylation of the *RASSF1A* gene in prostate cancer. *Oncogene* 21: 6835-6840, 2002.
35. Murray PG, Qiu GH, Fu L, Waites ER, Srivastava G, Heys D, Agathangelou A, Latif F, Grundy RG, Mann JR, Starczynski J, Crocker J, Parkes SE, Ambinder RF, Young LS and Tao Q: Frequent epigenetic inactivation of the *RASSF1A* tumour suppressor gene in Hodgkin's lymphoma. *Oncogene* 23: 1327-1332, 2004.
36. Kwong D, Lam KY, Guan X, Law S, Wong J and Sham J: Chromosomal aberrations in oesophageal squamous cell carcinoma among Chinese: gain of 12p predicts poor prognosis after surgery. *Hum Pathol* 35: 309-316, 2004.
37. Ko JM, Wong CP, Tang CM, Lau KW and Lung ML: Frequent loss of heterozygosity on multiple chromosomes in Chinese oesophageal squamous cell carcinomas. *Cancer Lett* 170: 131-138, 2001.
38. Kuroki T, Trapasso F, Yendamuri S, Matsuyama A, Alder H, Mori M and Croce CM: Promoter hypermethylation of *RASSF1A* in oesophageal squamous cell carcinoma. *Clin Cancer Res* 9: 1441-1445, 2003.
39. Hu Y, Lam KY, Wan TS, Fang W, Ma ES, Chan LC and Srivastava G: Establishment and characterization of HKESC-1, a new cancer cell line from human oesophageal squamous cell carcinoma. *Cancer Genet Cytogenet* 118: 112-120, 2000.
40. Tang JC, Wan TS, Wong N, Pang E, Lam KY, Law SY, Chow LM, Ma ES, Chan LC, Wong J and Srivastava G: Establishment and characterization of a new xenograft-derived human oesophageal squamous cell carcinoma cell line SLMT-1 of Chinese origin. *Cancer Genet Cytogenet* 124: 36-41, 2001.
41. Hu YC, Lam KY, Law SY, Wan TS, Ma ES, Kwong YL, Chan LC, Wong J and Srivastava G: Establishment, characterization, karyotyping, and comparative genomic hybridization analysis of HKESC-2 and HKESC-3: two newly established human oesophageal squamous cell carcinoma cell lines. *Cancer Genet Cytogenet* 135: 120-127, 2002.
42. Deng W, Tsao SW, Guan XY, Lucas JN, Si HX, Leung CS, Mak P, Wang LD and Cheung AL: Distinct profiles of critically short telomeres are a key determinant of different chromosome aberrations in immortalized human cells: whole-genome evidence from multiple cell lines. *Oncogene* 23: 9090-9101, 2004.
43. Herman JG and Baylin SB: Gene silencing in cancer in association with promoter hypermethylation. *N Engl J Med* 349: 2042-2054, 2003.