

REVIEW ARTICLE

Dietary, Anticancer and Medicinal Properties of the Phytochemicals in Chili Pepper (*Capsicum* spp.)

M.D.M. Chamikara^{1,2}, D.R.R.P. Dissanayake¹, M. Ishan^{1,3} and S.D.S.S. Sooriyapathirana^{1,*}

¹Department of Molecular Biology and Biotechnology, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka

²Current Address: Structural Biology and Biophysics Laboratory, Research School of Chemistry, Australian National University, Pauline Griffin Building, [11], Canberra ACT 0200, Australia

³Current Address: Regenerative Bioscience Center, Department of Animal and Dairy Science, University of Georgia, Athens, GA-30602, USA

Received: 01/04/2016; Accepted: 15/09/2016

Abstract: Chili pepper (*Capsicum* spp.) is important as a spice, flavour enhancer, vegetable and component in herbal medicine. The numerous phytochemicals and their medicinally important properties present in diverse germplasm of chili pepper have been characterized and documented. Capsaicinoids, carotenoids, vitamins, flavonoids such as anthocyanins are present as the major phytochemicals in chili pepper fruits. Capsaicinoids, pungent analogues of capsinoids, are the most important group of phytochemicals in which capsaicin and dihydrocapsaicin are prominent in providing the basis for pungency and medicinal properties. The detailed studies conducted on the phytochemicals in chili pepper fruits using mouse models and human cell lines have reported an array of anticancer effects on leukemia, myeloma and various carcinomas associated with the digestive tract. In addition, chili pepper possesses anti-inflammatory, antidiabetic, antimicrobial, anticholesteremic, anticlotting and antioxidant activities. Therefore, this review is compiled to provide a comprehensive assessment of the phytochemical profile and medicinal values of chili pepper. Collectively, numerous studies performed to date demonstrated the potential medicinal significance of chili pepper for development of herbal medicine and to plan further studies to elucidate other hitherto unknown medicinal values.

Keywords: *Capsicum*, Capsaicin, Capsaicinoids, Phytochemicals, Anticancer activity

INTRODUCTION

Chili pepper, (*Capsicum* spp.), is an essential element in our daily cuisine. It is the most highly consumed spice in the world with a mean annual consumption of 3.5 million metric tons of

dried fruit (FAO, 2015). Characteristic pungency and flavour of the chili pepper fruits provide unique hotness to meal preparations and a wide variety of foods utilize this pungency in combination with salt and other spices for flavour enhancement (Pino *et al.*, 2007). The genus *Capsicum* is highly diverse and contains 40 species (The Plant List, 2013). Of those, five species, *C. annuum* var. *annuum*, *C. chinense*, *C. frutescens*, *C. baccatum* varieties *pendulum* and *umblicatum* and *C. pubescens* are most frequently consumed by humans (Moscone *et al.*, 2007). The domestication of chili pepper has been dated to 6000 B.C., and archeological evidence (genus specific starch-granule-morphotypes) found in the Bahamas and Southern Peru imply that chili pepper had been a component in diets as early as in 7500 B.C. (Perry, 2007).

Apart from the culinary uses, chili pepper has also been used in medicinal preparations, providing nutritional and health advantages such as anticancer, anti-inflammatory, antidiabetic, anticholesteremic, anticlotting, analgesic and antimicrobial effects. Mayans used chili pepper to treat microbial diseases (Cichewicz and Thorpe, 1996) and many reports from India show that chili pepper was used to treat asthma (even the chili root extracts were used), gastrointestinal tract problems (Baruah *et al.*, 2014), extreme pain, toothache, wounds, ulcers, arthritis (Bhagowati and Changkija, 2009), headaches and night blindness (Deorani and Sharma, 2007). Treating hypersensitivity and rheumatism using chili pepper also was common

in Italian indigenous medical practices (Pieroni *et al.*, 2004).

The indigenous uses of chili pepper in folk medicinal applications and the safe and prolonged use of chili pepper as a spice in human history enable us to think that the phytochemicals present in chili pepper fruits promote and maintain the good health of human beings (Bosland, 1996). These phytochemicals and their nutritional and medicinal values have been characterized in numerous studies. A few reviews have been published on metabolism and bioavailability (Rollyson *et al.*, 2014) and anticancer effects (Cao *et al.*, 2015; Pawar *et al.*, 2011) of capsaicin and nutraceutical benefits of chili pepper in general (Milind and Sushila, 2012). However, the array of medicinal properties in chili pepper has not been reviewed together to realize its multiple roles. This paper

reviews the medicinal values of chili pepper on diverse ailments including cancer, diabetics, cholesteromia and microbial infections.

PHYTOCHEMICAL PROFILE

The phenolic compounds present in the chili pepper are attributed to many medicinally important properties such as anti-diarrheal, antimicrobial, antioxidant, antihyperglycemic, anti-lithogenic and antimutagenic activities (Adefegha and Oboh, 2013). Chili pepper fruits contain phenolic compounds such as flavonoids (Chu *et al.*, 2002), β -catenin, capsaicinoids (Vera-Guzman *et al.*, 2011), carotenoids (capsanthin, capsorubin and cryptocapsin), vitamins A, B, C and E (Kidmose *et al.*, 2006; Niizu and Rodriguez-Amaya, 2005; USDA Nutrient Data Laboratory, 2015), glycolipids, glycerolipids and esters (Bijttebier *et al.*, 2014).




Figure 1: The structures of the common capsaicinoids in chili pepper.

Chemical composition of chili pepper fruit extracts has been characterized using GC/MS spectrophotometry (Wesolowska *et al.*, 2011) and liquid chromatography-photodiode array-accurate mass mass spectrometry (LC-PDA-amMS) (Bijtebier *et al.*, 2014). There are five major capsaicinoids observed in chili pepper (capsaicin, homocapsaicin, dihydrocapsaicin, nordihydrocapsaicin, and homodihydrocapsaicin) which contribute to the pungency (Baruah *et al.*, 2014; Reyes-Escogido *et al.*, 2011) (Figure 1). Other capsaicinoids such as nonvamide and n-vanillyl decanamide are also present in significant levels. The total capsaicinoid content is in the range of 58.8% to 84.5% in acetone and hexane extracts of the fruits (Wesolowska *et al.*, 2011). The capsaicin [8-methyl-N-vanilloid-6-nonenamide] and dihydrocapsaicin [N-(4-hydroxy-3-methoxybenzyl)-8-methylnonenamide] (Surh, 2002), present at levels of 37% and 29% of the total capsaicinoids respectively (Wesolowska *et al.*, 2011), provide 90% of the pungency of chili pepper (Giuffrida *et al.*, 2013).

Within the genus *Capsicum*, *C. chinense* contains the highest level of capsaicin (Bosland and Baral, 2007). The contents of tannins, flavonoids, saponins, terpenoids and carotenoids are higher in *C. annuum* than those of other species but *C. frutescens* contains the highest levels of flavonoids (Rahim and Mat, 2012) and anthraquinone (Emmanuel-Ikpeme *et al.*, 2014) compared to the other domesticated species. Although the mean vitamin E content in chili pepper ranges from 1.98% to 2.10% (Wesolowska *et al.*, 2011), the species *C. annuum* contains a higher vitamin E content than *C. frutescens* (Emmanuel-Ikpeme *et al.*, 2014). The well-known chili pepper cultivar, bell pepper contains 20.73 g of dietary-fiber and 133 mg of vitamin C per 100 g of fresh fruit weight (Durucasu and Tokusoglu, 2007). It has been revealed that *C. annuum* would be a good source of Zn which is an essential micronutrient (Emmanuel-Ikpeme *et al.*, 2014).

Effects of drying and ripening on phytochemicals

Conventional sun drying retains capsaicin in the chili pepper fruits (Magied *et al.*, 2014), and it has been confirmed that drying under low temperature assures the high quality of chili pepper by preserving the phenolic compounds and vitamin C in the fruits (Moraes *et al.*, 2013).

Generally chili pepper has a higher amount of vitamin C (ascorbic acid) than many other vegetables showing its significant nutritional value (Kumar and Tata, 2009). It is also interesting to note that vitamin C is preserved during drying. The total phenolic and carotenoid contents increase during ripening, where the total phenolic content is doubled and the carotenoid content is increased by 10-fold. Although the total flavonoid content increases during initial ripening (i.e. yellowing), it is later reduced as fruits turn red (i.e. at complete ripening) (Shaha *et al.*, 2013). The content of ascorbic acid is increased during drying of green chili peppers, while total phenolic content is decreased, resulting in lowered overall antioxidant capacity (Ozgun *et al.*, 2011). However, Wangcharoen and Morasuk, (2008) reported that when chili pepper was dried at a temperature more than 100 °C, higher amounts of phenolic compounds were retained and hence resulting an increased antioxidant activity. It is also interesting to note that the antioxidant activity of dried chili pepper was maintained for as long as 18 months after harvesting (Ogiso *et al.*, 2008).

ANTIOXIDANT ACTIVITY

The phytochemicals present in chili pepper possess anticancer properties (Chung *et al.*, 2005), among them, vitamin C and phenolic compounds can act as antioxidants to counteract effects of reactive oxygen species (ROS) (Podsedek, 2007). ROS, which are generated from cellular metabolic processes, have been associated with several diseases such as cancer, diabetes, cardiovascular and neurological disorders (Sun-Hwa *et al.*, 2007). Detoxification of ROS can be done by several enzymes present in the body as well as antioxidants obtained through dietary components (Halliwell, 2006) such as chili pepper. The antioxidant activity of chili pepper (Rosa *et al.*, 2002) is only second to spinach among commonly consumed fruits and vegetables (Pellegrini *et al.*, 2003). The antioxidant activity of chili pepper has been measured and studied using several assaying systems such as DPPH (2,2-diphenyl-1-picrylhydrazyl assay), ABTS (2,2'-azino-bis[3-ethylbenzothiazoline-6-sulphonic acid] assay), DMPD (Dimethyl-4-phenylenediamine assay), CuPRAC (Cupric reducing antioxidant capacity assay) and FRAP (Ferric ion reducing antioxidant power assay) (Krishna *et al.*, 2010). Chili pepper contains both phenolic and

anthocyanin compounds which possess antioxidant activity (Borovsky *et al.*, 2004; Howard *et al.*, 2000; Marín *et al.*, 2004; Mennen *et al.*, 2005). Dietary phenolic compounds are advantageous as they donate H atoms to free radicals generated inside the body and neutralize them (Borovsky *et al.*, 2004). The vitamins C and E present in chili pepper are also effective antioxidants (Sun-Hwa *et al.*, 2007). Chili pepper is consumed in both dried and fresh forms, potentially influencing antioxidant levels, although studies have provided conflicting results regarding the effect of drying (Wangcharoen and Morasuk, 2008).

There are also differences for antioxidant capacity among different species and cultivars of chili pepper. Assessment of the antioxidant activity in Turkish pepper cultivars revealed an approximately 7-fold range in antioxidant activity (2.57 to 18.96 mmol of trolox per kg) (Frary *et al.*, 2008). Out of the five chili pepper types tested, Bell and Caribe (a type of yellow fruited chili pepper) were reported to contain the highest antioxidant activity. The correlation between the phenolic content and the antioxidant activity was estimated as 91.4% (Medina-Juarez *et al.*, 2012) indicating that majority of the antioxidant activity in chili pepper is due to the phenolic compounds. The highest level of phenolic content and the highest antioxidant activity compared to other species and cultivars were detected in bell pepper by Rahim and Mat, (2012), which was verified by Nadeem *et al.*, (2011). *C. chinense* is also reported to contain higher levels of antioxidants such as carotenoids, ascorbate and glutathione and the concentration of glutathione is increased during ripening (Castro-Concha *et al.*, 2012). Glutathione and ascorbate are the major antioxidants which contribute to Halliwell Asada Cycle, where detoxification of H₂O₂ produced in chloroplast takes place without the involvement of enzyme catalase (Foyer and Noctor, 2009).

ANTICANCER ACTIVITY

Application of *C. chinense* fruit extracts to the HepG2 cell lines (derived from a hepatocellular carcinoma) demonstrated the inhibition of cancer cell proliferation. This inhibitory activity was independently verified by using methylthiazol tetrazolium (MTT), lactate dehydrogenase leakage and nitrous oxide (NO) production assays (Amruthraj *et al.*, 2014). The phytochemicals present in chili pepper trigger

apoptosis of malignant cells. Capsaicin elicits the apoptosis pathway by repressing the plasma membrane NADH oxidoreductase enzyme in mitochondria (Morre *et al.*, 1995, 1996). Capsaicin also binds with the ATP generating coenzyme Q, inhibiting its activity and destabilizing the electron flow in the mitochondria, thus producing ROS and triggering apoptosis (Macho *et al.*, 1998, 1999, 2000; Wolvetang *et al.*, 1996). The ROS have been shown to disrupt the mitochondrial membrane and induce apoptosis pathways in pancreatic cancer cells (Zhang *et al.*, 2008). Capsaicin triggered the apoptosis pathway in human KB cancer cells (derived from *HeLa* cell line) by agitating the membrane potential of mitochondria and subsequently activating the caspase signaling pathway (Lin *et al.*, 2013). However, capsaicin could also inhibit NF-κB (nuclear factor kappa-light-chain-enhancer of activated B cells) pathway which can cause apoptosis and minimize carcinogenic effects (Han *et al.*, 2001; Patel *et al.*, 2002). The studies on human cell cultures with exogenous capsaicin exhibited apoptosis (Ghosh and Basu, 2010), autophagy (Choi *et al.*, 2009; Oh *et al.*, 2010) and inhibition of cell metabolism (Arora *et al.*, 2011). The carotenoid pigments present in bell pepper such as capsanthin, capsorubin and cryptocapsin possess high free radical scavenging activity (Matsufuji *et al.*, 1998).

The experiments conducted using derived human cutaneous squamous cell carcinoma (SCC cell lines) concluded that capsaicin has prophylactic and therapeutic potentials to the skin cancers through the induction of apoptosis (Hail and Lotan, 2002) and modulating the epidermal growth factor receptor (EGFR) (Hwang *et al.*, 2010). Capsaicin also down-regulates the expression of *Bcl-2* (B-cell lymphoma 2) (Jun *et al.*, 2007), and thereby induces the apoptosis in B16-F10 melanoma cells (derived from *Mus musculus* skin melanoma). Chemically induced skin cancers in mouse models were also shown to be repressed by capsaicin (Park and Surh, 1997).

Capsaicin is identified as a blocker for the interleukin-6 induced transcription factor STAT3 [signal transducer and activator of transcription 3 (acute-phase response factor)] (Yu *et al.*, 2009) which promotes tumor regenerative pathways. Thus, capsaicin could be used as a preventive and treatment drug for myeloma and other

cancers (Bhutani *et al.*, 2007). *In vitro* studies using T-cell leukemia cell cultures have reported that capsaicin inhibits the growth of leukemia cells by degrading the TAX protein (human T-cell leukemia virus type 1 transcriptional transactivator) and increasing the NF- κ B inhibitor alpha (I κ B α) which arrests the cell cycle and triggers apoptosis. This demonstrates the possibility of using capsaicin as a chemopreventive drug for leukemia (Zhang *et al.*, 2003). Capsaicin is also known to induce the apoptosis pathway in leukemia cells through oxidative stress (Ito *et al.*, 2004).

In glioma cancers of brain, capsaicin binds to the transient receptor potential vanilloid type I (TRPV I) receptor in glioma cells and induces apoptosis (Amantini *et al.*, 2007). Gil and Kang, (2008) have also reported that capsaicin induces apoptosis in human glioma cells by down-regulating *Bcl-2* expression. Capsaicin is also known to possess inhibitory effects against the numerous cancers in the digestive system. Capsaicin possesses anti carcinogenic activity against tongue cancers (Tanaka *et al.*, 2002) by inducing apoptosis through the stimulation of the expression of caspase-3 and caspase-9 activities (Ip *et al.*, 2012a). Wu *et al.*, (2006) reported that capsaicin triggers apoptosis via activating caspase-3 and generating the ROS in CE 81T/VGH cells (derived from esophagus epidermoid carcinoma). It has also been reported that capsaicin heightened the expression of proto-oncogenes (such as *c-myc* and *c-Ha-ras*), which trigger the apoptosis pathway, and tumor suppressor gene *p53* in SNU-1 cell line (derived from human stomach cancer cell) (Kim *et al.*, 1997). Jung *et al.*, (2001) reported that capsaicin induces apoptosis by activating the caspase-3 and down-regulating the *Bcl-2* in hepatocarcinoma cells SK-Hep-1 human liver cell line (derived from the ascetic fluid of a patient with adenocarcinoma). In pancreatic cancer cells,

capsaicin also mediates ROS resulted mitochondrial damages and induces apoptosis. In colorectal cancer cells capsaicin is known to suppress the β -catenin dependent signaling pathway. Capsaicin restricts the expression of transcription factor 4 (TCF-4) and inhibits the interaction between β -catenin and TCF-4 suggesting the potential use of chili pepper as a treatment for colorectal cancers (Lee *et al.*, 2012). Lee *et al.*, (2010) also reported that capsaicin represses the cell proliferation in colorectal cancer cell lines.

The anticancer role of capsaicin is also observed in lung associated carcinoma. Capsaicin stabilizes the mitochondrial related enzymes in lungs which minimize the cancer risk induced by using benzopyrene in mouse models (Anandakumar *et al.*, 2007). Jang *et al.*, (1989) have reported that capsaicin reduces the risk of lung tumor development induced by polycyclic aromatic hydrocarbons. According to Brown *et al.*, (2010) capsaicin can be used to treat human small cell lung cancer since it has anti-proliferative activity in both cell culture experiments and mouse models by inhibiting the *E2F* (a family of transcription factor genes in higher eukaryotes) responses and proliferative gene expression which triggers the anti-proliferative activity. The apoptosis inducing mechanisms in NPC cell lines (derived from human nasopharyngeal carcinoma) are reported in Ip *et al.*, (2012b) and it was found that capsaicin induces endoplasmic reticulum stress, caspase-3 activation and mitochondrial depolarization. In addition capsaicin has an inhibitory activity against the development of prostate cancer by down-regulating the expression of prostate specific antigen (PSA) (Mori *et al.*, 2006). The reported inhibitory effects of the capsaicinoids on diverse cancer cell lines are summarized in Table 1.

Table 1: The inhibitory effects of capsaicinoids on diverse cell lines mimicking the anticancer activity.

Cancer	Cell line	Effect/s	Reference
Leukemia	HPB-ATL-T (adult T-cell leukemia)	Inhibit growth.	Zhang <i>et al.</i> , (2003)
	NB4 –PL (neuroblastoma promyelocytic leukemia) and Kasumi-1 (myeloid leukemia)	Induce apoptosis by oxidative stress.	Ito <i>et al.</i> , (2004)
	HL-60 (human myelocytic leukemia)	Induce apoptosis by caspase-3-dependent mechanism.	Tsou <i>et al.</i> , (2006)
Multiple myeloma	U266 (human multiple myeloma) and MM.1S (immunoglobulin A lambda myeloma)	Inhibit the tumorigenesis by blocking STAT3 [signal transducer and activator of transcription 3 (acute-phase response factor)].	Bhutani <i>et al.</i> , (2007)
Cutaneous cell carcinoma	SRB-12 (scavenger receptor class B type 12 derived from an epidermal lesion of a patient) and COLO 16 (squamous-cell carcinoma)	Arrest cell cycle at G1 stage and induce apoptosis by mitochondrial depolarization.	Hail and Lotan, (2002)
Glioma	A172 (human glioblastoma)	Induce apoptosis by generating ROS.	Lee <i>et al.</i> , (2000)
	FLS (fibroblast-like synoviocyte) and FC1 (glioblastoma)	Induce apoptosis by mitochondrial depolarization.	Amantini <i>et al.</i> , (2007)
Tongue cancer	SCC-4 (squamous-cell carcinoma) human tongue cancer cells	Induce apoptosis by mitochondria dependent and independent mechanisms	Ip <i>et al.</i> , (2012a)
Nasopharyngeal carcinoma	NPC-TW 039 (human nasopharyngeal carcinoma)	Induce apoptosis by endoplasmic reticulum stress and mitochondrial depolarization.	Ip <i>et al.</i> , (2012b)
Esophageal carcinoma	CE 81T/VGH (human esophagus epidermoid carcinoma)	Arrest cell cycle at G0-G1 phase.	Wu <i>et al.</i> , (2006)
Gastric cancer	SNU-1. NIH/3T3 (a Korean stomach cancer)	Alter the expression of tumor forming genes and induce the apoptosis.	Kim <i>et al.</i> , (1997)
Pancreatic cancer	AsPC-1 and BxPC-3 (human pancreatic cancer)	Induce apoptosis by generating ROS.	Zhang <i>et al.</i> , (2008)
		Induce apoptosis by oxidative stress and mitochondrial damage.	Pramanik <i>et al.</i> , (2011)
Hepato carcinoma	SK-Hep-1 (sloan kettering hepatocarcinoma)	Induce apoptosis by caspase-3-dependent mechanism.	Jung <i>et al.</i> , (2001)
	HepG2 (human hepatoma)	Induce apoptosis by ROS disruption.	Huang <i>et al.</i> , (2009)
Colon carcinoma	HT-29 (human colon cancer)	Induce apoptosis by activating the peroxisome proliferator-activated receptor gamma.	Kim <i>et al.</i> , (2004)
	Colo 205 (human colon cancer)	Induce apoptosis by caspase-8 dependent mechanism.	Lu <i>et al.</i> , (2010)
Lung cancer	NCI-H69, NCI-H82 (small cell lung cancer)	Arrest cell cycle at G1.	Brown <i>et al.</i> , (2010)
Breast cancer	MCF-7 (Michigan Cancer Foundation-7) (human breast cancer)	Induce apoptosis by caspase-independent pathway.	Chou <i>et al.</i> , (2009)
	MCF-7 (Michigan Cancer Foundation-7), T47D, BT-474, SKBR-3 and MDA-MB231 (breast cancer)	Arrest cell cycle and induce apoptosis by altering the EGFR/HER-2 pathway.	Thoennissen <i>et al.</i> , (2010)
Prostate cancer	LNcaP (lymph node carcinoma of the prostate), PC-3 and DU-145 (prostate cancer)	Confer anti- proliferative activity by downregulating PSA.	Mori <i>et al.</i> , (2006)
	PC-3 (prostate cancer)	Induce apoptosis by ROS generation.	Sanchez <i>et al.</i> , (2007)
Human KB carcinoma	KB (derived from HeLa cell line)	Arrest cell cycle at G2/M phase and induce apoptosis.	Lin <i>et al.</i> , (2013)

ANTI-INFLAMMATORY AND PAIN RELIEVING ACTIVITIES

Chili pepper extracts contain anti-inflammatory (Kim *et al.*, 2003; Lee *et al.*, 2005) and anti-allergic properties (Lee *et al.*, 2005). Out of the numerous phytochemicals in chili pepper, capsaicin is reported mainly to confer the anti-inflammatory activity (Kim *et al.*, 2003) but the colour pigment anthocyanin has also been reported to possess significant anti-inflammatory activity (Wang *et al.*, 1999). Capsaicin overwhelmed the obesity induced inflammatory responses from macrophages in adipose tissue (Kang *et al.*, 2007) indicating the potential applications as an analgesic. Inhibition of the expression of pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α) and interleukin 1 β (IL-1 β) by capsaicin reduces the inflammatory responses induced by antigens (Spiller *et al.*, 2008). It was also found that anti-inflammatory activity was not triggered by vanilloid receptor-1 but by inhibiting the I κ B- α , which degrades the lipopolysaccharide stimulated peritoneal macrophages (Kim *et al.*, 2003). Capsaicin and nordihydrocapsiate (a type of capsinoid) inhibit the early T cell activation events such as NF- κ B activation (Sancho *et al.*, 2002). Capsaicin can be used to treat rheumatoid arthritis, osteoarthritis and peripheral arthritis (Cordell and Araujo, 1993) as it reduces the inflammatory heat. It can be used to relieve the pain from noxious chemical hyperalgesia as well (Arora *et al.*, 2011; Fraenkel *et al.*, 2004) because it has an ability to control the secretion of neurotransmitter relating to the pain (Lynn, 1990). The analgesic activity of capsaicin can be used to minimize neuropathic pain conditions (Sindrup and Jensen, 1999). When recovering from cancer treatments, capsaicin can be used as a temporary pain relieving agent for oral mucositis which results from chemo and radiation therapies (Berger *et al.*, 1995). Cruz, (2004) reported that capsaicin can reduce the bladder pain associated with patients who have hypersensitive disorders.

ANTIDIABETIC/ANTIHYPERTENSIVE ACTIVITIES

It is estimated that by 2025, the number of diabetic patients in the world could go up to 300 million (Sy *et al.*, 2005) and therefore more efficient and economical treatments are needed.

Chili pepper, with a significant potential as a candidate plant for new drugs, has shown to possess promising antidiabetic activity (Okumura *et al.*, 2012) as it contains inhibitors for α -amylase and α -glucosidase which are required for the degradation of polysaccharides and disaccharides. Chili pepper is also reported to lower the absorption of D-glucose in the intestine, thus it can be used to control carbohydrate digestion and post-prandial glucose level rise (Kwon *et al.*, 2007) (reviewed in Adefegha and Oboh, 2013). The effect of capsaicin on carbohydrate metabolism in exercising and resting states were determined using human subjects and it was found that capsaicin increases the plasma epinephrine and norepinephrine by increasing O₂ consumption. This implies an increase of carbohydrate oxidation due to capsaicin (Lim *et al.*, 1997). Domotor *et al.*, (2006) further confirmed using human subjects, that capsaicin increases the glucose adsorption and increases the level of blood glucagon concentration. The obesity induced insulin resistance which is a metabolic disorder that can be minimized by using capsaicin reducing the risk of having type-2 diabetes in the future. Capsaicin alters the gene expression, lessens the glucose intolerance (hyperglycemia) and reduces the fasting glucose levels in mouse models (Kang *et al.*, 2010). Capsaicin is known to increase the rate of carbohydrate metabolism in human subjects and in addition, the rat models were used to gain a deep insight on the antidiabetic effects of chili pepper. Administration of chili pepper to rats reduces blood glucose levels (Magied *et al.*, 2014; Okumura *et al.*, 2012) through the induction of thermogenesis reactions such as lipid and glucose metabolism leading to the lowering of fatty acid storage in the body. Induction of thermogenesis reactions are achieved by activating the sympathetic nervous system which promotes thermogenesis reactions (Okumura *et al.*, 2012). *C. frutescens* is always a center of attraction in this regard as it was shown to increase the blood insulin level in type-2 diabetes of rats (Islam and Choi, 2008). Furthermore *C. frutescens* was found to increase the affinity of insulin towards its receptor (glucose) implying the potential of using chili pepper as a treatment for diabetes (Anthony *et al.*, 2013; Patel *et al.*, 2012).

ANTICHOLESTEREMIC ACTIVITY AND THE EFFECTS ON LIPID METABOLISM

Chili pepper is reported to possess hypocholesteremic and hypolipidemic activities, thus it can be used to treat and prevent cardiovascular diseases (Kempaiah and Srinivasan, 2002; Srinivasan *et al.*, 2004). Capsaicin can induce the thermogenesis and lipid metabolism pathways (Ahuja *et al.*, 2006) by inhibiting the fatty acid storage in the body (Okumura *et al.*, 2012; Yoshioka *et al.*, 1998) showing its potential to use as a treatment to control obesity. Capsaicin stimulates catecholamine secretion leading to the release of β_3 adrenergic stimulation which generates thermogenesis reaction (Kawada *et al.*, 1986; Jimenez *et al.*, 2002; Watanabe *et al.*, 1987). Capsaicin specifically alters the lipid metabolism by lipid oxidation of the high fat diets in the gastrointestinal tract. Kempaiah *et al.*, (2005) reported that capsaicin inhibits low density lipoprotein oxidation induced by copper (*in vitro*) and ferrous (*in vivo*) ions. Ahuja *et al.*, (2006) reported that capsaicin and dihydrocapsaicin inhibit the copper-induced serum lipoprotein oxidation. Capsaicin has proven protective effects on hepatic lipid peroxidation in rats (Kempaiah *et al.*, 2005). Another study on rats revealed that administering chili pepper along with high fat diet affects HDL and LDL cholesterol, total glyceroids and total lipid contents, and thereby reduces serum cholesterol significantly compared to that of control treatments (Magied *et al.*, 2014). Similarly, the serum lipoprotein content was reported to be reduced in human subjects who had daily chili pepper diets (Ahuja and Ball, 2006). Chili pepper can be used to control lipid oxidation since it possesses higher antioxidant activity than most of the other vegetables (Embuscado, 2015). Experiments with human subjects also have shown that dietary capsaicin induces abdominal fat loss by increasing fat oxidation (Snitker *et al.*, 2009). Further studies with human subjects with higher Body Mass Index (BMI) demonstrated that consumption of capsaicinoids increased fat oxidation rate and energy expenditure in the body, suggesting it may be valuable to facilitate reduction in BMI (Inoue *et al.*, 2007).

ANTICLOTTING EFFECT

The phytochemicals present in chili pepper have proven healing effects on cardiovascular diseases

in rodents (Ogunlade *et al.*, 2012). Experiments conducted using rat models suggested that chili pepper can reduce the risk of cardiovascular diseases and arthritis (Kritchevsky, 1992). Capsaicin was shown to inhibit platelet aggregation (Adams, 2009; Hogaboam and Wallace, 1991) by stabilizing the membranes of red blood cells via interfering with the enzyme phospholipase A2 (PLA2) (Wang *et al.*, 1984). Mouse model studies revealed that capsaicin, which inhibits platelet formation by inhibiting the clotting factors VIII: C and IX (Adams *et al.*, 2009), is more effective than aspirin in averting the acute pulmonary thromboembolism (Wang *et al.*, 1985). Therefore, capsaicin may have the potential to be developed as a therapeutic and preventive agent of cardiovascular diseases.

ANTIMICROBIAL ACTIVITY

Herbal materials have long been appreciated because of their proven antimicrobial effects (Alavijeh *et al.*, 2012). In addition to the use of chili pepper as a spice, it has been widely used as an agent to preserve food (Omolo *et al.*, 2014). The antimicrobial activity of the compounds present in *Capsicum* species were reported in many studies (Careaga *et al.*, 2003; Cichewicz and Thorpe, 1996; Omolo *et al.*, 2014). In addition to the pungency related phytochemicals, the pigment anthocyanin in chili pepper also possesses antimicrobial activity (Zhao *et al.*, 2009). The antibacterial activities of chili pepper were reported against the noxious pathogens *Bacillus cereus*, *B. subtilis*, *Clostridium sporogenes*, *C. tetani*, *Streptococcus pyogenes* (Cichewicz and Thorpe, 1996), *Staphylococcus aureus* (Molina-Torres *et al.*, 1999), *Escherichia coli*, *Pseudomonas aeruginosa*, *Sarcina lutea*, *Candida albicans* (Soetarno *et al.*, 1997) and *Vibrio cholera* (Chatterjee *et al.*, 2010). The bactericidal activity of capsaicin against the gastric pathogen *Helicobacter pylori* was evaluated and the minimum inhibitory concentration (MIC) was found to be 10 $\mu\text{g/ml}$ (Jones *et al.*, 1997) and confirmed by Zeyrek and Oguz, (2005). Similarly MIC values of 25 $\mu\text{g/ml}$ against *B. subtilis*, 200-300 $\mu\text{g/ml}$ against *E. coli* (Molina-Torres *et al.*, 1999), 15 $\mu\text{l/g}$ against *Salmonella typhimurium* and 5-15 $\mu\text{l/g}$ against *P. aeruginosa* (Careaga *et al.*, 2003) were reported. Capsaicin also possesses inhibitory activity against numerous antibiotic resistant microbial strains (Zeyrek and Oguz, 2005). The acetonitrile extracts of phytochemicals chrysoeriol,

dihydrocapsaicin and capsaicin from *C. frutescence* were found to have antimicrobial activity against *Enterococcus faecalis*, *B. subtilis*, *S. aureus*, *P. aeruginosa*, *Klebsiella pneumoniae*, *E. coli* and *C. albicans* (Nascimento *et al.*, 2014). The mechanism of antimicrobial activity of capsaicin was evaluated through DNA microarray technology and it was found that capsaicin has a toxic effect against yeast cells and induces their pleiotrophic drug resistance network which expresses genes related to osmotic stress and membrane biosynthesis (Kurita *et al.*, 2002).

OTHER MISCELLANEOUS MEDICINAL VALUES

It has long been believed that chili pepper promotes proper functioning and good health of digestive tract. Capsaicin stimulates the secretion of saliva and gastric juice which is vital for efficient digestion of food and to overcome gastrointestinal abnormalities (Bosland, 1996). Capsaicin also increases the gastric blood flow by inducing the gastric sub mucosal arteriolar dilation (Chen *et al.*, 1992) which is important for efficient food absorption. Zeyrek and Oguz, (2005) reported that capsaicin inhibits the growth of *H. pylori*, hence chili pepper may have value as a treatment for ulcers in the stomach and duodenum. Chili pepper exhibited gastric ulcer protective activity (Das *et al.*, 2008) and capsaicin provided hepato-protective activity as it promoted recovery from CCl₄ induced liver injuries in rat (Hassan *et al.*, 2012). Finally, ripe red chili pepper has shown inhibitory activity against the angiotensin-1 converting enzyme which increases blood pressure, suggesting that chili pepper has anti-hypertension activity (Ranilla *et al.*, 2010).

CONCLUSION

The phytochemicals present in chili pepper such as capsaicin have the potential to provide numerous medicinal benefits. It is clear that, ethno-medicinal applications reported in the historical records are supported by the latest findings on the medicinal properties of chili pepper. The anticancer activity is conferred through the inhibition of cancer proliferation, induction of apoptosis and detention of cell cycle. These activities were confirmed using animal models and human cancer cell lines. In addition, analgesic, antihyperglycemic, antibacterial and antifungal effects of chili

pepper were also reported. The healing effects of chili pepper on other disease conditions such as strokes, hypercholesterolemia, aging, obesity, and ulcers associated with the digestive tract were well studied. The underlying phytochemicals and their associated molecular pathways have been characterized to show the mechanisms of action. The inter-genera diversity of the medicinal properties have been studied though mainly limited to the fruits of two main species, *C. annuum* and *C. frutescence* and, the famous cultivar bell pepper which is also a *C. annuum*. Further studies are essential to explore and characterize more efficient bio-active compounds from the underutilized *Capsicum* spp. in diverse geographical locations. The wealth of information that is currently available regarding the medicinal properties of chili pepper can be used to develop novel drugs to combat cancers and other important ailments mentioned.

REFERENCES

- Adams, M.J. (2009). Effect of capsaicin and dihydrocapsaicin on *in vitro* blood coagulation and platelet aggregation. *Thrombosis Research*. **124**: 721-723.
- Adefegha, S.A. and Oboh, G. (2013). Phytochemistry and mode of action of some tropical spices in the management of type-2 diabetes and hypertension. *African Journal of Pharmacy and Pharmacology*. **7**: 332-346.
- Ahuja, K.D. and Ball, M.J. (2006). Effects of daily ingestion of chilli on serum lipoprotein oxidation in adult men and women. *British Journal of Nutrition*. **96**: 239-242.
- Ahuja, K.D., Kunde, D.A., Ball, M.J. and Geraghty, D.P. (2006). Effects of capsaicin, dihydrocapsaicin, and curcumin on copper-induced oxidation of human serum lipids. *Journal of Agricultural and Food Chemistry*. **54**: 6436-6439.
- Alavijeh, P.K., Alavijeh, P.K. and Sharma, D. (2012). A study of antimicrobial activity of few medicinal herbs. *Asian Journal of Plant Science and Research*. **2**: 496-502.
- Amantini, C., Mosca, M., Nabissi, M., Lucciarini, R., Caprodossi, S., Arcella, A., Giangaspero, F. and Santoni, G. (2007). Capsaicin-induced apoptosis of glioma cells is mediated by TRPV1 vanilloid receptor and requires p38 MAPK activation. *Journal of Neurochemistry*. **102**: 977-990.
- Amruthraj, N.J., Raj, J.P.P., Saravanan, S. and Lebel, A. (2014). *In vitro* studies on anticancer activity of capsaicinoids from *capsicum chinense* against human hepatocellular carcinoma cells.

- International Journal of Pharmacy and Pharmaceutical Sciences*. **6**: 254-558.
- Anandakumar, P., Kamaraj, S., Jagan, S., Ramakrishnan, G., Vinodhkumar, R. and Devaki, T. (2007). Stabilization of pulmonary mitochondrial enzyme system by capsaicin during benzo (a) pyrene induced experimental lung cancer. *Biomed Pharmacotherapy*. **62**: 390-394.
- Anthony, O.E., Ese, A.C. and Lawrence, E.O. (2013). Regulated effects of *Capsicum frutescens* supplemented diet (C.F.S.D) on fasting blood glucose level, biochemical parameters and body weight in alloxan induced diabetic wistar rats. *British Journal of Pharmaceutical Research*. **3**: 496-507.
- Arora, R., Gill, N.S., Chauhan, G. and Rana, A.C. (2011). An overview about versatile molecule Capsaicin. *International Journal of Pharmaceutical Sciences and Drug Research*. **3**: 280-286.
- Baruah, S., Zaman, M.K., Rajbongshi, P. and Das S. (2014). A review on recent researches on *Bhut jolokia* and pharmacological activity of Capsaicin. *International Journal of Pharmaceutical Sciences Review and Research*. **15**: 89-94.
- Berger, A., Henderson, M., Nadoolman, W., Duffy, V., Cooper, D., Sabersli, L. and Bartoshuk, L. (1995). Oral capsaicin provides temporary relief for oral mucositis pain secondary to chemotherapy/radiation therapy. *Journal of Pain and Symptom Management*. **10**: 243-248.
- Bhagowati, R.R. and Chankija, S. (2009). Genetic variability and traditional practices in Naga King Chilli landraces of Nagaland. *Asian Agri-History*. **3**: 171-180.
- Bhutani, M., Pathak, A.K., Nair, A.S., Kunnumakkara, A.B., Guha, S., Sethi, G. and Aggarwal, B.B. (2007). Capsaicin is a novel blocker of constitutive and Interleukin-6-inducible STAT3 activation. *Cancer Therapy: Preclinical*. **13**: 3024-3032.
- Bijttebier, S., Zhani, K., D'Hondt, E., Noten, B., Hermans, N., Apers, S. and Voorspoels, S. (2014). Generic characterization of apolar metabolites in red chili peppers (*Capsicum frutescens* L.) by orbitrap mass spectrometry. *Journal of Agriculture and Food Chemistry*. **62**: 4812-4831.
- Borovsky, Y., Oren-Shamir, M., Ovadia, R., De Jong, W. and Paran I. (2004). The A locus that controls anthocyanin accumulation in pepper encodes a MYB transcription factor homologous to Anthocyanin2 of *Petunia*. *Theoretical and Applied Genetics*. **109**: 23-29.
- Bosland, P.W. (1996). Capsicum: innovative uses of an ancient crop. In: J Janick (Eds.), *Progress in New crops*, ASHS press, Arlington, Virginia, USA Pp.479-487.
- Bosland, P.W. and Baral, J.B. (2007). 'Bhut Jolokia' - The world's hottest known chili pepper is a putative naturally occurring interspecific hybrid. *Hort Science*. **42**: 222-224.
- Brown, K.C., Witte, T.R., Hardman, W.E., Luo, H., Chen, Y.C., Carpenter, A.B., Lau, J.K. and Dasgupta, P. (2010). Capsaicin displays anti-proliferative activity against human small cell lung cancer in cell culture and nude mice models via the E2F pathway. *Public Library of Science ONE* **5**: e10243.
- Cao, S., Chen, H., Xiang, S., Hong, J., Weng, L., Zhu, H. and Liu, Q. (2015). Anti-Cancer effects and mechanisms of capsaicin in chili peppers. *American Journal of Plant Sciences*. **6**: 3075-3081.
- Careaga, M., Fernández, E., Dorantes, L., Mota, L., Jaramillo, M.E. and Hernandez-Sanchez, H. (2003). Antibacterial activity of *Capsicum* extract against *Salmonella typhimurium* and *Pseudomonas aeruginosa* inoculated in raw beef meat. *International Journal of Food Microbiology*. **83**: 331-335.
- Castro-Concha, L.A., Canche-Chuc, I. and Miranda-Ham, M.D.L. (2012). Determination of antioxidants in fruit tissues from three accessions of habanero pepper (*Capsicum chinense* Jacq.). *Journal of the Mexican Chemical Society*. **56**: 15-18.
- Chatterjee, S., Asakura, M., Chowdhury, N., Neogi, S.B., Sugimoto, N., Haldar, S., Awasthi, S.P., Hinenoya, A., Aoki, S. and Yamasaki, S. (2010). Capsaicin, a potential inhibitor of cholera toxin production in *Vibrio cholera*. *FEMS Microbiology Letters*. **306**: 54-60.
- Chen, R.Y., Li, D.S. and Guth, P.H. (1992). Role of calcitonin gene-related peptide in capsaicin-induced gastric submucosal arteriolar dilation. *American Journal of Physiology*. **262**: H1350-H1355.
- Choi, C.H., Jung, Y.K. and Oh, S. (2009). Selective induction of catalase mediated autophagy by dihydrocapsaicin in lung cell lines. *Free Radical Biology and Medicine*. **49**: 245-257.
- Chou, C.C., Wu, Y.C., Wang, Y.F., Chou, M.J., Kuo, S.J. and Chen, D.R. (2009). Capsaicin-induced apoptosis in human breast cancer MCF-7 cells through caspase independent pathway. *Oncology Reports*. **21**: 665-671.
- Chu, Y.F., Sun, J., Wu, X. and Liu, R.H. (2002). Antioxidant and antiproliferative activities of common vegetables. *Journal of Agriculture and Food Chemistry*. **50**: 6910-6916.
- Chung, S.Y., Sung, M.K., Kim, N.H., Jang, J.O. and Go, E.J. (2005). P-glycoprotein by natural products in human breast cancer cells. *Archives of Pharmacal Research*. **28**: 823-828.
- Cichewicz, R.H. and Thorpe, P.A. (1996). The antimicrobial properties of chili peppers (*Capsicum* species) and their uses in Mayan medicine. *Journal of Ethnopharmacology*. **52**: 61-70.

- Cordell, G.A. and Araujo, O.E. (1993). Capsaicin: identification, nomenclature, and pharmacotherapy. *Annals of Pharmacotherapy*. **27**: 330-336.
- Cruz, F. (2004). Mechanisms involved in new therapies for overactive bladder. *Urology*. **63**: 65-73.
- Das, S., Deka, S. and Gohain, K. (2008). A preclinical study on gastric ulcer protective activity of world's hottest chilli *Capsicum frutescences*. *Journal of Clinical and Diagnostic Research*. **2**: 1024-1027.
- Deorani, S.C. and Sarma, G.D. (2007). Medicinal plants of Nagaland. *Bishen Singh/Mahendra Pal Singh, Dehradun, India*. p.71
- Domotor, A., Szolcsányi, J. and Mózsik, G. (2006). Capsaicin and glucose absorption and utilization in healthy human subjects. *European Journal of Pharmacology* **534**: 280-283.
- Durucasu, I. and Tokusoglu, O. (2007). Effect of grilling on luteolin (3', 4', 5, 7-tetrahydroxyflavone) content in sweet green bell pepper (*Capsicum annum*). *Pakistan Journal of Biological Sciences*. **10**: 3410-3414.
- Embuscado, M.E. (2015). Spices and herbs: Natural sources of antioxidants – a mini review. *Journal of Functional Foods*. dx.doi.org/10.1016/j.jff.2015.03.005.
- Emmanuel-Ikpeme, C., Henry, P. and Okiri, O.A. (2014). Comparative evaluation of the nutritional, phytochemical and microbiological quality of three pepper varieties. *Journal of Food and Nutrition Sciences*. **2**: 74-80.
- Food and Agriculture Organization (FAO). *Food and Agriculture Organization of the United States*. Available from: <http://www.fao.org/home/en>. (Accessed on August 01, 2015).
- Foyer, C.H. and Noctor, G. (2009). Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. *Antioxidants & Redox Signaling*. **11**: 861-905.
- Fraenkel, L., Bogardus, S.T., Concato, J. and Wittink, D.R. (2004). Treatment options in knee osteoarthritis: the patient's perspective. *Archives of Internal Medicine*. **164**: 1299-1304.
- Frary, A., Keceli, M.A., Okmen, B., Sigva, H.O., Yemenicioglu, A. and Doganlar, S. (2008). Water-soluble antioxidant potential of turkish pepper cultivars. *Hort Science*. **43**: 631-636.
- Ghosh, A.K. and Basu, S., (2010). Fas-associated factor 1 is a negative regulator in capsaicin induced cancer cell apoptosis. *Cancer Letters* **287**: 142-149.
- Gil, Y. and Kang, M. (2008). Capsaicin induces apoptosis and terminal differentiation in human glioma A172 cells. *Life Sciences*. **82**: 997-1003.
- Giuffrida, D., Dugo, P., Torre, P., Bignardi, C., Cavazza, A., Corradini, C., and Dugo, G. (2013). Characterization of 12 *Capsicum* varieties by evaluation of their carotenoid profile and pungency determination. *Food Chemistry*. **140**: 794-802.
- Hail, N. and Lotan, R. (2002). Examining the role of mitochondrial respiration in vanilloid-induced apoptosis. *Journal of the National Cancer Institute*. **94**: 1281-1292.
- Halliwell, B. (2006). Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Plant Physiology*. **141**: 312-322.
- Han, S.S., Keum, Y.S., Seo, H.J., Chun, K.S., Lee, S.S. and Surh, Y.J. (2001). Capsaicin suppresses phorbol ester-induced activation of NF- κ B/Rel and AP-1 transcription factors in mouse epidermis. *Cancer Letters*. **164**: 119-126.
- Hassan, M.H., Edfawy, M., Mansour, A. and Hamed, A.A. (2012). Antioxidant and antiapoptotic effects of capsaicin against carbon tetrachloride-induced hepatotoxicity in rats. *Toxicology and Health*. **28**: 428-438.
- Hogaboam, C.M. and Wallace, J.L. (1991). Inhibition of platelet aggregation by capsaicin. An effect unrelated to actions on sensory afferent neurons. *European Journal of Pharmacology*. **202**: 129-131.
- Howard, L.R., Talcott, S.T., Brenes, C.H. and Villalon, B. (2000). Changes in phytochemical and antioxidant activity of selected pepper cultivars (*Capsicum* species) as influenced by maturity. *Journal of Agricultural and Food Chemistry*. **48**: 1713-1720.
- Huang, S.P., Chen, J.C., Wu, C.C., Chen, C.T., Tang, N.Y., Ho, Y.T., Lo, C., Lin, J.P., Chung, J.G. and Lin, J.G. (2009). Capsaicin-induced apoptosis in human hepatoma HepG2 cells. *Anticancer Research*. **29**: 165-174.
- Hwang, M.K., Bode, A.M., Byun, S., Song, N.R., Lee, H.J. and Dong, Z. (2010). Cocarcinogenic effect of Capsaicin involves activation of EGFR signaling but not TRPV1. *Therapeutics, Targets, and Chemical Biology*. DOI: 10.1158/0008-5472.CAN-09-4393
- Inoue, N., Matsunaga, Y., Satoh, H. and Takahashi, M. (2007). Enhanced energy expenditure and fat oxidation in humans with high BMI scores by the ingestion of novel and non-pungent capsaicin analogues (Capsaicinoids). *Bioscience, Biotechnology, and Biochemistry*. **71**: 380-389.
- Ip, S.W., Lan, S.H., Huang, A.C., Yang, J.S., Chen, Y.Y., Huang, H.Y., Lin, Z.P., Hsu, Y.M., Yang, M.D., Chiu, C.F. and Chung, J.G. (2012a). Capsaicin induces apoptosis in SCC-4 human tongue cancer cells through mitochondria-dependent and -independent pathways. *Environmental Toxicology*. **27**: 332-341.
- Ip, S.W., Lan, S.H., Lu, H.F., Huang, A.C., Yang, J.S., Lin, J.P., Huang, H.Y., Lien, J.C., Ho, C.C., Chiu, C.F., Wood, W.G. and Chung, J.G. (2012b). Capsaicin mediates apoptosis in human nasopharyngeal carcinoma NPC-TW 039 cells

- through mitochondrial depolarization and endoplasmic reticulum stress. *Human and Experimental Toxicology* **31**: 539-549.
- Islam, M.S. and Choi, H. (2008). Dietary red chilli (*Capsicumfrutescens* L.) is insulinotropic rather than hypoglycemic in type 2 diabetes model of rats. *Phytotherapy Research* **22**: 1025-1029.
- Ito, K., Nakazato, T., Yamato, K., Miyakawa, Y., Yamada, T., Hozumi, N., Segawa, K., Ikeda, Y. and Kizaki, M. (2004). Induction of apoptosis in leukemic cells by homovanillic acid derivative, capsaicin, through oxidative stress: implication of phosphorylation of p53 at Ser-15 residue by reactive oxygen species. *Cancer Research*. **64**: 1071-1078.
- Jang, J.J., Kim, S.H. and Yun, T.K. (1989). Inhibitory effect of capsaicin on mouse lung tumor development. *In Vivo*. **3**: 49-53.
- Jimenez, M., Leger, B., Canola, K., Lehr, L., Arboit, P., Seydoux, J., Russell, A.P., Giacobino, J.P., Muzzin, P. and Preitner, F. (2002). β_1 / β_2 / β_3 adrenoreceptor knockout mice are obese and cold sensitive but have normal lipolytic responses to fasting. *FEBS Letters*. **530**: 37-40.
- Jones, N.L., Shabib, S. and Sherman, P.M. (1997). Capsaicin as an inhibitor of growth of the gastric pathogen *Helicobacter pylori*. *FEMS Microbiology Letters*. **146**: 223-227.
- Jun, H.S., Park, T., Lee, C.K., Kang, M.K., Park, M.S., Kang, H.I., Surh, Y.J. and Kim, O.H. (2007). Capsaicin induced apoptosis of B16-F10 melanoma cells through down-regulation of Bcl-2. *Food and Chemical Toxicology* **45**: 708-715.
- Jung, M.Y., Kang, H.J. and Moon, A., (2001). Capsaicin-induced apoptosis in SK-Hep-1 hepatocarcinoma cells involves Bcl-2 down-regulation and caspase-3 activation. *Cancer Letters*. **165**: 139-145.
- Kang, J.H., Kim, C.S., Han, I.S., Kawada, T. and Yu, R. (2007). Capsaicin, a spicy component of hot peppers, modulates adipokine gene expression and protein release from obese-mouse adipose tissues and isolated adipocytes, and suppresses the inflammatory responses of adipose tissue macrophages. *FEBS Letters* **581**: 4389-4396.
- Kang, J.H., Tsuyoshi, G., Han, I.S., Kawada, T., Kim, Y.M. and Yu, R. (2010). Dietary capsaicin reduces obesity-induced insulin resistance and hepatic steatosis in obese mice fed a high-fat diet. *Obesity*. **18**: 780-787.
- Kawada, T., Watanabe, T., Takaishi, T., Tanaka, T. and Iwai, K. (1986). Capsaicin induced beta-adrenergic action on energy metabolism in rats: influence of capsaicin on oxygen consumption, the respiratory quotient, and substrate utilization. *Proceedings of the Society for Experimental Biology and Medicine*. **183**: 250-256.
- Kempaiiah, R.K. and Srinivasan, K. (2002). Integrity of erythrocytes of hypercholesterolemic rats during spices treatment. *Molecular and Cellular Biochemistry*. **236**: 155-161.
- Kempaiiah, R.K., Manjunatha, H. and Srinivasan, K. (2005). Protective effect of dietary capsaicin on induced oxidation of low-density lipoprotein in rats. *Molecular and Cellular Biochemistry*. **275**: 7-13.
- Kidmose, U., Yang, R.Y., Thilsted, S.H., Christensen, L.P. and Brandt, K. (2006). Content of carotenoids in commonly Asian vegetables and stability and extractability during frying. *Journal of Food Composition and Analysis*. **19**: 562-571.
- Kim, C.S., Kawada, T., Kim, B.S., Han, I.S., Choe, S.Y., Kurata, T. and Yu, R. (2003). Capsaicin exhibits anti-inflammatory property by inhibiting IkB- α degradation in LPS-stimulated peritoneal macrophages. *Cell Signal*. **15**: 299-306.
- Kim, C.S., Park, W.H., Park, J.Y., Kang, J.H., Kim, M.O., Kawada, T., Yoo, H., Han, I.S. and Yu, R. (2004). Capsaicin, a spicy component of hot pepper, induces apoptosis by activation of the peroxisome proliferator-activated receptor gamma in HT-29 human colon cancer cells. *Journal of Medicinal Food*. **7**: 267-273.
- Kim, J.D., Kim, J.M., Pyo, J.O., Kim, S.Y., Kim, B.S., Yu, R. and Han, I.S. (1997). Capsaicin can alter the expression of tumor forming-related genes which might be followed by induction of apoptosis of a Korean stomach cancer cell line, SNU-1. *Cancer Letters*. **120**: 235-241.
- Krishna, A.G.G., Lokesh, B.R., Sugasini, D. and Kancheva, V.D. (2010). Evaluation of the antiradical and antioxidant properties of extracts from Indian red chili and black pepper by *in vitro* models. *Bulgarian Chemical Communications*. **42**: 62-69.
- Kritchevsky, D. (1992). Antioxidant vitamins in the prevention of cardiovascular disease. *Nutrition Today*. **27**: 30-33.
- Kumar, O.A. and Tata, S.S. (2009). Ascorbic acid contents in chili peppers (*Capsicum* L.). *Notulae Scientia Biologicae*. **1**: 50-52
- Kurita, S.K., Kitagawa, E., Kim, C.H., Momose, Y. and Iwahashi, H. (2002). Studies of the antimicrobial mechanism of Capsaicin using yeast DNA microarray. *Bioscience, Biotechnology, and Biochemistry*. **66**: 532-536.
- Kwon, Y.I., Apostolidis, E., Kim, Y.C. and Shetty, K. (2007). Health benefits of traditional corn, beans and pumpkin: *In vitro* studies for hyperglycemia and hypertension management. *Journal of Medicinal Food*. **10**: 266-275.
- Lee, J.J., Crosby, K.M., Pike, L.M., Yoo, K.S. and Lescobar, D.I. (2005). Impact of genetic and environmental variation of development of flavonoids and carotenoids in pepper (*Capsicum* spp.). *Scientia Horticulturae*. **106**: 341-352.
- Lee, S., Richardson, R.L., Dashwod, R.H. and Baek, S.J. (2012). Capsaicin represses transcriptional activity of β -catenin in human colorectal cancer

- cells. *The Journal of Nutritional Biochemistry*. **23**: 646-655.
- Lee, S.H., Krisanapun, C. and Baek, S.J. (2010). NSAID-activated gene-1 as a molecular target for capsaicin-induced apoptosis through a novel molecular mechanism involving GSK3 β , C/EBP β , and ATF3. *Carcinogenesis*. **31**: 719-728.
- Lee, Y.S., Nam, D.H. and Kim, J.A. (2000). Induction of apoptosis by capsaicin in A172 human glioblastoma cells. *Cancer Letters*. **161**: 121-130.
- Lim, K., Yoshioka, M., Kikuzato, S., Kiyonaga, A., Tanaka, H., Shindo, M. and Suzuki, M. (1997). Dietary red pepper ingestion increases carbohydrate oxidation at rest and during exercise in runners. *Medicine & Science in Sports & Exercise*. **29**: 355-361.
- Lin, C., Lu, W., Wang, C., Chan, Y. and Chen, M. (2013). Capsaicin induces cell cycle arrest and apoptosis in human KB cancer cells. *BMC Complementary and Alternative Medicine*. **13**: 46.
- Lu, H.F., Chen, Y.L., Yang, J.S., Yang, Y.Y., Liu, J.Y., Hsu, S.C., Lai, K.C. and Chung, J.G. (2010). Antitumor activity of capsaicin on human colon cancer cells in vitro and Colo 205 tumor xenografts in vivo. *Journal of Agricultural and Food Chemistry*. **58**: 12999-13005.
- Lynn, B. (1990). Capsaicin: Action on nociceptive C-fibres and therapeutic potential. *Pain*. **41**: 61-69.
- Macho, A., Blázquez, M.V., Navas, P. and Muñoz, E. (1998). Induction of apoptosis by vanilloid compounds does not require de novo gene transcription and activator protein 1 activity. *Cell Growth and Differentiation*. **9**: 277-286.
- Macho, A., Calzado, M.A., Muñoz-Blanco, J., Gomez-Diaz, C., Gajate, C., Mollinedo, F., Navas, P. and Muñoz, E. (1999). Selective induction of apoptosis by capsaicin in transformed cells: the role of reactive oxygen species and calcium. *Cell Growth and Differentiation*. **6**: 155-165.
- Macho, A., Lucena, C., Calzado, M.A., Blanco, M., Donnay, I., Appendino, G. and Muñoz, E. (2000). Phorboid 20-homovanillates induce apoptosis through a VR1-independent mechanism. *Chemistry and Biology*. **7**: 483-492.
- Magied, M.M.A., Salama, N.A.R. and Ali, M.R. (2014). Hypoglycemic and hypocholesterolemia effects of intragastric administration of dried red chili pepper (*Capsicum annum*) in alloxan-induced diabetic male albino rats fed with high-fat-diet. *Journal of Food and Nutrition Research*. **11**: 850-856.
- Marín, A., Ferreres, F., Tomás-Barberán, F.A. and Gil, M.I. (2004). Characterization and quantitation of antioxidant constituents of sweet pepper (*Capsicum annum* L.). *Journal of Agricultural and Food Chemistry*. **52**: 3861-3869.
- Matsufuji, H., Nakamuro, H., Chino, M. and Mitsuhashi, T. (1998). Antioxidant activity of capsaicin and the fatty acid esters in paprika (*Capsicum annum*). *Journal of Agricultural and Food Chemistry*. **46**: 3462-3472.
- Medina-Juárez, L.A., Molina-Quijada, D.M.A., Toro-Sánchez, C.L.D., González-Aguilar, G.A. and Gámez-Meza, N. (2012). Antioxidant activity of peppers (*Capsicum annum* L.) extracts and characterization of their phenolic constituents. *Interciencia*. **37**: 588-593.
- Mennen, L.I., Walker, R., Bennetau-Pelissero, C. and Scalbert, A. (2005). Risks and safety of polyphenol consumption. *The American Journal of Clinical Nutrition*. **81**: 326-329.
- Milind, P. and Sushila, K. (2012). A hot way leading to healthy stay. *International Research Journal of Pharmacy*. **3**: 21-25.
- Molina-Torres, J., García-Chávez, A. and Ramírez-Chávez, E. (1999). Antimicrobial properties of alkaloids present in flavouring plants traditionally used in Mesoamerica: affinin and capsaicin. *Journal of Ethnopharmacology*. **64**: 241-248.
- Moraes, I.C.F., Sobral, P.J.D.A., Branco, I.G., Re, T.B. and Gomide, C.A. (2013). Dehydration of "dedo de moça" pepper: kinetics and phytochemical concentration. *Food Science and Technology (Campinas)*. **33**: 134-141.
- Mori, A., Lehmann, S., O'Kelly, J., Kumagai, T., Desmond, J.C., Pervan, M., McBride, W.H., Kizaki, M. and Koeffler, H.P. (2006). Capsaicin, a component of red peppers, inhibits the growth of androgen independent, p53 mutant prostate cancer cells. *Cancer Research*. **66**: 3222-3229.
- Morre, D.J., Chueh, P.J. and Morre, D.M. (1995). Capsaicin inhibits preferentially the NADH oxidase and growth of transformed cells in culture. *Proceedings of the National Academy of Sciences*. **92**: 1831-1835.
- Morre, D.J., Sun, E., Geilen, C., Wu, L.Y., de-Cabo, R., Krasagakis, K., Orfanos, C.E. and Morre, D.M. (1996). Capsaicin inhibits plasma membrane NADH oxidase and growth of human and mouse melanoma lines. *European Journal of Cancer*. **32A**: 1995-2003.
- Moscone, E.A., Scaldaferrro, M.A., Grabielle, M., Cecchini, N.M., Sánchez García, Y., Jarret, R., Daviña, J.R., Ducasse, D.A., Barboza, G.E. and Ehrendorfer, F. (2007). The evolution of chili peppers (*Capsicum* – Solanaceae): A cytogenetic perspective. *Acta Horticulturae*. **745**: 137-170.
- Nadeem, M., Anjum, F.M., Khan, M.R., Saeed, M. and Riaz, A. (2011). Antioxidant Potential of Bell Pepper (*Capsicum annum* L.)—A Review. *Pakistan Journal of Food Science*. **21**: 45-51.
- Nascimento, P.L.A., Nascimento, T.C.E.S., Ramos, N.S.M., Silva, G.R., Gomes, J.E.G., Falcao, R.A.E., Moreira, K.A., Porto, A.L.F. and Silva, T.M.S. (2014). Quantification, Antioxidant and Antimicrobial Activity of Phenolics Isolated from Different Extracts of *Capsicum frutescens* (Pimenta Malagueta). *Molecules*. **19**: 5434-5447.

- Niizu, P. and Rodriguez-Amaya, D.B. (2005). New data on the carotenoid composition of raw salad vegetables. *Journal of Food Composition and Analysis*. **18**: 739-749.
- Ogiso, Y., Yabe, R.H., Kawamoto, Y., Kawamoto, T., Kato, K. and Yabe, T. (2008). An antioxidant of dried chili pepper maintained its activity through postharvest ripening for 18 months. *Bioscience, Biotechnology, and Biochemistry*. **72**: 3297-3300.
- Ogunlade, I., Alebiosu, A.A. and Osasona, A.I. (2012). Proximate, mineral composition, antioxidant activity, and total phenolic content of some pepper varieties (*Capsicum* species). *International Journal of Biological and Chemical Sciences*. **6**: 2221-2227.
- Oh, S., Choi, C.H. and Jung, Y.K. (2010). Autophagy induction by capsaicin in malignant human breast cells is modulated by p38 and ERK mitogen-activated protein kinase and retards cell death by suppressing endoplasmic reticulum stress mediated apoptosis. *Molecular Pharmacology*. **78**: 114-125.
- Okumura, T., Tsukui, T., Hosokawa, M. and Miyashita, K. (2012). Effect of caffeine and capsaicin on the blood glucose levels of obese/diabetic KK-A(y) mice. *Journal of Oleo Science*. **61**: 515-523.
- Omolo, M.A., Wong, Z.Z., Mergen, A.K., Hastings, J.C., Le, N.C., Reiland, H.A., Case, K.A. and Baumier, D.J. (2014). Antimicrobial Properties of Chili Peppers. *Infectious Diseases and Therapy*. **2**: 145 doi:10.4172/2332-0877.1000145.
- Ozgun, M., Ozcan, T., Akpınar-Bayazit, A. and Yilmaz-Ersan, L. (2011). Functional compounds and antioxidant properties of dried green and red peppers. *African Journal of Agricultural Research*. **6**: 5638-5644.
- Park, K.K. and Surh, Y.J. (1997). Effects of capsaicin on chemically-induced two-stage mouse skin carcinogenesis. *Cancer Letters*. **114**: 183-184.
- Patel, D.K., Prasad, S.K., Kumar, R. and Hemalatha, S. (2012). An overview on antidiabetic medicinal plants having insulin mimetic property. *Asian Pacific Journal of Tropical Biomedicine*. **2**: 320-330.
- Patel, P.S., Varney, M.L., Dave, B.J. and Singh, R.K. (2002). Regulation of constitutive and induced NF- κ B activation in malignant melanoma cells by capsaicin modulates interleukin-8 production and cell proliferation. *Journal of Interferon and Cytokine Research*. **22**: 427-435.
- Pawar, S.S., Bharude, N.V., Sonone, S.S., Deshmukh, R.S., Raut, A.K. and Umalkar, A.R. (2011). Chillies as food, spice and medicine: a perspective. *International Journal of Pharmacy and Biological Sciences*. **1**: 311-318.
- Pellegrini, N., Serafini, M., Colombi, B., Del Rio, D., Salvatore, S., Bianchi, M. and Brighenti F. (2003). Total antioxidant capacity of plant foods, beverages and oils consumed in Italy assessed by three different in vitro assays. *Journal of Nutrition*. **133**: 2812-2819.
- Perry, L. (2007). Starch fossils and the domestication and dispersal of chili pepper (*Capsicum* spp. L.) in the Americas. *Science*. **315**: 986-988.
- Pieroni, A., Quave, C.L. and Santoro, R.F. (2004). Folk pharmaceutical knowledge in the territory of the Dolomiti Lucane, inland southern Italy. *Journal of Ethnopharmacology*. **95**: 373-384.
- Pino, J., Gonzalez, M., Ceballos, L., Centurio'n-Yah, A.R., Trujillo-Aguirre, M.J., Latournerie-Moreno, L. and Sauri-Duch, E. (2007). Characterization of total capsaicinoids, colour and volatile compounds of Habanero chilli pepper (*Capsicum chinense* Jack.) cultivars grown in Yucatan. *Food Chemistry*. **104**: 1682-1686.
- Podsedek, A. (2007). Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *LWT - Food Science and Technology*. **40**: 1-11.
- Pramanik, K.C., Boreddy, S.R. and Sirivasatava, S.K. (2011). Role of mitochondrial electron transport chain complexes in capsaicin mediated oxidative stress leading to apoptosis in pancreatic cancer cells. *Public Library of Science ONE*. **6**: e20151
- Rahim, R.A. and Mat, I. (2012). Phytochemical contents of *Capsicum frutescens* (Chili Padi), *Capsicum annum* (Chili Pepper) and *Capsicum annum* (Bell Peper) aqueous extracts. *International Conference on Biological and Life Sciences*. **40**: 164-167.
- Ranilla, L.G., Kwon, Y.I., Apostolidis, E. and Shetty, K. (2010). Phenolic compounds, antioxidant activity and in vitro inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly used medicinal plants, herbs and spices in Latin America. *Bioresource Technology*. **101**: 4676-4689.
- Reyes-Escogido, M.L., Gonzalez-Mondragon, E.G. and Vazquez-Tzompantzi, E. (2011). Chemical and pharmacological aspects of capsaicin. *Molecules*. **16**: 1253-1270.
- Rollyson, W.D., Stover, C.A., Brown, K.C., Perry, H.E., Stevenson, C.D., McNeese, C.A., Ball, J.G., Valentovic, M.A. and Dasgupta, P. (2014) Bioavailability of capsaicin and its implications for drug delivery. *Journal of Controlled Release*. **196**: 96-105.
- Rosa, A., Deiana, M., Casu, V., Paccagnini, S., Appendino, G., Ballero, M. and Assunta Dessiä,M. (2002).Antioxidant activity of capsinoids. *Journal of Agricultural and Food Chemistry*. **50**: 7396-7401.
- Sanchez, A.M., Malagarie-Cazenave, S., Olea, N., Vara, D., Chiloeches, A. and Díaz- Laviada, I. (2007). Apoptosis induced by capsaicin in prostate PC-3 cells involves ceramide accumulation, neutral sphingomyelinase, and JNK activation. *Apoptosis*. **12**: 2013-2024.

- Sancho, R., Lucena, C., Macho, A., Calzado, M.A., Blanco-Molina, M., Minassi, A., Appendino, G., and Munoz, E. (2002). Immunosuppressive activity of capsaicinoids: capsiate derived from sweet peppers inhibits NF- κ B activation and is a potent anti-inflammatory compound *in vivo*. *European Journal of Immunology*. **32**: 1753-1763.
- Shaha, R.K., Rahman, S. and Asrul, A. (2013). Bioactive compounds in chilli peppers (*Capsicum annum* L.) at various ripening (green, yellow and red) stages. *Annals of Biological Research*. **4**: 27-34.
- Sindrup, S.H. and Jensen, T.S. (1999). Efficacy of pharmacological treatments of neuropathic pain: An update and effect related to mechanism of drug action. *Pain*. **83**: 389-400.
- Snitker, S., Fujishima, Y., Shen, H., Ott, S., Pi-Sunyer, X., Furuhashi, Y., Sato, H., and Takahashi, M. (2009). Effects of novel capsinoid treatment on fatness and energy metabolism in humans: possible pharmacogenetic implications. *The American Journal of Clinical Nutrition*. **89**: 45-50.
- Soetarno, S., Sukrasno, Yulinah, E. and Sylvia. (1997). Antimicrobial Activities of the Ethanol Extracts of *Capsicum* Fruits with Different Levels of Pungency. *Journal of Membrane Science*. **2**: 57-63.
- Spiller, F., Alves, M.K., Vieira, S.M., Carvalho, T.A., Leite, C.E., Lunardelli, A., Polomi, J.A., Cunha, F.Q. and Oliveira, J.R. (2008). Anti-inflammatory effects of red pepper (*Capsicum baccatum*) on carrageenan- and antigen-induced inflammation. *Journal of Pharmacy and Pharmacology*. **60**: 473-478.
- Srinivasan, K. Sambaiah, K. and Chandrasekhara, N. (2004). Spices as beneficial hypolipidemic food adjuncts: A Review. *Food Reviews International*. **20**: 187-220.
- Sun-Hwa, H., Jung-Bong, K., Jong-Sug, P., Shin-Woo, L. and Kang-Jim, C. (2007). A comparison of the carotenoid accumulation in *Capsicum* varieties that show different ripening colours: deletion of the capsanthin-capsorubin synthase gene is not a prerequisite for the formation of a yellow pepper *Journal of Experimental Botany*. **58**: 3135-3144.
- Surh, Y.J. (2002). More than spice: capsaicin in hot chili peppers makes tumor cells commit suicide. *Journal of the National Cancer Institute*. **94**: 1263-1265.
- Surh, Y.J. (2003). Cancer chemoprevention with dietary phytochemicals. *Reviews*. **3**: 768-780.
- Sy, G.Y., Cissé, A., Nongonierma, R.B., Sarr, M., Mbodj, N.A. and Faye, B. (2005). Hypoglycaemic and antidiabetic activity of acetonetic extract of *Vernonia colorata* leaves in normoglycaemic and alloxan-induced diabetic rats. *Journal of Ethnopharmacology*. **98**: 171-175.
- Tanaka, T., Kohno, H., Sakata, K., Yamada, Y., Hirose, Y., Sugie, S. and Mori, H. (2002). Modifying effects of dietary capsaicin and rotenone on 4-nitroquinoline 1-oxide-induced rat tongue carcinogenesis. *Carcinogenesis*. **23**: 1361-1367.
- The Plant List (2013). *Version 1.1*. Available from: URL: <http://www.theplantlist.org/> (Accessed on November 11, 2015).
- Thoennissen, N.H., O'Kelly, J., Lu, D., Iwanski, G.B., La, D.T., Abbassi, S., Leiter, A., Karlan, B., Mehta, R. and Koeffler, H.P. (2010). Capsaicin causes cell-cycle arrest and apoptosis in ER-positive and -negative breast cancer cells by modulating the EGFR/HER-2 pathway. *Oncogene*. **29**: 285-296.
- Tsou, M.F., Lu, H.F., Chen, S.C., Wu, L.T., Chen, Y.S., Kuo, H.M., Lin, S.S. and Chung, J.G. (2006). Involvement of Bax, Bcl-2, Ca²⁺ and caspase-3 in capsaicin-induced apoptosis of human leukemia HL-60 cells. *Anticancer Research*. **26**: 1965-1971.
- USDA Nutrient Data Laboratory. (2015). USDA national nutrient database for standard reference. Available from: URL: <http://ndb.nal.usda.gov/ndb/foods/show/> (Accessed on August 01, 2015).
- Vera-Guzman, A., Chavez-Servia, J.L., Carrillo-Rodriguez, J.C. and Lopez, M.G. (2011). Phytochemical evaluation of wild and cultivated pepper (*Capsicum annum* L. and *C. pubescens* Ruiz & Pav.) from Oaxaca, Mexico. *Chilean Journal of Agricultural Research*. **71**: 578-585.
- Wang, H.B., Nair, M.G., Strasburg, G. M., Chang, Y. C., Booren, A.M., Gray, J.I., DeWitt, D.L. (1999). Antioxidant and antiinflammatory activities of anthocyanins and their aglycon, cyanidin, from tart cherries. *Journal of Natural Products*. **62**: 294-296.
- Wang, J.P., Hsu, M.F. and Teng, C.M. (1984). Antiplatelet effect of capsaicin. *Thrombosis Research* **36**: 497-507.
- Wang, J.P., Hsu, M.F., Hsu, T.P. and Teng, C.M. (1985). Antihemostatic and antithrombotic effects of capsaicin in comparison with aspirin and indomethacin. *Thrombosis Research*. **37**: 669-679.
- Wangcharoen, W. and Morasuk, W. (2008). Antioxidant capacity changes in Chili Spur Pepper (*Capsicum annum* Linn. var. *acuminatum* Fingerh.) during drying process. *Asian Journal of Food and Agro-Industry*. **1**: 68-77.
- Watanabe, T., Kawada, T., Yamamoto, M. and Iwai, K. (1987). Capsaicin, a pungent principle of hot red pepper, evokes catecholamine secretion from the adrenal medulla of anesthetized rats. *Biochemical and Biophysical Research Communications*. **142**: 259-264.
- Wesolowska, A., Jadczyk, D. and Grzeszczuk, M. (2011). Chemical composition of the pepper fruit extracts of hot cultivars *Capsicum annum* L. *Acta Scientiarum Polonorum Horticulture*. **10**: 171-184.

- Wolvetang, E.J., Larm, J.A., Moutsoulas, P. and Lawen, A. (1996). Apoptosis induced by inhibitors of the plasma membrane NADH-oxidase involves Bcl-2 and calcineurin. *Cell Growth and Differentiation*. **7**: 1315-1325.
- Wu, C.C., Lin, J.P., Yang, J.S., Chou, S.T., Chen, S.C., Lin, Y.T. and Chung, J.G. (2006). Capsaicin induced cell cycle arrest and apoptosis in human esophagus epidermoid carcinoma CE 81T/VGH cells through the elevation of intracellular reactive oxygen species and Ca²⁺ productions and caspase-3 activation. *Mutation Research*. **601**: 71-82.
- Yoshioka, M., St-Pierre, S., Suzuki, M. and Tremblay, A. (1998). Effects of red pepper added to high-fat and high-carbohydrate meals on energy metabolism and substrate utilization in Japanese women. *British Journal of Nutrition*. **80**: 503-510.
- Yu, H., Pardoll, D. and Jove, R. (2009). STATs in cancer inflammation and immunity: a leading role for STAT3. *Nature Reviews Cancer*. **9**: 798-809.
- Zeyrek, Y.F. and Oguz, E. (2005). Invitro activity of capsaicin against *Heylicobacter pylori*. *Annals of Microbiology*. **55**: 125-127.
- Zhang, J., Nagasaki, M., Tanaka, Y. and Morikawa, S. (2003). Capsaicin inhibits growth of adult T-cell leukemia cells. *Leukemia Research*. **27**: 275-283.
- Zhang, R., Humphreys, I., Sahu, R.P., Shi, Y. and Sirivastava, S.K. (2008). *In vitro* and *in vivo* induction of apoptosis by capsaicin in pancreatic cancer cells is mediated through ROS generation and mitochondrial death pathway. *Apoptosis*. **13**: 1465-1478.
- Zhao, X.Y., Zhang, C., Guigas, C., Ma, Y., Corrales, M., Tauscher, B. and Hu, X. S. (2009). Composition, antimicrobial activity, and anti-proliferative active capacity of anthocyanin extracts of purple corn (*Zea mays* L.) from china. *European Food Research and Technology*. **228**: 759-765.
-