

Short-term Associations between Ambient Air Pollutants and Pediatric Asthma Emergency Department Visits

Matthew J. Strickland (1), Lyndsey A. Darrow (1), Mitchel Klein (1), W. Dana Flanders (2), Jeremy A. Sarnat (1), Lance A. Waller (3), Stefanie E. Sarnat (1), James A. Mulholland (4), Paige E. Tolbert (1)

Author Affiliations:

(1) Department of Environmental and Occupational Health, Rollins School of Public Health, Emory University

(2) Department of Epidemiology, Rollins School of Public Health, Emory University

(3) Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory University

(4) School of Civil and Environmental Engineering, Georgia Institute of Technology

Corresponding Author:

Matthew J. Strickland, PhD, MPH

Department of Environmental and Occupational Health

1518 Clifton Rd NE

Atlanta GA 30322

Tel: 404-712-8912 Fax: 404-727-8744 Email: mjstric@emory.edu

Sources of support: NIEHS R01ES012967, EPA STAR R82921301, and EPRI EP-P277231/C13172

Running title: Air pollution and pediatric asthma

Subject category number: 118 Air Pollution

Word Count: 3,891

Scientific Knowledge of the Subject:

Certain outdoor air pollutants can trigger asthma exacerbations. To advance understanding of the relationships between ambient air pollutant concentrations and asthma exacerbations among children, further characterization of the dose-response and pollutant lag effects are needed, as are investigations of pollutant species beyond the commonly measured criteria pollutants.

What this Study Adds to the Field:

In our large, population-based study we observed that both ozone and primary pollutants from traffic sources independently contributed to the burden of emergency department visits for pediatric asthma. These associations tended to be of the highest magnitude for concentrations on the day of the emergency department visit and were present at relatively low ambient concentrations.

This article has an online data supplement, which is accessible from this issue's table of content online at www.atsjournals.org

ABSTRACT

Rationale: Certain outdoor air pollutants cause asthma exacerbations in children. To advance understanding of these relationships, further characterization of the dose-response and pollutant lag effects are needed, as are investigations of pollutant species beyond the commonly measured criteria pollutants.

Objectives: Investigate short-term associations between ambient air pollutant concentrations and emergency department visits for pediatric asthma.

Methods: Daily counts of emergency department visits for asthma or wheeze among children age 5–17 were collected from 41 Metropolitan Atlanta hospitals during 1993–2004 (n = 91,386 visits). Ambient concentrations of gaseous pollutants and speciated particulate matter were available from stationary monitors during this time period. Rate ratios for the warm season (May–October) and cold season (November–April) were estimated using Poisson generalized linear models in the framework of a case-crossover analysis.

Measurements and Main Results:

Both ozone and primary pollutants from traffic sources were associated with emergency department visits for asthma or wheeze; evidence for independent effects of ozone and primary pollutants from traffic sources were observed in multipollutant models. These associations

tended to be of the highest magnitude for concentrations on the day of the emergency department visit and were present at relatively low ambient concentrations.

Conclusions: Even at relatively low ambient concentrations, ozone and primary pollutants from traffic sources independently contributed to the burden of emergency department visits for pediatric asthma.

Abstract word count: 217

Key words: Ambient Particulate Matter, Asthma, Minors, Ozone

INTRODUCTION

A broad literature supports associations between ambient air pollutant concentrations and asthma exacerbations (1-3). Children are thought to be particularly susceptible to ambient air pollutants, because their lungs and immune systems are not fully developed; they breathe more air per unit body weight and are typically more active than adults; and their peripheral airways are anatomically smaller than adults so that inflammation results in proportionally greater airway obstruction (4-6). To help advance understanding of the relationships between ambient air pollutant concentrations and asthma exacerbations in children, further characterization of the dose-response and pollutant lag effects are needed, as are investigations of pollutant species beyond the commonly measured urban air pollutants (3, 7). Further investigation of pollutant mixtures and effect modification may also provide insights (8, 9); for example, there have been reports of stronger pollution effects during the warm season (10-15), even though pediatric asthma rates peak during the cold season (16). To lessen concerns about uncontrolled confounding, aggressive control for variables such as meteorology and seasonal asthma trends is required.

In the present study, we analyzed data from SOPHIA (Study of Particles and Health in Atlanta) (14, 17-21), one of the largest single-city time-series studies of the health effects of urban air pollutants, to investigate short-term associations between ambient air pollutant concentrations and pediatric emergency department visits for asthma or wheeze in metropolitan Atlanta, USA,

during 1993–2004. Our study takes advantage of daily measurements of components of particulate matter less than 2.5 μm in aerodynamic diameter ($\text{PM}_{2.5}$), a resource not typically available to investigators, which enables us to distinguish among the various compounds that comprise $\text{PM}_{2.5}$. We investigated effect modification by season and potential confounding by ambient pollen concentrations and upper respiratory infections, because it is well known that these factors can trigger asthma exacerbations (9). The extent to which ambient pollen concentrations and circulating respiratory viruses confound associations between air pollutant concentrations and asthma exacerbations depends on the nature of the associations between short term changes in these risk factors and changes in air pollutant concentrations. We also present several characterizations of dose-response relationships, descriptions of pollution lag effects, and results from multipollutant models and models of pollutant interactions. Some of the results from this study have been previously reported in the form of an abstract (22).

METHODS

We obtained data on emergency department visits from metropolitan Atlanta hospitals during 1993–2004 (18). Using the International Classification of Diseases, 9th Revision, we defined emergency department visits for pediatric asthma as all visits with a code for asthma (493.0–493.9) or wheeze (786.09 before October 1, 1998; 786.07 after October 1, 1998) that did not have a code for an external injury or poisoning (E800–E999) among children age 5 to 17 years ($n = 91,386$). We also identified emergency department visits for acute respiratory infections (codes 460.0–466.0) that did not have a code for asthma or wheeze among children age 5 to 17 years ($n = 154,300$).

Daily concentrations of ambient 1-hour maximum carbon monoxide, nitrogen dioxide, and sulfur dioxide; 8-hour maximum ozone; and 24-hour average particulate matter less than 10 μm in diameter (PM_{10}), coarse particles between 2.5 and 10 μm in diameter ($\text{PM}_{10-2.5}$), $\text{PM}_{2.5}$, and the $\text{PM}_{2.5}$ components sulfate, elemental carbon, organic carbon, and water-soluble metals (comprised of water-soluble chromium, copper, iron, manganese, nickel, and vanadium) were obtained from several networks of ambient monitors (14, 21, 23). These pollutants reflect many of the predominant sources of air pollution in Atlanta (24-26). Measurement of $\text{PM}_{10-2.5}$, $\text{PM}_{2.5}$, and the $\text{PM}_{2.5}$ components began in August, 1998. Daily ambient airborne pollen concentrations were obtained from the Atlanta Allergy and Asthma Clinic.

In the framework of a case-crossover analysis, associations between ambient air pollutant concentrations and pediatric asthma emergency department visits were estimated via Poisson generalized linear models that accounted for overdispersion (27). The dependent variable was the hospital-specific daily count of pediatric asthma visits. In most analyses, the 3-day moving average pollutant concentration (the average of concentrations today [lag 0], yesterday [lag 1], and two days ago [lag 2]) was modeled linearly (14). To further describe associations we examined the 3-day moving average using quintiles, the 3-day moving average using a loess smoother (from a generalized additive model) (28, 29), and the 8-day moving average concentration (lags 0 through 7) constrained using a third degree polynomial (30). For all analyses we created separate warm season (May–October) and cold season (November–April) models. All models included a cubic polynomial for day-of-season, the moving average of dew point (lags 0-2), and the moving average of minimum temperature (lags 1 and 2); indicator variables for year, month, day of week, hospital, and lag 0 maximum temperature (for each

degree of Celsius); and interactions between month and year, month and lag 0 maximum temperature, and month and day of week. Additional terms for dew point and minimum temperature were included in distributed lag models to coincide with the pollutant averaging period. We investigated confounding by upper respiratory infections (the logarithm of the daily count of upper respiratory infections) and pollen concentrations (various lags of ambient ragweed, pine, oak, juniper, grass, and birch concentrations) by assessing whether their inclusion changed the pollutant regression coefficient estimates. We examined the sensitivity of our results to alternative model specifications and evaluated model misspecification by estimating associations with pollutant concentrations on the day after the emergency department visit. The sensitivity analyses include results from time-series models, which are based on models analogous to those we presented in earlier publications (14, 18, 20, 25, 31); “traditional” case-crossover models that include the three-way interactions between year, month, and day-of-week; traditional case-crossover models with bi-monthly (rather than monthly) time windows to further account for temporal trends; and our primary analytic approach without control for the daily count of emergency department visits for (non-asthma) respiratory infections. Generalized additive models were implemented using R 2.8.1; all other analyses were performed using SAS 9.2.

RESULTS

Mean counts of emergency department visits for asthma or wheeze among children age 5-17 years were 18.9/day during the warm season (May–October) and 22.8/day during the cold season (November–April). The seasonal difference in upper respiratory infections was more

pronounced, with mean counts of 21.5/day during the warm season and 34.6/day during the cold season. Descriptive statistics for ambient air pollutant concentrations and maximum temperature are shown in Table 1. All pollutants had some seasonal variability, although the mean concentrations of nitrogen dioxide, carbon monoxide, and PM_{2.5} elemental carbon (all markers of pollution from combustion engines) were similar between the warm and cold season.

Rate ratio estimates for associations between ambient air pollutant concentrations and pediatric asthma emergency department visits were similar regardless of whether pollen concentrations were included in the model as covariates; we examined various lags of pollen concentrations both continuously and using indicator variables for the top 5% and 10% of days. Therefore, covariates for pollen were not included in our final regression models. Results from our primary analysis, with pollutant concentrations characterized as three-day moving averages and rate ratios corresponding to interquartile range changes in pollutant concentrations, are presented in Table 2. Models for 11 pollutants were created for each season, and we observed 10 significant positive associations ($p < 0.05$) during the warm season and 1 during the cold season. Further examination of dose-response relationships included season-specific rate ratios for pollutant concentration quintiles, presented in Table 3, and loess dose-response curves for the 10 statistically significant warm season associations from Table 2, which are presented in Figure 1. Ozone was strongly associated with emergency department visits for pediatric asthma during the summer; the cold season ozone association was elevated but not significant. Within the cold season, we observed evidence for an effect during the more temperate months (November, March, and April) and no evidence for an effect during the coldest three months (December, January, and February) (see Table 2 footnote). As shown by both the quintile analysis and the

loess dose-response curves, evidence for a dose-response relationship with ozone was present at concentrations as low as 30 parts per billion. Several markers of pollution from combustion engines (carbon monoxide, nitrogen dioxide, $PM_{2.5}$ elemental carbon, and $PM_{2.5}$ water-soluble metals) were associated with emergency department visits for pediatric asthma during the warm season, as were sulfur dioxide, PM_{10} , $PM_{2.5}$, and $PM_{2.5}$ sulfate (a secondary pollutant derived largely from coal burning). $PM_{10-2.5}$ was associated with pediatric asthma emergency department visits during the cold season; the lower bound confidence interval for the $PM_{10-2.5}$ association was very close to 1.0, although the point estimates for $PM_{10-2.5}$ were elevated during both seasons.

Figure 2 displays rate ratio estimates for day-specific lags (lags 0-7) for associations between ambient air pollutant concentrations and pediatric asthma emergency department visits for the 10 statistically significant warm season associations from Table 2. The rate ratio and confidence interval summarizing the overall eight-day association are also presented for each pollutant. For 7 of the 10 associations, the lag-specific rate ratio of the highest magnitude was lag 0 (the pollutant concentration on the day of the emergency department visit). For the other 3 associations (PM_{10} , $PM_{2.5}$ elemental carbon, and $PM_{2.5}$ water-soluble metals), the lag-specific rate ratios were more uniformly distributed over the 8-day period. For all pollutants the confidence intervals are widest at lag 0, because the shape of the dose-response was constrained to follow a cubic polynomial (over lags 0-13), and the uncertainty of the estimates is greatest at the boundaries.

We created multipollutant models, including several two-pollutant models that paired ozone with another pollutant representing a particular source. A table of partial correlations between the various pollutants is available in the online data supplement. Warm season two-pollutant models are presented in Figure 3. The association between ozone and asthma emergency department visits observed in the single pollutant models persisted across the various two-pollutant models, and although the point estimates were attenuated, we observed evidence for an effect of primary traffic pollutants on asthma exacerbations independent of the effect of ozone. Carbon monoxide showed the strongest association of the three markers of pollution from traffic sources examined in two-pollutant models. Although significant associations were observed with warm season $PM_{2.5}$ sulfate and cold season $PM_{10-2.5}$ in single-pollutant models, these associations were not present in the two-pollutant models. We also explored two-pollutant models that allowed for interactions between ozone and each of the other pollutants; however, we did not observe evidence suggesting interaction in these models (results not shown).

We included the logarithm of the daily count of upper respiratory infections as an additional covariate in all of our models. As presented in Table 4, inclusion of this covariate tended to attenuate the estimated rate ratios, particularly during the warm season. Results from several alternative statistical models are also presented in Table 4. Broadly, these models supported the conclusions from our primary single-pollutant models. Asthma emergency department visits were associated with ambient ozone concentrations and with several traffic-related primary pollutants and $PM_{2.5}$ sulfate concentrations during the warm season and with $PM_{10-2.5}$ during the cold season. Table 4 also presents associations between lag -1 (tomorrow's) pollutant concentrations and pediatric asthma emergency department visits for the primary model

(controlling for the 3-day average concentration on lags 0, 1, and 2). We used these estimates, which were generally compatible with the null, as a diagnostic tool to assess model misspecification; a well-specified model should have a rate ratio estimate close to the null value for the lag -1 pollutant concentration.

DISCUSSION

In our large study we analyzed over 90,000 emergency department visits for pediatric asthma in relation to ambient air pollutant concentrations. We controlled tightly for meteorology and seasonal asthma trends, and we observed several positive, statistically significant associations between ambient air pollutant concentrations and the rate of pediatric asthma emergency department visits in Atlanta. Ozone was associated with emergency department visits for asthma during the warm season and during the temperate cold season months (November, March, and April). We also observed associations with several traffic-related primary pollutants during the warm season. These pollutants have been found to cause asthma exacerbations and airway inflammation in observational and experimental studies (32-36); as supported by both the quintile analysis and the smooth estimates of dose-response, we observed evidence that associations were present at relatively low ambient concentrations. Further, results from two-pollutant models support the conclusion that ambient concentrations of both ozone and traffic-related primary pollutants independently contribute to the burden of asthma exacerbations. Among the three markers of primary traffic pollution that we investigated in two-pollutant models, the rate ratio of the highest magnitude was for carbon monoxide. Since levels of carbon monoxide present in ambient air do not pose appreciable health risks, carbon monoxide concentrations are likely a surrogate for other pollutants emitted from combustion sources more

plausibly linked to asthma. Estimates from distributed lag models suggested there were both immediate and lagged effects for these pollutants, with the association of highest magnitude tending to occur on the day of the emergency department visit.

We also observed associations with lag 0-2 average concentrations of warm season sulfur dioxide, warm season $PM_{2.5}$ sulfate, warm season $PM_{2.5}$ organic carbon, and cold season coarse particles ($PM_{10-2.5}$). None of these warm season results were significant in two-pollutant models that also contained lag 0-2 ozone concentrations; however, many of the lag-specific point estimates (from the distributed lag models) were positive at lag 3 and longer, thereby raising the possibility that some pollutants might have shown effects had we created multipollutant models that spanned longer lag periods. We are particularly suspicious of the sulfur dioxide result, because local plume touchdowns strongly impact measured sulfur dioxide concentrations, and consequently it is challenging to develop a daily sulfur dioxide metric that could be considered representative of the urban airshed based on measurements from only 5 monitoring stations. Further, in previous epidemiological and experimental studies, ambient concentrations of sulfur dioxide and $PM_{2.5}$ sulfate have not been consistently associated with impaired pediatric respiratory function (6, 37, 38). Respiratory function decline and increased risks of asthma exacerbation associated with ambient $PM_{10-2.5}$ and $PM_{2.5}$ organic carbon concentrations have been reported in previous studies (39-42), although there have been relatively few investigations of these pollutants compared to the body of work on PM_{10} and $PM_{2.5}$. Both $PM_{10-2.5}$ and $PM_{2.5}$ organic carbon are comprised of several different compounds, with $PM_{10-2.5}$ concentrations in Atlanta being largely comprised of metal oxides and crustal material (43), and $PM_{2.5}$ organic carbon consisting of mixture of compounds of both primary and secondary origin (26, 44, 45).

Our tendency to find stronger associations during the warm season is consistent with previous findings (10-15), and although we are unclear about the underlying mechanism for these apparent seasonal differences, it may simply be that during the warm season a greater proportion of asthma exacerbations are caused by air pollution. Rates of emergency department visits for pediatric asthma increase by 60% during the cold season; this increase is largely attributable to exacerbations triggered by viral infections. If the additive effect of air pollution is similar year-round, then the attributable fraction (and, correspondingly, the rate ratio) will appear higher during the warm season, because there are fewer competing causes of asthma exacerbations during the warm season. Alternatively, it may be that children actually respond more severely to air pollutants during warmer temperatures, perhaps because of some unidentified synergism between the pollutant and a meteorological or physical factor. Additional contributions to the observed seasonal differences may include non-linear dose-response functions (e.g., air pollutant concentrations typical during warmer months may be on a steeper part of the dose-response curve) and behavior differences that impact personal pollutant exposures. For example, during the summer children are more likely to play outside, which may lead to a higher correlation between measured ambient concentrations and personal exposures, and consequently result in higher estimated effects of ambient pollutants.

We relied upon codes from hospital administrative databases to identify emergency department visits for pediatric asthma. Our definition was relatively broad and included codes for both asthma and wheeze among children age 5-17 years. We excluded children younger than 5 years from our analysis because young children frequently experience transient wheeze, and asthma

diagnoses may be suspect (46); however, even among children age 5 years and older, we observed significant hospital-to-hospital variability in the proportion of emergency visits coded as “asthma” as opposed to “wheeze.” Further, we observed variability in the coding of primary versus secondary diagnoses; for asthma, this typically occurred when a patient presented with both asthma symptoms and a respiratory infection. We conducted subanalyses limited to emergency department visits where asthma or wheeze was reported as the primary diagnosis and observed results similar to those from our primary analytic approach. Because co-morbidities were not coded completely and consistently across hospitals, we deemed these data to be of inadequate quality to support analyses where individual visits were stratified according to the presence or absence of a respiratory infection as a co-morbidity, even though at the aggregate-level the daily count of emergency department visits for respiratory infections was likely an adequate surrogate for the actual burden of respiratory infections in Atlanta. We controlled for the daily count of upper respiratory visits in our statistical models and found it to be an extremely strong predictor of the rate of emergency department visits for pediatric asthma; further, we observed evidence of confounding by respiratory infections, as control for this covariate tended to attenuate the rate ratio estimates, particularly during the warm season.

Although we chose our primary statistical model carefully, all statistical models are misspecified to some degree. Therefore, we reported results from sensitivity analyses using alternative model specifications. Our primary model is based on the case-crossover design, with implementation via Poisson time-series models that account for overdispersion, given that under certain formulations these approaches are nearly identical (the conditional logistic regression estimating equation that is used in time-stratified case-crossover analyses is equivalent to a Poisson time-

series model with an indicator variable for each stratum when disease is rare and there is a shared exposure such as air pollution) (27, 47). Traditionally, investigators have implemented the case-crossover design by matching either on day-of-week (48, 49) or temperature (50) within a given month; matching on both day-of-week and temperature is typically not feasible, because data become sparse with too many matching factors. To implement a case-crossover approach in a time-series framework requires terms for the main effects of year, month, and the matching-factor (e.g., a term for each day-of-week); terms for the two-way interactions between year and month, year and day-of-week, and month and day-of-week; and terms for the three-way interactions between year, month, and day-of-week. In developing our primary analytic approach, we explored case-crossover models with matching on year, month, and either day-of-week or lag 0 maximum temperature; however, regardless of the approach, we observed evidence of confounding by within-month trends, e.g., the increasing trend in asthma exacerbations during late August and September due to the “back-to-school” effect (51). To smoothly control for these within-month trends we included a cubic polynomial for day-of-season in the regression models. Given this cubic polynomial, inclusion of the three-way interaction terms no longer meaningfully changed the point estimates for the air pollutant effect. Therefore, we abandoned the three-way interactions (and thereby removed hundreds of parameters from the model), and instead implemented a case-crossover analysis by matching only on month and year. In addition to matching on these factors, we controlled tightly for both day-of-week and lag 0 maximum temperature; our base model included indicator variables for year, month, day-of-week, and lag 0 maximum temperature (for each degree of Celsius), as well as selected two-way interactions (between month and year; month and day-of-week; and month

and lag 0 maximum temperature) that we found to be highly predictive of the pediatric asthma emergency department visit rates.

Even though we controlled tightly for meteorology and temporal trends and employed a case-only analytic approach, confounding by an unmeasured or inadequately modeled risk factor that varied in a systematic way with short-term fluctuations in ambient air pollutant concentrations could have biased our results. Although we cannot dismiss the possibility of confounding by an unmeasured factor, we conducted extensive analyses to understand the relationships with meteorology, temporal trends, and ambient pollen concentrations in our data. Further, we investigated associations with the lag -1 pollutant concentration (the concentration on the day after the emergency department visit), while simultaneously controlling for the average concentration on lags 0-2, as an approach to evaluate model misspecification (14, 52), since we know that tomorrow's pollutant concentrations are not causally related to today's count of emergency department visits, and any association not due to chance must be biased.

Measurement error is inherent in all large epidemiologic studies of urban air pollution health effects. Although studies of personal exposures to air pollutants help to advance understanding of biological responses, from a regulatory standpoint the ambient concentrations are of greatest relevance. One prominent component of error in our study is, therefore, how well the population-weighted spatial average of measurements from urban monitoring stations approximates the ambient concentration across the entire metropolitan Atlanta area. The extent of this measurement error likely varies by pollutant, with primary pollutants (such as those from traffic sources) tending to have more measurement error than secondary pollutants (such as ozone and

PM_{2.5} sulfate) (18, 53). Indeed, in previous work, we observed associations between emergency department visits for cardiovascular disease and spatially heterogeneous pollutants (carbon monoxide and nitrogen dioxide) using measurements from several different air pollution monitors located within 20 miles of the Atlanta population center; however, we did not observe associations when measurements were used from a rural monitor located 38 miles away.

Conversely, we observed associations for the spatially homogeneous pollutants (ozone and PM_{2.5}) regardless of whether the measurements were from the rural or urban monitors (18). Also contributing to the measurement error issue is the number of air pollutant monitoring stations, which ranged from only one central monitor (for PM_{10-2.5} and PM_{2.5} water-soluble metals) to 11 monitors (for PM_{2.5}). Interpretation of two-pollutant models is complicated by these measurement error issues; the pollutant with the stronger estimated effect (e.g., ozone in our analyses) may not be the more harmful pollutant but may instead be the pollutant that has less measurement error (20). This measurement error also impacts the statistical power to detect effects. In our study we did not find evidence of synergism between ozone and any of the other air pollutants, perhaps due to issues involving measurement error and statistical power.

The findings from our large, population-based time-series study in Atlanta complement previous findings from multicity studies (13, 54). Whereas multicity designs offer a statistically powerful approach for investigating the health effects of ambient air pollutants, large single-city studies provide the opportunity for investigators to better understand and account for the nuances of local data. Further, the SOPHIA study, which has amassed data on over 10,000,000 emergency department visits in metropolitan Atlanta since 1993, affords ample statistical power to detect subtle health effects of ambient air pollutants, including the health effects of PM_{2.5} components.

In our study we observed evidence that ambient concentrations of ozone and primary pollutants from traffic sources independently contributed to the burden of emergency department visits for pediatric asthma. Further, these associations were present at relatively low ambient concentrations, reinforcing the need for continued evaluation of the EPA's National Ambient Air Quality Standards to ensure that the standards are sufficient to protect susceptible individuals.

ACKNOWLEDGMENTS

Jennifer Peel for insights into the statistical model building process and for her previous work on respiratory outcomes based on the SOPHIA database. Ted Russell for manuscript comments and assistance with characterization of the air quality data. Sarah Tinker for management and maintenance of the SOPHIA emergency department visit database.

REFERENCES

1. U.S. EPA. Air quality criteria for particulate matter (final). Washington, DC: U.S. Environmental Protection Agency 600/P-99/002aF-bF; 2004.
2. U.S. EPA. Air quality criteria for ozone and related photochemical oxidants (final). Washington, DC: United States Environmental Protection Agency EPA/600/R-05/004aF-cF; 2006.
3. World Health Organization. Health aspects of air pollution: Results from the WHO project "systematic review of air pollution in Europe". Copenhagen, Denmark: WHO Regional Office for Europe; 2004.
4. Bateson TF, Schwartz J. Children's response to air pollutants. *J Toxicol Environ Health A* 2008;71:238-243.
5. Selgrade MK, Plopper CG, Gilmour MI, Conolly RB, Foos BS. Assessing the health effects and risks associated with children's inhalation exposures--asthma and allergy. *J Toxicol Environ Health A* 2008;71:196-207.
6. Trasande L, Thurston GD. The role of air pollution in asthma and other pediatric morbidities. *J Allergy Clin Immunol* 2005;115:689-699.
7. National Research Council. Research priorities for airborne particulate matter: iv continuing research progress. Washington, D.C.: The National Academies Press; 2004.
8. Mauderly JL, Samet JM. Is there evidence for synergy among air pollutants in causing health effects? *Environ Health Perspect* 2009;117:1-6.

9. Eggleston PA. Complex interactions of pollutant and allergen exposures and their impact on people with asthma. *Pediatrics* 2009;123 Suppl 3:S160-167.
10. Villeneuve PJ, Chen L, Rowe BH, Coates F. Outdoor air pollution and emergency department visits for asthma among children and adults: A case-crossover study in Northern Alberta, Canada. *Environ Health* 2007;6:40.
11. Jalaludin B, Khalaj B, Sheppard V, Morgan G. Air pollution and ed visits for asthma in Australian children: A case-crossover analysis. *Int Arch Occup Environ Health* 2008;81:967-974.
12. Szyszkowicz M. Ambient air pollution and daily emergency department visits for asthma in Edmonton, Canada. *Int J Occup Med Environ Health* 2008;21:25-30.
13. Barnett AG, Williams GM, Schwartz J, Neller AH, Best TL, Petroeschovsky AL, Simpson RW. Air pollution and child respiratory health: A case-crossover study in Australia and New Zealand. *Am J Respir Crit Care Med* 2005;171:1272-1278.
14. Peel JL, Tolbert PE, Klein M, Metzger KB, Flanders WD, Todd K, Mulholland JA, Ryan PB, Frumkin H. Ambient air pollution and respiratory emergency department visits. *Epidemiology* 2005;16:164-174.
15. Mohr LB, Luo S, Mathias E, Tobing R, Homan S, Sterling D. Influence of season and temperature on the relationship of elemental carbon air pollution to pediatric asthma emergency room visits. *J Asthma* 2008;45:936-943.
16. Pendergraft TB, Stanford RH, Beasley R, Stempel DA, McLaughlin T. Seasonal variation in asthma-related hospital and intensive care unit admissions. *J Asthma* 2005;42:265-271.
17. Tolbert PE, Klein M, Metzger KB, Peel J, Flanders WD, Todd K, Mulholland JA, Ryan PB, Frumkin H. Interim results of the Study of Particulates and Health in Atlanta (SOPHIA). *J Expo Anal Environ Epidemiol* 2000;10:446-460.
18. Sarnat SE, Klein M, Sarnat JA, Flanders WD, Waller LA, Mulholland JA, Russell AG, Tolbert PE. An examination of exposure measurement error from air pollutant spatial variability in time-series studies. *J Expo Sci Environ Epidemiol* 2009.
19. Metzger KB, Klein M, Flanders WD, Peel JL, Mulholland JA, Langberg JJ, Tolbert PE. Ambient air pollution and cardiac arrhythmias in patients with implantable defibrillators. *Epidemiology* 2007;18:585-592.
20. Tolbert PE, Klein M, Peel JL, Sarnat SE, Sarnat JA. Multipollutant modeling issues in a study of ambient air quality and emergency department visits in Atlanta. *J Expo Sci Environ Epidemiol* 2007;17 Suppl 2:S29-35.
21. Metzger KB, Tolbert PE, Klein M, Peel JL, Flanders WD, Todd K, Mulholland JA, Ryan PB, Frumkin H. Ambient air pollution and cardiovascular emergency department visits. *Epidemiology* 2004;15:46-56.
22. Strickland MJ, Darrow LA, Klein M, Sarnat JA, Flanders WD, Sarnat SE, Waller LA, Tolbert PE. Acute associations between ambient air pollution and pediatric asthma emergency department visits in Atlanta, 1993-2004. *Epidemiology* 2009;20:S163.
23. Van Loy M, Bahadori T, Wyzga R, Hartsell B, Edgerton E. The Aerosol Research and Inhalation Epidemiology Study (ARIES): PM_{2.5} mass and aerosol component concentrations and sampler intercomparisons. *J Air Waste Manag Assoc* 2000;50:1446-1458.
24. Marmur A, Park SK, Mulholland JA, Tolbert PE, Russell AG. Source apportionment of PM_{2.5} in the Southeastern United States using receptor and emissions-based models: Conceptual differences and implications for time-series studies. *Atmos Environ* 2006;40:2533-2551.

25. Sarnat JA, Marmur A, Klein M, Kim E, Russell AG, Sarnat SE, Mulholland JA, Hopke PK, Tolbert PE. Fine particle sources and cardiorespiratory morbidity: An application of chemical mass balance and factor analytical source-apportionment methods. *Environ Health Perspect* 2008;116:459-466.
26. Zheng M, Cass GR, Ke L, Wang F, Schauer JJ, Edgerton ES, Russell AG. Source apportionment of daily fine particulate matter at Jefferson Street, Atlanta, GA, during summer and winter. *J Air Waste Manag Assoc* 2007;57:228-242.
27. Lu Y, Zeger SL. On the equivalence of case-crossover and time series methods in environmental epidemiology. *Biostatistics* 2007;8:337-344.
28. Dominici F, McDermott A, Hastie TJ. Improved semiparametric time series models of air pollution and mortality. *J Am Stat Assoc* 2004;99:938-948.
29. Flanders WD, Klein M, Tolbert PE. A new variance estimator for parameters of semiparametric generalized additive models. *J Agric Biol Envir Stat* 2005;10:246-257.
30. Schwartz J. The distributed lag between air pollution and daily deaths. *Epidemiology* 2000;11:320-326.
31. Darrow LA, Klein M, Sarnat JA, Mulholland JA, Strickland MJ, Sarnat SE, Russell AG, Tolbert PE. The use of alternative pollutant metrics in time-series studies of ambient air pollution and respiratory emergency department visits. *J Expo Sci Environ Epidemiol* 2009.
32. Moore K, Neugebauer R, Lurmann F, Hall J, Brajer V, Alcorn S, Tager I. Ambient ozone concentrations cause increased hospitalizations for asthma in children: An 18-year study in Southern California. *Environ Health Perspect* 2008;116:1063-1070.
33. Jansen KL, Larson TV, Koenig JQ, Mar TF, Fields C, Stewart J, Lippmann M. Associations between health effects and particulate matter and black carbon in subjects with respiratory disease. *Environ Health Perspect* 2005;113:1741-1746.
34. Gent JF, Triche EW, Holford TR, Belanger K, Bracken MB, Beckett WS, Leaderer BP. Association of low-level ozone and fine particles with respiratory symptoms in children with asthma. *JAMA* 2003;290:1859-1867.
35. Riedl M, Diaz-Sanchez D. Biology of diesel exhaust effects on respiratory function. *J Allergy Clin Immunol* 2005;115:221-228; quiz 229.
36. Hollingsworth JW, Kleeberger SR, Foster WM. Ozone and pulmonary innate immunity. *Proc Am Thorac Soc* 2007;4:240-246.
37. Reiss R, Anderson EL, Cross CE, Hidy G, Hoel D, McClellan R, Moolgavkar S. Evidence of health impacts of sulfate-and nitrate-containing particles in ambient air. *Inhal Toxicol* 2007;19:419-449.
38. Schlesinger RB. The health impact of common inorganic components of fine particulate matter (PM_{2.5}) in ambient air: A critical review. *Inhal Toxicol* 2007;19:811-832.
39. Brunekreef B, Forsberg B. Epidemiological evidence of effects of coarse airborne particles on health. *Eur Respir J* 2005;26:309-318.
40. Alexis NE, Lay JC, Zeman K, Bennett WE, Peden DB, Soukup JM, Devlin RB, Becker S. Biological material on inhaled coarse fraction particulate matter activates airway phagocytes in vivo in healthy volunteers. *J Allergy Clin Immunol* 2006;117:1396-1403.
41. Ostro B, Roth L, Malig B, Marty M. The effects of fine particle components on respiratory hospital admissions in children. *Environ Health Perspect* 2009;117:475-480.
42. Delfino RJ, Staimer N, Tjoa T, Gillen D, Kleinman MT, Sioutas C, Cooper D. Personal and ambient air pollution exposures and lung function decrements in children with asthma. *Environ Health Perspect* 2008;116:550-558.

43. Edgerton ES, Hartsell BE, Saylor RD, Jansen JJ, Hansen DA, Hidy GM. The Southeastern Aerosol Research and Characterization study: Part ii. Filter-based measurements of fine and coarse particulate matter mass and composition. *J Air Waste Manag Assoc* 2005;55:1527-1542.
44. Ding X, Zheng M, Yu L, Zhang X, Weber RJ, Yan B, Russell AG, Edgerton ES, Wang X. Spatial and seasonal trends in biogenic secondary organic aerosol tracers and water-soluble organic carbon in the Southeastern United States. *Environ Sci Technol* 2008;42:5171-5176.
45. Ding X, Zheng M, Edgerton ES, Jansen JJ, Wang X. Contemporary or fossil origin: Split of estimated secondary organic carbon in the southeastern united states. *Environ Sci Technol* 2008;42:9122-9128.
46. National Heart Lung and Blood Institute. Expert panel report 3: Guidelines for the diagnosis and management of asthma. NIH publication no. 07-4051; 2007.
47. Lu Y, Symons JM, Geyh AS, Zeger SL. An approach to checking case-crossover analyses based on equivalence with time-series methods. *Epidemiology* 2008;19:169-175.
48. Janes H, Sheppard L, Lumley T. Case-crossover analyses of air pollution exposure data: Referent selection strategies and their implications for bias. *Epidemiology* 2005;16:717-726.
49. Levy D, Lumley T, Sheppard L, Kaufman J, Checkoway H. Referent selection in case-crossover analyses of acute health effects of air pollution. *Epidemiology* 2001;12:186-192.
50. Schwartz J. The effects of particulate air pollution on daily deaths: A multi-city case crossover analysis. *Occup Environ Med* 2004;61:956-961.
51. Johnston NW, Johnston SL, Duncan JM, Greene JM, Keadze T, Keith PK, Roy M, Wasserman S, Sears MR. The September epidemic of asthma exacerbations in children: A search for etiology. *J Allergy Clin Immunol* 2005;115:132-138.
52. Dominici F, Peng RD, Bell ML, Pham L, McDermott A, Zeger SL, Samet JM. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* 2006;295:1127-1134.
53. Wade KS, Mulholland JA, Marmur A, Russell AG, Hartsell B, Edgerton E, Klein M, Waller L, Peel JL, Tolbert PE. Effects of instrument precision and spatial variability on the assessment of the temporal variation of ambient air pollution in Atlanta, Georgia. *J Air Waste Manag Assoc* 2006;56:876-888.
54. Atkinson RW, Anderson HR, Sunyer J, Ayres J, Baccini M, Vonk JM, Boumghar A, Forastiere F, Forsberg B, Touloumi G, Schwartz J, Katsouyanni K. Acute effects of particulate air pollution on respiratory admissions: Results from APHEA 2 project. *Am J Respir Crit Care Med* 2001;164:1860-1866.

Figure 1. Loess dose-response estimates (solid line) and twice-standard error estimates (dashed lines) from generalized additive models for associations between 3-day moving average air pollutant concentrations and emergency department visits for pediatric asthma. The reference (denominator) for the rate ratio is the estimated rate at the 5th percentile of the pollutant concentration. Estimates are presented for the 5th percentile through the 95th percentile of pollutant concentrations due to instability in the dose-response estimates at the distribution tails.

Figure 2. Constrained cubic polynomial distributed lag models. The rate ratio and 95% confidence interval displayed for each pollutant correspond to an interquartile range increase in the cumulative ambient pollutant concentration during the 8-day period of interest (lags 0 through 7). Point estimates and 95% confidence intervals are also presented graphically for the lag-specific rate-ratios. To enhance the stability of the distributed lag estimates, the cubic polynomial was fit to lags 0-13; however, the rate ratios and 95% confidence intervals presented in the Figure correspond to the effects of lags 0-7 only.

Figure 3. Warm season rate ratios and 95% confidence intervals for interquartile range increases in 3-day moving average ambient air pollutant concentrations for single-pollutant and two-pollutant models. Abbreviations: CO (carbon monoxide), O3 (ozone), NO2 (nitrogen dioxide), EC (PM_{2.5} elemental carbon), SO4 (PM_{2.5} sulfate).

*Time period limited to 1 August 1998 – 31 December 2004

Table 1. Descriptions (mean, standard deviation, interquartile range (IQR), percent missing and number of monitors) of population weighted ambient air pollutant concentrations and maximum temperature.*

	Overall (Jan-Dec) Mean \pm SD	Warm Season (May-Oct) Mean \pm SD	Cold Season (Nov-Apr) Mean \pm SD	IQR	Percent Missing	Number of monitors
8-h ozone (ppb)†	45.4 \pm 20.0	55.2 \pm 19.2	34.5 \pm 14.6	29.2	4	5
1-h nitrogen dioxide (ppb)†	23.3 \pm 9.7	22.0 \pm 9.4	24.5 \pm 9.8	12.9	<1	6
1-h carbon monoxide (ppm)†	0.9 \pm 0.5	0.9 \pm 0.4	1.0 \pm 0.6	0.6	5	4
1-h sulfur dioxide (ppb)†	10.8 \pm 9.5	9.6 \pm 8.7	12.0 \pm 10.2	11.5	0	5
24-h PM ₁₀ ($\mu\text{g}/\text{m}^3$)‡	23.8 \pm 11.5	27.6 \pm 11.6	20.0 \pm 10.0	14.6	1	9
24-h PM _{10-2.5} ($\mu\text{g}/\text{m}^3$)§	9.0 \pm 5.0	9.7 \pm 4.7	8.3 \pm 5.3	5.9	7	1
24-h PM _{2.5} ($\mu\text{g}/\text{m}^3$)§	16.4 \pm 7.4	18.4 \pm 7.6	14.3 \pm 6.5	9.2	<1	11
24-h PM _{2.5} sulfate ($\mu\text{g}/\text{m}^3$)§	4.6 \pm 3.1	5.9 \pm 3.5	3.2 \pm 1.8	3.5	1	6
24-h PM _{2.5} elemental carbon ($\mu\text{g}/\text{m}^3$)§	0.9 \pm 0.6	0.8 \pm 0.6	0.9 \pm 0.6	0.7	<1	6
24-h PM _{2.5} organic carbon ($\mu\text{g}/\text{m}^3$)§	4.9 \pm 3.4	4.8 \pm 2.7	5.0 \pm 4.0	3.4	<1	6
24-h PM _{2.5} water-soluble metals ($\mu\text{g}/\text{m}^3$)§	0.030 \pm 0.023	0.039 \pm 0.025	0.020 \pm 0.016	0.025	9	1
Maximum temperature ($^{\circ}\text{C}$)	22.4 \pm 8.4	28.5 \pm 4.6	16.1 \pm 6.6	13.0	0	1

* Because the means presented are population-weighted spatial averages, they may differ from the means of concentrations measured at urban central site monitors. For comparison, the mean central site concentrations were: ozone – 47.3ppb; nitrogen dioxide – 43.2 ppb; carbon monoxide – 1.6 ppm; sulfur dioxide – 15.0 ppb; PM₁₀ – 26.6 $\mu\text{g}/\text{m}^3$; PM_{10-2.5} – 9.0 $\mu\text{g}/\text{m}^3$; PM_{2.5} – 17.1 $\mu\text{g}/\text{m}^3$; PM_{2.5} sulfate – 4.9 $\mu\text{g}/\text{m}^3$; PM_{2.5} elemental carbon – 1.6 $\mu\text{g}/\text{m}^3$; PM_{2.5} organic carbon – 4.4 $\mu\text{g}/\text{m}^3$; PM_{2.5} water soluble metals – 0.030 $\mu\text{g}/\text{m}^3$.

† Measurements available during 1 January 1993 – 31 December 2004 (n=4383). Ozone was not measured December 1994–February 1995 and December 1995 – February 1995.

‡ Measurements available during 1 January 1996 – 31 December 2004 (n=3288).

§ Measurements available during 1 August 1998 – 31 December 2004 (n=2345).

Table 2. Rate ratios and 95% confidence intervals from Poisson general linear models for interquartile range increases in 3-day moving average population-weighted ambient air pollutant concentrations.*

	Overall (Jan-Dec)	Warm Season (May-Oct)	Cold Season (Nov-Apr)
Ozone†‡	1.062 (1.031, 1.093)	1.082 (1.043, 1.123)	1.044 (0.992, 1.098)
Nitrogen dioxide†	1.036 (1.018, 1.055)	1.066 (1.038, 1.095)	1.016 (0.992, 1.040)
Carbon monoxide†	1.023 (1.006, 1.041)	1.068 (1.034, 1.102)	1.005 (0.985, 1.025)
Sulfur dioxide†	1.012 (0.994, 1.030)	1.030 (1.002, 1.058)	1.001 (0.978, 1.025)
PM ₁₀ §	1.020 (1.003, 1.038)	1.026 (1.001, 1.051)	1.018 (0.994, 1.043)
PM _{10-2.5}	1.034 (1.011, 1.057)	1.025 (0.991, 1.059)	1.041 (1.010, 1.073)
PM _{2.5}	1.020 (1.002, 1.039)	1.043 (1.016, 1.070)	1.005 (0.978, 1.031)
PM _{2.5} sulfate	1.014 (0.995, 1.033)	1.027 (1.004, 1.049)	0.991 (0.953, 1.029)
PM _{2.5} elemental carbon	1.015 (0.997, 1.033)	1.041 (1.010, 1.072)	1.003 (0.981, 1.026)
PM _{2.5} organic carbon	1.008 (0.994, 1.021)	1.034 (1.007, 1.062)	1.000 (0.985, 1.016)
PM _{2.5} water-soluble metals	1.021 (1.000, 1.042)	1.029 (1.003, 1.055)	1.005 (0.968, 1.043)

*Interquartile ranges presented in Table 1

†Measurements available January, 1993 – December, 2004

‡ When the cold season was divided into the more temperate months (March, April, November) vs. the coldest months (December, January, February) we observed rate ratios of 1.107 (1.035, 1.184) for November/March/April and 0.968 (0.895, 1.047) for December/January/February.

§Measurements available January, 1996 – December, 2004

||Measurements available August, 1998 – December, 2004

Table 3. Rate ratios (RR) and 95% confidence intervals for quintiles of 3-day moving average population-weighted ambient air pollutant concentrations.

	Warm Season RR (95% CI)*	Cold Season RR (95% CI)*
Ozone (ppb)†		
Q2 (26.3 to < 38.7)	1.002 (0.953, 1.054)	1.039 (0.998, 1.081)
Q3 (38.7 to < 51.5)	1.016 (0.961, 1.074)	1.097 (1.037, 1.161)
Q4 (51.5 to < 67.0)	1.061 (0.999, 1.127)	1.151 (1.065, 1.243)
Q5 (67.0 to ≤ 147.5)	1.111 (1.038, 1.189)	1.150 (1.013, 1.306)
Nitrogen dioxide (ppb)†		
Q2 (28.0 to < 37.1)	1.033 (0.999, 1.069)	0.996 (0.964, 1.030)
Q3 (37.1 to < 46.0)	1.040 (1.000, 1.081)	0.984 (0.950, 1.020)
Q4 (46.0 to < 57.1)	1.087 (1.044, 1.131)	1.024 (0.985, 1.064)
Q5 (57.1 to ≤ 181.0)	1.087 (1.036, 1.140)	1.014 (0.973, 1.056)
Carbon monoxide (ppm)†		
Q2 (0.70 to < 1.01)	1.019 (0.986, 1.054)	1.010 (0.977, 1.045)
Q3 (1.01 to < 1.05)	1.046 (1.008, 1.086)	1.040 (1.005, 1.076)
Q4 (1.05 to < 2.30)	1.097 (1.049, 1.147)	1.005 (0.969, 1.042)
Q5 (2.30 to ≤ 7.70)	1.112 (1.054, 1.174)	1.021 (0.981, 1.064)
Sulfur dioxide (ppb)†		
Q2 (3.1 to < 7.0)	1.021 (0.988, 1.055)	0.968 (0.935, 1.002)
Q3 (7.0 to < 13.0)	1.041 (1.007, 1.077)	0.998 (0.964, 1.034)
Q4 (13.0 to < 24.2)	1.048 (1.010, 1.087)	0.982 (0.947, 1.017)
Q5 (24.2 to ≤ 149.0)	1.008 (0.967, 1.051)	0.987 (0.949, 1.026)
PM ₁₀ (µg/m ³)‡		
Q2 (16.0 to < 21.9)	1.014 (0.968, 1.061)	1.008 (0.978, 1.038)
Q3 (21.9 to < 28.0)	1.029 (0.981, 1.080)	0.996 (0.963, 1.030)
Q4 (28.0 to < 36.0)	1.027 (0.979, 1.078)	1.017 (0.977, 1.059)
Q5 (36.0 to ≤ 98.4)	1.059 (1.006, 1.116)	1.047 (0.991, 1.106)
PM _{10-2.5} (µg/m ³)§		
Q2 (5.0 to < 7.1)	0.975 (0.924, 1.028)	0.972 (0.930, 1.012)
Q3 (7.1 to < 9.3)	0.986 (0.934, 1.040)	1.006 (0.960, 1.054)
Q4 (9.3 to < 12.3)	0.964 (0.909, 1.022)	1.045 (0.996, 1.097)
Q5 (12.3 to ≤ 50.4)	1.005 (0.942, 1.072)	1.075 (1.015, 1.139)
PM _{2.5} (µg/m ³)§		
Q2 (10.0 to < 13.7)	0.993 (0.943, 1.047)	0.985 (0.952, 1.019)
Q3 (13.7 to < 17.6)	1.008 (0.956, 1.062)	0.979 (0.943, 1.017)
Q4 (17.6 to < 23.8)	1.018 (0.966, 1.073)	1.006 (0.960, 1.049)
Q5 (23.8 to ≤ 65.8)	1.052 (0.995, 1.112)	1.050 (0.997, 1.106)
PM _{2.5} sulfate (µg/m ³)§		
Q2 (2.2 to < 3.3)	1.032 (0.951, 1.120)	0.987 (0.950, 1.025)
Q3 (3.3 to < 4.6)	1.048 (0.968, 1.135)	1.045 (0.998, 1.094)
Q4 (4.6 to < 7.2)	1.061 (0.978, 1.150)	1.008 (0.944, 1.077)
Q5 (7.2 to ≤ 21.9)	1.082 (0.995, 1.177)	1.027 (0.931, 1.133)
PM _{2.5} elemental carbon (µg/m ³)§		
Q2 (0.78 to < 1.13)	0.981 (0.936, 1.028)	1.034 (0.994, 1.076)
Q3 (1.13 to < 1.55)	1.023 (0.977, 1.071)	1.027 (0.984, 1.072)
Q4 (1.55 to < 2.25)	1.050 (1.004, 1.107)	1.014 (0.971, 1.058)
Q5 (2.25 to ≤ 11.89)	1.056 (1.000, 1.115)	1.039 (0.990, 1.091)
PM _{2.5} organic carbon (µg/m ³)§		
Q2 (2.54 to < 3.41)	1.024 (0.976, 1.073)	0.983 (0.942, 1.025)
Q3 (3.41 to < 4.32)	1.070 (1.020, 1.122)	1.028 (0.983, 1.075)
Q4 (4.32 to < 5.82)	1.078 (1.024, 1.135)	0.993 (0.950, 1.038)
Q5 (5.82 to ≤ 25.93)	1.065 (1.006, 1.128)	1.017 (0.972, 1.063)
PM _{2.5} water-soluble metals (µg/m ³)§		
Q2 (0.0123 to < 0.0195)	1.026 (0.946, 1.112)	0.992 (0.953, 1.032)
Q3 (0.0195 to < 0.0276)	1.081 (0.999, 1.170)	1.004 (0.958, 1.053)
Q4 (0.0276 to < 0.0436)	1.113 (1.026, 1.206)	1.036 (0.978, 1.096)
Q5 (0.0436 to ≤ 0.202)	1.119 (1.028, 1.218)	1.009 (0.934, 1.090)

*Relative to the first quintile (concentrations less than the lower bound of the second quintile)

† Measurements available January, 1993 – December, 2004
‡ Measurements available August, 1998 – December, 2004
§ Measurements available August, 1998 – December, 2004

Table 4. Sensitivity analyses: Rate ratios (RR) and 95% confidence intervals for interquartile range increases in 3-day moving average population-weighted ambient air pollutant concentrations.

	Base model*	No URI control†	Alternative model specifications			Lag -1 pollution**
	RR (95% CI)	RR (95% CI)	Time-series‡ RR (95% CI)	Case-crossover§ RR (95% CI)	Bi-monthly windows RR (95% CI)	RR (95% CI)
Warm Season (May–October)						
Ozone	1.082 (1.043, 1.123)	1.106 (1.066, 1.147)	1.071 (1.025, 1.119)	1.092 (1.049, 1.137)	1.058 (1.014, 1.104)	1.015 (0.991, 1.039)
Nitrogen dioxide	1.066 (1.038, 1.095)	1.073 (1.045, 1.102)	1.058 (1.028, 1.088)	1.096 (1.065, 1.127)	1.065 (1.034, 1.098)	1.000 (0.981, 1.019)
Carbon monoxide	1.068 (1.034, 1.102)	1.073 (1.039, 1.107)	1.062 (1.030, 1.096)	1.090 (1.054, 1.127)	1.053 (1.016, 1.092)	1.009 (0.988, 1.031)
Sulfur dioxide	1.030 (1.002, 1.058)	1.033 (1.006, 1.062)	1.021 (0.992, 1.051)	1.009 (0.981, 1.039)	1.014 (0.984, 1.046)	1.007 (0.990, 1.025)
PM ₁₀	1.026 (1.001, 1.051)	1.043 (1.018, 1.068)	1.031 (1.005, 1.058)	1.057 (1.030, 1.085)	1.022 (0.994, 1.050)	1.024 (1.005, 1.043)
PM _{10-2.5}	1.025 (0.991, 1.059)	1.032 (0.998, 1.067)	1.028 (0.992, 1.065)	1.062 (1.026, 1.099)	0.984 (0.945, 1.024)	1.004 (0.983, 1.025)
PM _{2.5}	1.043 (1.016, 1.070)	1.060 (1.033, 1.088)	1.041 (1.012, 1.071)	1.043 (1.015, 1.072)	1.015 (0.984, 1.046)	1.024 (1.004, 1.045)
PM _{2.5} sulfate	1.027 (1.004, 1.049)	1.040 (1.017, 1.063)	1.025 (1.002, 1.048)	1.024 (1.000, 1.047)	1.003 (0.977, 1.029)	1.017 (1.001, 1.034)
PM _{2.5} elemental carbon	1.041 (1.010, 1.072)	1.051 (1.020, 1.083)	1.032 (1.000, 1.065)	1.057 (1.024, 1.090)	1.036 (0.984, 1.091)	1.000 (0.979, 1.020)
PM _{2.5} organic carbon	1.034 (1.007, 1.062)	1.047 (1.020, 1.075)	1.036 (1.005, 1.067)	1.050 (1.021, 1.079)	1.019 (0.987, 1.052)	1.000 (0.982, 1.018)
PM _{2.5} water-soluble metals	1.029 (1.003, 1.055)	1.039 (1.013, 1.065)	1.021 (0.996, 1.047)	1.030 (1.004, 1.057)	1.015 (0.986, 1.045)	0.992 (0.976, 1.009)
Cold Season (November–April)						
Ozone	1.044 (0.992, 1.098)	1.062 (1.010, 1.118)	1.013 (0.953, 1.077)	1.053 (0.996, 1.113)	1.019 (0.961, 1.080)	1.004 (0.973, 1.036)
Nitrogen dioxide	1.016 (0.992, 1.040)	1.023 (1.000, 1.048)	1.007 (0.982, 1.033)	1.020 (0.994, 1.047)	1.022 (0.995, 1.050)	0.999 (0.983, 1.014)
Carbon monoxide	1.005 (0.985, 1.025)	1.017 (0.997, 1.037)	1.002 (0.982, 1.023)	1.003 (0.981, 1.025)	1.005 (0.982, 1.028)	0.999 (0.986, 1.012)
Sulfur dioxide	1.001 (0.978, 1.025)	1.000 (0.977, 1.024)	1.009 (0.984, 1.034)	0.999 (0.975, 1.025)	1.014 (0.988, 1.040)	1.003 (0.990, 1.017)
PM ₁₀	1.018 (0.994, 1.043)	1.026 (1.001, 1.051)	1.009 (0.983, 1.035)	1.033 (1.006, 1.061)	1.020 (0.992, 1.049)	0.995 (0.976, 1.014)
PM _{10-2.5}	1.041 (1.010, 1.073)	1.036 (1.006, 1.068)	1.018 (0.985, 1.052)	1.073 (1.041, 1.107)	1.042 (1.004, 1.080)	1.001 (0.981, 1.020)
PM _{2.5}	1.005 (0.978, 1.031)	0.995 (0.969, 1.022)	0.999 (0.971, 1.028)	1.027 (0.998, 1.056)	1.012 (0.982, 1.044)	0.982 (0.961, 1.003)
PM _{2.5} sulfate	0.991 (0.953, 1.029)	0.976 (0.939, 1.014)	0.980 (0.941, 1.022)	1.026 (0.983, 1.070)	1.012 (0.968, 1.059)	0.971 (0.943, 1.000)
PM _{2.5} elemental carbon	1.003 (0.981, 1.026)	1.005 (0.983, 1.028)	0.998 (0.974, 1.022)	1.014 (0.992, 1.037)	1.004 (0.979, 1.030)	0.996 (0.981, 1.012)
PM _{2.5} organic carbon	1.000 (0.985, 1.016)	0.999 (0.984, 1.015)	0.998 (0.981, 1.105)	1.007 (0.992, 1.023)	1.002 (0.985, 1.020)	0.994 (0.983, 1.010)
PM _{2.5} water-soluble metals	1.005 (0.968, 1.043)	1.004 (0.967, 1.043)	0.985 (0.949, 1.023)	1.017 (0.978, 1.058)	0.994 (0.949, 1.042)	0.998 (0.974, 1.023)

* The “base model” is the primary statistical model as described in the Methods section. The results in this column are reproduced from Table 2.

† Model is identical to the base model, except there is no control for pediatric emergency department visits for upper respiratory infections.

‡ Model is does not contain terms for year; month; month*year interactions; month*lag 0 maximum temperature interactions; and month*day-of-week interactions. Instead cubic splines with monthly knots are included to control for long-term and seasonal trends. Daily average temperature and dew point were modeled using cubic splines with knots at the 25th and 75th percentiles. These models are analogous to those we presented in earlier publications (14, 18, 20, 24, 30).

§ Model contains three-way interactions between year, month, and day-of-week and does not contain month*lag 0 maximum temperature interactions.

|| Model is identical to the base model, except that bimonthly (twice-per-month) indicator variables are used instead of monthly indicator variables.

**Model is identical to the base model, save for the addition of tomorrow’s (lag -1) pollutant concentration, which has been included in the model. The results in this column are the rate ratio estimated for the lag -1 term.




Figure 1.




Figure 2.




Figure 3.

Short-term Associations between Ambient Air Pollutants and Pediatric Asthma
Emergency Department Visits

Matthew J. Strickland, Lyndsey A. Darrow, Mitchel Klein, W. Dana Flanders, Jeremy A. Sarnat, Lance A. Waller, Stefanie E. Sarnat, James A. Mulholland, Paige E. Tolbert

Online Data Supplement

Table E1. Partial spearman correlation coefficients* for daily population-weighted pollutant concentrations during the warm season (bottom) and cold season (top, shaded).

	Ozone	Nitrogen dioxide	Carbon monoxide	Sulfur dioxide	PM ₁₀	PM _{10-2.5}	PM _{2.5}	PM _{2.5} sulfate	PM _{2.5} element. carbon	PM _{2.5} organic carbon	PM _{2.5} water-soluble metals
Ozone		0.11	-0.02	0.04	-0.05	-0.05	-0.12	-0.02	-0.16	-0.08	-0.07
Nitrogen dioxide	0.42		0.59	0.36	0.46	0.25	0.37	0.11	0.51	0.42	0.41
Carbon monoxide	0.24	0.54		0.19	0.49	0.22	0.38	0.09	0.58	0.48	0.43
Sulfur dioxide	0.17	0.37	0.20		0.15	0.08	0.00	0.10	0.09	0.02	0.05
PM ₁₀	0.47	0.44	0.37	0.23		0.49	0.76	0.51	0.54	0.63	0.55
PM _{10-2.5}	0.15	0.27	0.27	0.17	0.51		0.26	0.15	0.31	0.29	0.33
PM _{2.5}	0.50	0.36	0.32	0.13	0.82	0.29		0.68	0.51	0.66	0.56
PM _{2.5} sulfate	0.45	0.25	0.18	0.16	0.71	0.19	0.81		0.19	0.31	0.39
PM _{2.5} elemental carbon	0.25	0.47	0.52	0.20	0.47	0.34	0.44	0.31		0.78	0.45
PM _{2.5} organic carbon	0.44	0.44	0.40	0.11	0.61	0.32	0.62	0.47	0.64		0.48
PM _{2.5} water-soluble metals	0.29	0.40	0.42	0.18	0.57	0.36	0.55	0.45	0.47	0.46	

*Correlation after adjusting for log of the daily count of emergency department visits for upper respiratory infections, same day maximum temperature (exact °C), lag 1-2 minimum temperature (cubic terms), lag 0-1-2 average dew point temperature (cubic terms), month, year, month*year interactions, weekday, holiday, maximum temperature*month interactions, month*weekday interactions.