

RESEARCH ARTICLE

Streptococcus pneumoniae Induces Autophagy through the Inhibition of the PI3K-I/Akt/mTOR Pathway and ROS Hypergeneration in A549 Cells

Pu Li¹, Jing Shi², Qiao He¹, Qin Hu¹, Yun Ying Wang¹, Li Jun Zhang¹, Wai Ting Chan³, Wei-Xian Chen^{1*}

1 Department of Laboratory medicine, the Second Hospital Affiliated to Chongqing Medical University, Chongqing, China, **2** Department of Laboratory medicine, the First Hospital Affiliated to Chongqing Medical University, Chongqing, China, **3** Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas, Ramiro de Maeztu 9, Madrid, Spain

* chenweixian75@163.com



OPEN ACCESS

Citation: Li P, Shi J, He Q, Hu Q, Wang YY, Zhang LJ, et al. (2015) *Streptococcus pneumoniae* Induces Autophagy through the Inhibition of the PI3K-I/Akt/mTOR Pathway and ROS Hypergeneration in A549 Cells. PLoS ONE 10(3): e0122753. doi:10.1371/journal.pone.0122753

Academic Editor: Jia Luo, University of Kentucky, UNITED STATES

Received: October 28, 2014

Accepted: February 12, 2015

Published: March 24, 2015

Copyright: © 2015 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This project was supported by research grants from National Natural Science Foundation of China (no. 81171628). Author WTC was supported by the Spanish Ministry of Economy and Competitiveness, grant CSD2008/00013. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

The present study focused on the action mechanism of *S. pneumoniae* (Sp) in inducing autophagy in human alveolar epithelial cells. Sp, a gram-positive extracellular bacterium, activates autophagy with considerably increased microtubule-associated protein light chain 3 (LC3) punctation in A549 cells. The accumulation of typical autophagosomes and conjugation of LC3 to phosphatidylethanolamine were observed in Sp-infected cells as an indication of autophagy. Using the pneumolysin (PLY) mutant, we successfully demonstrated that PLY is involved in initiating autophagy without affecting the expression levels of PI3K-III and Beclin1. PLY-mediated autophagy depends on the inhibition of the phosphoinositide 3-kinase/Akt/mammalian target of rapamycin (PI3K/Akt/mTOR) pathway. Furthermore, Sp could also lead to the reactive oxygen species (ROS) hypergeneration in A549 cells. Taken together, Sp infection-induced autophagy is PLY-mediated through ROS hypergeneration and mTOR inhibition. PI3K-I and rapamycin (autophagy inducers) enhanced bacterial clearance, whereas wortmannin (autophagy inhibitor) and acetylcysteine (ROS inhibitor) reduced intracellular bacteria clearance. Thus, Sp-induced autophagy represents a host-protective mechanism, providing new insight into the pathogenesis of respiratory tract Sp infection.

Introduction

Extracellular bacterium *S. pneumoniae* (Sp) is a major human respiratory tract pathogen with a redundant set of virulence factors against host clearance [1]. One of the most important toxins released by Sp is pneumolysin (PLY), which has various immunomodulatory effects, including induction of cytokine production, reactive oxygen species (ROS) accumulation, and activation

the classical pathway of complement [2–3]. Recent studies have shown that epithelial cells of the human respiratory tract and lung play a critical role in defending against host mucosal pathogens [4], but their role in fighting against Sp remains to be fully defined.

Autophagy is an intracellular process that delivers cytoplasmic components to the autophagosome and lysosome for degradation [5]. The autophagosome is the central organelle that eliminates intracellular pathogens and degrades cytoplasmic material to fuel starving cells [6]. The growing body of research has demonstrated that the autophagy pathway is a critical cellular process that strongly influences the functions of epithelial and immune cells [7]. Several signaling pathways have been implicated in regulating autophagy, including phosphoinositide 3-kinase/Akt/mammalian target of rapamycin (PI3K/Akt/mTOR) and ROS. Class I PI3Ks (PI3K-I) inhibits autophagy through triggers the target of mTOR (rapamycin) [8], whereas ROS upregulates autophagy under oxidative stress and inflammatory conditions, such as pathogenic microbe infections [9–10]. Thus, targeting critical autophagy regulators with a goal to promote autophagy in epithelial cells is an attractive new therapeutic strategy for mucosal pathogen infections [11–12].

Previous studies showed that the induction of autophagy can protect alveolar epithelial cells from respiratory pathogens infection, such as *Mycobacterium tuberculosis*, group A streptococci, and *Pseudomonas aeruginosa* [13–15], indicating that autophagy acts as an immune effector that mediates pathogen clearance [16]. However, most studies of bacterial autophagy only involve intracellular pathogens [17]. Until now, the role of autophagy in Sp pathogenesis has been completely unknown. Thus, we studied autophagy in Sp-infected A549 cells and, for the first time, revealed the induction of autophagy by pneumococcal PLY through inhibition of the PI3K/AKT/mTOR pathway via ROS. This observation could provide useful information for further understanding of the role of autophagy in respiratory pneumococcal infection and improve our knowledge of mucosal immunity against this pathogen.

Materials and Methods

Cells, bacteria, vectors, and cell transfection

A549 (human alveolar epithelial) cell lines and breast cancer cell line MCF7 were purchased from ATCC (USA) and maintained according to the supplier's instructions. Bacteria strains Sp strain 35A (st35A) wild-type (WT) was isolated and collected from the Department of Laboratory Medicine (The Second Hospital Affiliated to Chongqing Medical University, Chongqing, China). Corresponding PLY-negative mutants (mut-PLY) developed through insertion-duplication mutagenesis, as described previously [18], were cultivated prior to infection analyses under antibiotic pressure with 10 mg/L erythromycin and 50 mg/L kanamycin. The plasmid pMV158GFP, which harbors the gene encoding the green fluorescent protein under the control of a promoter inducible by maltose, was a gift from Manuel Espinosa (Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas, Spain) [19]. The pMV158GFP was transferred into Sp (Sp-GFP) according to the standard transfer assays as previously described [20]. The GFP-LC3 plasmid was kindly provided by Dr. Juan Chen (Chinese University of Hong Kong, China). The RFP-PLY plasmid was constructed by cloning the coding sequence of PLY (GenBank: X52474.1) to vector pHcRed1-N1/1 (Clontech, Cat. No. 632424), and the PCR primers for PLY were ATGGCAAATAAAGCAGTAAA (forward) and CTAGTCATTTTCTACCTTAT (reverse). A549 cells were transfected/co-transfected with GFP-LC3 and/or RFP-PLY plasmids using Lipofectamine 2000 reagent (Invitrogen, Grand Island, NY, USA) in serum-free Dulbecco's modified Eagle's medium (DMEM) (Invitrogen) following the manufacturer's instructions.

Proteins, antibodies, and reagents

Anti-PLY antibody was purchased from Abcam (ab71810, Cambridge, MA, USA), and sequestosome (p62) monoclonal antibody (H00008878-M01) was purchased from Abnova (Taipei, Taiwan). Phospho-PI3 kinase p85 (Tyr458)/p55 (Tyr199) antibody (4228), PI3 kinase class III antibody (3811), phospho-Akt (Ser473) (D9E) XP rabbit mAb (4060), phospho-ULK1 (Ser317) (D2B6Y) rabbit mAb (12753), mTOR rabbit mAb (2983), phospho-mTOR antibody (2974), phospho-Beclin-1 (Ser93/96) antibody (12476), and beclin-1 (D40C5) rabbit mAb (3495), rabbit monoclonal antibody against β -actin were all purchased from Cell Signaling Technology (Beverly, MA, USA). Rabbit polyclonal antibody against LC3B (light chain 3B, LC3B; L8918), rapamycin (R8781; Rapa, mTOR inhibitor), 3-methyladenine (M9281; 3-MA, PI3K-III inhibitor), and wortmannin (W1628; WT) were all obtained from Sigma Life Science (Shanghai, China). PIK-93 (S1489; Class I PI3 Kinase inhibitor) was purchased from Selleck Chemicals (Houston, TX, USA).

Infection experiments

The bacteria were grown for 15 hours in Todd-Hewitt medium supplemented with 1% yeast extract (THY medium; Oxoid, England) at 37°C and 5% CO₂ with shaking. The bacteria were then pelleted by centrifugation at 5000 rpm and resuspended in 10 mL of fresh THY broth, in which they were allowed to grow until the mid-logarithmic phase [21]. Thereafter, the optical density (OD) at 600 nm was measured, and the OD was adjusted to 0.25 (0.1OD = 1×10^8 cells/mL). Accuracy of the bacterial colony-forming units (CFUs) was verified by plate counts of serial dilutions.

The bacteria were washed with phosphate buffer saline (PBS) before being changed to antibiotic-free medium for infection. The bacteria then were added to cell cultures at a 1:30 multiplicity of infection (MOI). After 1 hour, the infected cells were washed 3 times with PBS, and antibiotics (100 μ g/mL gentamicin and 250 mg/mL

Ceftriaxone Sodium) were added to kill extracellular bacteria. The cells were further cultured for the indicated time (post infection; specific time was described in results section). The extracellular bacterial clearance and invading bacteria inside the cells were determined using the CFU assay [13].

Transmission electron microscopy analysis

Transmission electron microscopy (TEM) was employed to identify autophagosomes with modified Karnovsky's fixative [22]. A549 cells were infected with bacteria at an MOI of 1:30 for 1.5 h. The cells were then fixed with 2.5% glutaraldehyde. Ultrathin sections were stained with uranyl acetate and lead citrate and the sections were then analyzed by TEM. All sections were observed with an H7600 electron microscope (Hitachi, Japan).

ROS assay

Relative changes in intracellular ROS in A549 cells were monitored using a fluorescent probe, 20, 70-dichlorofluorescein diacetate (DCFH-DA) [23]. The cells were seeded and cultured on 96-well plates at a density of 1×10^4 cells/well. The cells were incubated with 10 μ mol/L of DCFH-DA for 30 min, washed 3 times with serum-free DMEM medium, and then infected with bacteria for 1 hour. Then the cells were subjected to fluorescence detection at the indicated times (30 min, 1 h, 2 h, and 3 h) in a FLX800TBI fluorescence analyzer (BioTek, USA) with excitation/emission set at 488/525 nm. The values were expressed as percentage of relative fluorescence compared to the control.

Western blotting

The cells were lysed and protease inhibitor cocktail added to obtain the whole protein to be quantified. The lysates with protein loading buffer were boiled for 5 min. The supernatants were collected and 40 µg of each sample were loaded onto 15% SDS-PAGE gels and electrophoresed for protein resolution. The separated proteins were then transferred to polyvinylidene fluoride membranes (0.45 µm pore size; Millipore, Billerica, MA) and blocked for 2 h at room temperature. The PVDF membranes were incubated overnight at 4°C with primary antibodies diluted at 1:1,000 in PBS with 5% bovine serum albumin buffer. After washing 3 times with tris-buffered saline (TBS) with 0.5% (v/v) Tween, the membranes were incubated for 2 h at room temperature with horseradish peroxidase-conjugated secondary antibody (1:2,000 dilution). The immunoreactive bands were visualized by enhanced chemiluminescence reagents (Millipore).

Statistical analysis

All experiments were performed in at least three independent assays. Data are presented as percentage change compared with the controls \pm SD from the three independent experiments. Group means were compared by one-way analysis of variance (ANOVA) with SPSS software. Statistical significances were analyzed by the two-tailed unpaired Student's *t*-test; *P*-values of less than 0.05 were considered statistically significant.

Results

Sp infection induces LC3 punctation in A549 cells

To investigate whether infection by Sp (st35A) could induce autophagy, GFP-LC3 vectors were transfected into A549 cells. The A549 cells were infected with st35A, a strain of Sp, in a dose- and time-dependent manner. We noticed that Sp infection with an MOI of 30:1 (bacteria: cell) induced remarkably high LC3 punctation in A549 cells (Fig 1A). Cellular apoptosis was also detected to exclude the possibility that autophagy was caused by the induction of A549 cell death. LC3 punctation reached its peak at 3 h after infection and no observable cell death was detected at this time (Fig 1B). The LC3 punctation, thereafter, gradually decreased, probably because of increased autophagic cell death caused by infection [5] (Fig 1C). A549 cells transfected with GFP-LC3 were also treated with the autophagy activator rapamycin (positive) and inhibitor wortmannin (negative) as controls (Fig 1A). To determine whether autophagy is a general phenomenon in Sp infection, other cell type was also investigated. We also noted that Sp infection induced LC3 punctation in breast cancer cell line MCF7 (Fig 1D).

Sp infection induces increased autophagosome and LC3-II formation and upregulated autophagic degradation

To exclude the possibility that the punctation flux was caused by transient overexpression of GFP-LC3 proteins, we further determined the newly formed autophagosomes with TEM. We observed that typical autophagosomes (with double membranes and vacuoles) were notably increased in st35A-infected A549 cells as compared with the control cells (Fig 2A). Thus, the morphological evidence further confirms that elevated autophagosome formation is consistent with the accumulation of LC3 punctation. Our results demonstrate that Sp infection-induced autophagy in A549 cells.

Autophagosomes accumulation might be caused by the blockage of autophagosome degradation rather than induction of autophagosome formation [24]. LC3-II is a well-established autophagosomes marker in mammalian cells. Therefore, we examined the change in LC3




Fig 1. Sp infection induced LC3 punctation in A549 cells. A549 cells were infected with Sp (st35A) for 1 h (MOI=30:1). Before infection, the cells were treated with rapamycin (1 mM, 10 h) and wortmannin (2 μM, 2 h). (A) A549 cells were transfected with GFP-LC3 plasmids for 24 h. Then, the cells were infected with Sp for 1 h (MOI=30:1). (B) The puncta in each cell were counted and cells with more than 10 punctae were considered GFP-LC3 puncta cells. Values are from 100 cells/sample (one-way ANOVA; Tukey's posthoc test). (C) A549 cells were infected at different times and with different MOIs. (D) Breast cancer cell line MCF7 cells were infected with Sp (st35A) for 4 h (MOI=30:1). The percentage of GFP-LC3 puncta cells was determined (one-way ANOVA; Tukey's posthoc test). Data are representative of three experiments with similar results. Sp: *S. pneumoniae*; Rapa: rapamycin; WT: wortmannin.

doi:10.1371/journal.pone.0122753.g001




Fig 2. Sp infection induces increased autophagosome and LC3-II formation and upregulated autophagic degradation. (A) After 3 h infection, cells were processed and examined by TEM. A1. Untreated cells (-). A2. Cells infected with Sp (st35A). A3. Cells treated with rapamycin. A4. Cells treated with wortmannin and then infected with Sp. A5. The number of autophagic vesicles in each cell was determined with 10 cells in each sample (one-way ANOVA; Tukey's posthoc test). (B) Sp infection induces LC3-II formation and upregulated autophagic degradation. B1. Western blot of LC3. B2. Western blot of P62. B3. Western blot of P-ULK1. Data are representative of three experiments with similar results. Sp: *S. pneumoniae*; Rapa: rapamycin; WT: wortmannin; p62: Sequestosome 1; p-ULK1: phosphorylated UNC-51-like kinase 1.

doi:10.1371/journal.pone.0122753.g002

patterns at the molecular level. Endogenous LC3-I transformation into phosphatidylethanolamine-conjugated (PE-conjugated) LC3-II was dramatically increased by Sp infection in a time-dependent manner (Fig 2B1). We monitored autophagic flux (refers to the whole process of autophagy, including autophagosome formation, maturation, fusion with lysosomes, subsequent breakdown and the release of macromolecules back into the cytosol) through the levels of endogenous p62 expression. Consistent with the induction of LC3-II accumulation, p62 showed a decrease by Sp infection in A549 cells (Fig 2B2). We also monitored phosphorylated UNC-51-like kinase 1 (p-ULK1). ULK1 can act as a convergence point for multiple signals that control autophagy [25]. Consistent with the induction of LC3-II accumulation, p-ULK1 showed an increase by Sp infection in A549 cells (Fig 2B3). Taken together, these findings firmly establish that Sp infection can specifically induce autophagy in A549 cells.

Pneumococcal PLY is involved in initiating autophagy

Extracellular bacterium Sp could be phagocytosed by phagocytic cells [13–21], indicating autophagy through intracellular pathways. However, a previous study reported that even without bacteria internalization, cells were found to show significant autophagy and autophagy induction did not require intracellular bacteria [14]. Thus, it could be inferred that particular virulence factors released by Sp, such as pneumococcal PLY, attributed to autophagy induction.

To determine PLY in inducing autophagy, we used the PLY-deficient strains (mut-PLY) and found that mut-PLY failed to induce LC3 punctation (Fig 3A). To further confirm whether intracellular PLY is involved in initiating autophagy, we constructed a RFP-PLY eukaryotic plasmid and transfected/cotransfected GFP-LC3 or/and RFP-PLY into A549 cells (Fig 3B). Autophagy typical markers for LC3 punctation, LC3 proteins, and p62 proteins were then examined as described above. We noticed that both the accumulation of LC3 punctation (Fig 3B and 3C) and the change in LC3 protein patterns (Fig 3D) were consistent with st35A-infected A549 cells. Thus, our results demonstrated that virulence factor PLY is involved in initiating autophagy in A549 cells.

Pneumococcus PLY induces autophagy by inhibiting the PI3K/AKT/mTOR pathway and it is mediated through ROS hypergeneration

The PI3K pathways act as a regulator of autophagy [8]. Class I PI3Ks (PI3K-I) triggers the target of the rapamycin signaling pathway (PI3K/AKT/mTOR), which inhibits autophagy. Class III PI3Ks (PI3K-III) mediates the recruitment of specific autophagic effectors to the sites of origin of autophagic membranes and accelerates autophagy. Thus, we examined the involvement of autophagic protein PI3Ks. We found that PI3Ks, upstream regulators of autophagy, are involved in autophagy. Expression of phosphorylated PI3K-I (p-PI3K-I) was downregulated with Sp infection in A549 cells (Fig 4A1).

To dissect this pathway further, we examined a major downstream autophagy-related protein Akt and mTOR. As with p-PI3K-I, Sp infection also resulted in a substantial downregulation of phosphorylated Akt (p-Akt) (Fig 4A2). However, Sp infection induces the downregulation of the PI3K/AKT/mTOR pathway without affecting the expression levels of PI3K-III and Beclin 1 (Fig 4A3). From the present results, our data identified that Sp-induced autophagy depends on the PI3K/Akt/mTOR inhibition pathway in A549 cells.

To identify whether pneumococcus PLY-induced autophagy is mediated through ROS hypergeneration, we detected the intracellular levels of ROS in A549 cells. ROS are generated during mitochondrial oxidative metabolism, as well as in cellular response to xenobiotics, cytokines, and bacterial invasion. Growing bodies of study show that autophagy activated by ROS




Fig 3. Pneumococcal PLY is involved in initiating autophagy. A549 cells were infected with Sp or mut-PLY for 1 h (MOI=30:1). Before infection, the cells were also treated with rapamycin (1 mM, 10 h) and wortmannin (2 μ M, 2 h). (A) mut-PLY failed to induce LC3 punctation. (B) RFP-PLY induced LC3 punctation in cotransfected cells. (C) The puncta in each cell were counted and cells with more than 10 punctae were considered GFP-LC3 puncta cells. Values are from 100 cells/sample (one-way ANOVA; Tukey's posthoc test). (D) Western blotting of LC3 and p62. Data are representative of three experiments with similar results; Rapa: rapamycin; WT: wortmannin; p62: Sequestosome 1; PLY: pneumolysin; mut-PLY: PLY-deficient strain.

doi:10.1371/journal.pone.0122753.g003

as a cytoprotective process [26–27]. On the other side, autophagy is stimulated to clear ROS to protect cells from bacterial toxin-induced damage [28]. However, the role of ROS generated by the host in pneumococcal containment during infection is unclear.

PLY induces the production of intracellular ROS in human neutrophils [2] and Sp may modulate the oxidative burst [29]. Therefore, the role of ROS in Sp-induced autophagy was examined. Sp infection increased the intensity of DCF fluorescence in A549 cells, showing that




Fig 4. Pneumococcus PLY induces autophagy by inhibiting the PI3K/AKT/mTOR pathway and it is mediated through ROS hypergeneration. (A). Sp infection induced autophagy through the PI3K/Akt/mTOR inhibition pathway. A549 cells were infected with Sp for 1 h. A1. Western blotting of phosphorylated PI3K-I. Sp (st35A)-induced autophagy depends on P-PI3K-I inhibition. A2. Western blotting of phosphorylated Akt and mTOR. Sp inhibits Akt/mTOR signaling in A549 cells. A3. The expression levels of PI3K-III and Beclin 1 were not affected by Sp infection. (B). Sp infection induces ROS hypergeneration. A549 cells were infected with Sp for 1 h and intracellular ROS hypergeneration was observed in A549 cells. B1. A time-dependent intracellular ROS hypergeneration was observed in A549 cells infected with st35A. B2. ROS was significantly increased when the cells were infected with st35A ($p < 0.005$). B3. The ROS-specific inhibitor NAC can upregulate PI3K/AKT/mTOR expression. Data are representative of three experiments with similar results. p-PI3K-I: phosphorylated PI3K-I; Rapa: rapamycin; Sp: *S. pneumoniae*; mut-PLY: PLY-deficient strain; ROS: reactive oxygen species; NAC: acetylcysteine (ROS inhibitor).

doi:10.1371/journal.pone.0122753.g004

intracellular ROS was increased within 30 min and peaked at 120 min (Fig 4B1). Thereafter, a decline in ROS generation was observed.

To analyze whether pneumococcus PLY-induced autophagy is mediated through ROS hypergeneration, the cells were also treated with mut-PLY and analyzed for cell ROS formation. The levels of ROS in mut-PLY-treated A549 cells were significantly reduced (Fig 4B2). However, acetylcysteine, the ROS specific inhibitor, can block this effect and upregulation of the PI3K/AKT/mTOR expression (Fig 4B3). These results confirm that the PLY-induced autophagy is mediated through ROS hypergeneration.

Autophagy regulates Sp clearance in A549 cells

Pneumococcus has long been considered an extracellular pathogen, and most autophagy studies thus far have only involved intracellular pathogens. To define the role of autophagy, we infected A549 cells with Sp-GFP to monitor internalization and bacterial clearance. After the cells were transfected with RFP-plasmid and then Sp infection, we utilized CFU to count the invading bacteria inside the cells. This approach showed bacterial internalization and the outline of cells (Fig 5), as well as the number of invading bacteria. To determine the role of autophagy in regulating phagocytosis of Sp and clearance, A549 cells were pre-treated with rapamycin, wortmannin, PI3K-I inhibitor, and ROS inhibitor respectively before infection. The intracellular Sp-GFP count decreased in rapamycin-treated and PI3K-I inhibitor-treated A549 cells, indicating that induction of autophagy can increase host defense against this pathogen. By contrast, the number of intracellular bacteria was increased by blocking autophagy with wortmannin in A549 cells. In particular, ROS-inhibitor treatment resulted in a marked increase in intracellular bacteria (Fig 5A and 5B). The data indicate that blocking autophagy with wortmannin or ROS inhibitor reduced Sp bacterial clearance. An autophagy activator, rapamycin, however, produced the opposite effect, improving bacterial clearance. Collectively, our observations indicate that autophagy might be a major benefit to host defenses by augmenting bacterial clearance.

Discussion

In the present study, we addressed that Sp infection specifically induces autophagy through the PI3K/AKT/mTOR autophagy pathway and ROS hypergeneration in a human pulmonary epithelial cell line (A549). Moreover, our observations showed that the autophagy was initiated by PLY, an important virulence factor of Sp. Bacterial pore-forming toxins have been shown to induce autophagy through mTOR [30]. Several pathogens, such as intracellular bacteria and viruses, could induce autophagy, although the pathways involved and the impact on infection outcomes vary with intracellular pathogens [24, 31–33].

Sp is traditionally considered an extracellular pathogen, and many virulence factors, such as PLY and capsular polysaccharide, might contribute to the extracellular pathogenic features of the bacterium [2, 34]. To our knowledge, prior to the current study, it was not known whether Sp-infection could induce autophagy. This is the first to demonstrate the induction of autophagy by Sp through the inhibition of the PI3K/Akt/mTOR pathway and ROS hypergeneration, as well as its physiological significance in relation to improved bacterial clearance.

Under stress or infection conditions, endogenous LC3-I proteins can conjugate with PE to form LC3-II [35]. Contrary to the cytoplasmic localization of LC3-I, LC3-II associates with both the outer and inner membranes of the autophagosome, thereby being a typical marker of autophagy formation. We identified LC3 punctation and confirmed the significant elevated autophagosome formation in st35A-infected A549 cells (Fig 1A; $p < 0.005$). Since p62 accumulates when autophagy is inhibited, and decreased levels can be observed when autophagy is




Fig 5. Autophagy regulates Sp clearance in A549 cells. (A) A549 cells were transfected with RFP-plasmid for 24 h and then infected with Sp-GFP for 1 h (MOI = 30:1). Before infection, the cells were also treated with rapamycin (1 mM, 10 h), wortmannin (2 μ M, 2 h), PI3K-I inhibitor, or ROS inhibitor. The fluorescence images are displayed to confirm the internalization of bacteria. (B) Number of internalized bacteria per cell. CFU experiments were utilized to count the invading bacteria inside the cells. The data are representative of 100 cells (one-way ANOVA; Tukey's posthoc test). Data are representative of three experiments with similar results. Rapa: rapamycin; WT: wortmannin; Sp: *S. pneumoniae*; ROS: reactive oxygen species; Sp-GFP: Sp + pMV158GFP plasmid; CFU: colony-forming unit.

doi:10.1371/journal.pone.0122753.g005

induced, p62 could be used as a marker to monitor autophagic flux. However, there is not always a clear correlation between increases in LC3-II and decreases in p62. Thus, we used LC3-II turnover in combination with p62 to monitor autophagy flux (Fig 2B). The increased autophagosome formation was also shown to be a result of autophagy induction rather than of blocked lysosome degradation. Rapamycin, a widely used inducer for autophagy induction, caused similar autophagy. However, wortmannin, a typical autophagy inhibitor, reduced autophagy following Sp infection.

Among the main mechanisms in the regulation of autophagy are the PI3Ks [8]. There are three distinct classes of PI3Ks; however, only Class I and III PI3Ks have been shown to play a role in autophagy [36]. The Class I and III PI3Ks have opposing roles in autophagy. PI3K-III

mediates the recruitment of specific autophagic effectors to the origin sites of autophagic membranes and, thereby, plays a positively role in canonical autophagy. By contrast, PI3K-I trigger the AKT/mTOR signaling pathway to inhibit autophagy.

In this work, we demonstrated that the crucial upstream autophagy regulator PI3K-I was inhibited by Sp infection (Fig 4A). To evaluate its key downstream autophagy-related protein, we examined the phosphorylated Akt and mTOR. We noticed that protein levels of the phosphorylated Akt and mTOR were decreased significantly (Fig 4B; $p < 0.005$), which were consistent with PI3K-I expression patterns. To determine additional virulence factors in inducing autophagy, we used the PLY-deficient strain (mut-PLY) and found that whereas the mut-PLY strain failed to induce autophagy, the Sp strain did induce it (Fig 3). Our observations indicate that virulence factor PLY might involve autophagy induction. Taken together, our studies indicate that Sp infection specifically induces autophagy and that this induction depends on the PI3K/AKT/mTOR autophagy pathway.

Having successfully demonstrated that Sp infection caused autophagy through the PI3K-I/AKT/mTOR pathway in A549 cells, we addressed the problem of how Sp induced the autophagic phenomenon. PLY has been characterized as a potent virulence factor released by Sp [37], which could induce cell death by pore formation and toxin-induced apoptosis, but although its cytolytic effects are well understood, less is known about the interactions between this potent toxin and the phenomenon of autophagy [3]. Therefore, the capacity of this pneumococcal factor to induce autophagy was analyzed. In this study, PLY-negative mutant (mut-PLY) was developed through insertion-duplication mutagenesis; The RFP-PLY plasmid was constructed and transfected to A549 cells, then PLY was overexpressed intracellularly. We did not observe significant apoptosis-changes in cell morphology.

In our analyses, transfection of A549 with the plasmid RFP-PLY resulted in a significant induction of autophagy (Fig 3; $p < 0.005$) and ROS hypergeneration (Fig 4), while the corresponding PLY-negative mutants led to a blockade of autophagy induction. Tattoli *et al* found that infection of epithelial cells with *Shigella* and *Salmonella* triggers acute intracellular amino acid starvation due to host membrane damage. Pathogen-induced amino acid starvation caused downregulation of mTOR activity, resulting in the induction of autophagy [38]. This study could be the most probable explanation for autophagy is induced in a PLY-dependent manner.

ROS has been shown to activate autophagy to protect cells from nutrient starvation, cell death, and invading pathogens. Some studies show that ROS activate autophagy by regulating mTOR activity. In malignant gliomas, ROS disrupt mitochondrial membrane potential and induce autophagy through inhibiting Akt/mTOR signaling [39]. In glioma cells, ROS induce autophagy by inhibiting mTOR in a BNIP3-dependent manner (BNIP3: BCL2/adenovirus E1B 19 kDa protein-interacting protein 3) [40]. ROS can also inhibit PTEN (phosphatase and tensin homolog located on chromosome 10) [41], which has been shown to activate autophagy through the downregulation of the PI3K-I/Akt pathway. However, many unknowns still exist regarding the downstream signals of ROS in autophagy regulation. In this study, we detected the hypergeneration of ROS and downregulation of PI3K-I/Akt/mTOR during Sp infection. Taken together, we conclude that pneumococcal PLY-induced autophagy is mediated through ROS hypergeneration.

Recently, the role of autophagy in bacterial infections disease has garnered increasing interest [42–43], and it has been shown to play crucial roles in host defense, especially in immunological cells. Moreover, autophagy has a direct impact on immunity and inflammatory response within the whole organism. Intracellular bacteria, such as *Salmonella* and *Mycobacterium tuberculosis*, have all been shown to fuse with or be directly engulfed by autophagosomes, offering an alternative method for the bacteria destruction by autophagy [44–45]. Although most studies of bacterial-specific autophagy involve the capture of intracellular bacteria,

autophagy can similarly eliminate extracellular bacteria *P. aeruginosa*, which have the capacity of autophagy induction in the alveolar macrophage cell line MH-S [14, 33].

Our studies have also defined that human pulmonary epithelial cell lines A549 can capture and destroy the invading Sp (Fig 5). PI3K is a classic pathway to influence the whole process of autophagy [8], whereas ROS can upregulate autophagy when host cells were infected with pathogenic microbe [9–10]. Thus, we targeted critical autophagy regulators with its inhibitors to observe the autophagy impact on Sp infections. Treatment with the autophagy inhibitor wortmannin or ROS inhibitor resulted in a marked increase in bacterial load whereas treatment with the autophagy inducer rapamycin or PI3K-I inhibitor enhanced Sp clearance (Fig 5B). Because airway epithelial cells and alveolar epithelial cells play a pivotal role in orchestrating both the innate immune response and the transition to adaptive immunity [46–47], autophagy in epithelial cells could impact the consequence of Sp infection. Previous reports have showed that Sp could be killed in a lysosome dependent manner [48], but host cells may use more than one strategy to eliminate invading pathogens. Based on our data, the mechanism of autophagy also plays a role in limiting Sp infection. In fact, lysosome system overlaps autophagy. More researches should be taken out in this field. Our results suggest that autophagy can mediate the removal of Sp in alveolar epithelial cells. Thus, Sp could serve as a model pathogen for investigating the impact of autophagy on crucial cellular events, such as bacterial entry and clearance. But Sp infection-induced autophagy in vivo need further assessment, however, this observation could provide useful information for further understanding the role of autophagy in respiratory pneumococcal infection and ultimately lead to new therapeutic targets.

Acknowledgments

We thank Prof. M. Espinosa for the gift of the plasmid pMV158GFP. We also thank Dr. Juan Chen kindly provided the GFP-LC3 plasmid.

Author Contributions

Conceived and designed the experiments: WC PL JS. Performed the experiments: PL JS Q. He. Analyzed the data: Q. Hu YYW. Contributed reagents/materials/analysis tools: LJZ WTC YYW. Wrote the paper: WC PL JS WTC.

References

1. McCullers JA. Insights into the interaction between influenza virus and pneumococcus. *Clin Microbiol Rev.* 2006; 19: 571–582. PMID: [16847087](#)
2. Martner A1, Dahlgren C, Paton JC, Wold AE. Pneumolysin released during *Streptococcus pneumoniae* autolysis is a potent activator of intracellular oxygen radical production in neutrophils. *Infect Immun.* 2008; 76: 4079–4087. doi: [10.1128/IAI.01747-07](#) PMID: [18559434](#)
3. Molloy S. Pneumolysin: stimulating protection. *Nat Rev Microbiol.* 2011; 9: 4. PMID: [21204311](#)
4. Eddens T, Kolls J K. Host defenses against bacterial lower respiratory tract infection. *Curr Opin Immunol.* 2012; 24: 424–430. doi: [10.1016/j.coi.2012.07.005](#) PMID: [22841348](#)
5. Choi AM, Ryter SW, Levine B. Autophagy in human health and disease. *N Engl J Med.* 2013; 368: 1845–1846. doi: [10.1056/NEJMc1303158#SA1](#) PMID: [23656659](#)
6. Codogno P, Mehrpour M, Proikas-Cezanne T. Canonical and non-canonical autophagy: variations on a common theme of self-eating? *Nat Rev Mol Cell Biol.* 2011; 13: 7–12. doi: [10.1038/nrm3249](#) PMID: [22166994](#)
7. Patel KK, Stappenbeck TS. Autophagy and intestinal homeostasis. *Annu Rev Physiol.* 2013; 75: 241–262. doi: [10.1146/annurev-physiol-030212-183658](#) PMID: [23216414](#)
8. FOF, Rusten TE, Stenmark H. Phosphoinositide 3-kinases as accelerators and brakes of autophagy. *FEBS J.* 2013; 280: 6322–6337. doi: [10.1111/febs.12486](#) PMID: [23953235](#)
9. Huang J, Lam GY, Brumell JH. Autophagy signaling through reactive oxygen species. *Antioxid Redox Signal.* 2011; 14: 2215–2231. doi: [10.1089/ars.2010.3554](#) PMID: [20874258](#)

10. Kroemer G, Marino G, Levine B. Autophagy and the integrated stress response. *Mol Cell*. 2010; 40: 280–293. doi: [10.1016/j.molcel.2010.09.023](https://doi.org/10.1016/j.molcel.2010.09.023) PMID: [20965422](https://pubmed.ncbi.nlm.nih.gov/20965422/)
11. Shoji-Kawata S, Sumpster R, Leveno M, Campbell GR, Zou Z, Kinch L, et al. Identification of a candidate therapeutic autophagy-inducing peptide. *Nature*. 2013; 494: 201–206. doi: [10.1038/nature11866](https://doi.org/10.1038/nature11866) PMID: [23364696](https://pubmed.ncbi.nlm.nih.gov/23364696/)
12. Patel AS, Morse D, Choi AM. Regulation and functional significance of autophagy in respiratory cell biology and disease. *Am J Respir Cell Mol Biol*. 2013; 48: 1–9. doi: [10.1165/rcmb.2012-0282TR](https://doi.org/10.1165/rcmb.2012-0282TR) PMID: [22984088](https://pubmed.ncbi.nlm.nih.gov/22984088/)
13. Nakagawa I, Amano A, Mizushima N, Yamamoto A, Yamaguchi H, Kamimoto T, et al. Autophagy defends cells against invading group A Streptococcus. *Science*. 2004; 306: 1037–1040. PMID: [15528445](https://pubmed.ncbi.nlm.nih.gov/15528445/)
14. Yuan K, Huang C, Fox J, Laturnus D, Carlson E, Zhang B, et al. Autophagy plays an essential role in the clearance of *Pseudomonas aeruginosa* by alveolar macrophages. *J Cell Sci*. 2012; 125: 507–515. doi: [10.1242/jcs.094573](https://doi.org/10.1242/jcs.094573) PMID: [22302984](https://pubmed.ncbi.nlm.nih.gov/22302984/)
15. Guo XG, Ji TX, Xia Y, Ma YY. Autophagy protects type II alveolar epithelial cells from *Mycobacterium tuberculosis* infection. *Biochem Biophys Res Commun*. 2013; 432: 308–313. doi: [10.1016/j.bbrc.2013.01.111](https://doi.org/10.1016/j.bbrc.2013.01.111) PMID: [23396060](https://pubmed.ncbi.nlm.nih.gov/23396060/)
16. Puleston DJ, Simon AK. Autophagy in the immune system. *Immunology*. 2014; 141: 1–8. doi: [10.1111/imm.12165](https://doi.org/10.1111/imm.12165) PMID: [23991647](https://pubmed.ncbi.nlm.nih.gov/23991647/)
17. Deretic V. Autophagy in immunity and cell-autonomous defense against intracellular microbes. *Immunol Rev*. 2011; 240:92–104. doi: [10.1111/j.1600-065X.2010.00995.x](https://doi.org/10.1111/j.1600-065X.2010.00995.x) PMID: [21349088](https://pubmed.ncbi.nlm.nih.gov/21349088/)
18. Berry AM, Yother J, Briles DE, Hansman D, Paton JC. Reduced virulence of a defined pneumolysin-negative mutant of *Streptococcus pneumoniae*. *Infect Immun*. 1989; 57: 2037–2042. PMID: [2731982](https://pubmed.ncbi.nlm.nih.gov/2731982/)
19. Nieto C, Espinosa M. Construction of the mobilizable plasmid pMV158GFP, a derivative of pMV158 that carries the gene encoding the green fluorescent protein. *Plasmid*. 2003; 49: 281–285. PMID: [12749839](https://pubmed.ncbi.nlm.nih.gov/12749839/)
20. Priebe SD, Lacks SA. Region of the streptococcal plasmid pMV158 required for conjugative mobilization. *J Bacteriol*. 1989; 171: 4778–4784. PMID: [2768188](https://pubmed.ncbi.nlm.nih.gov/2768188/)
21. Lüttge M, Fulde M, Talay SR, Nerlich A, Rohde M, Preissner KT, et al. *Streptococcus pneumoniae* induces exocytosis of Weibel-Palade bodies in pulmonary endothelial cells. *Cell Microbiol*. 2012; 14: 210–225. doi: [10.1111/j.1462-5822.2011.01712.x](https://doi.org/10.1111/j.1462-5822.2011.01712.x) PMID: [21999205](https://pubmed.ncbi.nlm.nih.gov/21999205/)
22. Burghardt RC, Droleskey R. Transmission electron microscopy. *Curr Protoc Microbiol*. 2006; Chapter 2, Unit 2B 1. doi: [10.1002/9780471729259.mc02b01s03](https://doi.org/10.1002/9780471729259.mc02b01s03) PMID: [18770588](https://pubmed.ncbi.nlm.nih.gov/18770588/)
23. Kuster GM, Pimentel DR, Adachi T, Ido Y, Brenner DA, Cohen RA, et al. Alpha-adrenergic receptor-stimulated hypertrophy in adult rat ventricular myocytes is mediated via thioredoxin-1-sensitive oxidative modification of thiols on Ras. *Circulation*. 2005; 111: 1192–1198. PMID: [15723974](https://pubmed.ncbi.nlm.nih.gov/15723974/)
24. Deretic V, Saitoh T, Akira S. Autophagy in infection, inflammation and immunity. *Nat Rev Immunol*. 2013; 13: 722–737. doi: [10.1038/nri3532](https://doi.org/10.1038/nri3532) PMID: [24064518](https://pubmed.ncbi.nlm.nih.gov/24064518/)
25. Young AR, Chan EY, Hu XW, Köchl R, Crawshaw SG, High S, et al. Starvation and ULK1-dependent cycling of mammalian Atg9 between the TGN and endosomes. *J Cell Sci*. 2006; 119: 3888–3900. PMID: [16940348](https://pubmed.ncbi.nlm.nih.gov/16940348/)
26. Nair RR, Emmons MF, Cress AE, Argilagos RF, Lam K, Kerr WT, et al. HYD1-induced increase in reactive oxygen species leads to autophagy and necrotic cell death in multiple myeloma cells. *Mol Cancer Ther*. 2009; 8: 2441–2451. doi: [10.1158/1535-7163.MCT-09-0113](https://doi.org/10.1158/1535-7163.MCT-09-0113) PMID: [19671765](https://pubmed.ncbi.nlm.nih.gov/19671765/)
27. Huang J, Canadien V, Lam GY, Steinberg BE, Dinauer MC, Magalhaes MA, et al. Activation of antibacterial autophagy by NADPH oxidases. *Proc Natl Acad Sci U S A*. 2009; 106: 6226–6231. doi: [10.1073/pnas.0811045106](https://doi.org/10.1073/pnas.0811045106) PMID: [19339495](https://pubmed.ncbi.nlm.nih.gov/19339495/)
28. Yuan H, Perry CN, Huang C, Iwai-Kanai E, Carreira RS, Glembotski CC, et al. LPS-induced autophagy is mediated by oxidative signaling in cardiomyocytes and is associated with cytoprotection. *Am J Physiol Heart Circ Physiol*. 2009; 296: H470–479. doi: [10.1152/ajpheart.01051.2008](https://doi.org/10.1152/ajpheart.01051.2008) PMID: [19098111](https://pubmed.ncbi.nlm.nih.gov/19098111/)
29. Barbuti G, Moschioni M, Fumarulo R, Censini S, Montemurro P. *Streptococcus pneumoniae* modulates the respiratory burst response in human neutrophils. *FEMS Immunol Med Microbiol*. 2010; 60: 57–62. doi: [10.1111/j.1574-695X.2010.00716.x](https://doi.org/10.1111/j.1574-695X.2010.00716.x) PMID: [20618848](https://pubmed.ncbi.nlm.nih.gov/20618848/)
30. Shin JH, Park SJ, Kim ES, Jo YK, Hong J, Cho DH. Sertindole, a potent antagonist at dopamine D(2) receptors, induces autophagy by increasing reactive oxygen species in SH-SY5Y neuroblastoma cells. *Biol Pharm Bull*. 2012; 35: 1069–1075. PMID: [22791154](https://pubmed.ncbi.nlm.nih.gov/22791154/)
31. Wileman T. Autophagy as a defence against intracellular pathogens. *Essays Biochem*. 2013; 55: 153–163. doi: [10.1042/bse0550153](https://doi.org/10.1042/bse0550153) PMID: [24070478](https://pubmed.ncbi.nlm.nih.gov/24070478/)

32. Choy A, Roy CR. Autophagy and bacterial infection: an evolving arms race. *Trends Microbiol.* 2013; 21: 451–456. doi: [10.1016/j.tim.2013.06.009](https://doi.org/10.1016/j.tim.2013.06.009) PMID: [23880062](https://pubmed.ncbi.nlm.nih.gov/23880062/)
33. Junkins RD, Shen A, Rosen K, McCormick C, Lin TJ. Autophagy enhances bacterial clearance during *P. aeruginosa* lung infection. *PLoS One.* 2013; 8: e72263. doi: [10.1371/journal.pone.0072263](https://doi.org/10.1371/journal.pone.0072263) PMID: [24015228](https://pubmed.ncbi.nlm.nih.gov/24015228/)
34. van der Poll T, Opal SM. Pathogenesis, treatment, and prevention of pneumococcal pneumonia. *Lancet.* 2009; 374: 1543–1556. doi: [10.1016/S0140-6736\(09\)61114-4](https://doi.org/10.1016/S0140-6736(09)61114-4) PMID: [19880020](https://pubmed.ncbi.nlm.nih.gov/19880020/)
35. Ichimura Y, Imamura Y, Emoto K, Umeda M, Noda T, Ohsumi Y. In vivo and in vitro reconstitution of Atg8 conjugation essential for autophagy. *J Biol Chem.* 2004; 279: 40584–40592. PMID: [15277523](https://pubmed.ncbi.nlm.nih.gov/15277523/)
36. Petiot A, Ogier-Denis E, Blommaert EF, Meijer AJ, Codogno P. Distinct classes of phosphatidylinositol 3'-kinases are involved in signaling pathways that control macroautophagy in HT-29 cells. *J Biol Chem.* 2000; 275: 992–998. PMID: [10625637](https://pubmed.ncbi.nlm.nih.gov/10625637/)
37. Mitchell AM, Mitchell TJ. *Streptococcus pneumoniae*: virulence factors and variation. *Clin Microbiol Infect.* 2010; 16: 411–418. doi: [10.1111/j.1469-0691.2010.03183.x](https://doi.org/10.1111/j.1469-0691.2010.03183.x) PMID: [20132250](https://pubmed.ncbi.nlm.nih.gov/20132250/)
38. Yang F, Chen WD, Deng R, Li DD, Wu KW, Feng GK, et al. Hirsutanol A induces apoptosis and autophagy via reactive oxygen species accumulation in breast cancer MCF-7 cells. *J Pharmacol Sci.* 2012; 119: 214–220. PMID: [22786562](https://pubmed.ncbi.nlm.nih.gov/22786562/)
39. Zhang H, Kong X, Kang J, Su J, Li Y, Zhong J, et al. Oxidative stress induces parallel autophagy and mitochondria dysfunction in human glioma U251 cells. *Toxicol Sci.* 2009; 110: 376–388. doi: [10.1093/toxsci/kfp101](https://doi.org/10.1093/toxsci/kfp101) PMID: [19451193](https://pubmed.ncbi.nlm.nih.gov/19451193/)
40. Byun YJ, Kim SK, Kim YM, Chae GT, Jeong SW, Lee SB. Hydrogen peroxide induces autophagic cell death in C6 glioma cells via BNIP3-mediated suppression of the mTOR pathway. *Neurosci Lett.* 2009; 461: 131–135. doi: [10.1016/j.neulet.2009.06.011](https://doi.org/10.1016/j.neulet.2009.06.011) PMID: [19539716](https://pubmed.ncbi.nlm.nih.gov/19539716/)
41. Cho SH, Lee CH, Ahn Y, Kim H, Kim H, Ahn CY, et al. Redox regulation of PTEN and protein tyrosine phosphatases in H(2)O(2) mediated cell signaling. *FEBS Lett.* 2004; 560: 7–13. PMID: [15017976](https://pubmed.ncbi.nlm.nih.gov/15017976/)
42. Haspel JA, Choi AM. Autophagy: a core cellular process with emerging links to pulmonary disease. *Am J Respir Crit Care Med.* 2011; 184: 1237–1246. doi: [10.1164/rccm.201106-0966CI](https://doi.org/10.1164/rccm.201106-0966CI) PMID: [21836133](https://pubmed.ncbi.nlm.nih.gov/21836133/)
43. Mostowy S. Autophagy and bacterial clearance: a not so clear picture. *Cell Microbiol.* 2013; 15: 395–402. doi: [10.1111/cmi.12063](https://doi.org/10.1111/cmi.12063) PMID: [23121192](https://pubmed.ncbi.nlm.nih.gov/23121192/)
44. Castillo EF, Dekonenko A, Arko-Mensah J, Mandell MA, Dupont N, Jiang S, et al. Autophagy protects against active tuberculosis by suppressing bacterial burden and inflammation. *Proc Natl Acad Sci U S A.* 2012; 109: E3168–3176. doi: [10.1073/pnas.1210500109](https://doi.org/10.1073/pnas.1210500109) PMID: [23093667](https://pubmed.ncbi.nlm.nih.gov/23093667/)
45. Conway KL, Kuballa P, Song JH, Patel KK, Castoreno AB, Yilmaz OH, et al. Atg16l1 is required for autophagy in intestinal epithelial cells and protection of mice from *Salmonella* infection. *Gastroenterology.* 2013; 145: 1347–1357. doi: [10.1053/j.gastro.2013.08.035](https://doi.org/10.1053/j.gastro.2013.08.035) PMID: [23973919](https://pubmed.ncbi.nlm.nih.gov/23973919/)
46. Juncadella IJ, Kadl A, Sharma AK, Shim YM, Hochreiter-Hufford A, Borish L, et al. Apoptotic cell clearance by bronchial epithelial cells critically influences airway inflammation. *Nature.* 2013; 493: 547–551. doi: [10.1038/nature11714](https://doi.org/10.1038/nature11714) PMID: [23235830](https://pubmed.ncbi.nlm.nih.gov/23235830/)
47. Proud D, Leigh R. Epithelial cells and airway diseases. *Immunol Rev.* 2011; 242: 186–204. doi: [10.1111/j.1600-065X.2011.01033.x](https://doi.org/10.1111/j.1600-065X.2011.01033.x) PMID: [21682746](https://pubmed.ncbi.nlm.nih.gov/21682746/)
48. Gradstedt H, Iovino F, Bijlsma JJ. *Streptococcus pneumoniae* invades endothelial host cells via multiple pathways and is killed in a lysosome dependent manner. *PLoS One.* 2013; 8: e65626. doi: [10.1371/journal.pone.0065626](https://doi.org/10.1371/journal.pone.0065626) PMID: [23785439](https://pubmed.ncbi.nlm.nih.gov/23785439/)