

REVIEW SERIES

Chronic obstructive pulmonary disease • 6: The aetiology of exacerbations of chronic obstructive pulmonary disease

A J White, S Gompertz, R A Stockley

Thorax 2003;58:73–80

Exacerbations of COPD are thought to be caused by interactions between host factors, bacteria, viruses, and changes in air quality to produce increased inflammation in the lower airway. The evidence for this and the potential mechanisms by which they result in the characteristic symptoms of exacerbations is reviewed. A better understanding of the causes and processes is needed for the appropriate use of existing treatments and the development of new ones. Future studies need to define populations clearly, stratify for known confounding factors, and should aim to identify clinical correlates so that clinical practice can be modified appropriately.

adoption in research or clinical practice. However, recently a consensus statement defined exacerbation as “a sustained worsening of the patient’s condition, from the stable state and beyond normal day to day variations that is acute in onset and necessitates a change in regular medication in a patient with underlying COPD”.¹³ It is likely that this definition will become generally accepted in the future.

FACTORS DETERMINING SUSCEPTIBILITY TO EXACERBATIONS

Predictors for frequent exacerbations include various clinical factors such as daily cough, wheeze, and sputum production as well as frequency of exacerbations in the previous year.⁴ A prospective study of 5887 patients published recently showed that COPD smokers with chronic bronchitis had exacerbations 1.6–1.9 times more frequently than those without.⁶ This suggests that factors predisposing to chronic bronchitis also predispose to exacerbation, possibly due to reduced mucociliary clearance¹⁴ that facilitates bacterial invasion of mucosa.¹⁵ Indeed, excess production of mucus, which characterises chronic bronchitis, has been shown to provide a site for bacterial adherence¹⁶ and is associated with increased mortality from bacterial infection.¹⁷

Secretory leukoprotease inhibitor (SLPI) is the main proteinase inhibitor found in the larger airways and submucosal glands¹⁸ and also has antiviral and antibacterial properties.¹⁹ During the stable clinical state the level of SLPI in the sputum has been found to be significantly lower in patients with frequent exacerbations (three or more exacerbations in the preceding 12 months) than in those with infrequent exacerbations.²⁰ In addition, saliva and serum levels of lysozyme (a cationic protein with antibacterial properties with an important role in the first line of host defences) have been studied in patients with chronic bronchitis. Patients with two or more exacerbations over 12 months had significantly lower serum levels of lysozyme, lysozyme activity, and ability of saliva to aggregate non-typeable *Haemophilus influenzae* than those with less than two exacerbations.²¹ Although deficiencies in other host defence mechanisms may also predispose to infection, their role in COPD is yet to be determined.

The role of inflammatory mediators is less clear. Bhowmik *et al* found a relationship between stable state interleukin (IL)-8 and IL-6 concentrations in induced sputum samples and exacerbation frequency.²² However, there was no stratification for the presence or absence of confounding

The natural history of chronic obstructive pulmonary disease (COPD) is one of a progressive decline in ventilatory function, exercise capacity, and health status that is punctuated with varying frequency by exacerbations of symptoms.¹ Patients often consult their primary care physician at the time of exacerbation,² providing a contact early in the course of their disease, and as ventilatory function declines exacerbations lead to an increasing need for medical care and hospitalisation.^{1,3} Patients who suffer the most exacerbations have significantly lower health status⁴ and there is also evidence that exacerbation frequency predicts accelerated decline in lung function,^{5,6} although this remains controversial.^{1,7}

The heterogeneity of COPD is being appreciated increasingly,^{8,9} although this is rarely taken into account in studies of exacerbations which are heterogeneous in their own right. For this and other reasons, conclusions regarding the aetiology, as well as the inflammatory processes involved and their effects, remain unresolved.

DEFINITIONS

It is perhaps surprising that there is no consistently used definition of acute exacerbation of COPD (AECOPD), either in clinical practice or in research. In 1968 Fisher wrote “the value of future work on these lines would be increased if agreement could be reached on the definition of an exacerbation”¹⁰ but, unfortunately, inconsistencies continue. Although there have been attempts to gain consensus,^{11,12} these have been imprecise and have not resulted in universal

See end of article for authors’ affiliations

Correspondence to:
Professor R A Stockley
Department of Respiratory
Medicine, First Floor,
Nuffield House, Queen
Elizabeth Hospital,
Edgbaston, Birmingham
B15 2TH, UK;
r.a.stockley@bham.ac.uk

factors known to influence one or both of these including spontaneous sputum production, bacterial colonisation,²³ severity of airflow obstruction,²⁴ smoking status,^{25, 26} bronchiectasis,²⁷ or treatment with inhaled steroids, all of which may influence the inflammatory status. On the other hand, Gompertz *et al* found no association between IL-8 levels and exacerbation frequency when all these factors were taken into account,²⁰ which suggests that the high IL-8 concentrations observed by Bhowmik *et al*²² were reflecting another risk factor for exacerbations. However, the observation that plasma fibrinogen levels are also associated with increased rates of hospital admission with exacerbation, as well as with reduced lung function and risk of COPD,²⁸ may support a role for IL-6 which is the primary cytokine regulating fibrinogen expression.

CHARACTERISATION OF COPD EXACERBATIONS

The most widely quoted clinical criteria used in the characterisation of AECOPD are those described by Anthonisen *et al* in what remains the best placebo controlled trial of antibiotics for such episodes.²⁹ In that study exacerbations were divided into three groups: type 1 exacerbations were characterised by increased breathlessness, increased sputum volume, and new or increased sputum purulence; type 2 included any two of these symptoms; and type 3 consisted of any one of the symptoms together with at least one additional feature, including sore throat or nasal discharge within the last five days; unexplained fever; increased wheeze; increased cough; or a 20% increase in respiratory or heart rate compared with baseline. These criteria have been used as a benchmark ever since, and all proposed aetiologies of exacerbation need to establish their relationship to these key features.

PATHOPHYSIOLOGY OF EXACERBATIONS OF COPD

The stable clinical state is characterised by varying degrees of inflammation affecting the large and small airways as well as the alveoli, resulting in mucus hypersecretion, airway narrowing and alveolar destruction, respectively. The main mediators directly responsible for this damage are proteinases released by inflammatory cells, particularly neutrophils, that are found in abundance in the bronchial secretions of patients with COPD, especially as the forced expiratory volume in 1 second (FEV₁) declines.^{24, 26, 30, 31} The neutrophils are recruited from the circulation in response to chemoattractants, particularly IL-8 and LTB₄ via adhesion molecules (such as E selectin and ICAM-1) expressed on the vascular endothelium. These receptors are thought to be upregulated in COPD³⁰ and bind to their counterparts on the neutrophil (such as MAC-1) that are also upregulated.³² The activated neutrophils degranulate releasing neutrophil elastase and other proteinases (cathepsin G, proteinase 3, and matrix metalloproteinases) that are thought to be responsible for various aspects of tissue damage, as well as myeloperoxidase.¹⁸ This latter protein has a green colouration that is useful for monitoring neutrophil influx clinically (see below).³³ A proportion of patients also have increased numbers of bronchial eosinophils which may contribute to inflammation.³⁴⁻³⁶

With this background of mild inflammation there is a general belief that exacerbations are episodes where the inflammatory process is enhanced, although the processes involved and their effects are poorly understood. However, over the past two years in particular there has been an increased interest in the mechanisms involved in exacerbations of COPD,^{2, 20, 22, 37-44} although interpretation and comparison between studies has been hampered by loose definitions, differences in methodology, study populations, and patient characterisation.

Understanding the aetiology of exacerbations requires that potential mechanisms of these episodes are related to the key

Box 1 Relationship of exacerbation symptoms to potential underlying mechanisms

(1) Increased breathlessness

- Increased metabolic/catabolic state
- Airway narrowing
 - Mucosal damage, increased irritability/bronchospasm
 - Inflammatory cell infiltration into airway tissues
 - Airway oedema
 - Increased airway secretion and viscosity/mucus plugging
- Increased ventilation/perfusion (V/Q) mismatch

(2) Increased sputum production

- Mucous gland hypertrophy
- Goblet cell hyperplasia
- Goblet cell degranulation

(3) New or increased sputum purulence

- Eosinophil recruitment
- Neutrophil recruitment

symptoms outlined by Anthonisen *et al*.²⁹ The three main symptoms and how they might arise are outlined in box 1. The evidence for the relevant processes occurring during exacerbations is reviewed.

Increased metabolic/catabolic state

A systemic inflammatory response is capable of causing increased breathlessness without changes in respiratory tract pathology. The increased metabolic state associated with pyrexia increases oxygen demand and may increase hydrogen ion concentrations stimulating the respiratory centre to increase tidal volume and respiratory rate.⁴⁵ In those with normal lungs this effect may go unnoticed, but in patients with COPD the increase in ventilatory drive may lead to an increased sensation of breathlessness. Undoubtedly, a systemic effect is present during exacerbations as indicated by the rise in C reactive protein (CRP)² and IL-6.²² Whether treatment to reduce the metabolic drive (such as antipyretics) improves the sensation of dyspnoea during these episodes has yet to be proved.

Airway narrowing

Several potential mechanisms of exacerbations may exert their influence by reducing the calibre of the airways. Such a process would limit airflow further, result in increased residual volume, and hence increase the work of breathing.

In a longitudinal study of patients with moderate to severe COPD, patients recorded symptoms and peak expiratory flow rates (PEFR) in a daily diary and some also measured FEV₁ and forced vital capacity (FVC) using a handheld spirometer.⁴⁶ Significant falls in PEFR, FEV₁, and FVC were observed during the exacerbations. The recovery time was related to the magnitude of the fall, and the reduction in PEFR was greater in patients with increased dyspnoea. These findings are supported by intervention studies that have used FEV₁ as an outcome measure and demonstrated significant increases in PEFR and/or FEV₁ on recovery from exacerbation,⁴⁷⁻⁴⁹ and bronchodilator treatment has been shown to speed the resolution of airflow obstruction.⁵⁰⁻⁵⁴

One potential mechanism for this change in airflow would be bronchospasm which could be mediated by endothelin-1. In addition to promoting bronchial hyperresponsiveness, endothelin-1 may stimulate mucus secretion, promote airway oedema and smooth muscle proliferation, as well as upregulate production of cytokines. The concentration of this peptide, which is produced by the bronchial epithelium, alveolar macrophages and pulmonary endothelium, is increased in sputum

in stable COPD⁵⁵ and a recent study has shown that concentrations of endothelin-1 are increased in induced sputum during AECOPD, suggesting it may play an important role.⁴¹

Infiltration of the airway wall with inflammatory cells could also contribute to airflow limitation. This has been described extensively in stable COPD where increased numbers of CD8+ lymphocytes^{56, 57} and neutrophils²⁴ are found. During exacerbation, biopsy studies have shown that the inflammation becomes more marked with recruitment of eosinophils (see below) and an increase in the population of CD4+ lymphocytes.⁴⁴ Furthermore, sputum studies have shown increased neutrophil content and increased protein leakage from serum to sputum which also occurs during mucoid exacerbations that may lead to oedema of the airway wall.³⁹

An increase in mucus production would lead to an increase in sputum production that characterises many AECOPD. The presence of sputum in the airways would be expected to reduce the airway calibre and this effect would be enhanced if the viscosity of the sputum also increased. Such secretions would be harder to clear and result in plugging of smaller airways and hence increased breathlessness. There are numerous potential mechanisms which could cause increased sputum production and these are outlined below. Support for this feature comes from studies of mucolytic drugs that have been shown to reduce exacerbation frequency by 29%⁵⁸ and speed up resolution of symptoms when associated with bacteria.⁵⁹

Changes in ventilation/perfusion (V/Q) matching

The relationship of ventilation to perfusion has been studied in patients admitted to hospital with AECOPD using a multiple inert gas elimination technique.^{60, 61} Gas exchange deteriorates during exacerbations through worsening V/Q matching. Proportionately more blood perfuses poorly ventilated lung units than in the stable clinical state, and the resulting reduction in arterial oxygen tension is compounded by increased oxygen consumption. The relationship of these changes to airway wall oedema and mucus plugging has yet to be determined.

Eosinophilic inflammation

A potential mechanism which may explain many of the features of AECOPD is eosinophil recruitment. Biopsy studies have shown 30 fold higher numbers of eosinophils in bronchial mucosa during exacerbations⁶² and serum levels of eosinophilic cationic protein are higher in patients with exacerbations than in those with stable COPD.⁶³ Although this suggests an "asthmatic profile", the observed eosinophils are not degranulated (as they would be in asthma) and are not associated with increased IL-5 expression.⁶⁴ A recent study suggests that the chemokine associated with the influx of eosinophils is RANTES, which is secreted by both epithelial cells and the subepithelial lymphocytes.⁴⁴ The mechanism may be indirect and relate to increased TNF- α positive cells⁶² and sputum concentrations^{37, 42} during exacerbations. TNF- α increases the expression of RANTES, providing a potential mechanism for eosinophil recruitment.⁴⁴ The results suggest that at least a subset of patients experience eosinophilic exacerbations of bronchitis. The relative importance of the eosinophilia remains to be determined, but several eosinophil products may cause inflammatory damage to the airway (eosinophil peroxidase, major basic protein, eosinophil cationic protein, metalloproteinases, platelet activating factor, and cysteinyl leukotrienes)⁶⁵ and, together with histamine, can also cause bronchospasm. The presence of eosinophils in airway secretions can contribute to sputum purulence which is a feature of some exacerbations.

In stable COPD bronchial eosinophilia has been shown to be a marker for steroid responsiveness.^{66, 67} Thus, the eosinophilia associated with some exacerbations may explain the effects of

long term inhaled steroids on reducing exacerbations in COPD⁶⁸ and the beneficial effect of steroids seen in some exacerbations.^{47, 49}

Neutrophilic inflammation

A major finding in airway secretions during exacerbations is an increase in neutrophils⁶⁹ that is also associated with the presence or change in sputum purulence.³³ Neutrophil degranulation results in release of elastase and other proteinases which may cause epithelial damage, reduce ciliary beat frequency,⁷⁰ stimulate mucus secretion by goblet cells,⁷¹ and increase the permeability of the bronchial mucosa resulting in airway oedema and protein exudation into the airway.³⁹ These changes, especially in the small airways, may adversely affect airflow leading to increased breathlessness, as well as the mucus secretion and purulence characteristic of some exacerbations. The cytokine most associated with the neutrophilic inflammatory changes seen at exacerbation is LTB₄³⁹ which increases, whereas IL-8 (the other cytokine regularly associated with chronic bronchitis) is unchanged^{22, 39} except in the most severe episodes.³⁸

The increase in TNF- α in the sputum during exacerbations^{37, 42} may increase expression of endothelial adhesion molecules facilitating cell migration as well as activating neutrophils directly.²⁶ Furthermore, granulocyte/macrophage colony stimulating factor (GM-CSF) is increased in bronchoalveolar lavage fluid during exacerbations.⁶⁹ This cytokine stimulates differentiation of granulocytes and macrophages and can activate them directly,⁶⁹ providing another mechanism whereby neutrophils—as well as eosinophils and macrophages—can contribute to inflammatory changes within the airways. Studies of bronchial neutrophilic inflammation during exacerbations show resolution usually within 5 days following treatment, and this parallels clinical recovery.³⁹ Figure 1 outlines the changes which occur in neutrophilic inflammation during AECOPD.

ROLE OF BACTERIA IN EXACERBATIONS OF COPD

The association of bacteria with COPD has been appreciated for many years.⁷² Bacteria are frequently isolated from the sputum of patients when stable, and recent studies using the protected brush to obtain lower airway samples have shown that approximately one third of patients are colonised at any one time.^{73–76} *Haemophilus influenzae* and *Streptococcus pneumoniae* represent the majority of the organisms, although *Moraxella catarrhalis*, *H parainfluenzae*, Gram negative enteric bacilli, and *Pseudomonas* spp also account for a proportion. The role of bacteria in disease progression has been investigated extensively but remains relatively unclear.

There are a number of potential mechanisms by which bacteria may affect symptoms in COPD. Strains of *H influenzae*, *P aeruginosa*, and *S pneumoniae* have been shown to stimulate mucus hypersecretion in vitro,⁷⁷ and the first two organisms have been shown to inhibit ciliary beat frequency.⁷⁸ Furthermore, *H influenzae* can cause direct epithelial damage⁷⁹ and its endotoxin has been shown to increase epithelial expression of the pro-inflammatory cytokines IL-6, IL-8, and TNF- α in vitro,⁸⁰ providing potential mechanisms to upregulate inflammation. In support of this concept, patients with chronic bronchitis and COPD with a positive bacterial culture of *H influenzae* have higher concentrations of TNF- α in their sputum.²³ In a study of patients with stable chronic bronchitis, airway bacterial load has been shown to correlate directly with markers of neutrophilic inflammation, irrespective of the pathogen isolated.⁸¹ Similarly, the presence of potentially pathogenic microbes isolated from bronchoalveolar lavage fluid is strongly associated with increased neutrophils and concentrations of TNF- α .⁷⁶ Eradication or reduction of bacteria is associated with resolution of inflammation.




Figure 1 Neutrophilic inflammation during exacerbations of COPD. (A) Phagocytosis/macrophage activation (release of LTb4 and TNF α). (B) Epithelial activation (in response to TNF α and bacterial products, release of TNF α and IL-8). (C) Activation of endothelium (due to TNF α , increase in adhesion molecules). (D) Neutrophil migration (follows from binding to adhesion molecules and IL-8/LTB4 chemotactic gradient, elastase release and tissue damage). (E) Neutrophil elastase release in airway (epithelial damage and protein leakage; increased mucus secretion; reduced mucociliary clearance; associated release of IL-8 and LTb4 resulting in amplification of chemotactic signal). LTb4=leukotriene B4; IL-8=interleukin 8; TNF α =tumour necrosis factor α ; ICAM-1=intercellular adhesion molecule 1.

The background of colonisation and inflammation in the stable state has led to uncertainty concerning the role of bacteria during exacerbations. Some authors suggest that there is no compelling evidence that bacteria play a role in acute exacerbations.⁸² The majority of microbiological studies aimed at establishing the role of bacteria have used cultures of spontaneously produced sputum and compared the exacerbation with the stable clinical state. Because the same organisms are frequently found both during an exacerbation and in the stable clinical state in the same patients, interpretation of their role can be difficult. Nevertheless, the proportion of patients with positive bacterial cultures and a high bacterial load increases during exacerbations in most,^{2, 10, 83} although not in all studies.⁸⁴ Newer molecular techniques have recently shown that colonisation is not stable and there is frequent turnover of discrete strains of *H influenzae* that elicit specific host responses to outer membrane proteins.⁸⁵ It is therefore likely that a change in the strain but not the organism may be responsible for the symptoms of an exacerbation, and studies which have failed to discriminate between different bacterial strains³⁷ may miss evidence of a new infection.

To refine our understanding, several studies have used the protected specimen brush method to derive more information about the bacteriology of the lower airways during exacerbations. Monso *et al* compared patients with an exacerbation to matched patients in the stable clinical state and showed marked differences, both in pathogen recovery (52% *v* 25% in the stable group) and bacterial load (24% of those with an exacerbation having a concentration of bacteria in excess of 10 000 cfu/ml *v* 5% of the stable group).⁷⁴ Other bronchoscopic studies have found a similar proportion of exacerbations to be associated with bacteria.^{86, 87} The most common organisms isolated are *H influenzae*, *S pneumoniae*, and *M catarrhalis*.⁷⁴ However, studies have shown that exacerbations in more severe patients are more likely to be associated with recovery of Gram negative enterobacteriae and *Pseudomonas* spp.⁸⁷⁻⁸⁹ Although these data suggest a relationship with decline in lung function, the prior use of antibiotics may have influenced some of the results.

Further evidence to support the role of bacteria in AECOPD relate to studies of the immune response. Strain specific antibodies to surface outer membrane proteins of *H influenzae* have been shown to develop in relation to exacerbations.⁹⁰ Similarly, patients who experience an exacerbation of COPD with a

positive sputum culture for *M catarrhalis* have been shown to mount an antibody response to the homologous strain,⁹¹ and this has also recently been shown for *H parainfluenzae*.⁹² This, together with sputum and protected brush studies, provides compelling evidence that bacteria cause a proportion of exacerbations of COPD.

The inconclusive results of placebo controlled trials of antibiotics for exacerbations raises some further concerns about the importance of bacteria. Some studies have failed to show a significant difference in the outcome of patients treated with or without antibiotics.^{93, 94} However, when Anthonisen and co-workers studied a larger numbers of patients characterised by specific symptoms it was possible to identify a subgroup of patients who benefited from antibiotics—namely, those with all three symptoms (see above).²⁹ Furthermore, meta-analysis indicates that antibiotics have a positive effect.⁹⁵ Because bacterial infections are mucosal and some are likely to resolve spontaneously, the difference between antibiotic and placebo treatment may be difficult to detect, particularly in exacerbations that have not been characterised and when a proportion will not be bacterial in origin. When relapse rate was used as an outcome measure, a recent retrospective study indicated that patients treated with antibiotics had significantly lower rates of relapse than those not receiving them,⁹⁶ again supporting a role for bacteria in at least some episodes.

As already emphasised, exacerbations are not a single entity and recent data indicate that use of a simple sputum colour chart to divide episodes into those with mucoid or purulent sputum identifies a major difference in bacterial isolation from spontaneous sputum samples.² A positive culture was obtained from 84% of sputum samples from patients with a purulent exacerbation compared with 38% from mucoid exacerbations. In the stable clinical state the proportion of patients with a positive culture was similar in the two groups (approximately 35%). The presence of green sputum was 94.4% sensitive and 77% specific for the yield of a high bacterial load, identifying a distinct subset of patients in whom bacteria are strongly associated with the exacerbation. Resolution of sputum purulence was shown to relate to resolution of symptoms.⁹⁷ Conversely, there was little evidence of a bacterial aetiology for exacerbations associated with mucoid sputum and these patients improved without antibiotics. The mixed nature of exacerbations would therefore minimise the benefit of antibiotics unless stratification is undertaken.




Figure 2 Potential mechanisms of viral induced exacerbations of COPD. M2=type 2 muscarinic receptors; RANTES=regulated on activation, normal T cell expressed and secreted; PAF=platelet activating factor; MBP=major basic protein; EPO=eosinophil peroxidase; ECP=eosinophil cationic protein.

ROLE OF VIRUSES IN EXACERBATIONS OF COPD

Lower respiratory tract viral infections in patients with COPD may also cause direct damage to the airway epithelium resulting in loss of ciliated epithelium, increased mucus production, sloughing of necrotic cells into the airway lumen, together with increased plasma exudation.⁹⁸ In addition to causing airway narrowing,⁴⁸ the altered environment may promote secondary bacterial infection with the further potential to promote inflammatory changes as outlined above. Viral infections have been shown to increase airway responsiveness and this may be due to a change in the balance of bronchoconstricting and bronchodilating substances in the airway.^{98, 99} Indeed, viral infections have been shown to increase levels of endothelin-1, precipitating acute episodes of asthma, and endothelin-1 also increases during exacerbations of COPD.⁴¹ Furthermore, viruses can also increase release of acetylcholine by direct and indirect effects upon M2 muscarinic receptors leading to increased bronchoconstriction.¹⁰⁰ Thus, there are several mechanisms by which viruses can cause AECOPD and these are outlined in fig 2.

Several pathological findings in COPD provide indirect evidence for the role of viruses in AECOPD. Eosinophilia has been an important feature in biopsies and it has been suggested that this may be via the increased expression of RANTES,⁴⁴ which might be a direct effect of the virus on the epithelial cell. All of the inflammatory processes, together with the systemic inflammatory response associated with viral infections, would be expected to increase the symptoms of breathlessness in COPD.

Direct evidence of viral infection is, however, necessary to establish the contribution of viruses to exacerbations. Table 1 summarises some of the more important studies of the viral aetiology of exacerbations of COPD. Most of the evidence is indirect and depends on serological assessment.^{84, 101-105} Seroconversion to specific viruses is not consistently associated with clinical deterioration and therefore seroconversion at the time of an exacerbation does not provide incontrovertible evidence of aetiology. Studies conducted over short periods can be influenced by the effect of epidemics and probably account for the variation in some of the viruses implicated.

Culture of rhinovirus has been undertaken in several studies with varying results. The largest was a longitudinal study of 150 patients over 8 years and positive culture of rhinovirus was associated with 4.8% of 1030 exacerbation episodes compared with 0.5% during illness free periods.¹⁰⁴ This finding has been supported by other shorter studies,¹⁰¹ although isolation of the virus has been achieved in up to 23% of episodes using induced sputum and nasal/throat samples assessed by polymerase chain reaction (PCR) which increases the sensitivity of rhinovirus detection.¹⁰⁶

As suggested previously, viral infection may have a role in predisposing to secondary bacterial infection in patients susceptible because of their underlying lung disease. One study specifically examined the relationship between viral infection and bacteria in COPD; viral infections were found to be associated with increased isolation of *H influenzae* and *S pneumoniae* from sputum cultures.¹⁰⁷ Furthermore, there was a higher incidence of *H influenzae* antibody seroconversion in the virus infected group, suggesting that this is a potential sequence of events.

Despite the research that has been published, the contribution of viruses to AECOPD remains to be determined and probably varies considerably with epidemics and immunisation policy. Undoubtedly, there is evidence that a proportion of episodes are associated with viral infection, particularly influenza viruses, as vaccination has shown benefit.¹⁰⁸ However, viral infections of the respiratory tract are a common cause of morbidity in otherwise healthy individuals, and whether or not patients with COPD are more susceptible to viral infections remains controversial. Some studies have shown significantly more viral infections in patients with COPD,^{102, 109} while others have found no difference from healthy subjects.^{104, 110}

ROLE OF ATYPICAL ORGANISMS IN EXACERBATIONS OF COPD

Mycoplasma pneumoniae and *Chlamydia* spp are intracellular bacteria that share some of the characteristics of viruses. The mechanism by which they cause exacerbations is likely to be similar to viral infections and, likewise, the investigations into the role of these organisms have depended largely on serological diagnosis.^{101, 104, 111-113} Mycoplasma infections predominantly affect young adults and immunity is achieved by

Table 1 Summary of main findings from viral studies

	No of exacerbations	Rhinovirus		Influenza virus		Parainfluenza virus		RSV		Adenovirus	
		Ex* (%)	Stable† (%)	Ex (%)	Stable (%)	Ex (%)	Stable (%)	Ex (%)	Stable (%)	Ex (%)	Stable (%)
Smith ¹⁰⁴	1030	4.8	0.5	4.9	0.2	2.8	0.7	0.8	0.6	0.7	0.5
Seemungal ¹⁰⁶	43	23.3	0								
Buscho ¹⁰⁶	166	2.7	0.55	10.2	6.4	6	7	0	0.7	1.2	0.7
Lamy ¹⁰³	49			28.6	1	24.5	0.5	6	1	0	0.75
Greenberg ¹¹⁰	323	4.9		2.5		6.2		2.5		0.3	
Stenhouse ¹⁰⁵	64			9.4	1.5	0	0	3.1	0		
Gump ⁸⁴	116			12.9	0.2	7.8	0.2	4.3	0.05	2.6	0

*Viral infection determined from seroconversion PCR or viral isolation in the exacerbated state.

†Viral infection determined from seroconversion PCR or viral isolation in the stable clinical state.

The figures in the table indicate the proportion of patients in the stable and exacerbated state from which each type of virus was detected or seroconversion occurred.

the time most individuals reach the age where AECOPD becomes a clinical problem. It is therefore not surprising that most studies have found that few exacerbations are associated with evidence implicating this organism,^{101–104} although a recent study implicated *M pneumoniae* in as many as 14% of exacerbations.¹¹⁴ There has, however, been more debate regarding the role of *Chlamydia pneumoniae*. Two studies of acute exacerbations (one in hospitalised patients and one in outpatients) found *C pneumoniae* in approximately 5% of episodes, and serological evidence suggesting previous infection in approximately three quarters of patients (similar to a matched population).^{111–113} However, two other recent studies have found *C pneumoniae* to be associated with 24% and 34% of exacerbations.^{112–115} The implications of these findings remain uncertain and further studies are required to resolve these issues.

AIR POLLUTION

It has been suggested that AECOPD could be induced by increases in air pollution. The effects of diesel particulates, sulphur dioxide (SO₂), ozone (O₃), and nitrogen dioxide (NO₂) have been studied and potential mechanisms by which airway inflammation is enhanced have been proposed. Recent bronchoscopic studies have shown that exposure of healthy volunteers to diesel exhaust results in increases in neutrophils^{116–117} and methylnhistamine.¹¹⁶ In vitro studies have shown that diesel exhaust particles simulated production of pro-inflammatory cytokines such as GM-CSF and IL-8, both of which may explain the enhanced neutrophilic inflammation.¹¹⁸ Ozone exposure has been shown to be associated with markers of nasal eosinophil inflammation in non-atopic children,¹¹⁹ and SO₂ and NO₂ have been shown to enhance the airway response to inhaled allergens.¹²⁰ Thus, potential mechanisms exist whereby changes in air pollution can cause exacerbations of respiratory symptoms in COPD.

Evidence to support a role for air pollution has been based on epidemiological studies that have implicated increases of SO₂, NO₂, PM₁₀, and black smoke particulate matter in changes in chronic respiratory symptoms and increased respiratory mortality in patients with COPD.^{121–126} These city based studies have investigated excess hospital admissions at times of increased atmospheric pollution and concluded that this may account for approximately 6–9% of admissions, depending on the time of year.¹²⁵ Patients with COPD have also been shown to be at increased risk of death associated with urban particle air pollution, although much of this may be cardiovascular rather than respiratory.¹²⁶

CONCLUSIONS

The evidence suggests that bacteria, viruses, and changes in air quality interact with host factors and with each other to produce increased inflammation in the lower airway. This leads to the development of the characteristic symptoms associated with exacerbations of COPD. The appreciation that exacerbations have several causes has not resulted in a change of clinical practice^{127–128} and, in particular, antibiotics remain overused, contributing to global antibiotic resistance. Patients are usually treated empirically with increased β₂ agonists, anticholinergic agents, oral steroids, and antibiotics. The appropriate use of these treatments and the development of new ones can only be established once a better understanding of the nature and causes of exacerbations has been achieved. Future studies need to define populations clearly, stratify for known confounding factors, and should aim to identify clinical correlates so that clinical practice can be modified appropriately.

Authors' affiliations

A J White, S Gompertz, R A Stockley, Department of Respiratory Medicine, Queen Elizabeth Hospital, Edgbaston, Birmingham B15 2TH, UK

REFERENCES

- 1 Fletcher C, Peto R. The natural history of chronic airflow obstruction. *BMJ* 1977;1:1645–8.
- 2 Stockley RA, O'Brien C, Pye A, et al. Relationship of sputum colour to nature and outpatient management of acute exacerbations of COPD. *Chest* 2001;117:1638–45.
- 3 Garcia-Aymerich J, Monso E, Marrades M, et al. Risk factors for hospitalization for a chronic obstructive pulmonary disease exacerbation: Efram study. *Am J Respir Crit Care Med* 2001;164:1002–7.
- 4 Seemungal TA, Donaldson GC, Paul EA, et al. Effect of exacerbation on quality of life in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;157:1418–22.
- 5 Kanner RE, Renzetti AD Jr, Klauber MR, et al. Variables associated with changes in spirometry in patients with obstructive lung diseases. *Am J Med* 1979;67:44–50.
- 6 Kanner RE, Anthonisen NR, Connett JE. Lower respiratory illnesses promote FEV₁ decline in current smokers but not ex-smokers with mild chronic obstructive pulmonary disease. Results from the lung health study. *Am J Respir Crit Care Med* 2001;164:358–64.
- 7 Burrows B, Bloom JW, Traver GA, et al. The course and prognosis of different forms of chronic airways obstruction in a sample from the general population. *N Engl J Med* 1987;317:1309–14.
- 8 O'Brien C, Guest PJ, Hill SL, et al. Physiological and radiological characterisation of patients diagnosed with chronic obstructive pulmonary disease in primary care. *Thorax* 2000;55:635–42.
- 9 Wedzicha JA. The heterogeneity of chronic obstructive pulmonary disease. *Thorax* 2000;55:631–2.
- 10 Fisher M, Akhtar AJ, Calder MA, et al. Pilot study of factors associated with exacerbations in chronic bronchitis. *BMJ* 1969;4:187–92.
- 11 COPD Guidelines Group of the Standards of Care Committee of the British Thoracic Society. BTS guidelines for the management of chronic obstructive pulmonary disease. *Thorax* 1997;52(Suppl 5):S1–28.
- 12 Pauwels R, Buist AS, Calverley PM, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: NHLBI/WHO global initiative for chronic obstructive lung disease (GOLD) workshop summary. *Am J Respir Crit Care Med* 2001;163:1256–76.
- 13 Rodriguez-Roisin R. Towards a consensus definition for COPD exacerbations. *Chest* 2000;117:398–401S.
- 14 Mossberg B, Afzelius B, Camner P. Mucociliary clearance in obstructive lung diseases. Correlations to the immotile cilia syndrome. *Eur J Respir Dis Suppl* 1986;146:295–301.
- 15 Wilson R, Roberts D, Cole P. Effects of bacterial products on human ciliary function in vitro. *Thorax* 1984;40:125–31.
- 16 Kubiet M, Ramphal R. Adhesion of nontypeable Haemophilus influenzae from blood and sputum to human tracheobronchial mucins and lactoferrin. *Infect Immun* 1995;63:899–902.
- 17 Prescott E, Lange P, Vestbo J. Chronic mucus hypersecretion in COPD and death from pulmonary infection. *Eur Respir J* 1995;9:1333–8.
- 18 Stockley RA. Cellular mechanisms in the pathogenesis of COPD. *Eur Respir J* 1996;9:264–9.
- 19 Tomee JFC, Koeter GH, Hiemstra PS, et al. Secretory leukoprotease inhibitor: a native antimicrobial protein presenting a new therapeutic option? *Thorax* 1998;53:114–6.
- 20 Gompertz S, Bayley DL, Hill SL, et al. Relationship between airway inflammation and the frequency of exacerbations in patients with smoking related COPD. *Thorax* 2001;56:36–41.
- 21 Taylor DC, Cripps AW, Clancy RL. A possible role for lysozyme in determining acute exacerbation in chronic bronchitis. *Clin Exp Immunol* 1995;102:406–16.
- 22 Bhowmik A, Seemungal TA, Sapsford RJ, et al. Relation of sputum inflammatory markers to symptoms and lung function changes in COPD exacerbations. *Thorax* 2000;55:114–20.
- 23 Bresser P, Out TA, van Alphen L, et al. Airway inflammation in nonobstructive and obstructive chronic bronchitis with chronic haemophilus influenzae airway infection. Comparison with noninfected patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2000;162:947–52.
- 24 Di Stefano A, Capelli A, Lusuardi M, et al. Severity of airflow limitation is associated with severity of airway inflammation in smokers. *Am J Respir Crit Care Med* 1998;158:1277–85.
- 25 Hill AT, Bayley D, Campbell EJ, et al. Airways inflammation in chronic bronchitis: the effects of smoking and alpha-1 antitrypsin deficiency. *Eur Respir J* 2000;15:886–90.
- 26 Keatings VM, Collins PD, Scott DM, et al. Differences in interleukin-8 and tumor necrosis factor-alpha in induced sputum from patients with chronic obstructive pulmonary disease or asthma. *Am J Respir Crit Care Med* 1996;153:530–4.
- 27 Gompertz S, O'Brien C, Bayley D, et al. Comparison of stable state bronchial inflammation in chronic bronchitis and previously unsuspected bronchiectasis. *Eur Respir J* 1999;14:165S.
- 28 Dahl M, Tybjaerg-Hansen A, Vestbo J, et al. Elevated plasma fibrinogen associated with reduced pulmonary function and increased risk of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;164:1008–11.

- 29 **Anthonisen NR**, Manfreda J, Warren CP, *et al*. Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;**106**:196–204.
- 30 **Stanescu D**, Sanna A, Veriter C, *et al*. Airways obstruction, chronic expectoration, and rapid decline of FEV1 in smokers are associated with increased levels of sputum neutrophils. *Thorax* 1996;**51**:267–71.
- 31 **Thompson AB**, Daughton D, Robbins RA, *et al*. Intraluminal airway inflammation in chronic bronchitis: characterisation and correlation with clinical parameters. *Am Rev Respir Dis* 1989;**140**:1527–37.
- 32 **Noguera A**, Batle S, Miralles C, *et al*. Enhanced neutrophil response in chronic obstructive pulmonary disease. *Thorax* 2001;**56**:432–7.
- 33 **Stockley RA**, Bayley D, Hill SL, *et al*. Assessment of airway neutrophils by sputum colour: correlation with airways inflammation. *Thorax* 2001;**56**:366–72.
- 34 **Lacoste J-Y**, Bousquet J, Chanez P, *et al*. Eosinophilic and neutrophilic inflammation in asthma, chronic bronchitis, and chronic obstructive pulmonary disease. *J Allergy Clin Immunol* 1993;**92**:537–48.
- 35 **O'Connell JM**, Baird LI, Campbell AH. Sputum eosinophilia in chronic bronchitis and asthma. *Respiration* 1978;**35**:65–72.
- 36 **Pesci A**, Balbi B, Majori M, *et al*. Inflammatory cells and mediators in bronchial lavage of patients with chronic obstructive pulmonary disease. *Eur Respir J* 1998;**12**:380–6.
- 37 **Aaron S**, Angel JB, Lunau M, *et al*. Granulocyte inflammatory markers and airway infection during acute exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;**163**:349–55.
- 38 **Crooks SW**, Bayley DL, Hill SL, *et al*. Bronchial inflammation in acute bacterial exacerbations of chronic bronchitis: the role of leukotriene B4. *Eur Respir J* 2000;**15**:274–80.
- 39 **Gompertz S**, O'Brien C, Bayley D, *et al*. Changes in bronchial inflammation during acute exacerbations of chronic bronchitis. *Eur Respir J* 2001;**17**:1112–9.
- 40 **Hill AT**, Campbell EJ, Bayley DL, *et al*. Evidence for excessive bronchial inflammation during an acute exacerbation of chronic obstructive pulmonary disease in patients with alpha₁-antitrypsin deficiency (P1Z). *Am J Respir Crit Care Med* 1999;**160**:1968–75.
- 41 **Roland M**, Bhowmik A, Sapsford RJ, *et al*. Sputum and plasma endothelin-1 levels in exacerbations of chronic obstructive pulmonary disease. *Thorax* 2001;**56**:30–5.
- 42 **Sethi S**, Muscarella BS, Evans N, *et al*. Airway inflammation and etiology of acute exacerbations of chronic bronchitis. *Chest* 2000;**118**:1557–65.
- 43 **Wedzicha JA**, Seemungal TA, MacCallum PK, *et al*. Acute exacerbations of chronic obstructive pulmonary disease are accompanied by elevations of plasma fibrinogen and serum IL-6 levels. *Thromb Haemost* 2000;**84**:210–5.
- 44 **Zhu J**, Qiu YS, Majumdar S, *et al*. Exacerbations of Bronchitis: bronchial eosinophilia and gene expression for interleukin-4, interleukin-5, and eosinophil chemoattractants. *Am J Respir Crit Care Med* 2001;**164**:109–16.
- 45 **Bone RC**. The sepsis syndrome. Definition and general approach to management. *Clin Chest Med* 1996;**17**:175–81.
- 46 **Seemungal TA**, Donaldson GC, Bhowmik A, *et al*. Time course and recovery of exacerbations in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2000;**161**:1608–13.
- 47 **Davies L**, Angus RM, Calverley PM. Oral corticosteroids in patients admitted to hospital with exacerbations of chronic obstructive pulmonary disease: a prospective randomised controlled trial. *Lancet* 1999;**354**:456–60.
- 48 **Smith CB**, Kanner RE, Golden CA, *et al*. Effect of viral infections on pulmonary function in patients with chronic obstructive pulmonary diseases. *J Infect Dis* 1980;**141**:271–80.
- 49 **Thompson WH**, Nielson CP, Carvalho P, *et al*. Controlled trial of oral prednisone in outpatients with acute COPD exacerbation. *Am J Respir Crit Care Med* 1996;**154**:407–12.
- 50 **Fernandez A**, Lazaro A, Garcia A, *et al*. Bronchodilators in patients with chronic obstructive pulmonary disease on mechanical ventilation. Utilization of metered-dose inhalers. *Am Rev Respir Dis* 1990;**141**:164–8.
- 51 **Lloberes P**, Ramis L, Montserrat JM, *et al*. Effect of three different bronchodilators during an exacerbation of chronic obstructive pulmonary disease. *Eur Respir J* 1988;**1**:536–9.
- 52 **McCroory DC**, Brown CD. Inhaled short-acting beta₂-agonists versus ipratropium for acute exacerbations of chronic obstructive pulmonary disease (Cochrane Review). *Cochrane Database Syst Rev* 2001;**2**:CD002984.
- 53 **Moayyedi P**, Congleton J, Page RL, *et al*. Comparison of nebulised salbutamol and ipratropium bromide with salbutamol alone in the treatment of chronic obstructive pulmonary disease. *Thorax* 1995;**50**:834–7.
- 54 **Patrick DM**, Dales RE, Stark RM, *et al*. Severe exacerbations of COPD and asthma. Incremental benefit of adding ipratropium to usual therapy. *Chest* 1990;**98**:295–7.
- 55 **Chalmers GW**, Macleod KJ, Sriiriam S, *et al*. Sputum endothelin-1 is increased in cystic fibrosis and chronic obstructive pulmonary disease. *Eur Respir J* 1999;**13**:1288–92.
- 56 **O'Shaughnessy TC**, Ansari TW, Barnes NC, *et al*. Inflammation in bronchial biopsies of subjects with chronic bronchitis: inverse relationship of CD8+ T lymphocytes with FEV1. *Am J Respir Crit Care Med* 1997;**155**:852–7.
- 57 **Saetta M**, Di Stefano A, Turato G, *et al*. CD8+ T-lymphocytes in the peripheral airways of smokers with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;**157**:822–6.
- 58 **Poole PJ**, Black PN. Oral mucolytic drugs for exacerbations of chronic obstructive pulmonary disease: systematic review. *BMJ* 2001;**322**:1271–4.
- 59 **Marchioni CF**, Polu JM, Taytard A, *et al*. Evaluation of efficacy and safety of erdosteine in patients affected by chronic bronchitis during an infective exacerbation phase and receiving amoxicillin as basic treatment (ECOBES, European Chronic Obstructive Bronchitis Erdosteine Study). *Int J Clin Pharmacol Ther* 1995;**33**:612–8.
- 60 **Barbera JA**, Roca J, Ferrer A, *et al*. Mechanisms of worsening gas exchange during acute exacerbations of chronic obstructive pulmonary disease. *Eur Respir J* 1997;**10**:1285–91.
- 61 **Rodriguez-Roisin R**. Pulmonary gas exchange in acute respiratory failure. *Eur J Anaesthesiol* 1994;**11**:5–13.
- 62 **Saetta M**, Di Stefano A, Maestrelli P, *et al*. Airway eosinophilia in chronic bronchitis during exacerbations. *Am J Respir Crit Care Med* 1994;**150**:1646–52.
- 63 **Fiorini G**, Crespi S, Rinaldi M, *et al*. Serum ECP and MPO are increased during exacerbations of chronic bronchitis with airway obstruction. *Biomed Pharmacother* 2000;**54**:274–8.
- 64 **Saetta M**, Di Stefano A, Maestrelli P, *et al*. Airway eosinophilia and expression of interleukin-5 protein in asthma and in exacerbations of chronic bronchitis. *Clin Exp Allergy* 1995;**26**:766–74.
- 65 **Gompertz S**, Stockley RA. Inflammation: role of the neutrophil and the eosinophil. *Semin Respir Infect* 2000;**15**:14–23.
- 66 **Chanez P**, Vignola AM, O'Shaughnessy TC, *et al*. Corticosteroid reversibility in COPD is related to features of asthma. *Am J Respir Crit Care Med* 1997;**155**:1529–34.
- 67 **Fujimoto K**, Kubo K, Yamamoto H, *et al*. Eosinophilic inflammation in the airway is related to glucocorticoid reversibility in patients with pulmonary emphysema. *Chest* 1999;**115**:697–702.
- 68 **Burge PS**, Calverley PM, Jones PW, *et al*. Randomised, double blind, placebo controlled study of fluticasone propionate in patients with moderate to severe chronic obstructive pulmonary disease: the ISOLDE trial. *BMJ* 2000;**320**:1297–303.
- 69 **Balbi B**, Bason C, Balleari E, *et al*. Increased bronchoalveolar granulocytes and granulocyte/macrophage colony-stimulating factor during exacerbations of chronic bronchitis. *Eur Respir J* 1997;**10**:846–50.
- 70 **Smallman LA**, Hill SL, Stockley RA. Reduction of ciliary beat frequency in vitro by sputum from patients with bronchiectasis: a serine proteinase effect. *Thorax* 1984;**39**:663–7.
- 71 **Nadel JA**. Role of neutrophil elastase in hypersecretion during COPD exacerbations, and proposed therapies. *Chest* 2000;**117**:386–9S.
- 72 **May JR**. The bacteriology of chronic bronchitis. *Lancet* 1953;**2**:534–7.
- 73 **Cabello H**, Torres A, Celis R, *et al*. Distal airway bacterial colonisation in healthy subjects and chronic lung disease: a bronchoscopic study. *Eur Respir J* 1997;**10**:1137–44.
- 74 **Monso E**, Ruiz J, Manterola J, *et al*. Bacterial infection in chronic obstructive pulmonary disease: a study of stable and exacerbated outpatients using the protected specimen brush. *Am J Respir Crit Care Med* 1995;**152**:1316–20.
- 75 **Monso E**, Rosell A, Bonet G, *et al*. Risk factors for lower airway bacterial colonization in chronic bronchitis. *Eur Respir J* 1999;**13**:338–42.
- 76 **Soler N**, Ewig S, Torres A, *et al*. Airway inflammation and bronchial microbial patterns in patients with stable chronic obstructive pulmonary disease. *Eur Respir J* 1999;**14**:1015–22.
- 77 **Adler KB**, Hendley DD, Davis GS. Bacteria associated with obstructive pulmonary disease elaborate extracellular products that stimulate mucin secretion by explants of guinea pig airways. *Am J Pathol* 1986;**125**:501–14.
- 78 **Wilson R**, Roberts D, Cole P. Effects of bacterial products on human ciliary function in vitro. *Thorax* 1984;**40**:125–31.
- 79 **Read RC**, Wilson R, Rutman A, *et al*. Interaction of nontypeable *Haemophilus influenzae* with human respiratory mucosa in vitro. *J Infect Dis* 1991;**163**:549–58.
- 80 **Khair OA**, Devalia JL, Abdelaziz MM, *et al*. Effect of *Haemophilus influenzae* endotoxin on the synthesis of IL-6, IL-8, TNF-alpha and expression of ICAM-1 in cultured human bronchial epithelial cells. *Eur Respir J* 1994;**7**:2109–16.
- 81 **Hill AT**, Campbell EJ, Hill SL, *et al*. Association between airway bacterial load and markers of airway inflammation in patients with stable chronic bronchitis. *Am J Med* 2000;**109**:288–95.
- 82 **Hirschman JV**. Do bacteria cause exacerbations of COPD. *Chest* 2000;**118**:193–203.
- 83 **Scott KW**. Bacteriology of the sputum in exacerbation and remission of chronic bronchitis. *Br J Dis Chest* 1973;**67**:43–8.
- 84 **Gump DW**, Phillips CA, Forsyth BR, *et al*. Role of infection in chronic bronchitis. *Am Rev Respir Dis* 1976;**113**:465–74.
- 85 **Murphy TF**, Sethi S, Klingman KL, *et al*. Simultaneous respiratory tract colonisation by multiple strains of nontypeable *Haemophilus influenzae* in chronic obstructive pulmonary disease: implications for antibiotic therapy. *J Infect Dis* 1999;**180**:404–9.
- 86 **Fagon JY**, Chastre J, Trouillet JL, *et al*. Characterization of distal bronchial microflora during acute exacerbation of chronic bronchitis. Use of the protected specimen brush technique in 54 mechanically ventilated patients. *Am Rev Respir Dis* 1990;**142**:1004–8.
- 87 **Soler N**, Torres A, Ewig S, *et al*. Bronchial microbial patterns in severe exacerbations of chronic obstructive pulmonary disease (COPD) requiring mechanical ventilation. *Am J Respir Crit Care Med* 1998;**157**:1498–505.
- 88 **Eller J**, Ede A, Schaberg T, *et al*. Infective exacerbations of chronic bronchitis: relation between bacteriological etiology and lung function. *Chest* 1998;**113**:1542–8.

- 89 **Miravittles M**, Espinosa C, Fernandez-Laso E, *et al*. Relationship between bacterial flora in sputum and functional impairment in patients with acute exacerbations of COPD. Study Group of Bacterial Infection in COPD. *Chest* 1999;**116**:40–6.
- 90 **Yi K**, Sethi S, Murphy TF. Human immune response to nontypeable *Haemophilus influenzae* in chronic bronchitis. *J Infect Dis* 1997;**176**:1247–52.
- 91 **Chapman A**, Musher DM, Jonsson S, *et al*. Development of bactericidal antibody during *Branhamella catarrhalis* infection. *J Infect Dis* 1985;**151**:878–82.
- 92 **Mitchell J**, Hill SL. Immune response to *Haemophilus parainfluenzae* in patients with chronic obstructive lung disease. *Clin Diagn Lab Immunol* 1999;**7**:25–30.
- 93 **Nicotra MB**, Rivera M, Awe RJ. Antibiotic therapy of acute exacerbations of chronic bronchitis. A controlled study using tetracycline. *Ann Intern Med* 1982;**97**:18–21.
- 94 **Sachs AP**, Koeter GH, Groenier KH, *et al*. Changes in symptoms, peak expiratory flow, and sputum flora during treatment with antibiotics of exacerbations in patients with chronic obstructive pulmonary disease in general practice. *Thorax* 1995;**50**:758–63.
- 95 **Saint S**, Bent S, Vittinghoff E, *et al*. Antibiotics in chronic obstructive pulmonary disease exacerbations. A meta-analysis. *JAMA* 1995;**273**:957–60.
- 96 **Adams SG**, Melo J, Luther M, *et al*. Antibiotics are associated with lower relapse rates in outpatients with acute exacerbations of COPD. *Chest* 2000;**117**:1345–52.
- 97 **Woolhouse I**, Hill SL, Stockley RA. Symptom resolution assessed using a patient directed diary card during exacerbations of chronic bronchitis. *Thorax* 2001;**56**:947–53.
- 98 **Hegeler RG**, Hayashi S, Hogg JC, *et al*. Mechanisms of airway narrowing and hyperresponsiveness in viral respiratory tract infections. *Am J Respir Crit Care Med* 1995;**151**:1659–64.
- 99 **Melbye H**, Kongerud J, Vorland L. Reversible airflow limitation in adults with respiratory infection. *Eur Respir J* 1994;**7**:1239–45.
- 100 **Jacoby DB**, Fryer AD. Interaction of viral infections with muscarinic receptors. *Clin Exp Immunol* 1999;**29**:59–64.
- 101 **Buscho RO**, Saxtan D, Shultz PS, *et al*. Infections with viruses and *Mycoplasma pneumoniae* during exacerbations of chronic bronchitis. *J Infect Dis* 1978;**137**:377–83.
- 102 **Carilli AD**, Gohd RS, Gordon W. A virological study of chronic bronchitis. *N Engl J Med* 1964;**270**:123.
- 103 **Lamy ME**, Pouthier-Simon F, Debacker-Willame E. Respiratory viral infections in hospital patients with chronic bronchitis. Observations during periods of exacerbation and quiescence. *Chest* 1973;**63**:336–41.
- 104 **Smith CB**, Golden CA, Kanner RE, *et al*. Association of viral and *Mycoplasma pneumoniae* infections with acute respiratory illness in patients with chronic obstructive pulmonary diseases. *Am Rev Respir Dis* 1980;**121**:225–32.
- 105 **Stenhouse AC**. Viral antibody levels and clinical status in acute exacerbations of chronic bronchitis: a controlled prospective study. *BMJ* 1968;**3**:287–90.
- 106 **Seemungal TA**, Harper-Owen R, Bhowmik A, *et al*. Detection of rhinovirus in induced sputum at exacerbation of chronic obstructive pulmonary disease. *Eur Respir J* 2000;**16**:677–83.
- 107 **Smith CB**, Golden C, Klauber MR, *et al*. Interactions between viruses and bacteria in patients with chronic bronchitis. *J Infect Dis* 1976;**134**:552–61.
- 108 **Poole PJ**, Chacko E, Wood-Baker RW, *et al*. Influenza vaccine for patients with chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2000;CD002733.
- 109 **Monto AS**, Bryan ER. Susceptibility to rhinovirus infection in chronic bronchitis. *Am Rev Respir Dis* 1978;**118**:1101–3.
- 110 **Greenberg SB**, Allen M, Wilson J, *et al*. Respiratory viral infections in adults with and without chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000;**162**:167–73.
- 111 **Beatty CD**, Grayston JT, Wang SP, *et al*. Chlamydia pneumoniae, strain TWAR, infection in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1991;**144**:1408–10.
- 112 **Mogulkoc N**, Karakurt S, Isalska B, *et al*. Acute purulent exacerbation of chronic obstructive pulmonary disease and Chlamydia pneumoniae infection. *Am J Respir Crit Care Med* 1999;**160**:349–53.
- 113 **Verkooyen RP**, Van Lent NA, Mousavi Joulandan SA, *et al*. Diagnosis of Chlamydia pneumoniae infection in patients with chronic obstructive pulmonary disease by micro-immunofluorescence and ELISA. *J Med Microbiol* 1997;**46**:959–64.
- 114 **Lieberman D**, Lieberman D, Ben Yaakov M, *et al*. Infectious etiologies in acute exacerbation of COPD. *Diagn Microbiol Infect Dis* 2001;**40**:95–102.
- 115 **Karnak D**, Beng Sun S, Beder S, *et al*. Chlamydia pneumoniae infection and acute exacerbation of chronic obstructive pulmonary disease (COPD). *Respir Med* 2001;**95**:811–6.
- 116 **Nordenhall C**, Pourazar J, Blomberg A, *et al*. Airway inflammation following exposure to diesel exhaust: a study of time kinetics using induced sputum. *Eur Respir J* 2000;**15**:1046–51.
- 117 **Rudell B**, Blomberg A, Helleday R, *et al*. Bronchoalveolar inflammation after exposure to diesel exhaust: comparison between unfiltered and particle trap filtered exhaust. *Occup Environ Med* 1999;**56**:527–34.
- 118 **Ohtoshi T**, Takizawa H, Okazaki H, *et al*. Diesel exhaust particles stimulate human airway epithelial cells to produce cytokines relevant to airway inflammation in vitro. *J Allergy Clin Immunol* 1998;**101**:778–85.
- 119 **Kopp MV**, Ulmer C, Ihorst G, *et al*. Upper airway inflammation in children exposed to ambient ozone and potential signs of adaptation. *Eur Respir J* 1999;**14**:854–61.
- 120 **Devalia JL**, Ruzsnaik C, Herdman MJ, *et al*. Effect of nitrogen dioxide and sulphur dioxide on airway response of mild asthmatic patients to allergen inhalation. *Lancet* 1994;**344**:1668–71.
- 121 **Anderson HR**, Spix C, Medina S, *et al*. Air pollution and daily admissions for chronic obstructive pulmonary disease in 6 European cities: results from the APHEA project. *Eur Respir J* 1997;**10**:1064–71.
- 122 **Garcia-Aymerich J**, Tobias A, Anto JM, *et al*. Air pollution and mortality in a cohort of patients with chronic obstructive pulmonary disease: a time series analysis. *J Epidemiol Community Health* 2000;**54**:73–4.
- 123 **Schwartz J**. Air pollution and hospital admissions for the elderly in Detroit, Michigan. *Am J Respir Crit Care Med* 1994;**150**:648–55.
- 124 **Schwartz J**. PM₁₀, ozone, and hospital admissions for the elderly in Minneapolis-St Paul, Minnesota. *Arch Environ Health* 1994;**49**:366–74.
- 125 **Sunyer J**, Saez M, Murillo C, *et al*. Air pollution and emergency room admission for chronic obstructive pulmonary disease: a 5-year study. *Am J Epidemiol* 1993;**137**:701–5.
- 126 **Sunyer J**, Schwartz J, Tobias A, *et al*. Patients with chronic obstructive pulmonary disease are at increased risk of death associated with urban particle air pollution: a case-crossover analysis. *Am J Epidemiol* 2000;**151**:50–6.
- 127 **Davies L**, Hadcroft J, Mutton K, *et al*. Antimicrobial management of acute exacerbations of chronic airflow limitation. *Q J Med* 2001;**94**:373–8.
- 128 **Smith JA**, Redman P, Woodhead MA. Antibiotic use in patients admitted with acute exacerbations of chronic obstructive pulmonary disease. *Eur Respir J* 1999;**13**:835–8.

THORAX

Chronic obstructive pulmonary disease • 6: The aetiology of exacerbations of chronic obstructive pulmonary disease

A J White, S Gompertz and R A Stockley

Thorax 2003 58: 73-80

doi: 10.1136/thorax.58.1.73

Updated information and services can be found at:
<http://thorax.bmj.com/content/58/1/73>

References

These include:

This article cites 116 articles, 48 of which you can access for free at:
<http://thorax.bmj.com/content/58/1/73#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Air pollution](#) (110)
[Environmental issues](#) (253)
[Inflammation](#) (1020)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>