

# *Drosophila mojoless*, a Retroposed GSK-3, Has Functionally Diverged to Acquire an Essential Role in Male Fertility

Rasika Kalamegham,\*† David Sturgill,\* Esther Siegfried,†<sup>1</sup> and Brian Oliver\*

\*Laboratory of Cellular and Developmental Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland; and †Department of Biochemistry, Molecular Biology and Microbiology, The Pennsylvania State University

Retroposition is increasingly recognized as an important mechanism for the acquisition of new genes. We show that a *glycogen synthase kinase-3* gene, *shaggy* (*sgg*), retroposed at least 50 MYA in the *Drosophila* genus to generate a new gene, *mojoless* (*mjl*). We have extensively analyzed the function of *mjl* and examined its functional divergence from the parental gene *sgg* in *Drosophila melanogaster*. Unlike *Sgg*, which is expressed in many tissues of both sexes, *Mjl* is expressed specifically in the male germ line, where it is required for male germ line survival. Our analysis indicates that *mjl* has acquired a specific function in the maintenance of male germ line viability. However, it has not completely lost its ancestral biochemical function and can partially compensate for loss of the parental gene *sgg* when ectopically expressed in somatic cells. We postulate that *mjl* has undergone functional diversification and is now under stabilizing selection in the *Drosophila* genus.

## Introduction

New genes arise from exon shuffling (Long, Deutsch, et al. 2003), gene splits and joins (Wang et al. 2004), or duplication (Li et al. 2005) followed by diversification of function (Long, Betran, et al. 2003). Duplication can occur at the DNA level or via an RNA intermediate. RNA-mediated gene duplication, or retroposition, occurs by a retrovirus-like integration mechanism. Relatively recent retropositions can be identified based on simple criteria (Betran and Long 2002): retroposed genes show strong sequence homology with a parent gene in the genome, the coding sequence of the newly retroposed gene is not interrupted by introns (due to the mechanism of transmission), the PolyA tract from the mRNA intermediate is encoded in the genome, and the terminal repeats associated with the integration event are evident. These newly created genes have been under intense study, but older retroposed genes have not yet been extensively characterized. One might expect that some of the features of retroposed genes, such as PolyA tracts, would be unconstrained by evolution and eventually be lost. Such genes might ultimately acquire introns and become more difficult to distinguish from genes arising from strictly DNA-mediated duplication (Betran and Long 2002). Retroposed genes have several different fates: one member of the duplicated pair can become nonfunctional and silenced (nonfunctionalization), there could be a partitioning of the ancestral function between the 2 copies (subfunctionalization), or 1 of the copies could acquire new functions (neofunctionalization) (Lynch and Conery 2000).

We have identified a new retroposed glycogen synthase kinase-3 (GSK-3)-encoding gene, derived from one of the alternatively spliced mRNAs from the *shaggy* (*sgg*) locus. In *Drosophila melanogaster*, *sgg* locus is on the X chromosome and the retroposed duplicate, and *mojoless* (*mjl*) is on the third chromosome. We show that this

retroposition event occurred near the root of the *Drosophila* genus and that *sgg* and *mjl* have diverged rapidly.

This gene pair is of particular interest because of the highly pleiotropic function of the parental *sgg* gene. GSK-3 proteins are kinases essential for metabolic regulation, cell fate determination, stem cell maintenance, and nuclear import in a wide range of eukaryotes (Jope and Johnson 2004). *Sgg*, the *Drosophila* homolog of GSK-3 $\beta$  is probably best known for its role as the key negative regulator of the Wingless (*wg*) signaling pathway, but is also intimately involved in diverse developmental processes, such as establishment of anterior–posterior polarity in the embryo, patterning of the embryonic cuticle as well as various adult structures (wing, eye, haltere etc.), and regulating circadian rhythms (Siegfried et al. 1994; Stanewsky 2002). *Sgg* is maternally loaded and has been detected ubiquitously throughout development (Bourouis et al. 1990). In addition, it has multiple splice variants ranging in size from 10 to 110 kilo basepair (kb) (Bourouis et al. 1990). These encode for at least 7 predicted protein isoforms (FlyBase 2003). Multiple isoforms of *sgg* mRNA show no obvious temporal pattern of expression and in situ hybridization using common probes show uniform expression in embryos (Bourouis et al. 1990; Siegfried et al. 1990).

On the other hand, *mjl* is intronless and it encodes for a single transcript with highly specific expression in the male germ line. Altered expression patterns are expected to promote neofunctionalization, and indeed, *mjl* knockdown results in male sterility due to ablation of the male germ line. The loss of *mjl* causes a germ line-specific defect with no obvious effects in the soma. This indicates that the retrocopy has now acquired a critical novel function, which probably explains why it has been retained in the genome. However, this neofunctionalization is not complete. Ectopically *Mjl* was capable of partially rescuing an *sgg* mutant. These results are consistent with retention of ancestral biochemical function (as a gene family member), but evolution of a new spatial expression pattern along with a specific novel role.

## Methods

### Fly Stocks and Crosses

All flies were raised and maintained on standard cornmeal media (Tucson *Drosophila* Stock Center, Tucson, AZ)

<sup>1</sup> Present address: Department of Biology, University of Pittsburgh at Johnstown.

Key words: germ cell, male sterile, gene duplication, GSK-3, adaptive evolution, retroposition, male germ line.

E-mail: rasika@nidk.nih.gov.

*Mol. Biol. Evol.* 24(3):732–742, 2007

doi:10.1093/molbev/msl201

Advance Access publication December 18, 2006

at 25 °C, except the ones used for heat-shock regimens and the dsRNA-mediated interference (RNAi) studies, as noted. Enhancer trap 3914 expresses LacZ in all the somatic cells of the gonad (Asaoka et al. 1998). The nos-GAL4 line drives expression of Upstream Activating Sequence (UAS) transgenes in germ lines of both males and females (Schulz et al. 2004). The hsp-70-GAL4 line drives expression of UAS transgenes ubiquitously following heat shock (FlyBase 2003). The armadillo (arm)-GAL4 line was used to induce expression of transgenes in cells where Sgg is required in embryos (FlyBase 2003).

The coding region of *mjl* was polymerase chain reaction (PCR) amplified from expressed sequence tag (EST) clone GH16447 (Celniker et al. 2002) with primers designed to include ~120 bp upstream of the start of transcription (forward primer [FP]: 5'-CAG TTC CAC ACG CAT ACG CAC-3'; reverse primer [RP]: 5'-GTG CGG TTC AAA TCA AGA CC-3'). The 1.8-kb fragment was cloned into the pUAST vector (Brand and Perrimon 1993) to create the *UAS-mjl* transgene. To generate the *UAS-mjl-RNAi* transgene, two 700-bp fragments encoding the C-terminal region of *mjl* were PCR amplified (fragment 1: FP 5'-GCGAATTCGATCGTCAAGGTTATGGGCATCTC-3'; RP 5'-GCACTAGTGCTATTGCACTGCCACCATCAT-3'; fragment 2: FP 5'-GCGGTACCGATCGTCAAGGTTATGGGCACTC-3'; RP 5'-GCACTAGTGCTATTGCACTGCCACCATCAT-3'). The fragments were then cloned into a modified pUAST vector (Han K, personal communication) containing a 500-bp heterologous intron inserted between the insertion sites. Transgenic lines were generated by *P* element-mediated transformation using the stock *y w;Δ2-3 Sb/TM6, Ubx* (Robertson et al. 1988). Eight and 10 independent lines of insertion were established for the *UAS-mjl* and *UAS-mjl<sup>RNAi</sup>* transgenes, respectively.

#### Phylogenetic Analysis

Sequence alignments were performed using RevTrans 1.4 (Wernersson and Pedersen 2003), which aligns coding sequence in translated amino acid space. Alignments were visualized using Jalview (Clamp et al. 2004). *Ka/Ks* was calculated pairwise using the Nei-Gojobori method (Nei and Gojobori 1986) and the modified Nei-Gojobori method (Zhang et al. 1998) in MEGA3.1 with Jukes-Cantor correction and transition/transversion *R* value of 2. For comparison, *Ka/Ks* was also calculated with a maximum likelihood approach using the yn00 program within the phylogenetic analysis by maximum likelihood package, implementing the method developed by Yang and Nielsen (2000). Phylogenetic trees were constructed using the minimum evolution method in MEGA3.1, and significance was measured via bootstrapping with 1,000 replicates (Kumar et al. 2004) and Neighbor-Joining to draw the trees, with the same substitution model as the modified Nei-Gojobori calculation.

#### Antibody Generation

To minimize cross-reaction of the antibody with Sgg, the divergent C-terminal region of Mjl was fused to bacterial glutathione-S-transferase (GST) and the recombinant protein was purified and then used to raise antibodies. A 763-bp C-terminal region fragment was amplified by

PCR (FP: 5'-GACGAATTCATGTTATGGGCACT-3'; RP: 5'-GCACTGGAGTTAGTTTTTCATCTC-3'). The fragment was cloned into the pGEX-4T-2 vector (Amersham-Pharmacia, Piscataway, NJ) to produce an in-frame GST fusion protein. The protein was purified on a GST affinity column, using the manufacturer's protocol (Amersham-Pharmacia), and injected into rabbits (Pocono Rabbit Farm, Canadensis, PA). The antibody was affinity purified against the fusion protein (Brent et al. 1987).

#### Immunoblots

Standard molecular biology techniques were used for protein transfer onto nitrocellulose membranes. Samples were homogenized in 50 μl of 2× Gel buffer (8 M urea, 2 M thio-urea, 3% sodium dodecyl sulfate (SDS), 75 mM dithiothreitol, 25 mM Tris pH 6.8, 0.1% bromophenol blue) using a motorized homogenizer. Samples were spun for 2 min in a microfuge to pellet cell debris and loaded onto a 12% SDS-polyacrylamide gel. Protein transfer was confirmed by Ponceau-S staining. Blots were blocked with 5% milk in Tris-buffered saline (TBS) (20 mM Tris, 150 mM NaCl) for 2–3 h at room temperature on a rocker, incubated with rabbit polyclonal anti-Mjl antibody overnight at 4 °C in 5% milk in TBS-Tween (20 mM Tris, 150 mM NaCl, 0.1% Tween) at a final concentration of 1:2000, washed, incubated in horseradish peroxidase-conjugated goat anti-rabbit antibody for 2–3 h at room temperature (Jackson Immuno Research Laboratories Inc., West Grove, PA), and developed using a commercial Enhanced chemiluminescence kit (Amersham-Pharmacia).

#### Rescue of *sgg* Mutant Embryos

Virgin females of the genotype *sgg<sup>M11-1</sup> FRT<sup>101</sup>/FM7; arm-GAL4/arm-GAL4* were mated to *ovo<sup>D1</sup> FRT<sup>101</sup>/Y; FLP recombine (FLP)<sup>38</sup>/FLP<sup>38</sup>* males. The progeny of this cross were heat shocked at 37 °C for 1 h during the crawling third instar larval stage to induce FLP-directed recombination (Siegfried et al. 1992). Virgin females containing mosaic germ lines as well as 1 copy of the *arm-GAL4* chromosome were mated to homozygous *UAS-mjl<sup>4B</sup>*, *UAS-sgg<sup>WDT7A</sup>*, or Oregon-R males. Embryo sorting and cuticle preparations were performed (Steitz et al. 1998). Embryos were scored as (+/–) for rescue, based on presence or absence of denticle bands. Five hundred embryos were screened for each genotype. Embryos were viewed and photographed in dark field.

#### RNAi Treatments

Virgin nos-GAL4 females were crossed to various RNAi transgenic males and maintained at 29 °C (*GAL4* is more effective at higher temperature). The progeny of the crosses were collected and newly eclosed males and females were individually mated to their respective wild-type counterparts. The crosses were maintained at 29 °C for up to 10 days and vials were examined for evidence of larvae. As a control, the experiment was repeated with virgin *hsp-70-GAL4* females instead of nos-GAL4 females. The progenies were heat shocked at 38 °C for 15 min a day. The mating experiments were repeated at 25 °C, as above.

## Immunofluorescence Staining of Testes

Testes from newly eclosed males (<24 h) were dissected and fixed (Fabrizio et al. 2003). Samples were blocked in 3% bovine serum albumin in phosphate buffered saline with 1% Triton X-100 overnight at 4 °C. Antibodies were used at the following concentrations: rabbit polyclonal anti-Mjl antibody at 1:20; chicken anti-Vasa at 1:1,000 (Burnett and Howard 2003). Goat anti-rabbit Alexa 488 and anti-chicken Alexa 555 (Invitrogen Molecular Probes, Carlsbad, CA) secondary antibodies were used at a final concentration of 1:250. After the final wash, testes were counterstained with 4',6-diamidino-2-phenylindole (DAPI) (50 µg/ml) and mounted (50% glycerol with 1,4-diazabicyclo[2.2.2]octane), and images were captured on a LSM510 confocal laser scanning microscope (Zeiss, Thornwood, NY).

## Results

### *mjl* Arose in *Drosophila* by Retroposition

The *mjl* (*CG11338*) locus is on the third chromosome and encodes a novel GSK-3 protein sharing 53% amino acid identity with Sgg, the well-studied X-linked *D. melanogaster* GSK-3-encoding gene (fig. 1A). This gene was previously named *gasket* for GSK-3 based on its sequence homology (FlyBase 2003). We have subsequently renamed the gene *mjl* based on its loss-of-function phenotype, which is male sterility.

The greatest level of conservation between Sgg and Mjl is within the canonical kinase domain (69% identity, outlined in red box in fig. 1A) and the least conserved region is the C-terminal region (11% similarity, outlined in green box in fig. 1A). Mjl and Sgg are the only predicted GSK-3 proteins we found encoded in the *D. melanogaster* genome (Adams et al. 2000; Celniker et al. 2002; FlyBase 2003). The *mjl* transcript is intronless, unlike *sgg*, which has 9 introns, indicating that it arose by retroposition of *sgg*. To determine which of the 11 *sgg* mRNA isoforms might have given rise to *mjl*, we examined the regions specific to those transcripts and found that *mjl* is most similar to *sgg-RH*. This suggests that an ancient version of *sgg-RH* mRNA retroposed to give rise to *mjl* (fig. 1C). There is no evidence of a PolyA tract or short repeats in *mjl*, implying that it is not a newly retroposed gene.

In order to investigate how long ago in the history of the genus *mjl* retroposed, we examined the genomic sequence of newly sequenced *Drosophila* species. We found that *mjl* was present in each species examined (fig. 1B; *D. melanogaster*, *Drosophila sechellia*, *Drosophila mauritiana*, *Drosophila simulans*, *Drosophila yakuba*, *Drosophila ananassae*, *Drosophila pseudoobscura*, *Drosophila mojavensis*, and *Drosophila virilis*). Interestingly, we have been unable to identify a homologue of *mjl* in the genomes of non-*Drosophila* insects (*Anopheles gambiae*, *Apis mellifera*, and *Tribolium castaneum*) or indeed any other published genomes. We also examined the linkage of *sgg* and *mjl* in the various species, and where available, the data indicate that *sgg* is X linked, whereas *mjl* is on the third chromosome (Muller element E). Collectively, these results demonstrate that *mjl* arose as a retroposed *sgg*, an event that

was probably *Drosophila* specific. Based on estimates of molecular clocks, we can date this event to at least 50 MYA (Russo et al. 1995; Tamura et al. 2004).

### Mjl Evolution

We examined the sequence conservation of *sgg* and *mjl* within the genus *Drosophila*. The predicted Sgg proteins are more conserved as a group than the predicted Mjl proteins, especially at their N termini (fig. 2). Residues that are crucial for functioning of the kinase domain, however, are well conserved within both groups of orthologs. This suggests that the *sgg* genes are under greater evolutionary constraint on their divergence. The observation that the N-terminal region outside the canonical kinase domain is strictly conserved within the predicted Sgg proteins and not within the Mjl proteins suggests that this domain may not be integral to function in the Mjl proteins. The C-terminal regions of both predicted Sgg and Mjl proteins show poor conservation indicating that this region is experiencing the least functional constraint within both groups of proteins (fig. 2).

We next examined nonsynonymous and synonymous substitution rates ( $Ka/Ks$ ) among orthologs of *sgg* and *mjl* (table 1). Maximum likelihood methods for comparing the rate of substitution at synonymous sites in a gene sequence ( $Ks$ ) and comparing it with the rate of substitution at nonsynonymous sites ( $Ka$ ) are effective approaches to characterizing evolutionary forces (Nei and Kumar 2000; Hurst 2002). Orthologs of *mjl* have experienced on average a greater number of nonsynonymous substitutions per site than *sgg* (table 1). The average synonymous substitution rate ( $Ks$ ) is much higher for *mjl* orthologs than for *sgg* orthologs (table 1). Most saliently, the fact that the  $Ka/Ks$  ratio is much less than 0.5 for both genes (table 1) implies that both genes are under strong selective constraints and therefore are functional.

### Mjl Has Retained GSK-3 Family Function

The greatest level of conservation between the Mjl and Sgg proteins is within the canonical GSK-3 kinase domain (69% identity, outlined in red in fig. 1A). All the residues crucial for GSK-3 function, as deduced from the mammalian crystal structure studies, are conserved in Mjl, such as R96, R180, and K205, which form a positively charged pocket to attract the primed substrate; Y216 within the activation loop whose phosphorylation increases catalytic activity >200-fold; R96 whose phosphorylation is important to align the  $\beta$ -strand domain and the  $\alpha$ -helical domain; S9 whose phosphorylation inhibits GSK-3 activity; and K85, which forms a salt bridge with E97 at the active site (Dajani et al. 2001; ter Haar et al. 2001). This indicates that the canonical kinase domain in the duplicate gene copy, *mjl*, is under strong selective pressure in order to retain its biochemical activity. However, the other regulatory domains within the coding sequence and perhaps upstream and downstream sequences are not subject to the same level of conservation, suggesting that *mjl* has started functionally diverging from *sgg* in its ability to recognize and bind to substrates, a key step in neofunctionalization (Zhang et al. 2004).

Sequence conservation suggests that the kinase domain is the most constrained, but does this conservation




FIG. 1.—Mjl is a unique GSK-3 family member related to Sgg. (A) Sequence alignment between Sgg and Mjl proteins from *Drosophila melanogaster*. Identical residues are shaded blue. The red box marks the canonical kinase domain and the green box marks the divergent C-terminal region used to generate anti-Mjl antibody. (B) Phylogenetic tree of *Drosophila* Sgg and Mjl proteins generated using MEGA3.1 software. The Sgg proteins all cluster together as do the Mjl proteins, indicating that they are distinct from one another. (C) Model depicting retroposition of *sgg-RH* mRNA from X to an autosome (the third chromosome in *D. melanogaster*) giving rise to *mjl*.

mean that Mjl and Sgg have the potential to recognize some of the same substrates? If Mjl and Sgg share biochemical activities, then ectopic expression of *mjl* should mimic *sgg* overexpression. Previous work has shown that overexpression of *sgg* along the presumptive wing margin disrupts the endogenous *wg* signaling pathway (Steitz et al. 1998).

Overexpression of *sgg* in the wing imaginal disc results in a loss of bristles and marginal tissue in the adult wing (compare panels A and B in fig. 3). When *mjl* is similarly expressed there is a comparable disruption of the wing margin (fig. 3C). The observation that ectopic expression of *sgg* and *mjl* result in similar phenotypes suggests that Mjl is also




FIG. 2.—Alignments of Mjl and Sgg orthologs. Sequence alignment of various *Drosophila* (A) Mjl and (B) Sgg proteins generated using ClustalW. Residues in blue are identical and the red box outlines the canonical kinase domain.

capable of interacting with and disrupting the endogenous wg signaling pathway. To confirm this hypothesis, we modulated the levels of 2 known components of the wg signaling pathway, *Drosophila* Axin (dAxin) and Arm. Expressing *mjl* in the wing imaginal disc in the presence of an extra copy of *arm* (fig. 3D) or a single copy of dAxin (fig. 3E) suppressed the wing phenotype. Similar types of suppression are seen with overexpression of *sgg* and modulation of gene dosages of either *arm* or *d-Axin* (Steitz 2000). Therefore, we conclude that Mjl is capable of interacting with the canonical wg signaling pathway. These suggest that any neofunctionalization of Mjl protein function is subtle or that any dramatic subfunctionalization does not include portions of the proteins involved in wg function.

We also addressed the question of whether Mjl and Sgg have overlapping biological activity by attempting

to express *sgg* mutant embryos with ectopic expression of *mjl*. Because *sgg* is a maternal effect gene, mutant embryos were derived from mothers with homozygous mutant germ lines or germ line clone females (see Methods). Mutant embryos are not viable and exhibit a distinct perturbation of segmentation that is revealed in a loss of all denticles or a “naked” phenotype (fig. 4, compare panels A and B). Zygotic expression of *sgg* in these embryos can partially restore segmentation (fig. 4C) (Steitz et al. 1998). In a similar fashion, ectopic expression of *mjl* also partially restores segmentation (fig. 4D). Again, this suggests that any differences in Mjl and Sgg protein function in this spatiotemporal context are subtle. This does not mean that there is no evidence of functional divergence. Indeed, the extent of *sgg* mutant rescue by the *mjl* transgene is less than the rescue by the *sgg* transgene. A rescue of 43% was seen for the *sgg*

**B**

D.melanogaster Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.simulans Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.yakuba Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.ananassae Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.pseudoobscura Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.mojavensis Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.virilis Sgg	MSGRRPTSSFAEGNKQSPSLVLLGGVKTCSDGSKIITVVVTPGGGTDRVQEVSYTDTKVIGNGSFGVVFOAKLCD75	
D.melanogaster Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.simulans Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.yakuba Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.ananassae Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.pseudoobscura Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.mojavensis Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.virilis Sgg	TGELVAIKKVLQDRRFKNRELQIMRRLHHCNIVKLLYFFYSSGEKRDEVFLNLVLEYIPETVYKVAROYAKTKQT150	
D.melanogaster Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.simulans Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.yakuba Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.ananassae Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.pseudoobscura Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.mojavensis Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.virilis Sgg	IPINFIRLYMYQLFRSLAYIHS LGICHRDIKPNQLLDPE TAVLKLCDFGSAKQLLHGEPNVS YICSRYYRAPEL225	
D.melanogaster Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.simulans Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.yakuba Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.ananassae Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.pseudoobscura Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.mojavensis Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.virilis Sgg	IFGAINYTTKIDVWSAGCVLAE LLLGQPIFPGDSGVDQLVEV I KVLGTP TREQIREMNPNYTEFKFPQIKSHPWQ300	
D.melanogaster Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.simulans Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.yakuba Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.ananassae Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.pseudoobscura Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.mojavensis Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.virilis Sgg	KVFRIRTPTEA INLVSLLEYTPSARI TPLKACAHPPFDELRMEGNHTLPNGRDMPP LFNFTHELS IQPSLVPQ375	
D.melanogaster Sgg	LLPKHLQNASGPG -GNRPSAGGAAS I AASGSTSVSS TSGASVEGSAQPO - - - - - SQGTAAAAGSGSGGA 439	
D.simulans Sgg	LLPKHLQNASGPG -GNRPSAGGAAS I AASGSTSVSS TSGASVEGSAQPO - - - - - SQGTAAAAGSGSGGA 439	
D.yakuba Sgg	LLPKHLQNASGPG -ANRSAGGAASAAASGSTSVSS TSGASAEGSAQPOQ - - - - - SQGTAAAAGSGSGGA 440	
D.ananassae Sgg	LLPKHLQNAAGGAAANRSAG -ASAGAASGSTSVSS TSGASAEGSAQPOQAASA TAAAAASASGAAAGTSGSGS 449	
D.pseudoobscura Sgg	LLPKHLANRRSSGGGP - - - - - AAAGAVSGSTSVSS TSGASADGAA LSO - - - - - AGGTSGS SVVSGSGAG 435	
D.mojavensis Sgg	LLPKHLQNAHAASANRASAGGPA GAAASGSTNVSSAGSGTSGDGAAGSQQ - - - - - QQQAPASAAAGTGAG 441	
D.virilis Sgg	LLPKHLQNAHAASANRASAGGPA GAAASGSTNVSSAGSGTSGDGAAGSQQ - - - - - QQQAPASAAAGTGAG 441	
D.melanogaster Sgg	TAG - - - - - TGGASAGG -PGSGNNSSSGGASGAPSAVAAAGGANA AVAGGAGGGGAGAAATAA - - ATATGA IG 502	
D.simulans Sgg	TAG - - - - - TGGASAGG -PGSGNNSSSGGASGAP -TVAAG -ANA AVAG - - -GGGAGAAATAA - - ATATGA IG 497	
D.yakuba Sgg	TAG - - - - - TGGASAGGVPGSGNNSSSGGVSGAPTAVTAG -ANA AVAGGAGGGGAGAAATAA - - ATAGA IG 503	
D.ananassae Sgg	GAGTAAAGVASGGAAAGTSGSGGNNSSSGGGGGGGVPAPSA GNAQQSAGSGGGGGGGGAAAAATGATASGA IA 524	
D.pseudoobscura Sgg	GAGSSVG - - -GPA TGSAGGQ -NNSNPGSGAPSAVAAA PAAQANAGGPAAA GGGGGGGGAGAA - - ATAGS IA 504	
D.mojavensis Sgg	GAA - - - - - NVGGVQGNNSSSNNSGNPTTV TAAAGAPNA QQS NAAAAG -SGGGGGGAAATAA - - ASAG -A IA 503	
D.virilis Sgg	GAA - - - - - NVGGVQGNNSSSNNSGNPTTV TAAAGAPNA QQS NAAAAG -SGGGGGGAAATAA - - ASAG -A IA 503	
D.melanogaster Sgg	ATNAGGANVTGSQNSALNSSGSGSGSNGEAAAGSGSGSGSG - - - - - GNGGNDAGDSGA IAS - -GGGAAE TE 570	
D.simulans Sgg	ATNAGGANVTGSQNSALNSSGSGSGSNGEAAAGSGSGSGSG - - - - - GNGGNDAGDSGAAAS -GGGAAE TE 565	
D.yakuba Sgg	ATNAGGANVTGSQNSALNSSGSGSGSNGEAAAGSGSGSGSGSGGGNGCNGGNDAGDSGAAASGAGGGAPE TE 578	
D.ananassae Sgg	ATNAGGANVTGSQNSALNS - - - - - TGNGNGSGSNGNGNGGGS - - - - - TGSNGEADDESAGAAETPAAAAENE 591	
D.pseudoobscura Sgg	ATNAGGANVTGSQNSALNSSG - - - - - TANGNGSGSNGNGET TAA - - - - - VGTG -GSAATPAAAGGEENGATAAAAA 571	
D.mojavensis Sgg	TSNAGGANITDSQNSAFNSSGNDAEASAAGDEGGGGSSNG - - - - - SCSGNDAAAE - - - - - 557	
D.virilis Sgg	TSNAGGANITDSQNSAFNSSGNDAEASA - - - - - 533	
D.melanogaster Sgg	AAASG	575
D.simulans Sgg	AAASG	570
D.yakuba Sgg	AAASG	583
D.ananassae Sgg	AAASG	596
D.pseudoobscura Sgg	AAAA -	575
D.mojavensis Sgg	- - - - -	
D.virilis Sgg	- - - - -	

FIG. 2. (Continued)

transgene as opposed to 10% for the *mjl* transgene (fig. 4E). These results are consistent with functional divergence, but could also be due to technical factors, such as degree of overexpression and position effects, which are inherent in all transgenic analysis.

Taken together the above data demonstrate that Mjl has at least partially retained its ability to substitute for Sgg and is indeed capable of genetically interacting with components of the canonical wg pathway. Functional divergence at the level of the proteins is subtle.

**Table 1**  
**Estimation of Codon Substitution Events in *mjl* and *sgg***

	<i>mjl</i>			<i>sgg</i>		
	<i>Ka</i>	<i>Ks</i>	<i>Ka/Ks</i>	<i>Ka</i>	<i>Ks</i>	<i>Ka/Ks</i>
PAML <sup>a</sup>	0.19	3.51	0.06	0.09	0.61	0.14
PAML <sup>b</sup>	0.19	1.51	0.13	0.08	0.60	0.14
MEGA	0.22	0.84	0.24	0.09	0.46	0.19

NOTE.—PAML, phylogenetic analysis by maximum likelihood.

<sup>a</sup> Yang–Nielsen method.

<sup>b</sup> Nei–Gojobori method.

### *mjl* Is Expressed in the Male Germ Line


Functional diversification often involves new spatial patterns of expression. Therefore, we explored the spatial expression pattern of *mjl* gene products by querying EST data, northern blotting, reverse transcriptase (RT)–PCR, western blotting, and antibody detection methods.

Multiple *mjl* ESTs have been identified in testis cDNA libraries, but not in ovary cDNA libraries (Andrews et al. 2000; Stapleton et al. 2002). One *mjl* EST from a head library exists (Stapleton et al. 2002). Although Mjl might also be expressed in other tissues, these EST data suggest that *mjl* expression is highly testes biased. To further explore *mjl* mRNA expression, we performed Northern blot analysis and RT–PCR experiments and confirmed that the expected 1.9-kb *mjl* transcript is expressed in the testis (data not shown). We have not detected *mjl* in females or in any other nongonadal tissues including head, either by RT–PCR or immunostaining techniques, thus confirming that *mjl* mRNA shows a highly testes-biased expression pattern.

To further characterize Mjl expression pattern in vivo, the divergent C-terminal portion of Mjl was used to produce an antibody that, predictably, does not cross-react with Sgg (data not shown). On western blots of protein extracts from adult males, male abdomens, and testes, a single band of the predicted 56-kDa mobility was detected with anti-Mjl antibody (fig. 5A). No protein was detected in extracts from females (fig. 5A).

To determine what cell types express Mjl in the testis and where it is localized within cells, we performed immunostaining experiments. In *D. melanogaster*, the apical tip of the testis forms the germinal proliferation center (Fuller 1993). At the very tip of the testis a tight cluster of 12–16 nondividing somatic cells form a structure called the hub, which forms the niche (Kiger et al. 2001; Tulina and Matunis 2001). Closely attached to the hub is a layer of germ line stem cells (GSCs) interspersed with somatic stem cells.

Mjl was readily detected in adult testis, but not in ovary (fig. 5, compare panels B and C). The anti-Vasa antibody is a good marker of germ cells (Rongo et al. 1997), and anti-Mjl staining was detected in Vasa-positive cells of the testis (fig. 5B and E). Anti-Mjl staining revealed a cytoplasmic subcellular localization, as evidenced by the lack of overlap with DAPI nuclear counter staining (fig. 5E). This was expected because all GSK-3s are cytoplasmic (Woodgett 1990). Mjl is expressed in the GSCs, which immediately surround the hub, the gonialblasts, and the developing spermatogonia as evidenced by overlapping staining of anti-Mjl with anti-Vasa (fig. 5E). To examine if Mjl is expressed



### G The "Destruction Complex"




FIG. 3.—Ectopic Mjl shows Sgg-like genetic interactions. Wings from adults of the following genotypes are shown: (A) Wild type, (B) *vg-GAL4/UAS-sgg*<sup>WDT7A</sup>, (C) *vg-GAL4/UAS-mjl*<sup>14B</sup>, (D) *+/DpY*<sup>armYD35</sup>; *vg-GAL4/UAS-mjl*, (E) *vg-GAL4/UAS-mjl; dAxin*<sup>1 [3] SO44230/+</sup>, and (F) *vg-GAL4/UAS-mjl*<sup>14B/UAS-mjl</sup><sup>RNAi10.2</sup>. Ectopic expression of *mjl* in the wing margin results in a notched wing phenotype, which is similar to the phenotype seen with *sgg* overexpression. This phenotype can be suppressed by the reduction of dAxin levels (*dAxin* heterozygous) or by the increase of Arm levels (*DpY*<sup>arm</sup> duplication of *arm* on the Y), indicating that Mjl interacts with components of the wg pathway. Arm (yellow circle) is phosphorylated and targeted for degradation by a collection of proteins that comprise the “destruction complex.” These proteins include Sgg (red box), dAxin (blue box), and antigen-presenting cells (green box). Once Arm is phosphorylated, it is targeted for degradation via the ubiquitin–proteasome pathway. In the presence of wg signal, Arm escapes phosphorylation translocates into the nucleus where it activates wg target genes.

in somatic cells within the male gonad, we stained Enhancer trap line 3914 with anti-β-galactosidase and anti-Mjl antibodies. Enhancer trap line 3914 (Asaoka et al. 1998) marks the somatic cyst cells and hub cells in the gonad and Mjl does not colocalize with these cells (fig. 5D). We also immunostained wild-type testes with anti-Fas III antibody, which strongly outlines the somatic hub cells at the apical tip (Le Bras and Van Doren 2006), and Mjl is clearly excluded from those cells (data not shown). These data indicate that Mjl expression is restricted to the male germ line. This is highly significant in light of the fact that Sgg is expressed in multiple somatic tissues in both sexes (Bourouis et al. 1990; Siegfried et al. 1990). Mjl evolving a tissue-specific expression pattern is highly reminiscent of a model for evolution of new protein function following restriction of expression to specific tissue types (Hughes 1994).




FIG. 4.—Mjl expression rescues the segmentation defect of *sgg* mutant embryos. Cuticle phenotypes of (A) wild-type embryo (B) *sgg* mutant embryo derived from a germ line clone, (C) *sgg* mutant embryo ectopically expressing a *sgg* rescue construct (*sgg; arm-GAL4/UAS-sgg<sup>WDT7A</sup>*), (D) *sgg* mutant embryo ectopically expressing a *mjl*-rescuing construct (*sgg; arm-GAL4/UAS-mjl<sup>4B</sup>*). (E) Comparison of efficiency of *sgg* mutant embryo rescue by the *UAS-sgg* transgene and *UAS-mjl* transgene.

*mjl* Is Required for Male Germ Line Survival

Although evolutionary conservation is a strong indicator of function, we were interested in directly testing for *mjl* genetic function. We utilized RNAi to disrupt *mjl* mRNA expression (Kennerdell and Carthew 2000). We generated *mjl* transgenic lines that express dsRNA under control of the GAL4/UAS system (see Methods). The *UAS-mjl<sup>RNAi</sup>* constructs strongly downregulate ectopically expressed *mjl* mRNA and Mjl protein and do not affect *sgg* function, mRNA expression, or Sgg protein levels (fig. 3F, additional data not shown).

Our studies on the endogenous expression pattern of Mjl establish that it is expressed in the male germ line. Hence, we used the nos-GAL4 driver to drive expression of *UAS-mjl<sup>RNAi</sup>* in the germ line. We also used a heat-shock GAL4 driver and obtained similar results (data not shown) indicating that the RNAi construct did not have overt off-target effects. The nos-GAL4 driver has been shown to activate expression in germ cells at the apical region of the testes, including stem cells, gonialblasts, spermatogonia, and early spermatocytes (Schulz et al. 2004).




FIG. 5.—Mjl is enriched in testis and is required for male germ line viability. (A) Western blots containing samples as indicated. The broken line signifies that intervening lanes are not shown although all lanes correspond to a single blot. Wild-type male gonads (B) and female adult gonads (C) immunostained for anti-Mjl (green) and anti-Vasa (red). Nuclei are counterstained with DAPI (blue). Overlap of Vasa and Mjl can be seen (yellow). (D) Testis from line 3914 immunostained for anti-Mjl (green) and anti-β-galactosidase (red). The asterisk marks the hub. (E) Wild-type and (F) *UAS-mjl<sup>RNAi11.1</sup>* testis immunostained for anti-Mjl (green) and anti-Vasa (red). Nuclei counterstained with DAPI (blue).

*UAS-mjl<sup>RNAi</sup>/+nos-GAL4/+* males were mated to wild-type virgin females and the progeny were examined. The Mjl-depleted males were sterile. Female siblings as well as control males lacking either one of the UAS/GAL4 binary system components were fertile. These observations indicate that Mjl is crucial for maintaining male fertility in *D. melanogaster*.

To explore the cause of male sterility, testes of Mjl-depleted males were dissected and examined. Mjl-depleted

males have grossly abnormal testes morphology when compared with wild-type males. The testes of Mjl-depleted males are stunted, as is seen in testes lacking a germ line (compare panels in fig. 5E and F). In a wild-type testis, Vasa-positive cells (germ line lineage) fill the testis (fig. 5E), whereas Mjl-depleted testes lack all germ cells (fig. 5F). The DAPI counterstain highlights the outer layer of sheath cells, which are the only cells left in the testes of Mjl-depleted adults. The Mjl-depleted testes often showed a complete loss of Vasa-positive cells, but a few isolated Vasa-positive cells were occasionally observed. These results showed that sterility was due to the loss of the entire germ line lineage in the adult testes of Mjl-depleted flies and that even when a few germ cells survive they are incapable of completing gametogenesis. This strongly suggests that Mjl has assumed an essential role for spermatogenesis in *D. melanogaster*.

Our analysis also indicates that Mjl has acquired a specific expression pattern and new function in the male germ line. Unlike *sgg*, which is required in a plethora of cell types and developmental stages, Mjl is only required in the male germ line. This can be inferred from the fact that global RNA interference of Mjl (using hs-GAL4) caused a very specific defect loss of the germ line in males. Taken together, our data argue for neofunctionalization of Mjl.

## Discussion

Retroposition is a means by which organisms can expand their genomes (Brosius 1991). Here we demonstrate that a retroposed gene, *mjl*, has acquired an important function in *Drosophila*. The retroposition itself appears to have occurred near, or in, the last common ancestor of the *Drosophila* genus. The crucial function of *mjl* in maintaining male fertility is probably the reason it has been faithfully retained in every species of the genus examined.

### Divergence of *mjl* and *sgg*

The most highly conserved domain between Mjl and Sgg is the kinase domain, which appears to be under strict evolutionary constraint. Structural conservation extends into functional relevancy, for when Mjl is placed in Sgg-centric contexts within *D. melanogaster*, it is capable of producing Sgg-like phenotypes. Thus, the core protein function is conserved in the retroposed copy following gene duplication. Does this highlight a lack of functional diversification or highlight the conserved core function of GSK-3 family members? The fact that Mjl, which was derived from just one of several isoforms of GSK-3 encoded by *sgg* over 50 MYA, rescues *sgg* mutants so well is striking, but not without precedent. Rat GSK-3 can also rescue *sgg* mutants (Siegfried et al. 1992). Clearly, the Sgg and Mjl proteins are under selection in the *Drosophila* genus. Therefore, we suggest that such cross-rescue is more a reflection of the large error in measuring phenotypes in a lab setting. Although we did see that altered rescue efficacy is consistent with Sgg being better adapted for *sgg* functions than Mjl, these transgenic experiments are inherently fraught with technical difficulties.

The best evidence for functional divergence is in the spatial expression (testes specific) as well as its sexually di-

morphic expression (male-specific) pattern of Mjl and the largely uniform expression of Sgg. The GSK-3 kinase domain is used in multiple signaling pathways (Cohen and Frame 2001; Frame and Cohen 2001; Eldar-Finkelman 2002), but is not known to function in the male germ line. Mjl, on the other hand, is required only for maintaining male fertility. These data argue for *mjl* having undergone neofunctionalization following duplication (Hughes 1994; Lynch and Conery 2000; Zhang et al. 2004).

It is curious that the *Ks* values for Mjl and Sgg are not similar, as expected under nearly neutral models of DNA sequence evolution (Ohta 2002). It is possible that this is a reflection of selection that is independent of amino acid coding (Xing and Lee 2006). It is well documented that *sgg* “pre-mRNAs” undergo differential splicing giving rise to various functional isoforms, whereas the *mjl* pre-mRNAs are devoid of introns and therefore do not undergo posttranscriptional splice modifications (Bourouis et al. 1990). Higher *Ks* rates in Mjl might be the result of relaxed selection at sites that have a regulatory role in processing in *sgg* pre-mRNAs.

### Selection or Chance

One of the more fascinating aspects of retroposition is the highly directional patterns of both distribution in the genome and the expression pattern of the daughter gene. In both *Drosophila* and mammals, the parental gene in a retroposed pair is usually located on the X chromosome and the daughter gene is autosomal (Betran et al. 2002; Emerson et al. 2004). As a general rule, retroposed genes in both mammals (Hisano et al. 2003; Rohozinski and Bishop 2004; Luo et al. 2006; Vinckenbosch et al. 2006) and *Drosophila* (Betran et al. 2002) tend to be expressed in testes. This preference is startling. In *D. melanogaster*, genes are preferential retroposed from the X chromosome onto autosomes (40% observed vs. 23.3% assuming random retroposition), where the majority (91%) of the retroposed genes evolved a testes-specific expression pattern (Betran et al. 2002). This pattern of expression along with growing evidence that the X chromosome is a disfavored location for genes with male-biased function (Parisi et al. 2003; Wu and Xu 2003) suggests that the 2 observations may be linked. Retroposition patterns may indicate that important male functions are being transferred to more favorable long-term locations in the genome (Betran et al. 2002, 2004).

In contrast to such functional movements, it is also possible that the testis is a permissive environment for gene expression (Vinckenbosch et al. 2006). Male-biased expression of retroposed genes is therefore a function of chance in this model. This is not well supported by global expression studies in *D. melanogaster*, which show a complex but highly specific pattern of testes gene expression (Andrews et al. 2000; Parisi et al. 2003, 2004). However, if genes retropose more frequently in the male germ line and then are preferential inserted due to the open chromatin structure of genes being highly expressed in the male germ line, then testis expression would also be expected (Khil et al. 2005).

If testes-biased expression of retroposed genes is neutral, then expression within the male germ line should not

be evolutionarily stable. Indeed, a recent report shows that retroposed genes in primates initially tend to be transcribed in testes, but over time they tend to “come out” of the testes and be expressed more broadly (Vinckenbosch et al. 2006). The *mjl* gene does not fit this pattern, because it is an older retroposed gene, but its expression remains highly testes biased, and in fact, may be expressed specifically in the male germ line of *D. melanogaster*. Our data support the idea that retroposition may be one of the modes employed by genes with male-biased functions to emigrate from the disfavored X chromosome to an autosomal location.

### Acknowledgments

We thank K. Howard, H. Lin, G. Thomas, K. Han, and A. Clark for reagents and flies. We thank M. Long, J. Dean, and members of the Oliver lab for critical reading of the manuscript and helpful discussions. We owe great thanks to providers of genomic sequence data: *mjl* and *sgg* sequences are from draft annotations (Iyer VN, Pollard DA, Eisen MB, personal communication) of assemblies generated by Agencourt (Smith D, personal communication) and Washington University (Wilson R, personal communication) genome sequencing centers. This research was supported, in part, by the Intramural Research Program of the National Institutes of Health and the National Institutes of Diabetes & Digestive & Kidney Diseases.

### Literature Cited

- Adams MD, Celniker SE, Holt RA, et al. (195 co-authors). 2000. The genome sequence of *Drosophila melanogaster*. *Science*. 287:2185–2195.
- Andrews J, Bouffard GG, Cheadle C, Lu J, Becker KG, Oliver B. 2000. Gene discovery using computational and microarray analysis of transcription in the *Drosophila melanogaster* testis. *Genome Res.* 10:2030–2043.
- Asaoka M, Sano H, Obara Y, Kobayashi S. 1998. Maternal Nanos regulates zygotic gene expression in germline progenitors of *Drosophila melanogaster*. *Mech Dev.* 78:153–158.
- Betran E, Emerson JJ, Kaessmann H, Long M. 2004. Sex chromosomes and male functions: where do new genes go? *Cell Cycle.* 3:873–875.
- Betran E, Long M. 2002. Expansion of genome coding regions by acquisition of new genes. *Genetica.* 115:65–80.
- Betran E, Thornton K, Long M. 2002. Retroposed new genes out of the X in *Drosophila*. *Genome Res.* 12:1854–1859.
- Bourouis M, Moore P, Ruel L, Grau Y, Heitzler P, Simpson P. 1990. An early embryonic product of the gene shaggy encodes a serine/threonine protein kinase related to the CDC28/cdc2+ subfamily. *EMBO J.* 9:2877–2884.
- Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development.* 118:401–415.
- Brent R, Kingston RE, Seidman JG, Struhl K, Ansel FM, Chanda VB, Moore DD. 1987. Current protocols in molecular biology. Hoboken (NJ): Wiley and Sons.
- Brosius J. 1991. Retroposons—seeds of evolution. *Science.* 251:753.
- Burnett C, Howard K. 2003. Fly and mammalian lipid phosphate phosphatase isoforms differ in activity both in vitro and in vivo. *EMBO Rep.* 4:793–799.
- Celniker SE, Wheeler DA, Kronmiller B, et al. (32 co-authors). 2002. Finishing a whole-genome shotgun: release 3 of the *Drosophila melanogaster* euchromatic genome sequence. *Genome Biol.* 3:RESEARCH0079.
- Clamp M, Cuff J, Searle SM, Barton GJ. 2004. The Jalview Java alignment editor. *Bioinformatics.* 20:426–427.
- Cohen P, Frame S. 2001. The renaissance of GSK3. *Nat Rev Mol Cell Biol.* 2:769–776.
- Dajani R, Fraser E, Roe SM, Young N, Good V, Dale TC, Pearl LH. 2001. Crystal structure of glycogen synthase kinase 3 beta: structural basis for phosphate-primed substrate specificity and autoinhibition. *Cell.* 105:721–732.
- Eldar-Finkelman H. 2002. Glycogen synthase kinase 3: an emerging therapeutic target. *Trends Mol Med.* 8:126–132.
- Emerson JJ, Kaessmann H, Betran E, Long M. 2004. Extensive gene traffic on the mammalian X chromosome. *Science.* 303:537–540.
- Fabrizio JJ, Boyle M, DiNardo S. 2003. A somatic role for eyes absent (*eya*) and sine oculis (*so*) in *Drosophila* spermatocyte development. *Dev Biol.* 258:117–128.
- FlyBase. 2003. The FlyBase database of the *Drosophila* genome projects and community literature. *Nucleic Acids Res.* 31:172–175.
- Frame S, Cohen P. 2001. GSK3 takes centre stage more than 20 years after its discovery. *Biochem J.* 359:1–16.
- Fuller MT. 1993. Spermatogenesis. In: Bate M, Martinez-Arias A, editors. The development of *Drosophila*. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press. p. 71–148.
- Hisano M, Yamada S, Tanaka H, Nishimune Y, Nozaki M. 2003. Genomic structure and promoter activity of the testis haploid germ cell-specific intronless genes, *Tact1* and *Tact2*. *Mol Reprod Dev.* 65:148–156.
- Hughes AL. 1994. The evolution of functionally novel proteins after gene duplication. *Proc Biol Sci.* 256:119–124.
- Hurst LD. 2002. The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends Genet.* 18:486.
- Jope RS, Johnson GV. 2004. The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem Sci.* 29:95–102.
- Kennerdell JR, Carthew RW. 2000. Heritable gene silencing in *Drosophila* using double-stranded RNA. *Nat Biotechnol.* 18:896–898.
- Khil PP, Oliver B, Camerini-Otero RD. 2005. X for intersection: retrotransposition both on and off the X chromosome is more frequent. *Trends Genet.* 21:3–7.
- Kiger AA, Jones DL, Schulz C, Rogers MB, Fuller MT. 2001. Stem cell self-renewal specified by JAK-STAT activation in response to a support cell cue. *Science.* 294:2542–2545.
- Kumar S, Tamura K, Nei M. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* 5:150–163.
- Le Bras S, Van Doren M. 2006. Development of the male germline stem cell niche in *Drosophila*. *Dev Biol.* 294(1):92–103.
- Li WH, Yang J, Gu X. 2005. Expression divergence between duplicate genes. *Trends Genet.* 21:602–607.
- Long M, Betran E, Thornton K, Wang W. 2003. The origin of new genes: glimpses from the young and old. *Nat Rev Genet.* 4:865–875.
- Long M, Deutsch M, Wang W, Betran E, Brunet FG, Zhang J. 2003. Origin of new genes: evidence from experimental and computational analyses. *Genetica.* 118:171–182.
- Luo C, Lu X, Stubbs L, Kim J. 2006. Rapid evolution of a recently retroposed transcription factor YY2 in mammalian genomes. *Genomics.* 87:348–355.
- Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science.* 290:1151–1155.
- Nei M, Gojobori T. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol Biol Evol.* 3:418–426.

- Nei M, Kumar S. 2000. Molecular evolution and phylogenetics. Oxford: Oxford University Press.
- Ohta T. 2002. Near-neutrality in evolution of genes and gene regulation. *Proc Natl Acad Sci USA*. 99:16134–16137.
- Parisi M, Nuttall R, Edwards P, et al. (12 co-authors). 2004. A survey of ovary-, testis-, and soma-biased gene expression in *Drosophila melanogaster* adults. *Genome Biol*. 5:R40.
- Parisi M, Nuttall R, Naiman D, Bouffard G, Malley J, Andrews J, Eastman S, Oliver B. 2003. Paucity of genes on the *Drosophila* X chromosome showing male-biased expression. *Science*. 299:697–700.
- Robertson HM, Preston CR, Phillis RW, Johnson-Schlitz DM, Benz WK, Engels WR. 1988. A stable genomic source of P element transposase in *Drosophila melanogaster*. *Genetics*. 118:461–470.
- Rohozinski J, Bishop CE. 2004. The mouse juvenile spermatogonial depletion (jsd) phenotype is due to a mutation in the X-derived retrogene, mUtp14b. *Proc Natl Acad Sci USA*. 101:11695–11700.
- Rongo C, Broihier HT, Moore L, Van Doren M, Forbes A, Lehmann R. 1997. Germ plasm assembly and germ cell migration in *Drosophila*. *Cold Spring Harbor Symp Quant Biol*. 62:1–11.
- Russo CA, Takezaki N, Nei M. 1995. Molecular phylogeny and divergence times of drosophilid species. *Mol Biol Evol*. 12:391–404.
- Schulz C, Kiger AA, Tazuke SI, Yamashita YM, Pantalena-Filho LC, Jones DL, Wood CG, Fuller MT. 2004. A misexpression screen reveals effects of bag-of-marbles and TGF beta class signaling on the *Drosophila* male germ-line stem cell lineage. *Genetics*. 167:707–723.
- Siegfried E, Chou TB, Perrimon N. 1992. Wingless signaling acts through zeste-white 3, the *Drosophila* homolog of glycogen synthase kinase-3, to regulate engrailed and establish cell fate. *Cell*. 71:1167–1179.
- Siegfried E, Perkins LA, Capaci TM, Perrimon N. 1990. Putative protein kinase product of the *Drosophila* segment-polarity gene zeste-white3. *Nature*. 345:825–829.
- Siegfried E, Wilder EL, Perrimon N. 1994. Components of wingless signalling in *Drosophila*. *Nature*. 367:76–80.
- Stanewsky R. 2002. Clock mechanisms in *Drosophila*. *Cell Tissue Res*. 309:11–26.
- Stapleton M, Carlson J, Brokstein P, et al. (13 co-authors). 2002. A *Drosophila* full-length cDNA resource. *Genome Biol*. 3: RESEARCH0080.
- Steitz MC. 2000. An investigation of Zeste white-3 inactivation by wingless signalling. University Park (PA): Department of Biochemistry, Microbiology and Molecular Biology. The Pennsylvania State University. p. 173.
- Steitz MC, Wickenheisser JK, Siegfried E. 1998. Overexpression of zeste white 3 blocks wingless signaling in the *Drosophila* embryonic midgut. *Dev Biol*. 197:218–233.
- Tamura K, Subramanian S, Kumar S. 2004. Temporal patterns of fruit fly (*Drosophila*) evolution revealed by mutation clocks. *Mol Biol Evol*. 21:36–44.
- ter Haar E, Coll JT, Austen DA, Hsiao HM, Swenson L, Jain J. 2001. Structure of GSK3beta reveals a primed phosphorylation mechanism. *Nat Struct Biol*. 8:593–596.
- Tulina N, Matunis E. 2001. Control of stem cell self-renewal in *Drosophila* spermatogenesis by JAK-STAT signaling. *Science*. 294:2546–2549.
- Vinckenbosch N, Dupanloup I, Kaessmann H. 2006. Evolutionary fate of retroposed gene copies in the human genome. *Proc Natl Acad Sci USA*. 103:3220–3225.
- Wang W, Yu H, Long M. 2004. Duplication-degeneration as a mechanism of gene fission and the origin of new genes in *Drosophila* species. *Nat Genet*. 36:523–527.
- Wernersson R, Pedersen AG. 2003. RevTrans: multiple alignment of coding DNA from aligned amino acid sequences. *Nucleic Acids Res*. 31:3537–3539.
- Woodgett JR. 1990. Molecular cloning and expression of glycogen synthase kinase-3/factor A. *EMBO J*. 9:2431–2438.
- Wu CI, Xu EY. 2003. Sexual antagonism and X inactivation—the SAXI hypothesis. *Trends Genet*. 19:243–247.
- Xing Y, Lee C. 2006. Can RNA selection pressure distort the measurement of Ka/Ks? *Gene*. 370:1–5.
- Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution rates under realistic evolutionary models. *Mol Biol Evol*. 17:32–43.
- Zhang J, Dean AM, Brunet F, Long M. 2004. Evolving protein functional diversity in new genes of *Drosophila*. *Proc Natl Acad Sci USA*. 101:16246–16250.
- Zhang J, Rosenberg HF, Nei M. 1998. Positive Darwinian selection after gene duplication in primate ribonuclease genes. *Proc Natl Acad Sci USA*. 95:3708–3713.

Koichiro Tamura, Associate Editor

Accepted December 12, 2006