

The Effect of Increasing Volume of Exercise on Activation Pattern of Vastus Medialis and Lateralis and its Correlation With Anterior Knee Pain in Karate Elites

Afsaneh Safar Cherati,^{1,2} Sara Lotfian,^{1,2,*} Aliashraf Jamshidi,³ Mohammad Ali Sanjari,⁴ and Mohammad Razi¹

¹Rasoul Akram Medical Complex, Iran University of Medical Sciences, Tehran, IR Iran

²Sports Medicine Research Center, Tehran University of Medical Sciences, Tehran, IR Iran

³Rehabilitation Research Center, Department of Physical Therapy, Faculty of Rehabilitation Science, Iran University of Medical Sciences, Tehran, IR Iran

⁴Biomechanics Lab, Rehabilitation Research Center, Department of Rehabilitation, Department of Rehabilitation Basic Sciences, Iran University of Medical Sciences, Tehran, IR Iran

*Corresponding author: Sara Lotfian, Rasoul Akram Medical Complex, Iran University of Medical Sciences, Tehran, IR Iran. Tel: +98-2164352446, Fax: +98-2166517309, E-mail: saralotfiyan@yahoo.com

Received 2015 December 08; Revised 2016 January 08; Accepted 2016 January 17.

Abstract

Background: The effects of exercise volume on the pattern of muscle activity is one of the most important factors in training management and injury risk reduction. In the lower limb, the quadriceps muscle which plays a determining role in performing the stance and other karate techniques could be injured in intensive exercise and may induce anterior knee pain in athletes.

Objectives: The aim of this study was to determine the relationship between training volume and muscle activity of vastus medialis and vastus lateralis and its association with anterior knee pain in karate elites.

Patients and Methods: Male and female athletes from national junior and cadet karate team (14 to 18 years) were invited to participate in the study at the beginning and the end of the training camps. Studies involved measurement of electromyographic muscle activity of vastus medialis and vastus lateralis in both lower extremities with surface electromyography device and assessment of movement by electrogoniometry. Muscle activity was recorded in three tests of dachi, walking up and walking down stairs. Simultaneously, anterior knee pain was evaluated using visual analogue scale and anterior knee pain scale questionnaire.

Results: Eight athletes of a total number of 23 reported increased ratings of pain in their right knees. No differences in muscle activity were observed in tests of Dachi and stairs between the groups with and without pain. Comparing Dachi task pattern at the beginning and end of training camps, there was no significant difference in pattern of biomechanical movement; however, reducing the amount of muscle activity in early and late phases of tasks was observed in electromyographic assessment.

Conclusions: The results showed that performing the same task after a six-week training period, less muscle activity was required in all phases in two groups of tasks, including karate-specific movement (dachi) and activities of daily living (up or down stairs).

Keywords: Vastus Medialis, Vastus Lateralus, Exercise Volume, Karate

1. Background

A comprehensive understanding of muscle activity pattern during execution of different maneuvers and techniques is one of the most important factors in set-out of exercise volume and reducing the risk of injury in professional athletes. Among the lower limb muscles, quadriceps has a main role in implementation of karate techniques and can be susceptible to overtraining addressed by anterior knee pain. Anterior knee pain is a common debilitating complaint in athletes and one of the most important differential diagnoses is patellofemoral pain syndrome (PFPS) (1, 2).

Although PFPS is the most common cause of knee pain, many studies have been conducted to investigate the underlying pathomechanics (1). Several causes are involved in the pathogenesis of the syndrome, which is mainly divided into two categories: internal and external factors (1). Imbalanced forces exerted on patella by VMO and VL, improper activation time, tightness of lateral retinaculum, hypermobile patella and abnormal patellar tracking due to structural malalignment in lower extremity are the most important internal factors. Kinematic errors, increase in the duration and frequency of physical activity and training volume are external factors (1). One of the

highly regarded basic pathomechanics in PFPS is delay in the activation of vastus medialis (VM) to vastus lateralis (VL) which in theory could induce laterally pulled patella and anterior knee pain (3, 4).

Surface electromyography (sEMG) is a non-invasive technique for measuring electrical activity of muscles. Moreover, it reveals the onset and amplitude of muscle activity during voluntary dynamic contraction (5). Although most of the previous studies investigated the difference in onset timing between the VM and VL, some sEMG studies have shown that the ratio of the electrical activity of the VM to VL is different in healthy subjects versus patients with patellofemoral pain (6-11).

The role of overload on VM/VL activity was evaluated by Herrington et al. They assessed sEMG activity of muscles during different load (12) but few prospective studies have been conducted on PFPS in athletes.

PFPS is common in karate mainly due to overstress on knee. Proper execution of techniques in this sport supplements muscle balance and strengths (13-15).

Allegorically, stand or dachi indicates lower part of the body position, which is used to carry the upper part of body. For correct technique, dachi should be done harmonically and strongly. Many techniques in karate overload the patellofemoral joint; therefore, an increase in the volume of training can lead to the development of anterior knee pain. Due to the negative impact of pain on athletic performance, identification and correction of pathomechanical factors could improve the function of professional athletes.

A prospective study by Witvrouw et al. showed that one of the risk factors of patellofemoral syndrome is rescheduling muscle reflex of VM (16). However, there are also studies with inconsistent results concerning VM/VL ratio and the timing onset of the muscles, including imbalance in the electric activity and abnormal recruitment pattern, with greater delay and lower amplitude of activation of the VM (17-26). The meta-analyses by Chester (20) and Wong (21) demonstrated heterogeneous finding regarding any correlation between timing of activation of VMO to VL and anterior knee pain.

One of the major obstacles in investigating the role of imbalance between activation of VM and VL in PFPS is the temporal priority of knee pain or delay of VM activation. Hodges et al. found that pain is the source not the consequence of muscle imbalance. Thus, anterior knee pain control is fundamental during exercise (27).

The main objective of this study was to determine the relationship between training volume and VM and VL activity and its correlation to anterior knee pain in elite karate athletes. It deems that amplitude and pattern of muscle recruitment could be changed regarding training volume.

Furthermore, it would be aimed to investigate other functional parameters of electromyographic properties of VM and VL as well as time domain index and electrogoniometry.

2. Objectives

Regarding limited evidence in karate, this type of study deems essential for improving training programs and reducing the risk of injury in elites. Understanding the factors in the development of knee pain and altered muscle activity during high-volume training in professional athletes could be helpful in definition of prevention and treatment strategies.

3. Patients and Methods

3.1. Subjects

All junior and cadet athletes (14-18 years old) of Iranian national karate team were recruited to participate in this quasi-experimental study. They were required to provide informed consent.

The subjects were included if they had no history of significant injury to the lower extremities; no history of patellar subluxation or meniscal or ligamentous injury; and normal radiological findings. They were excluded if the athlete left the team due to being crossed out. A total of 29 subjects, including 15 boys and 14 girls, took part in the first session of assessment; however, after 6 weeks of training, 23 of them-10 boys and 13 girls- had the criteria to complete the study.

3.2. Procedure

After arrival at the laboratory, data of acute and recent knee pain was recorded by a 10-cm visual analog scale (VAS) and anterior knee pain scale (AKPS) questionnaire, respectively (28, 29). The skin was prepared by shaving, if necessary, and the subject started the session by a 10-minute warm-up by a stationary bike and quadriceps and hamstring exercises and a few performance trials. It was explained to the athletes to keep the intensity of warm-up including bike-riding at a comfortable level and keep their heart rate less than 100 bpm. Quadriceps and hamstring exercises consisted of hip and knee flexion and extensions with no resistance and each exercise was repeated 10 times. Each participant practiced dachi and stair-up and down tasks 2 or 3 times, intended for familiarization with the process. Then the skin was cleaned by isopropyl alcohol and surface electrodes were set according to SENIAM recommendations (30). Silver/silver chloride disposable

electrodes (Skintact, Austria) were placed with an inter-electrode distance of 30 mm on VM and VL muscles of right and left thigh.

Electrogoniometer was positioned on the lateral aspect of right knee and centered on lateral femoral condyle. It was attached to the lower limb by a double-sided adhesive stick. Movement of right knee was continuously measured by the electrogoniometer. The sEMG data was synchronously sampled using a Mega receiver (ME 6000, Mega Electronics Ltd., Kuopio, Finland). Obtaining data from electrogoniometer, the points of changes in flexion and extension of right knee were determined on the plot. The intervals between the points were defined as phases of tasks. Signals were collected from 6 channels, including 4 channels for sEMG electrodes and 2 channels for electrogoniometer. Signals of sEMG were pre-amplified 10 times and raw data was collected without filtering at a sampling frequency of 1000 Hz. The data was recorded in Megawin software, version 2.4.

Data was collected under three different conditions: 1, Zenkutsu-dachi: with the left foot forward, with hearing the command, the subjects moved forward at a self-selected pace (31). Data was recorded simultaneously; 2, climb up and down the stairs: subjects were instructed to come up and down three steps with a height of 20 cm for each step, placing only one foot on each step, starting with the left foot. The stepping rate was adjusted at 96 steps per minute which is, approximately, the usual stepping rate (32). Ascending and descending trials were performed separately; 3, maximal voluntary isometric contraction (MVIC) of quadriceps, which was an isometric contraction done after dynamic tasks. It was performed in sitting position with full knee extension. The subject was requested to keep their muscle contracted as forcefully as possible for 5 seconds. They were encouraged by continuous verbal commands (Prepare! Go! Hold! Hold! Relax!).

At the end of each test signal accuracy and absence of noise were reviewed. The tests were repeated to record three sets of acceptable signals. All tests have been taken by one person (second author S.L (sports medicine specialist)).

3.3. Data Analysis

During later off-line analysis, raw data was first visually checked in Megawin 2.4 to select one of three repetitions of each task with minimal noise and best signal accuracy. The selected file was saved as a binary file in MATLAB for further analysis.

The tasks were divided into different phases. In Dachi three phases of movement were determined which is discussed in another article. The phases correspond to the beginning of the movement, maximum flexion and return to

extension respectively (31). Step up and down phases are shown in Figure 1. The frequency spectrum was assessed in these phases. For each individual muscle maximal mean root mean square (RMS) of activity (50 millisecond time-constant) was used to identify peak sEMG. Mean RMS of amplitude of muscle activity of each one of the four muscles was calculated. Then it was normalized to the maximal RMS data obtained during MVIC, as it is the preferred method of normalization (33, 34). To investigate the level of activity of the muscles, VM/VL ratio was measure in all tasks as:

- If it was more than 1, means VM had more activity during the specified phase of task.
- If it was equal to 1, VM and VL had the same level of activity.
- If it was less than 1, means VM had less activity during the specified phase of task.

3.4. Statistical Analysis

Data was analyzed using the statistical software package SPSS (version 14). To compare variables with normal distribution parametric tests and to compare variables with non-normal distribution, non-parametric test of Wilcoxon signed rank test was used with statistical significance set as 0.05.

4. Results

The mean age of subjects was 16.10 ± 1.26 years and the mean BMI was 20.86 ± 0.40 . Their BMI did not change during the study because the athletes should stay in their weight categories. The training course lasted 6 weeks, with a training volume equivalent to 16.5 hours per week.

Acute pain score, measured by VAS, increased significantly ($P < 0.001$), from a mean (SD) of 0.61 (1.29) to 1.87 (1.89). However, the changes in AKPS score were not significant ($P = 0.080$). Therefore, sEMG data was compared between participants with and without increase in acute knee pain. No significant difference was found in all 4 muscles activity between these groups, at the beginning and at the end of training course. In intra-group comparison of participants who experienced acute knee pain there was no significant change in muscle amplitude in different tasks.

Comparing the movement pattern of dachi, there was no change in motion at the beginning and the end of the camp. It was expected because of their proficiency and longtime of training karate (31). Normalized muscle activity in the task of dachi was compared at the beginning and end of 6 weeks of training. The results are presented in Table 1, which shows a decrease in total muscle activity that is




Figure 1. Phasing EMG and Electro-Goniometer Signals During Walk Up The Stairs in MATLAB

significant in right VL and left VM. When comparing amplitude of muscle activity in the first phase of the movement (dachi), decrease in muscle activity was statistically significant in right VM ($P = 0.002$), right VL ($P = 0.007$) and left VM ($P = 0.042$) except for left VL ($P = 0.346$), after 6 weeks of training.

Table 1. Comparison of Normalized Muscle Activity in Dachi at the Beginning and End of Training^a

Muscle Activity	At the Beginning	After 6 Weeks	P Value
RVM	1.77 (1.37)	1.29 (1.30)	0.078
RVL	1.56 (1.34)	0.98 (0.81)	0.010*
LVM	0.93 (0.65)	0.72 (0.45)	0.026*
LVL	1.04 (0.74)	0.78 (0.49)	0.089

Abbreviations: LVL, left vastus lateralis; LVM, left vastus medialis; RVM, right vastus medialis; RVL, right vastus lateralis.

^aValues are expressed as mean (SD).

Total muscle activity during the tasks of stair-up and

down was compared at the beginning and end of 6 weeks of training. The results are presented in Tables 2 and 3 respectively.

Table 2. Comparison of Normalized Muscle Activity in Stair Up at the Beginning and End of Training^a

Muscle Activity	At the Beginning	After 6 Weeks	P Value
RVM	0.67 (0.66)	0.40 (0.33)	0.050*
RVL	0.59 (0.65)	0.36 (0.26)	0.011*
LVM	0.59 (0.57)	0.28 (0.13)	0.004*
LVL	0.48 (0.33)	0.30 (0.13)	0.036*

^aValues are expressed as mean (SD).

Muscle activity was also compared before and after the training period in weight-bearing phases of stair tasks, which were separated according to data of electrogoniometry. Weight-bearing phases of right lower limb were 2 and 4 ones in stair-up and 1 and 3 in stair-down task (Figure 1). Decrease in normalized amplitude of muscle during

Table 3. Comparison of Normalized Muscle Activity in Stair Down at the Beginning and End of Training^a

Muscle Activity	At the Beginning	After 6 Weeks	P Value
RVM	0.56 (0.68)	0.34 (0.37)	0.039*
RVL	0.48 (0.50)	0.30 (0.31)	0.012*
LVM	0.51 (0.47)	0.26 (0.27)	0.005*
LVL	0.47 (0.43)	0.34 (0.35)	0.026*

^aValues are expressed as mean (SD).

weight-bearing phases was significant in right VL in stair-up ($P = 0.033$) and stair-down ($P = 0.023$). Muscle activity of right VM decreased in the specified phases of stair-up ($P = 0.036$) and stair-down ($P = 0.036$). For the left lower limb weight-bearing phases consisted of 1 and 3 ones in stair-up and 2 and 4 in stair-down task. Decrease of muscle activity was statistically significant in left VM in these phases of stair-up ($P = 0.039$) and stair-down ($P = 0.013$).

VM/VL ratio in right and left quadriceps was compared at the beginning and end of 6 weeks of training which is presented in **Table 4**. As it is shown, the changes of ratio was not statistically significant.

5. Discussion

This study aimed to investigate correlation between training volume and activity of the VM and VL and its contribution to anterior knee pain in karate elites.

Although several studies have evaluated the electromyographic activity of VMO and VL, few of them settled activity level and muscle contraction in their analysis. In the present study, the activity level was appraised as well as time domain index. Goniometric measurement was applied for evaluation of dynamic movement in athletes.

To our knowledge, this is the first time that the electromyographic activity of VMO and VL were addressed during a sport-specific task in karate, so the result could be applied for proper pattern of muscle recruitment and improving performance in elites.

One of the important variables in this study is the muscle activity in different phases of dachi. Decrease of muscle activity voltage in all phases of the dachi after six weeks of training was demonstrated and this reduction in the start and finish was significant. Regarding the first phase of movement, it would be hypothesized that after six weeks of training, the explosive start needs low levels of muscle activity and at the end of motion, control becomes possible with less muscle activity. Since changes in VM/VL ratio was not significant, it could be concluded that this reduction was uniform in both parts of quadriceps.

In the Irish study the effect of open and close kinetic chain exercises on VL to VMO activity was compared and it was demonstrated that lounge and squatting are effective in strengthening of VMO (35). As lounge is similar to dachi in karate, their results are consistent with our study.

According to Collado's study if there is no significant impairment in physical examination, the most important cause of anterior knee pain is overuse (1). This is a common phenomenon at national athletes who experience high volume of training.

Comparing two groups with and without increase in VAS Score, no difference was observed before and after six weeks of training in different phases of dachi. This inconsistent response might be due to short time follow up in our study. Previous studies showed that closed kinetic exercises improve sport performance and strength better than open kinetic knee exercises but open kinetic chain exercises could be applicable and effective in anterior knee pain rehabilitation. However, regarding the sample size, the clinical interpretation of the results should be done cautiously.

Hodges et al. found out that by inducing pain in the infrapatellar fat pad, muscle activity of VL decreased significantly during stair-up test (27). However in our study the decline in muscle activity was not different between the two with and without pain groups. However, regarding differences in methodology of the two studies, comparing deems difficult.

There are some limitations in our study. While this study had a sample containing the total available population, the sample size is still small. In order to decrease the effect of small sample size in future study, recruitment of more subjects can be helpful. Since this study was conducted with a six-week interval, it is proposed to perform biomechanical tests with more frequency and longer follow-up.

The advantages of the present study are subject population that is all professional athletes. The other strong point is the precise signal processing, evaluation of several functional parameters and electromyographic data analysis in MATLAB software.

Since prospective studies in this area, especially in athletes are limited, it is suggested to conduct basic and applied studies including kinematics and kinetics of specific movements of karate, featuring a variety of techniques of movement, kicks and hand strikes. Finally, long term follow up study of athletes can determine the impact of the training camp of the national team. The main challenge that which training program could improve the athlete's performance is also worth exploring.

The results showed that after a six-week training period for performing the same task, less muscle activity was

Table 4. Comparison of VM/VL Ratio at the Beginning and End of Training^a

Tasks	Right Quadriceps			Left Quadriceps		
	At the Beginning	After 6 Weeks	P Value	At the Beginning	After 6 Weeks	P Value
Dachi	1.29 (0.63)	1.31 (0.70)	0.951	0.97 (0.42)	1.50 (1.08)	0.078
Stair-up	1.34 (0.95)	1.20 (0.47)	0.685	1.33 (0.71)	0.95 (0.27)	0.058
Stair-down	1.40 (0.86)	1.19 (0.48)	0.330	1.22 (0.64)	0.97 (0.50)	0.144

^aValues are expressed as mean (SD).

required in all phases in both sport-specific movement (dachi) and activities of daily living (up or down stairs).

Acknowledgments

Sports sciences research institute of Iran and Iran Karate federation.

Footnotes

Authors' Contribution: Study concept and design, Sara Lotfian; Afsaneh Safar Cherati; acquisition of data, Sara Lotfian; analysis and interpretation of data, Sara Lotfian, Aliashraf Jamshidi, Mohammad Ali Sanjari; drafting of the manuscript, Sara Lotfian, Mohammad Razi; critical revision of the manuscript for important intellectual content, Sara Lotfian, Afsaneh Safar Cherati; statistical analysis, Sara Lotfian, Aliashraf Jamshidi, Mohammad Ali Sanjari; administrative, technical, and material support, Sara Lotfian, Afsaneh Safar Cherati; study supervision, Afsaneh Safar Cherati, Mohammad Razi.

Conflict of Interests: None declared.

References

- Collado H, Fredericson M. Patellofemoral pain syndrome. *Clin Sports Med.* 2010;**29**(3):379–98. doi: [10.1016/j.csm.2010.03.012](https://doi.org/10.1016/j.csm.2010.03.012). [PubMed: [20610028](https://pubmed.ncbi.nlm.nih.gov/20610028/)].
- Brunkner P, Brunkner K. Khan's clinical sports medicine. Australia: McGraw-Hill; 2012.
- Smith TO, Bowyer D, Dixon J, Stephenson R, Chester R, Donell ST. Can vastus medialis oblique be preferentially activated? A systematic review of electromyographic studies. *Physiother Theory Pract.* 2009;**25**(2):69–98.
- Cerny K. Vastus medialis obliquus/vastus lateralis muscle activity ratios for selected exercises in persons with and without patellofemoral pain syndrome. *Phys Ther.* 1995;**75**(8):672–83. [PubMed: [7644571](https://pubmed.ncbi.nlm.nih.gov/7644571/)].
- Drost G, Stegeman DF, van Engelen BG, Zwarts MJ. Clinical applications of high-density surface EMG: a systematic review. *J Electromyogr Kinesiol.* 2006;**16**(6):586–602. doi: [10.1016/j.jelekin.2006.09.005](https://doi.org/10.1016/j.jelekin.2006.09.005). [PubMed: [17085302](https://pubmed.ncbi.nlm.nih.gov/17085302/)].
- Souza DR, Gross MT. Comparison of vastus medialis obliquus: vastus lateralis muscle integrated electromyographic ratios between healthy subjects and patients with patellofemoral pain. *Phys Ther.* 1991;**71**(4):310–6. [PubMed: [2008454](https://pubmed.ncbi.nlm.nih.gov/2008454/)] discussion 317–20.
- Tang SF, Chen CK, Hsu R, Chou SW, Hong WH, Lew HL. Vastus medialis obliquus and vastus lateralis activity in open and closed kinetic chain exercises in patients with patellofemoral pain syndrome: an electromyographic study. *Arch Phys Med Rehabil.* 2001;**82**(10):1441–5. doi: [10.1053/apmr.2001.26252](https://doi.org/10.1053/apmr.2001.26252). [PubMed: [11588751](https://pubmed.ncbi.nlm.nih.gov/11588751/)].
- Qi Z, Ng GYF. EMG analysis of vastus medialis obliquus/vastus lateralis activities in subjects with patellofemoral pain syndrome before and after a home exercise program. *J Phys Ther Sci.* 2007;**19**(2):131–7.
- Pal S, Draper C, Fredericson M, Gold GE, Delp SL, Beaupre GS, et al. Patellar maltracking correlates with vastus medialis activation delay in patellofemoral pain patients. *Am J Sports Med.* 2011;**39**(3):590–8.
- Miao P, Xu Y, Pan C, Liu H, Wang C. Vastus medialis oblique and vastus lateralis activity during a double-leg semisquat with or without hip adduction in patients with patellofemoral pain syndrome. *BMC Musculoskelet Disord.* 2015;**16**:289. doi: [10.1186/s12891-015-0736-6](https://doi.org/10.1186/s12891-015-0736-6). [PubMed: [26459411](https://pubmed.ncbi.nlm.nih.gov/26459411/)].
- Chang W, Huang W, Lai P. Muscle Activation of Vastus Medialis Oblique and Vastus Lateralis in Sling-Based Exercises in Patients with Patellofemoral Pain Syndrome: A Cross-Over Study. *Evid Based Complement Alternat Med.* 2015;**2015**.
- Herrington L, Pearson S. Does level of load affect relative activation levels of vastus medialis oblique and vastus lateralis? *J Electromyogr Kinesiol.* 2006;**16**(4):379–83.
- Zetaruk MN, Zurakowski D, Violan MA, Micheli LJ. Safety recommendations in Shotokan karate. *Clin J Sport Med.* 2000;**10**(2):117–22. [PubMed: [10798793](https://pubmed.ncbi.nlm.nih.gov/10798793/)].
- Critchley GR, Mannion S, Meredith C. Injury rates in Shotokan karate. *British J Sports Med.* 1999;**33**(3):174–7.
- Kordi R, Maffulli N, Wroble RR, Wallace WA. Combat sports medicine. Springer Science & Business Media; 2009.
- Witvrouw E, Lysens R, Bellemans J, Cambier D, Vanderstraeten G. Intrinsic risk factors for the development of anterior knee pain in an athletic population a two-year prospective study. *Am J Sports Med.* 2000;**28**(4):480–9.
- McKenzie K, Galea V, Wessel J, Pierrynowski M. Lower extremity kinematics of females with patellofemoral pain syndrome while stair stepping. *J Orthop Sports Phys Ther, and JOSPT.* 2010;**40**(10):625–32.
- Voight ML, Wieder DL. Comparative reflex response times of vastus medialis obliquus and vastus lateralis in normal subjects and subjects with extensor mechanism dysfunction. An electromyographic study. *Am J Sports Med.* 1991;**19**(2):131–7. [PubMed: [2039064](https://pubmed.ncbi.nlm.nih.gov/2039064/)].
- Karst GM, Willett GM. Onset timing of electromyographic activity in the vastus medialis obliquus and vastus lateralis muscles in subjects with and without patellofemoral pain syndrome. *Phys Ther.* 1995;**75**(9):813–23. [PubMed: [7659741](https://pubmed.ncbi.nlm.nih.gov/7659741/)].
- Chester R, Smith TO, Sweeting D, Dixon J, Wood S, Song F. The relative timing of VMO and VL in the aetiology of anterior knee pain: a systematic review and meta-analysis. *BMC Musculoskeletal Disord.* 2008;**9**:64. doi: [10.1186/1471-2474-9-64](https://doi.org/10.1186/1471-2474-9-64). [PubMed: [18452611](https://pubmed.ncbi.nlm.nih.gov/18452611/)].

21. Wong YM. Recording the vastii muscle onset timing as a diagnostic parameter for patellofemoral pain syndrome: fact or fad?. *Phys Ther Sport*. 2009;**10**(2):71–4. doi: [10.1016/j.ptsp.2009.02.001](https://doi.org/10.1016/j.ptsp.2009.02.001). [PubMed: [19376476](https://pubmed.ncbi.nlm.nih.gov/19376476/)].
22. Cavazzuti L, Merlo A, Orlandi F, Campanini I. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Gait Posture*. 2010;**32**(3):290–5. doi: [10.1016/j.gaitpost.2010.06.025](https://doi.org/10.1016/j.gaitpost.2010.06.025). [PubMed: [20727760](https://pubmed.ncbi.nlm.nih.gov/20727760/)].
23. Coqueiro KR, Bevilacqua-Grossi D, Berzin F, Soares AB, Candolo C, Monteiro-Pedro V. Analysis on the activation of the VMO and VLL muscles during semisquat exercises with and without hip adduction in individuals with patellofemoral pain syndrome. *J Electromyogr Kinesiol*. 2005;**15**(6):596–603. doi: [10.1016/j.jelekin.2005.03.001](https://doi.org/10.1016/j.jelekin.2005.03.001). [PubMed: [15878289](https://pubmed.ncbi.nlm.nih.gov/15878289/)].
24. Peng HT, Kernozeck TW, Song C. Muscle activation of vastus medialis obliquus and vastus lateralis during a dynamic leg press exercise with and without isometric hip adduction. *Phys Ther Sport*. 2013;**14**(1):44–9.
25. Bevilacqua-Grossi D, Felicio LR, Simoes R, Coqueiro KRR, Monteiro-Pedro V. Electromyographic activity evaluation of the patella muscles during squat isometric exercise in individuals with patellofemoral pain syndrome. *Revista Brasileira de Medicina do Esporte*. 2005;**11**(3):159–63.
26. Bevilacqua-Grossi D, Felicio LR, Silverio GWP. Onset of electrical activity of patellar stabilizer muscles in subjects with patellofemoral pain. *Acta Ortopédica Brasileira*. 2009;**17**(5):297–9.
27. Hodges PW, Mellor R, Crossley K, Bennell K. Pain induced by injection of hypertonic saline into the infrapatellar fat pad and effect on coordination of the quadriceps muscles. *Arthritis Rheum*. 2009;**61**(1):70–7. doi: [10.1002/art.24089](https://doi.org/10.1002/art.24089). [PubMed: [19116977](https://pubmed.ncbi.nlm.nih.gov/19116977/)].
28. Singer B, Singer K. Anterior knee pain scale. *J Physiother*. 2009;**55**(2):140.
29. Watson CJ, Propp M, Ratner J, Zeigler DL, Horton P, Smith SS. Reliability and responsiveness of the lower extremity functional scale and the anterior knee pain scale in patients with anterior knee pain. *J Orthop Sports Phys Ther*. 2005;**35**(3):136–46. doi: [10.2519/jospt.2005.35.3.136](https://doi.org/10.2519/jospt.2005.35.3.136). [PubMed: [15839307](https://pubmed.ncbi.nlm.nih.gov/15839307/)].
30. Hermens HJ, Freriks B. Seniam Available from: <http://seniam.org>.
31. Lotfian S, Cherati AS, Jamshidi A, Sanjari MA. Assessment of Correlation between Electrogoniometer Measurements and Sports-Specific Movement in Karate Elites. *Asian J Sports Med*. 2014;**5**(2):115.
32. Cowan SM, Bennell KL, Hodges PW, Crossley KM, McConnell J. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Arch Phys Med Rehabil*. 2001;**82**(2):183–9. doi: [10.1053/apmr.2001.19022](https://doi.org/10.1053/apmr.2001.19022). [PubMed: [11239308](https://pubmed.ncbi.nlm.nih.gov/11239308/)].
33. Burden A. How should we normalize electromyograms obtained from healthy participants? What we have learned from over 25 years of research. *J Electromyogr Kines*. 2010;**20**(6):1023–35.
34. Nicholson WR. The importance of normalization in the interpretation of surface electromyography: A proof of principle. *J Manipulative Physiol Ther*. 2000;**23**(5):369–70. doi: [10.1016/S0161-4754\(00\)90218-5](https://doi.org/10.1016/S0161-4754(00)90218-5).
35. Irish SE, Millward AJ, Wride J, Haas BM, Shum GL. The effect of closed-kinetic chain exercises and open-kinetic chain exercise on the muscle activity of vastus medialis oblique and vastus lateralis. *J Strength Cond Res*. 2010;**24**(5):1256–62. doi: [10.1519/JSC.0b013e3181cf749f](https://doi.org/10.1519/JSC.0b013e3181cf749f). [PubMed: [20386128](https://pubmed.ncbi.nlm.nih.gov/20386128/)].