

The Association between *Helicobacter pylori* Infection and Body Weight among Children

Jun Sik Choi, Kyung Og Ko, Jae Woo Lim, Eun Jeong Cheon, Gyung Min Lee, and Jung Min Yoon

Department of Pediatrics, Konyang University College of Medicine, Daejeon, Korea

Purpose: We performed to reveal the association between the *Helicobacter pylori* infection and body weight among children.

Methods: Out retrospective study included patients who underwent the *H. pylori* immunoglobulin G testing at Konyang University Hospital between March 2011 and June 2014. These patients were classified as seropositive (28 boys, 27 girls; mean age: 9.89±3.28 years) or seronegative (55 boys, 54 girls; mean age: 9.84±3.02 years). Next, we compared various characteristics between the seropositive and negative groups, as well as between obese children (body weight ≥90th percentile) and non-obese children (body weight <90th percentile). Furthermore, we compared the change in body weight after 2 months of treatment with amoxicillin, clarithromycin and omeprazole among the 55 seropositive children (14 treated children and 41 non-treated children).

Results: There were no differences in the weights and laboratory data for the 55 seropositive children and 109 seronegative children (weight; 40.96±18.11 kg vs. 36.85±13.72 kg, respectively; $p=0.14$). And, there was no difference in the prevalence of *H. pylori* infection among the 29 obese and 135 non-obese children ($p=0.581$). However, after 2 months of eradication, the 14 treated patients exhibited a significant weight gain (+0.91±0.52 kg), compared to the 41 non-treated patients (-0.29±1.16 kg, $p=0.025$).

Conclusion: Our findings present that obesity was not associated with the *H. pylori* infection, although *H. pylori* eradication led to significant increase in body weight.

Key Words: Helicobacter, Obesity, Body weight, Disease eradication, Child

INTRODUCTION

The *Helicobacter pylori* infection is one of the causes in the gastritis, gastric and duodenal ulcer, and stomach cancer. In addition several studies have

described that the infection of *H. pylori* is associated with various extra-gastric conditions, such as diabetes mellitus (DM) [1-4], dyslipidemia [5-7], cardiovascular disease [8,9] and ischemic cerebrovascular disease [10,11]. But, the association between

Received : September 30, 2015, Revised : December 31, 2015, Accepted : January 14, 2016

Corresponding author: Jung Min Yoon, Department of Pediatrics, Konyang University College of Medicine, 158 Gwanjeodong-ro, Seo-gu, Daejeon 35365, Korea. Tel: +82-42-600-9230, Fax: +82-42-600-9090, E-mail: jmyoon@kyuh.ac.kr

Copyright © 2016 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

infection of *H. pylori* and obesity is unclear, as various studies have reported conflicting findings regarding whether obesity is associated with this infection. For example, one study has demonstrated the positive association between the *H. pylori* positivity and body mass index (BMI) in individuals who were > 15 years old [12], and another study has demonstrated that the prevalence of *H. pylori* infection was 57.2% among obese adults and 27.0% among non-obese adults [13]. In contrast, several studies have reported that obesity was not associated with the *H. pylori* infection [14,15]. Nevertheless, several studies have demonstrated that *H. pylori* eradication leads to increased BMI, owing to recovery of gastric acid secretion and improved pancreatic function [16-18]. However, almost all of these studies examined adult patients, except one study that reported *H. pylori* infection was associated with DM among children. Therefore, we performed this study to evaluate the association between the *H. pylori* infection and obesity among children and to examine the relationship between the eradication of *H. pylori* and weight gain.

MATERIALS AND METHODS

We retrospectively reviewed the medical records of all children who underwent *H. pylori* immunoglobulin G (IgG) test with IMMULITE 2000 kits (DPC Lab Diagnostic Products Corporation, Los Angeles, CA, USA) at Konyang University Hospital between March 2011 and June 2014. They had several gastrointestinal symptoms—recurrent abdominal pain, heartburn, dyspepsia, nausea, vomiting, and weight loss. A total of 55 children had a positive result, and we randomly selected 109 IgG-negative age- and sex-matched children (approximately double the number of positive children) to serve as the controls. We also classified the included patients as obese (body weight \geq 90th percentile, 29 patients) and non-obese (body weight < 90th percentile, 135 patients) to investigate the difference in seropositivity of *H. pylori* between two groups.

Fourteen patients who exhibited serious symp-

toms and positive results from rapid urease test (RUT), biopsy and urea breath test (UBT) in the 55 seropositive patients were treated with amoxicillin, clarithromycin, and omeprazole for 2 weeks. Twelve of the 14 treated patients tested positive in the RUT or via histologic examination, and 2 of these patients tested positive in the UBT. The other 41 seropositive patients were treated using symptomatic care, and we compared the change in body weight for the treated and non-treated groups via outpatient follow-up after 2 months of treatment. Fifteen patients in the non-treated group were treated using a proton pump inhibitor or H2 receptor blocker.

All statistical analyses were performed by IBM SPSS Statistics ver. 19.0 (IBM Co., Armonk, NY, USA), and differences with a *p*-value of < 0.05 were considered statistically significant. Inter-group comparisons were performed by the dependent Student *t*-test or chi-square test, as appropriate. All data was presented as mean \pm standard deviation or number (%).

RESULTS

The association between the infection of *H. pylori* and obesity

When we compared the seropositive children (28 boys, 27 girls; mean age: 9.89 ± 3.28 years) and the seronegative children (55 boys, 54 girls; mean age: 9.84 ± 3.02 years), we did not observe any significant differences in their laboratory data. The seropositive patients were slightly heavier than the seronegative patients (40.96 ± 18.11 kg vs. 36.85 ± 13.72 kg), although there was no statistically significant differences (Table 1). The obese group (29 patients) included 11 seropositive children (37.9%), and the non-obese group (135 patients) included 44 seropositive children (32.6%); this difference was not statistically significant (Table 2).

Does eradication of *H. pylori* lead to increased body weight?

There was no significant difference in the initial body weights of the treated and non-treated groups.

Table 1. The Demographic and Laboratory Characteristics of the Seropositive and Seronegative Groups

Characteristic	<i>Helicobacter pylori</i> immunoglobulin G		p-value
	Seropositive (n=55)	Seronegative (n=109)	
Sex			0.96
Male	28 (50.9)	55 (50.5)	
Female	27 (49.1)	54 (49.5)	
Age (yr)	9.89±3.28	9.84±3.02	0.93
Weight (kg)	40.96±18.11	36.85±13.72	0.14
White blood cell count (/μL)	7,554.55±2,238.29	7,053.21±2,346.49	0.19
Hemoglobin (g/dL)	13.08±0.84	12.94±0.97	0.35
Hematocrit (%)	38.79±2.51	38.77±5.27	0.97
Platelet (/μL)	289,890.91±70,775.77	282,692.42±117,861.17	0.97
Glucose (mg/dL)	94.15±9.90	104.25±78.25	0.68
Aspartate aminotransferase (IU/L)	25.13±7.21	25.56±7.63	0.34
Alanine aminotransferase (IU/L)	14.67±6.69	15.25±9.23	0.73

Values are presented as number of cases (%) or mean±standard deviation.

Table 2. The Prevalence of *Helicobacter pylori* Infection in the Obese and Non-Obese Groups

<i>H. pylori</i> infection	Body weight		p-value
	Obese group (≥90th percentile)	Non-obese group (<90th percentile)	
<i>H. pylori</i> IgG positive	11 (37.9)	44 (32.6)	0.581
<i>H. pylori</i> IgG negative	18 (62.1)	91 (67.4)	
Total	29 (100.0)	135 (100.0)	

Values are presented as number of cases (%).

IgG: immunoglobulin G.

Table 3. The Change in Body Weight after *Helicobacter pylori* Eradication in Obese Group

Body weight	<i>H. pylori</i> eradication		p-value
	Treatment (n=14)	No treatment (n=15)	
Initial weight (kg)	40.65±20.07	41.10±22.99	0.991
Change in body weight (kg)	+0.91±0.52	-0.29±1.16	0.025

Values are presented as mean±standard deviation.

By dependent Student t-test.

After 2 months of eradication treatment using omeprazole, amoxicillin, and clarithromycin, the 14 treated patients exhibited an increased body weight (+0.91±0.52 kg). In contrast, 15 non-treated patients who were successfully followed-up exhibited a decreased body weight (-0.29±1.16 kg), and this difference was statistically significant ($p=0.025$) (Table 3). However, 26 of the 41 non-treated children were lost to follow-up.

Thirteen of the 14 treated patients underwent upper endoscopy and were diagnosed with nodular gastritis (6 patients), erythematous gastritis (3 patients), hemorrhagic gastritis (1 patient), and atrophic gastritis (1 patient). Two patients exhibited normal upper endoscopy findings.

DISCUSSION

This retrospective study aimed to reveal the relationships between infection of *H. pylori* and obesity among pediatric patients. Our findings indicate that obesity was not associated with *H. pylori* infection, although triple eradication therapy led to an increase in body weight.

The association between *H. pylori* infection and BMI remains unclear, as is the relationship between eradication of *H. pylori* and change in body weight. For example, several studies have described that the prevalence of *H. pylori* infection was higher in patients with a high BMI [13,19,20]. Arslan et al. [13] have also described that the prevalence of *H. pylori* infection is higher in their obese group by impaired immunity. In contrast, similar to our findings, other studies have described that *H. pylori* infection is not associated with BMI or being overweight [14,15]. Ioannou et al. [14] have described that *H. pylori* infection is not associated with BMI because it did not change serum leptin levels in their study. However, one study have described that *H. pylori* infection was negatively associated with the BMI among 4-15-year-old individuals, but was positively associated with BMI in >15-year-old individuals [12]. These discrepancies are likely caused by a complex group of factors, as being obese or overweight is the result of multiple factors, including—age, personal habits, socioeconomic environment, culture, hormonal change and chronic diseases like metabolic syndrome. Furthermore, most of the previous studies evaluated adult populations, and only small number of studies evaluated children. In our study, the infection of *H. pylori* is not associated with overweight, although we cannot comment on the reason for this absence of a relationship, as we did not evaluate many of the factors that are associated with being overweight or obese.

Interestingly, many studies have described that the relationship of *H. pylori* infection and metabolic disease, such as DM and dyslipidemia, among adults [2-7]. For example, one study described that the prevalence of *H. pylori* infection was higher in the

DM patients [2], and another study described that *H. pylori* infection was associated with elevated serum lipid levels [6]. Therefore, additional studies are needed regarding the relationship between these metabolic diseases and *H. pylori* infection among children.

In this study, we observed that triple therapy of *H. pylori* induced weight gain, and previous studies have reported similar findings [16,17,21-23]. Kamada et al. [22] described that the *H. pylori* eradication led to increased obesity, but also improved the patients' dyspeptic symptoms and quality of life. Furthermore, Loffeld [24] reported that eradication of *H. pylori* led to increased body weight and appetite, due to changes in the patients' ghrelin and leptin levels. The *H. pylori* eradication led to decreased gastric leptin levels and improved pancreatic function, which might lead to increase in BMI [17,21]. Ghrelin—a gut-brain peptide hormone that regulates appetite—enhances the gastric emptying and gastrointestinal motility [25-27], and two studies have demonstrated that *H. pylori* eradication led to increased ghrelin levels [28,29]. Based on the results of these reports, we hypothesize that these diverse mechanisms (relieving gastric symptoms, hormonal changes, and improved quality of life owing to regular meals) may be responsible for the post-eradication weight gain that we observed in this study.

The use of *H. pylori* immunoglobulin as a diagnostic marker has high sensitivity and specificity in adults, although this method is rarely used in children, because of its low sensitivity [30]. Therefore, we only performed *H. pylori* eradication in the 14 patients who also tested positive in the RUT, histological examination or UBT. Thus, the remaining 41 seropositive patients with no other warning signs were only treated using supportive care.

This study has some limitations. Our study was performed at the single institution, and our findings may not be representative of a broader patient population. Second, we only evaluated a small number of patients in the eradication and control groups, and our control group may have been affected by selection bias. Third, the most commonly used index to

determine obesity is BMI, although we were unable to reveal the association between the BMI and infection of *H. pylori* because height was not recorded for a large number of our patients. Therefore we categorized patients using their age-adjusted body weight percentiles (<90th and ≥90th percentile). Fourth, our findings cannot support a causal mechanism for the increase in body weight, and further studies should evaluate the relevant factors, which include serum lipid levels, ghrelin levels, leptin levels and BMI.

In conclusion, we confirmed that *H. pylori* infection was not associated with obesity among children, although *H. pylori* eradication led to increasing in body weight. However, the limitations of our study preclude us from commenting on the potential causative factors that influenced these results. Therefore, large-scale well-designed the prospective studies are needed to evaluate the association of obesity with *H. pylori* infection and eradication.

REFERENCES

1. Arslan D, Kendirci M, Kurtoglu S, Kula M. Helicobacter pylori infection in children with insulin dependent diabetes mellitus. *J Pediatr Endocrinol Metab* 2000;13: 553-6.
2. Senturk O, Canturk Z, Cetinarslan B, Ercin C, Hulagu S, Canturk NZ. Prevalence and comparisons of five different diagnostic methods for Helicobacter pylori in diabetic patients. *Endocr Res* 2001;27:179-89.
3. Quatrini M, Boarino V, Ghidoni A, Baldassarri AR, Bianchi PA, Bardella MT. Helicobacter pylori prevalence in patients with diabetes and its relationship to dyspeptic symptoms. *J Clin Gastroenterol* 2001;32: 215-7.
4. Gulcelik NE, Kaya E, Demirbas B, Culha C, Koc G, Ozkaya M, et al. Helicobacter pylori prevalence in diabetic patients and its relationship with dyspepsia and autonomic neuropathy. *J Endocrinol Invest* 2005;28: 214-7.
5. Chimienti G, Russo F, Lamanuzzi BL, Nardulli M, Messa C, Di Leo A, et al. Helicobacter pylori is associated with modified lipid profile: impact on lipoprotein(a). *Clin Biochem* 2003;36:359-65.
6. Laurila A, Bloigu A, Näyhä S, Hassi J, Leinonen M, Saikku P. Association of Helicobacter pylori infection with elevated serum lipids. *Atherosclerosis* 1999;142: 207-10.
7. Hoffmeister A, Rothenbacher D, Bode G, Persson K, März W, Nauck MA, et al. Current infection with Helicobacter pylori, but not seropositivity to Chlamydia pneumoniae or cytomegalovirus, is associated with an atherogenic, modified lipid profile. *Arterioscler Thromb Vasc Biol* 2001;21:427-32.
8. Jafarzadeh A, Esmaeeli-Nadimi A, Nemati M, Tahmasbi M, Ahmadi P. Serum concentrations of Helicobacter pylori IgG and the virulence factor CagA in patients with ischaemic heart disease. *East Mediterr Health J* 2010;16:1039-44.
9. Niccoli G, Franceschi F, Cosentino N, Giupponi B, De Marco G, Merra G, et al. Coronary atherosclerotic burden in patients with infection by CagA-positive strains of Helicobacter pylori. *Coron Artery Dis* 2010; 21:217-21.
10. Grau AJ, Buggle F, Lichy C, Brandt T, Becher H, Rudi J. Helicobacter pylori infection as an independent risk factor for cerebral ischemia of atherothrombotic origin. *J Neurol Sci* 2001;186:1-5.
11. Diomedi M, Stanzione P, Sallustio F, Leone G, Renna A, Misaggi G, et al. Cytotoxin-associated Gene-A-positive Helicobacter pylori strains infection increases the risk of recurrent atherosclerotic stroke. *Helicobacter* 2008;13:525-31.
12. Kopáčová M, Bures J, Koupil I, Rejchrt S, Vorisek V, Seifert B, et al; European Society for Primary Care Gastroenterology. Body indices and basic vital signs in Helicobacter pylori positive and negative persons. *Eur J Epidemiol* 2007;22:67-75.
13. Arslan E, Atilgan H, Yavaşoğlu I. The prevalence of Helicobacter pylori in obese subjects. *Eur J Intern Med* 2009;20:695-7.
14. Ioannou GN, Weiss NS, Kearney DJ. Is Helicobacter pylori seropositivity related to body mass index in the United States? *Aliment Pharmacol Ther* 2005;21:765-72.
15. Cho I, Blaser MJ, François F, Mathew JP, Ye XY, Goldberg JD, et al. Helicobacter pylori and overweight status in the United States: data from the Third National Health and Nutrition Examination Survey. *Am J Epidemiol* 2005;162:579-84.
16. Azuma T, Suto H, Ito Y, Muramatsu A, Ohtani M, Dojo M, et al. Eradication of Helicobacter pylori infection induces an increase in body mass index. *Aliment Pharmacol Ther* 2002;16 Suppl 2:240-4.
17. Furuta T, Shirai N, Xiao F, Takashima M, Hanai H. Effect of Helicobacter pylori infection and its eradication on nutrition. *Aliment Pharmacol Ther* 2002;

- 16:799-806.
18. Kamada T, Hata J, Kusunoki H, Ito M, Tanaka S, Kawamura Y, et al. Eradication of *Helicobacter pylori* increases the incidence of hyperlipidaemia and obesity in peptic ulcer patients. *Dig Liver Dis* 2005;37:39-43.
 19. Rosenstock SJ, Jørgensen T, Andersen LP, Bonnevie O. Association of *Helicobacter pylori* infection with life-style, chronic disease, body-indices, and age at menarche in Danish adults. *Scand J Public Health* 2000;28:32-40.
 20. Thjodleifsson B, Olafsson I, Gislason D, Gislason T, Jögi R, Janson C. Infections and obesity: a multinational epidemiological study. *Scand J Infect Dis* 2008;40:381-6.
 21. Azuma T, Suto H, Ito Y, Ohtani M, Dojo M, Kuriyama M, et al. Gastric leptin and *Helicobacter pylori* infection. *Gut* 2001;49:324-9.
 22. Kamada T, Sugiu K, Haruma K. Development of obesity and hyperlipidemia after eradication of *Helicobacter pylori*. *Nihon Rinsho* 2005;63 Suppl 11:536-8.
 23. Fujiwara Y, Higuchi K, Arafa UA, Uchida T, Tominaga K, Watanabe T, et al. Long-term effect of *Helicobacter pylori* eradication on quality of life, body mass index, and newly developed diseases in Japanese patients with peptic ulcer disease. *Hepatogastroenterology* 2002;49:1298-302.
 24. Loffeld RJ. *Helicobacter pylori*, obesity and gastro-oesophageal reflux disease. Is there a relation? A personal view. *Neth J Med* 2005;63:344-7.
 25. Akamizu T, Kangawa K. Translational research on the clinical applications of ghrelin. *Endocr J* 2006;53:585-91.
 26. Williams DL, Cummings DE. Regulation of ghrelin in physiologic and pathophysiologic states. *J Nutr* 2005;135:1320-5.
 27. Chen CY, Asakawa A, Fujimiya M, Lee SD, Inui A. Ghrelin gene products and the regulation of food intake and gut motility. *Pharmacol Rev* 2009;61:430-81.
 28. Kawashima J, Ohno S, Sakurada T, Takabayashi H, Kudo M, Ro S, et al. Circulating acylated ghrelin level decreases in accordance with the extent of atrophic gastritis. *J Gastroenterol* 2009;44:1046-54.
 29. Gunji T, Matsuhashi N, Sato H, Fujibayashi K, Okumura M, Sasabe N, et al. *Helicobacter pylori* infection is significantly associated with metabolic syndrome in the Japanese population. *Am J Gastroenterol* 2008;103:3005-10.
 30. de Oliveira AM, Rocha GA, Queiroz DM, Mendes EN, de Carvalho AS, Ferrari TC, et al. Evaluation of enzyme-linked immunosorbent assay for the diagnosis of *Helicobacter pylori* infection in children from different age groups with and without duodenal ulcer. *J Pediatr Gastroenterol Nutr* 1999;28:157-61.