

## Letter to the Editor

# Soluble A $\beta$ oligomer-induced synaptopathy: c-Jun N-terminal kinase's role

### Dear Editor,

Among the neurodegenerative diseases, Alzheimer disease (AD) is the most common and severe age-related dementia for which there is currently no available treatment. Many studies support the assumption that AD is a spine pathology (Selkoe, 2002; Sivanesan et al., 2013) and that soluble amyloid- $\beta$  (A $\beta$ ) oligomers are causative of AD synaptopathy. Diverse lines of evidence indicate that A $\beta$  oligomers induce formation of pore-like structures on the membrane (Arispe et al., 1993; Lashuel et al., 2002) and interfere with glutamatergic transmission. The A $\beta$  oligomers result in a decreased number of AMPA receptors (AMPA-r) and NMDA receptors (NMDA-r), as well as PSD-95 at the postsynaptic membrane, and thus reduce the strength and plasticity of excitatory synapses (Chapman et al., 1999; Walsh et al., 2002). However, the underlying intracellular mechanisms regulating synaptic changes are only partially known. By understanding the pathophysiological mechanisms leading to synaptic dysfunction and the progression of this dysfunction, better interference in the pathogenesis of AD can be achieved.

We present an *in vitro* model to study the temporal sequence of dendritic spine modifications induced by soluble A $\beta$  oligomers, and to analyse the intracellular signalling pathways leading to AD synaptopathy. This model allows synaptic alterations to be followed in living neurons before and after treatment and reduces bias due to cell variability. This model also permits testing of pharmaceuticals that are designed to reverse the biochemical and structural alterations of synapses induced by A $\beta$  oligomers.


Brainbow hippocampal neurons, which express fluorescent proteins, were used to visualize dendritic spines and study synaptic plasticity (Figure 1A and B). To obtain isolated cells, fluorescent neurons were

seeded on a layer of non-fluorescent neurons (ratio: 1/16) (Figure 1A and B). In this way we avoided the need for transfection, infection protocols, and low density cultures that are not well tolerated by neurons. Neurons were treated with a subtoxic dose (Figure 1E) of soluble A $\beta$ 1–42 oligomers in order to induce synaptic changes without any signs of neuronal death (Figure 1E). The preparation of synthetic A $\beta$ 1–42 that were used to induce *in vitro* synaptic dysfunction had been previously characterized. Oligomeric assemblies were only observed in peptide preparations after a 24-h incubation at 4°C (referred to as oligomers). Immediately after dissolution, the majority of A $\beta$ 1–42 remained as unassembled monomeric structures (Figure 1C and D and Supplementary Figure S1).

Subtoxic concentrations of soluble A $\beta$  oligomers induced alterations in the postsynaptic density (PSD) composition of dendritic spines, while monomers had no effect on synaptic plasticity (Supplementary Figure S1F). Exposing the neurons for 3 h to 1  $\mu$ M A $\beta$  oligomers induced changes in the PSD region, leading to a 68% and 61% drop of GluN2A and GluN2B subunits of NMDA-r, respectively; a decrease of 70% and 65% of GluA1 and GluA2 subunits of AMPA-r, respectively; a 53% loss of PSD-95; and a 76% loss of drebrin (Figure 1F and Supplementary Figure S2A). To assess dendritic spine modifications *in vitro*, we analysed changes in spine density and morphology in neurons exposed to soluble A $\beta$  oligomers. Application of soluble A $\beta$  oligomers (1  $\mu$ M) for 3 h caused a 25% decrease in total spine number compared with that before A $\beta$  application (Figure 1J and K). The decrease involved all types of spines in a proportional manner. The number of mushroom, stubby, and thin spines decreased by 32%, 25%, and 22%, respectively (Figure 1J and L). Moreover, A $\beta$

oligomer treatment induced a 67% reduction in new spine formation compared with control conditions, and led to spine shrinkage (Supplementary Figure S2E). The number of mushroom spines that became stubby or thin was significantly increased by A $\beta$  oligomer treatment, while the number of spines that became mushroom was decreased (Supplementary Figure S2E). The morphological changes were consistent with the observed biochemical alterations since thin and stubby spines have a less extended PSD region and lower contents of glutamate receptors as well as postsynaptic markers, in comparison with mushroom spines (Tackenberg et al., 2009).

To analyse the pathways involved in A $\beta$  oligomer-induced synaptopathy, we evaluated synaptic changes induced after 30 min and 3 h of A $\beta$  oligomer exposure (1  $\mu$ M) and correlated them to the activation of two stress signalling pathways, c-Jun N-terminal kinase (JNK) and caspase-3. After 30 min there was no sign of molecular changes: NMDA-r and AMPA-r subunits, PSD-95, and drebrin levels were unaffected by the A $\beta$  oligomer treatment (Figure 1F). However, JNK was already activated at this stage, as indicated by a 2.36-fold increase of the P-JNK/JNK ratio compared with control conditions (Figure 1G and Supplementary Figure S2B). There was no indication of caspase-3 cleavage after 30 min of A $\beta$  oligomer exposure (Figure 1H and Supplementary Figure S2C). Exposing neurons to soluble A $\beta$  oligomers for 3 h induced a biochemical perturbation of PSD (Figure 1F) with a 4.33-fold increase of caspase-3 cleavage (Figure 1H) (Li et al., 2010; D'Amelio et al., 2011), while JNK activity remained elevated by 2.26 folds (Figure 1G). Our results showed that JNK activation was triggered by A $\beta$  oligomers before PSD alterations were induced and JNK activation persisted up to 3 h, at which



**Figure 1** JNK inhibition prevents Aβ oligomer-induced synaptopathy. **(A and B)** *In vitro* model to study synaptic plasticity. **(A)** A Brainbow mouse fluorescent hippocampal neuron (green, 20×) seeded on a layer of non-fluorescent neurons in a ratio 1:16 (merge with bright field). **(B)** Higher magnification of a dendrite portion and 3D reconstruction showing that dendritic spines are easily visualized and classified in mushroom, stubby, and thin subtypes. **(C and D)** Transmission electron micrographs of monomeric **(C)** and oligomeric **(D)** assemblies. **(E)** The effect of crescent doses of synthetic soluble Aβ oligomers (0.01, 0.1, 1, 10 μM) on neuron viability was assessed with the MTT assay. Soluble Aβ oligomers at 0.01, 0.1, and 1 μM did not affect cell viability after 3 h of treatment. However, Aβ oligomers led to neuronal death at 10 μM (one-way ANOVA, Dunnett's *post hoc*

stage PSD changes and caspase-3 cleavage appeared. These results suggest that JNK activation likely occurs at the onset of AD synaptopathy.

In order to define JNK's role in A $\beta$  oligomer-induced synaptopathy *in vitro*, neurons were pre-treated with the specific cell permeable JNK inhibitor peptide D-JNKI1 and subsequently exposed to A $\beta$  oligomer stimulation. D-JNKI1 completely protected neurons against A $\beta$  oligomer-induced synaptic changes, abolishing modifications of the composition of PSD (GluN2A, GluN2B, GluA1, GluA2, and PSD-95 returned to control levels) (Figure 11 and Supplementary Figure S2D), as well as spine alterations (Supplementary Figure S2E) and loss (Figure 1J–L). Importantly, D-JNKI1 also prevented caspase-3 cleavage (Figure 11 and Supplementary Figure S2D). These results suggest that the JNK pathway triggers synaptic dysfunction *in vitro* following A $\beta$  oligomer exposure.

To better understand the relevance of JNK and caspase-3 pathways in synaptopathy, we compared the effect of D-JNKI1 with z-DEVD-FMK, a caspase-3 inhibitor that partially prevents dendritic spine dysfunction in Tg2576 mice (D'Amelio et al., 2011). In our experimental conditions, z-DEVD-FMK completely inhibited A $\beta$  oligomer-induced loss of AMPA-r, but only partially (about 50%) restored NMDA-r subunits and PSD-95 to control levels (Supplementary Figure S3). Caspase-3 inhibition by z-DEVD-FMK was less efficient in protecting neurons against synaptic changes than JNK inhibition that led to a complete recovery of GluN2A, GluN2B, GluA1, GluA2, PSD-95,

and drebrin levels. Our results demonstrate that JNK plays a pivotal role in A $\beta$  oligomer-induced synaptopathy and acts upstream of caspase-3 (D'Amelio et al., 2011).

To our knowledge this is the first demonstration identifying JNK as a key modulator in the degeneration of excitatory synapses. JNK, therefore, represents an innovative target as its inhibition in neurons *in vitro* completely protects synapse degeneration, without interfering with A $\beta$  oligomerization.

[Supplementary material is available at *Journal of Molecular Cell Biology* online. This study was supported by San Paolo 2008–2437, Banca Intesa San Paolo Grant 2009–2013 to M.S., Marie Curie Industry-Academia Partnerships and Pathways (IAPP), CPADS (Cell permeable peptides as drug delivery system) and CARIPLO (Cassa di risparmio delle provincie Lombarde) Foundation 2009–2425, ADF (Alzheimer's Drugs Discovery Foundation) USA Grant, Swiss National Science Foundation Grant (31003A\_125379). We thank Alessandro Rossi (IRCCS-Istituto di Ricerche Farmacologiche 'Mario Negri') for peptide synthesis, Dr Gardoni and Prof. Di Luca (Department of Pharmaceutical Sciences University of Milan) for their advices and criticisms. The kind gift of FMOc amino acids by Flamma Spa (Bergamo, Italy) is gratefully acknowledged. There is no actual or potential conflict of interest.]

Alessandra Sclip<sup>1</sup>, Andrea Arnaboldi<sup>1</sup>, Isabella Colombo<sup>1</sup>, Pietro Veglianesi<sup>1</sup>, Laura Colombo<sup>1</sup>, Massimo Messa<sup>1</sup>,

Simona Mancini<sup>1</sup>, Sara Cimini<sup>1</sup>, Federica Morelli<sup>1</sup>, Xanthi Antoniou<sup>1</sup>, Egbert Welker<sup>2</sup>, Mario Salmona<sup>1</sup>, and Tiziana Borsello<sup>1,\*</sup>

<sup>1</sup>IRCCS-Istituto di Ricerche Farmacologiche 'Mario Negri', Via La Masa 19, 20156 Milano, Italy  
<sup>2</sup>Département de Neurosciences Fondamentales, University of Lausanne, 1005 Lausanne, Switzerland

\*Correspondence to: Tiziana Borsello, E-mail: tiziana.borsello@marionegri.it

## References

- Arispe, N., Rojas, E., and Pollard, H.B. (1993). Alzheimer disease amyloid beta protein forms calcium channels in bilayer membranes: blockade by tromethamine and aluminum. *Proc. Natl Acad. Sci. USA* 90, 567–571.
- Chapman, P.F., White, G.L., Jones, M.W., et al. (1999). Impaired synaptic plasticity and learning in aged amyloid precursor protein transgenic mice. *Nat. Neurosci.* 2, 271–276.
- D'Amelio, M., Cavallucci, V., Middei, S., et al. (2011). Caspase-3 triggers early synaptic dysfunction in a mouse model of Alzheimer's disease. *Nat. Neurosci.* 14, 69–76.
- Lashuel, H.A., Hartley, D., Petre, B.M., et al. (2002). Neurodegenerative disease: amyloid pores from pathogenic mutations. *Nature* 418, 291.
- Li, Z., Jo, J., Jia, J.M., et al. (2010). Caspase-3 activation via mitochondria is required for long-term depression and AMPA receptor internalization. *Cell* 141, 859–871.
- Selkoe, D.J. (2002). Alzheimer's disease is a synaptic failure. *Science* 298, 789–791.
- Sivanesan, S., Tan, A., and Rajadas, J. (2013). Pathogenesis of abeta oligomers in synaptic failure. *Curr. Alzheimer Res.* 10, 316–323.
- Tackenberg, C., Ghori, A., and Brandt, R. (2009). Thin, stubby or mushroom: spine pathology in Alzheimer's disease. *Curr. Alzheimer Res.* 6, 261–268.
- Walsh, D.M., Klyubin, I., Fadeeva, J.V., et al. (2002). Naturally secreted oligomers of amyloid beta protein potentially inhibit hippocampal long-term potentiation in vivo. *Nature* 416, 535–539.

test, \* $P < 0.05$ ,  $n = 6$ ). (F–H) Synthetic soluble A $\beta$  oligomers induced synaptic activation of JNK before synaptopathy. (F) Western blot performed on the triton insoluble fraction (TIF) of neurons treated with A $\beta$  oligomers (1  $\mu$ M) for 30 min and for 3 h. See for more detail in Supplementary Figure S2A. (G) Western blot showing P-JNK and JNK levels in the TIF of neurons exposed to A $\beta$  oligomers (1  $\mu$ M) for 30 min and 3 h. See for more detail in Supplementary Figure S2B. (H) Western blot of caspase-3 activation in the TIF of neurons exposed to A $\beta$  oligomers (1  $\mu$ M) for 30 min and 3 h. See for more detail in Supplementary Figure S2C. (I–L) D-JNKI1 prevented A $\beta$ -induced synaptopathy *in vitro*. (I) Western blot and relative quantification were performed on the TIF of neurons treated with D-JNKI1 (2  $\mu$ M, 30 min before A $\beta$  oligomer application) and A $\beta$  oligomers (1  $\mu$ M, 3 h), either alone or combined. D-JNKI1 prevented A $\beta$  oligomer-induced protein loss from the PSD and caspase-3 activation in the postsynaptic compartment. See also Supplementary Figure S2D. (J) Time-lapse photographs showing neurons treated with D-JNKI1 (2  $\mu$ M, 30 min before A $\beta$  oligomer treatment) alone, A $\beta$  oligomers (1  $\mu$ M, 3 h) alone, or combined, recorded at time 0 (t0h) and 3 h after the treatment (t3h). Red arrows represent spines that disappeared; green arrows represent newly formed spines. (K and L) Quantification of dendritic spine density. (K) At the starting point (t0h), the total spine density was not significantly different between groups. Graphs show that D-JNKI1 (2  $\mu$ M, 30 min before A $\beta$  oligomer application) was able to prevent synthetic soluble A $\beta$  oligomer (1  $\mu$ M, 3 h)-induced reduction in spine density (K) and restore the number of mushroom, stubby, and thin spines to control levels (L) (two-way ANOVA, Bonferroni *post hoc* test, \*\*\* $P < 0.001$  CTR vs. A $\beta$  oligomers, #### $P < 0.001$  A $\beta$  oligomers vs. A $\beta$  oligomers + D-JNKI1, CTR  $n = 20$  cells, D-JNKI1  $n = 14$  cells, A $\beta$  oligomers  $n = 19$  cells, A $\beta$  oligomers + D-JNKI1  $n = 14$  cells). The number of spines recorded after 3 h of treatment (t3h) was normalized to the number of spines recorded at time 0 (t0h).