



## The first outbreak of brucellosis in the region of Šabac

### Prva epidemija bruceloze u šabačkom okrugu

Ljiljana Marković-Denić\*, Vesna Škodrić Trifunović†, Vladimir Žugić†,  
Dragana Radojčić‡, Goran Stevanović§

\*School of Medicine, Institute of Epidemiology, Belgrade, Serbia; Clinical Center of Serbia, †Institute of Pulmonary Diseases and Tuberculosis, ‡Institute of Infectious and Tropical Diseases, Belgrade, Serbia; §Institute of Public Health, Šabac, Serbia

#### Abstract

**Background/Aim.** In Serbia brucellosis is a primary disease of the animals in the southern parts of the country. The aim of this study was to describe the first outbreak of human and animal brucellosis in the region of Šabac, Serbia. **Methods.** An epidemiological investigation was conducted to identify a source of outbreak and the ways of transmission of brucellosis infection in human population. A descriptive and analytical epidemiological methods (cohort study) were used. Additional data included monthly reports of the infectious diseases from the Institutes of Public Health and data from the Veterinary Specialistic Institute in Šabac. The serological tests for human brucellosis cases were performed in the Laboratory of the Military Medical Academy; laboratory confirmation of animal brucellosis cases was obtained from the reference laboratory of the Faculty of Veterinary Medicine, Belgrade. **Results.** Twelve cases of brucellosis were recorded from February 9 to September 1, 2004. Total attack rate was 8.1% (7.5% of males, 14.2% of females). Relative risk (RR) of milk consumption was 8.9 (95% confidence interval: 1.63-13.38), and RR for direct contact with animals was 14 (95% confidence interval: 3.5-55.6). The prevalence of seropositive animals in 33 villages of the Mačva region accounted for 0.8%. Regarding animal species, sheep were predominant – 264 (95.7%). Out of a total number of seropositive animals, ELISA results were positive in 228 (88.7%) of them. **Conclusion.** As contact epidemics generally last longer, it is probable that the implemented measures of outbreak control did reduce the length of their duration.

#### Key words:

brucellosis; serbia; disease outbreaks; epidemiologic methods; serologic test; humans; animals.

#### Apstrakt

**Uvod/Cilj.** Bruceloza je prvenstveno oboljenje životinja u južnom delu Srbije. Cilj ovog rada bio je da prikaže prvu epidemiju humane i animalne bruceloze na području Šapca. **Metode.** Sprovedeno je epidemiološko istraživanje u cilju otkrivanja izvora i puteva prenošenja bruceloze u humanoju populaciji. Korišćen je deskriptivni i analitički (kohortna studija) epidemiološki metod. Analizirane su mesečne prijave zaraznih bolesti Zavodu za javno zdravlje Šabac i Veterinarskom specijalističkom institutu u Šapcu. Serološki test za humanu brucelozu rađen je na Vojnomedicinskoj akademiji, a potvrda animalne bruceloze na Veterinarskom fakultetu u Beogradu. **Rezultati.** Dvanaest slučajeva humane bruceloze otkriveno je u periodu od 9. februara do 1. septembra 2004. god. Stopa javljanja iznosila je 8,1% (7,5% kod muškaraca i 14,2% kod žena). Relativni rizik od konzumacije mleka iznosio je 8,9 (interval poverenja: 1,63–13,38), a za kontakt sa životinjama 14 (interval poverenja: 3,5–55,6). Prevalencija seropozitivnih životinja u 33 sela mačvanskog regiona iznosila je 0,8%. Najviše je bilo obolelih ovaca – 264 (95,7%). Od ukupnog broja seropozitivnih životinja, ELISA test bio je pozitivan kod njih 228 (88,7%). **Zaključak.** Iako kontaktne epidemije obično traju dugo, primenjene mere suzbijanja verovatno su doprinele kraćem trajanju ove epidemije.

#### Ključne reči:

bruceloza; srbija; epidemije; epidemiološki metodi; serološki testovi; ljudi; životinje.

#### Background

Brucellosis is one of the most common zoonotic infections of domestic and wild animals caused by members of the genus *Brucella*<sup>1</sup>. The humans become infected by di-

rect contact with infected animals or inhalation of infected aerosols, or through the consumption of raw animal products such as unpasteurized milk or cheese. In humans, ovine and caprine brucellosis caused by *B. melitensis* is the most important clinically apparent disease. Reported inci-

dence and prevalence of the disease vary widely from country to country. According to annual reports from the Centers for Disease Control and Prevention (CDC), the eradication campaign resulted in elimination of cattle brucellosis and substantial decline in the incidence of human disease in the USA. During 1993–2002, 46 states reported at least one case, mainly attributed to *B. melitensis*<sup>2</sup>. Mexico is one of the most important reservoirs of human brucellosis in Latin America<sup>3</sup>. The disease remains a major problem in the Mediterranean region, Western Asia and parts of Africa<sup>4</sup>. North European countries declared as “officially bovine and/or ovine and canine brucellosis free”<sup>5</sup>. Incidence of disease was reported high in the Balkan Peninsula, Greece, Macedonia, Albania and Bosnia-Herzegovina<sup>4,6</sup>. In Serbia, brucellosis used to be a primary disease of animals in the territories of Kosovo and Metohija and southern parts of the country.

The aim of this study was to describe the first outbreak of human and animal brucellosis in the region of Šabac, Mačva, Serbia.

### Methods

An epidemiological investigation was conducted to identify a source of outbreak and the ways of transmission of the brucellosis infection in human population. In addition, the sources of data included monthly reports on infectious diseases from the Institute of Public Health, Šabac<sup>7</sup> as well as the official reported data about infectious diseases for Serbia<sup>8</sup>. A descriptive and analytical epidemiological method (cohort study) were used. Relative risk (RR) and 95% confidence interval (CI) were calculated. Moreover, data were obtained from the Veterinary Specialistic Institute in Sabac, e.g., decisions on verified household infection and implemented measures were used<sup>9</sup>.

Brucellosis was diagnosed on the basis of clinical symptoms and signs compatible with brucellosis, supported by laboratory results, as it was recommended by the World Health Organisation<sup>10</sup>. A regional health center sent blood samples to the laboratory of the Military Medical Academy in Belgrade. In Serbia, PCR test, as a gold standard for diagnosis of brucellosis, was not in use during the period of the study. Consequently, a rapid agglutination reaction – BAB test (in Serbian: *brza serumska aglutinacija* – spead serra agglutination) for brucella antibodies with antigen solution prepared from brucella species of interest in region of Serbia was used. Detection of antibodies to brucellas was done in some patients by Wright serological reaction. This method was performed in the Institute of Public Health of Serbia in Belgrade.

The veterinarian investigation was conducted simultaneously with epidemiological investigation of human population. As serological test was positive in a animal sample, a laboratory confirmation of animal brucellosis cases was obtained from the reference laboratory of the Faculty of Veterinary Medicine, Belgrade, using the enzyme-linked immunosorbent assay (ELISA).

### Results

Twelve cases of human brucellosis were recorded during the outbreak, which developed in the Šabac district from February 9 to September 1, 2004. Figure 1 shows the distribution of patients presented with human brucellosis. The epidemic curve suggested common and persistent source of contamination. The largest number of patients was recorded in March. Ten patients were males and two patients females, and their age ranged from 20 to 59 years. A severe clinical picture was manifested in eight patients (fever, muscular pain, headache, spleno- and hepatomegaly), so they were hospitalized. A prompt response to antibiotics was seen in all the cases but one. No case had any former history of brucellosis.




Fig. 1 – The distribution of patients with brucellosis in Šabac district (February 9 to September 1, 2004)

All the cases were positive by BAB test with titers ranging from 1/80 to 1/1280. These findings with positive antibodies tests in some patients were diagnostic criteria for *Brucella* infections. Isolation of *Brucella spp* organisms were not done routinely in human material, but it was done in animals.

The first (index) case in the outbreak was performed in a sheep and goat farm in the village Miokus near Šabac. The patient mentioned that, during his work in the farm, he used to drink freshly milked milk. Upon joining the veterinarian in the outbreak inspection, it was found that sheep from the aforementioned farm in Miokus had been sold in three more villages before the first case of brucellosis was diagnosed. Laboratory analyses of blood cultures collected from sheep, goats and cows of these farms revealed infected 93 head of cattle. On the basis of these results, epidemiologic cohort study was initiated in the families with the established infection of animals. A total number of the subjects exposed to epidemic, *ie* family members where animal infection had been verified were 148. Considering that 12 persons were affected during the epidemic, a total attack rate was 8.1% (7.5% of males, 14.2% of females). The highest attack rate was in the age group from 20 to 29 years (33.3%). All the cases and family members were interviewed to discover the mode of transmission. With the exception of the index case in the outbreak, other patients could not confirm a consumption of non-boiled milk, so RR of milk consumption was 8.9



excretions. By veterinary service regulations, the sheep were slaughtered already at the beginning of March.

Detection of a larger number of infected animals in the villages of Mačva suggests that the mode of transmission was most probably from animals having been bought in other part of Serbia or neighboring countries. The prevalence of the infected animals in this region (0.8%) was significantly lower than the prevalence of seropositive animals in Kosovo, where in 26 out of 29 districts 6.3% of seropositive sheep, 7.2% of goats and 0.6% of bovine cattle were recorded<sup>17</sup>. There is an increase in the number of reported outbreaks in the ruminants of Bosnia and Herzegovina<sup>18</sup>. An outbreak of human brucellosis in southern Croatia in 2004, the region with no brucellosis history, was

accounted for illegal transport of goats and sheep from Bosnia and Herzegovina<sup>19</sup>. The possible illegal trafficking of animals and their products across the borders of surrounding countries could be responsible for the persistence of foci of brucellosis in the area<sup>20</sup>.

### Conclusion

This outbreak shows selling of animals without any adequate control measures higher risk of spread of disease in animals. Contact with animals without any appropriate preventive measures may lead to infection of people, and consequently, to epidemic.

### R E F E R E N C E S

1. *World Health Organization*. Brucellosis in human and animals. Geneva: World Health Organization, WHO/CDS/EPR/2006.7; 2006 [cited 2009 Jan 7]. Available from: [www.cdc.gov/eid/content/15/2/pdfs/314.pdf](http://www.cdc.gov/eid/content/15/2/pdfs/314.pdf)
2. *Centers for Disease Control and Prevention*. Summary of notifiable diseases – United States, 2006. [cited 2008 March 21]. Available from: <http://www.cdc.gov/mmwr/summary.html>
3. Luna-Martínez JE, Mejía-Terán C. Brucellosis in Mexico: current status and trends. *Vet Microbiol* 2002; 90(1–4): 19–30.
4. Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV. The new global map of human brucellosis. *Lancet Infect Dis* 2006; 6(2): 91–9.
5. Godfroid J, Käsbobrer A. Brucellosis in the European Union and Norway at the turn of the twenty-first century. *Vet Microbiol* 2002; 90(1–4): 135–45.
6. Taleski V, Zerva L, Kantardžiev T, Cvetnic Z, Erski-Biljic M, Nikolovski B, et al. An overview of the epidemiology and epizootology of brucellosis in selected countries of Central and Southeast Europe. *Vet Microbiol* 2002; 90(1–4): 147–55.
7. *Institute of Public Health Šabac*. Monthly reports of the infectious diseases. Šabac: Institute of Public Health; 2004. (Serbian)
8. *Institute of Public Health of Serbia*. Infectious diseases in Serbia 1994–2005. Belgrade: Institute of Public Health of Serbia; 2005. (Serbian)
9. *Veterinary Specialist Institute Šabac*. Šabac-decisions about the infected household and measures undertaken (No 323 – 432 / 04 - 05 from April 23, 2004. to No 323 – 845 / 04–05 from July 26, 2004.) Šabac:Veterinary Specialist Institute; 2004. (Serbian)
10. *World Health Organization*. WHO Recommended Surveillance Standards. 2nd ed. WHO/CDS/CSR/ISR/99.2; Geneva: World Health Organization; 1999.
11. *Institute of Public Health of Serbia*. Infectious diseases in Serbia, 2003. Belgrade: Institute of Public Health of Serbia; 2004. (Serbian)
12. *Institute of Public Health of Serbia*. Infectious diseases in Serbia, 2001. Belgrade: Institute of Public Health of Serbia; 2002. (Serbian)
13. Méndez Martínez C, Pérez Jiménez A, Cortés Blanco M, Salmoral Chamizo E, Mohedano Mohedano E, Plata C, et al. Brucellosis outbreak due to unpasteurized raw goat cheese in Andalucía. (Spain), January - March 2002. *Eurosurveillance* 2003; 8: 164–8.
14. Castell Monsalve J, Rullán JV, Peiró Callizo EF, Nieto-Sandoval Alcolea A. Epidemic outbreak of 81 cases of brucellosis following the consumption of fresh cheese without pasteurization. *Rev Esp Salud Publica* 1996; 70(3): 303–11.
15. Park MY, Lee CS, Choi YS, Park SJ, Lee JS, Lee HB. A sporadic outbreak of human brucellosis in Korea. *J Korean Med Sci* 2005; 20(6): 941–6.
16. Landau Z, Green L. Chronic brucellosis in workers in a meat-packing plant. *Scand J Infect Dis* 1999; 31(5): 511–2.
17. Jackson R, Pite L, Kennard R, Ward D, Stack J, Domi X, et al. Survey of the seroprevalence of brucellosis in ruminants in Kosovo. *Vet Rec* 2004; 154(24): 747–51.
18. Šerić-Haračić S, Salman M, Fejzić N, Čavajuga S. Brucellosis of ruminants in Bosnia and Herzegovina: disease status, past experiences and initiation of a new surveillance strategy. *Bosn J Basic Med Sci* 2008; 8(1): 27–33.
19. Punda-Polić V, Cvetinić Z. Human brucellosis in Croatia. *Lancet Infect Dis* 2006; 6(9): 540–1.
20. Andikou I, Maipa V, Alamanos Y. Epidemiology of human brucellosis in a defined area of north western Greece. *Epidemiol Infect* 2005; 133(5): 905–10.

Received on May 19, 2009.  
Revised on September 20, 2009.  
Accepted on December 4, 2009.