



THE CEYLON MEDICAL JOURNAL

Established 1887

The Official Publication of the
Sri Lanka Medical Association
Volume 56, No.4, December 2011
Quarterly ISSN 0009-0875

All communications
should be addressed to
The Editors, CMJ

Editors Emeritus

Chris G Uragoda MD, FRCP
Colvin Goonaratna FRCP, PhD

Editors

Janaka de Silva DPhil, FRCP
Anuruddha Abeygunasekera MS, FRCS

Assistant Editors

Dennis Aloysius MBBS, FCGP
D N Atukorala MD, FRCP
Sarath Gamini de Silva MD, FRCP
S A S Goonewardena MS, FRCS
Dulani Gunasekara MD, FRCP
A Pathmeswaran MBBS, MD
Lalini Rajapakse MD, MSc
Channa Ranasinha MRCP, DTM & H
Udaya Ranawaka MD, MRCP
Koliitha Sellahewa MD, FCCP
B J C Perera MD, FRCP
Harshalal R Seneviratne DM, FRCOG
Shalini Sri Ranganathan MD, PhD

International Advisory Board

Kamran Abbasi MBChB, MRCP
London, UK

Raja Bandaranayake FRACS, PhD
Sydney, Australia

Peush Sahni MS, PhD
New Delhi, India

R K Tandon MD, PhD
New Delhi, India

Zulfiqar Ahmed Bhutta FRCPCH, PhD
Karachi, Pakistan

Continued overleaf

Chronic kidney disease of unknown aetiology in the North Central Province of Sri Lanka: trying to unravel the mystery

The 'battle' over establishing the cause of chronic kidney disease of unknown aetiology (CKDu) in the North Central Province (NCP), the seat of Sri Lanka's ancient civilisation, has been highlighted in many newspapers, television and radio programmes in recent years. Whilst impartiality in media reports sounds good in principle, and is appropriate for many subjects, it can misrepresent science unless genuine equipoise exists [1]. Because media reports about science can affect an individual's health and environment, the media also have a duty by the public to report science accurately, in a manner that fairly represents the weight of evidence. To enable this, scientists must take responsibility in allowing high quality evidence to develop over time, come to a consensus and make statements with a good degree of certainty.

Ever since CKDu became an issue in the NCP, many experts have worked to both investigate and treat patients as well as to identify associated risk factors and possibly its cause. In 2008, the World Health Organisation together with the Epidemiology Unit of the Ministry of Healthcare and Nutrition developed a draft research proposal to provide a framework for a programme of research activities, the main aim of which was to estimate the prevalence and identify the determinant(s) of CKDu. This research programme discouraged other scientists exploring the issue to the detriment of scientific enquiry and the wellbeing of residents in the NCP. As we approach the end of 2011, there is still little scientific evidence of the determinants of kidney disease or its aetiology.

Given the widespread discussion and debate in the media recently, it is timely that the available, credible, scientific evidence on CKDu (published in peer reviewed journals) is collated and analysed, and the difficulties faced in establishing causality are discussed.

Bibliographic computerised searches using the Ovid search engine located 12 articles on CKDu published in peer-reviewed scientific journals. In 2009, Athuraliya *et al* described CKDu patients as being predominantly young males from low socio-economic farming communities; patients presented with non-specific symptoms, mild proteinuria and had bilateral echogenic small kidneys on ultrasound examination [2]. However, this year, the same group reported the occurrence of proteinuric-chronic kidney disease in the NCP to be strongly associated with an age over 60 years and farming; specific risk factors were not identified in 87% of the patients [3]. A possible association between CKDu and exposure to organophosphate pesticides was reported in 2006 [4]. A later study found evidence of clustering suggesting an environmental aetiology, and reported that a family history of renal disease, using ayurvedic medications and snakebite were significant predictors of

Samiran Nundy FRCS, FRCP
New Delhi, India

N Medappa MD
New Delhi, India

Jane Smith BA, MSc
London, UK

Anita KM Zaidi MMBS, SM
Karachi, Pakistan

David Warrell MD, FRCP
Oxford, UK

**Advisory Board for
Statistics and Epidemiology**

Lalini Rajapakse MD, MSc

Kumudu Wijewardene MBBS, MD

A Pathmeswaran MBBS, MD

Published by

The Sri Lanka Medical
Association
Wijerama House
6, Wijerama Mawatha
Colombo 7
Sri Lanka

Tel: +94 11 2693324

Fax: +94 11 2698802

Internet home page

<http://www.sljol.info/index.php/CMJ/index>

e-mail: office@cmj.slma.lk

Printed by

Ananda Press
82/5, Sir Ratnajothi Saravanamuttu
Mawatha, Colombo 13
Sri Lanka

Tel: +94 11 2435975

Fax: +94 11 2385039

e-mail: anpress@sltnet.lk

For advertising

Please contact:

Mr. Anthony

Saatchi & Saatchi

79, C W W Kannangara Mawatha
Colombo 7

Tel: +94 11 2671026

+94 772514858

© The Ceylon

Medical Journal

**This journal is indexed by BIOSIS,
Elsevier SCOPUS, EMBASE, CABI, and
Index Medicus/Medline**

CKDu (5); being a farmer, using pesticides and drinking well water were also associated with CKDu [5]. Nanayakkara *et al* reported evidence of familial clustering of patients suggestive of a polygenic inheritance pattern comparable to that associated with multifactorial diseases [6]. They also demonstrated that renal tubular damage occurred at a very early stage of CKDu [6]. Another recent study found that hypertension, diabetes mellitus, urinary tract infections, drinking well water in the fields, smoking and pesticide spraying were significantly associated with microalbuminuria [7]; pesticide spraying appeared less common in patients with CKDu, and this may have been due to a selection bias where patients with CKDu may not have been involved in recent spraying activity due to ill health.

From the time an environmental aetiology was first suggested several investigators have attempted to identify the agent(s) responsible with a particular focus on medical and geomedical studies. Much attention has been drawn to cadmium (Cd). In 2008, Bandara *et al* provided evidence of NCP residents being exposed to high levels of Cd from reservoir water, irrigated rice, lotus rhizomes, freshwater fish and milk from cows reared in Cd-contaminated pastures [8]. In 2010, they claimed that dietary Cd was the cause of CKDu [9]. Their findings have been challenged; Chandrajith *et al* [10] reported that the Cd content in water sources and rice from affected regions was much lower than the levels found by Bandara *et al* [8, 9], and also showed that urinary Cd excretion was low in both CKDu patients and asymptomatic individuals from endemic and non-endemic (Kandy) areas. Nanayakkara *et al* too demonstrated low urinary Cd excretion in both CKDu patients and their unaffected relatives compared to Japanese controls [6]. More recently, Chandrajith *et al* investigating water sources in CKDu affected regions found moderate to high levels of fluoride (F1) in the water [10]. They also found evidence of a very low Na⁺:Ca²⁺ ratio in water in these regions compared to water in non-endemic regions (Huruluwewa and Wellawaya), and suggested that renal tubular damage due to F1 was dependent on the low Na⁺:Ca²⁺ ratio in drinking water [11]. A study investigating food samples, cultivated and consumed by people in regions where CKDu is prevalent, for ochratoxin A, found levels to be below the statutory maximum, and concluded that the toxin was unlikely to be a potential risk factor [12]. However, a more recent study has detected higher rates of urinary ochratoxins in CKDu patients in the NCP and their unaffected relatives than in Japanese controls, suggesting that exposure to the toxin may be common in the region [13].

Epidemiologically, establishing causality is difficult. Sir Bradford Hill outlined the minimum conditions that are required to establish causality, and these form the basis of modern epidemiologic research [14]. The conditions are:

- 1) Temporal relationship: the causative factor or agent must necessarily always precede the outcome.
- 2) Strength of the association: the strength or the size of the association is measured by appropriate statistical tests (relative risks, odds ratios and *p*-values). The larger the risk estimate or smaller the *p*-value, the greater the evidence for causality provided that other conditions are also fulfilled. The strength of the association is also dependent on the epidemiological study design used, with randomized controlled trials and cohort studies providing the strongest evidence, and cross sectional studies providing the weakest.
- 3) Dose response relationship: an increase in the amount of exposure should increase the risk or vice versa. If a dose-response relationship is present, it is strong evidence for a causal relationship. However, the absence of a dose-response relationship does not rule out a causal relationship, because

a threshold may exist above which a relationship may develop. If a specific factor is the cause of a disease, the incidence of the disease should decline when exposure to the factor is reduced or eliminated.

- 4) Consistency: The association is consistent when results are replicated in studies in different settings using different methods. Hence, numerous experiments have to be done before meaningful statements can be made about the causal relationship between two or more factors.
- 5) Plausibility: The association should agree with currently accepted understanding of pathological processes. There needs to be some theoretical basis for causality either in humans or in animals. However, research that disagrees with established theory is not necessarily false; it may, in fact, force a reconsideration of accepted beliefs and principles.
- 6) Consideration of alternate explanations: To establish causality, it is not only necessary to determine an association, but equally important is to rule out other explanations. It is always necessary to consider multiple hypotheses before making conclusions about the causal relationship between any two items.
- 7) Experiment: The condition should be modifiable (prevented or ameliorated) by an appropriate experimental regimen.
- 8) Specificity: Specificity is established when a single putative cause produces a specific effect. This is considered by some to be the weakest of all the criteria and may not be applicable to diseases of multifactorial origin. When specificity of an association is found, it provides additional support for a causal relationship but the absence of specificity in no way negates a causal relationship.
- 9) Coherence: The association should be compatible with existing theory and knowledge. As with plausibility, research that disagrees with established theory and knowledge are not automatically false. They may, in fact, force a reconsideration of accepted beliefs and principles.

The *minimum* criteria listed above makes it daunting to establish the cause of CKDu in the NCP of Sri Lanka; research done to date does not come even close to fulfilling these criteria. Mere associations, though encouraging, should not be considered to be of causal importance without documented evidence of proof. We advocate caution by the scientific community and the media when using such assertions that impact on policy, the livelihoods of the farming community and the economy of the country without scientific validity and biological plausibility.

The cause of CKDu is likely to be multifactorial. At this point in time there is insufficient evidence to pinpoint a cause(s). Both the wellbeing of residents of the NCP

and the enormous drain on health system resources and the economy demand that resolving the issue is a national priority. Well designed and adequately powered studies into CKDu should be strongly encouraged and adequately funded.

References

1. Science journalism: a delicate balancing act (Editorial). *Lancet* 2011; **378**: 374.
2. Athuraliya NTC, Abeysekera TDJ, Amerasinghe PH, Kumarasiri R, Dissanayake V. Prevalence of chronic kidney disease in two tertiary care hospitals: high proportion of cases with uncertain aetiology. *Ceylon Medical Journal* 2009; **54**: 23-5.
3. Athuraliya NTC, Abeysekera TDJ, Amerasinghe PH, *et al.* Uncertain etiologies of proteinuric-chronic kidney disease in rural Sri Lanka. *Kidney International* 2011 (Advance online publication 10 August 2011; doi: 10.1038/ki.2011.258).
4. Peiris-John RJ, Wanigasuriya JK, Wickremasinghe AR, Dissanayake WP, Hittarage A. Exposure to acetylcholinesterase-inhibiting pesticides and chronic renal failure. *Ceylon Medical Journal* 2006; **51**: 42-3.
5. Wanigasuriya KP, Peiris-John RJ, Wickremasinghe R, Hittarage A. Chronic renal failure in North Central Province of Sri Lanka: an environmentally induced disease. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2007; **101**: 1013-7.
6. Nanayakkara S, Senevirathna ST, Karunaratne U, *et al.* Evidence of tubular damage in the very early stage of chronic kidney disease of uncertain etiology in the North Central Province of Sri Lanka: a cross-sectional study. *Environmental Health and Preventive Medicine* 2011 (Advance online publication 27 June 2011; DOI: 10.1007/s12199-011-0224-z).
7. Wanigasuriya KP, Peiris-John RJ, Wickremasinghe R. Chronic kidney disease of unknown aetiology in Sri Lanka: is cadmium a likely cause? *BMC Nephrology* 2011; **12**: 32-8.
8. Bandara JMRS, Senevirathna DMAN, Dasanayake DMRSB, *et al.* Chronic renal failure among farm families in cascade irrigation systems in Sri Lanka associated with elevated dietary cadmium levels in rice and freshwater fish (Tilapia). *Environmental Geochemistry and Health* 2008; **30**: 465-78.
9. Bandara JMRS, Wijewardena HVP, Liyanage J, Upul MA, Bandara JMUA. Chronic renal failure in Sri Lanka caused by elevated dietary cadmium: Trojan horse of the green revolution. *Toxicology Letters* 2010; **198**: 33-9.
10. Chandrajith R, Nanayakkara S, Itai K, *et al.* Chronic kidney diseases of uncertain etiology (CKDu) in Sri Lanka: geographic distribution and environmental implications. *Environmental Geochemistry and Health* 2011; **33**: 267-78.
11. Chandrajith R, Dissanayake CB, Ariyaratna T, Herath HMJMK, Padmasiri JP. Dose-dependent Na and Ca in

- fluoride-rich drinking water-another major cause of chronic renal failure in tropical arid regions. *Science of the Total Environment* 2011; **409**: 671-5.
12. Wanigasuriya KP, Peiris H, Ileperuma N, Peiris-John RJ, Wickremasinghe R. Could ochratoxin A in food commodities be the cause of chronic kidney disease in Sri Lanka? *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2008; **102**: 726-8.
13. Desalegn B, Nanayakkara S, Harada KH, *et al.* Mycotoxin detection in urine samples from patients with chronic kidney disease of uncertain etiology in Sri Lanka. *Bulletin of Environmental Contamination and Toxicology* 2011; **87**: 6-10.
14. Hill AB. The Environment and Disease: Association or Causation? *Proceedings of the Royal Society of Medicine* 1965; **58**: 295-300.

A R Wickremasinghe, Department of Public Health, Faculty of Medicine, University of Kelaniya, Sri Lanka, **R J Peiris-John**, Section of Epidemiology and Biostatistics, School of Population Health, Faculty of Medical and Health Sciences, University of Auckland, New Zealand, **K P Wanigasuriya**, Department of Medicine, Faculty of Medical Sciences, University of Sri Jayewardenepura, Sri Lanka.

Correspondence: ARW, e-mail <arwicks@sltnet.lk>. Competing interests: none declared.