



Cite this: *RSC Adv.*, 2019, 9, 33130

# Distribution, contents and health risk assessment of heavy metal(loid)s in fish from different water bodies in Northeast China

XiaoLi Huang,<sup>ab</sup> DongLi Qin,<sup>id</sup><sup>b</sup> Lei Gao,<sup>b</sup> Qirui Hao,<sup>b</sup> Zhongxiang Chen,<sup>b</sup> Peng Wang,<sup>b</sup> Shizhan Tang,<sup>b</sup> Song Wu,<sup>b</sup> Haifeng Jiang<sup>b</sup> and Wei Qiu<sup>\*ac</sup>

This study aimed at investigating the accumulation and potential risks of eight metal(loid)s in fish from natural and culturing water samples in Northeast China. Chromium (Cr), nickel (Ni), copper (Cu), zinc (Zn), arsenic (As), cadmium (Cd), lead (Pb) and mercury (Hg) contents in 16 fish species (155 samples) and sediments of their habitats were analyzed. In general, the concentrations of these eight metal(loid)s in most fish samples are lower than the guideline levels and legal limits, and the Pb and Hg level in 0.65% of samples were exceeded the quality standards in China. The Cr, As, Cd and Pb levels in most fish samples are less than those reported in previous studies. Nonetheless, Hg levels in these fish samples are significantly higher than those reported in previous studies conducted in other regions. Different from the wild fish, significant positive correlations are found between Cr, Ni, As and Cd concentrations in cultured fish and those in pond sediment ( $P < 0.05$ ), which might be due to the closed static water environment and concentrated feeding operations. Cu, Zn, As and Hg concentrations differed significantly among wild species, while Cu and Zn concentrations differed significantly among cultured species ( $P < 0.05$ ), which might be because of the different feeding and foraging habitats. The target hazard quotients (THQs) at high exposure levels of target metal(loid)s in the studied fish were below 1 (except for Hg), and the carcinogenic risk indices of Cr, As, and Cd were less than  $10^{-4}$ . The levels of metal(loid)s (except for Hg) in the studied fishes fell within an acceptable range, but more attention should be paid to the potential carcinogenic risks.

Received 9th July 2019  
 Accepted 11th September 2019

DOI: 10.1039/c9ra05227e

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## 1 Introduction

Increasing human activities have caused the increasing release of heavy metals into the hydrosphere.<sup>1,2</sup> This is especially the case for the surface water, which turns into a sink for heavy metals from the wastes (*e.g.*, emission, wastewater and waste solids). Studies have found high levels of heavy metals being released into freshwater ecosystems because of the effects of urbanization and industrialization.<sup>3</sup> Some rivers and lakes have encountered different levels of ecological risks due to heavy metal pollution.<sup>4-7</sup> Fishes exposed to elevated metal levels can absorb the bioavailable metals (*e.g.*, cadmium (Cd), nickel (Ni), lead (Pb), and mercury (Hg)) directly from an aquatic environment<sup>8</sup> and regulate metal concentrations where bioaccumulation will occur in tissues to some extent.<sup>9</sup> Researchers have found evidence that supports the biomagnification of

some heavy metals (*e.g.*, chromium (Cr), copper (Cu), arsenic (As), Cd, Pb and Hg) with trophic levels in marine and freshwater ecosystems.<sup>10-14</sup> Bioaccumulation and biomagnification can cause heavy metal burdens in the muscle tissues of fish to increase.<sup>8,15</sup> Excessive levels of heavy metals could harm aquatic organisms, causing the escalation of heavy metal levels in fish tissues to dangerous levels.<sup>16</sup> The consumption of polluted fish could potentially result in serious food poisoning among the general public.

In general, heavy metal levels in a fish are related to its living environment, feeding behavior and foraging habitats.<sup>17,18</sup> There are numerous studies on the relationship between heavy metal pollution in the environment and in fish. For instance, sediment acts as an important sink for heavy metals, which affects the heavy metal bioconcentration by affecting bioavailability.<sup>19</sup> Because of the biogeochemical characteristics and pathways of trophic transfer, the metal burdens in food web components are different, which lead to the difference in individual fish taxa by functional feeding groups.<sup>20,21</sup> Previous studies have suggested that the foraging habitat is a strong predictor for variations in heavy metal concentrations in fish.<sup>21</sup> While heavy metal contamination of freshwater and marine biota in China has been well described,<sup>10,12,17,18,22</sup> such investigations are rather

<sup>a</sup>State Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology, 150090, Harbin, Heilongjiang, China. E-mail: [qiuweihit@126.com](mailto:qiuweihit@126.com)

<sup>b</sup>Heilongjiang River Fisheries Research Institute, Chinese Academy of Fishery Sciences, 150070, Harbin, Heilongjiang, China

<sup>c</sup>School of Municipal & Environmental Engineering, Harbin Institute of Technology, 150090, Harbin, Heilongjiang, China



limited in Northeast China. Some heavy metals (*i.e.*, As, Cd, Pb and Hg) are of major environmental concern as they can cause severe health implications for humans and other living organisms. Moreover, some elements (*e.g.*, Cr, Ni, Cu and zinc (Zn)) that play essential roles in life activities become toxic in excess amounts.

Northeast China is a water-rich area with satisfactory water resource conditions and long fishing history, especially the Amur River basin, which contains a rich freshwater ecosystem with various fish species from the frigid-zone, temperate zone and even the tropical zone. In Northeast China, aquatic environments such as rivers, lakes, artificial reservoirs and ponds provide all types of aquaculture. However, many major cities and chemical enterprises in Northeast China are located near rivers and lakes, where the released waste threatens these aquatic ecosystems. Recently, two serious environmental contamination accidents have occurred in the Songhua River, which is the third-largest river in China and one of the biggest tributaries of the Amur River.<sup>23,24</sup> Studies have found elevated heavy metal concentrations in the water and sediment in some rivers and lakes in Northeast China.<sup>25–27</sup> Studies have been conducted to investigate the trace elements in the tissues of fish from fish farms<sup>28</sup> and individual rivers or areas in Northeast China.<sup>29</sup> Nonetheless, little is known about the heavy metal levels in fish from Northeast China, particularly for various species with different feeding habits, and the relationship between heavy metals and the environment.

This study aims to develop a large-scale analysis to compare eight metal(loid) levels (including Cr, Ni, Cu, Zn, As, Cd, Pb and Hg) in fish from different water types in Northeast China, investigate the relationship between heavy metals and the environment as well as evaluate the potential health risks associated with fish consumption.

## 2 Materials and methods

### 2.1 Study area and sample collection

In the period from May to September of 2015, a total of 155 fish samples representing 16 species were randomly collected from various fishery areas, including the trunk stream of Amur River (Amur R.), Songhua River (Songhua R.), Ussuri River (Ussuri R.), Second Songhua River (SSonghua R.) and Nen River (Nen R.), Chagan Lake (CGL), Songhua Lake (SHL), Xingkai Lake (XKL), Jingpo Lake (JPL) and some ponds at city sites (Harbin, Jiamusi, Zhaodong, Changchun, Jilin and Songyuan) (Fig. 1). Sediment samples were also collected from the same localities. Basic biological information on the sampled fish, *e.g.*, species, number, feeding habits, age, total length and body weights, are displayed in Table 1. The dorsal muscle tissues of fish samples were taken and homogenized. Fish samples were put into a clean polythene bag and immediately stored at  $-20\text{ }^{\circ}\text{C}$  before the subsequent heavy metal analysis. A total of 235 sediment samples were extracted from the upper layer (0–10 cm) of the sampling sites (five sampling sites in each sampling area) using polyvinyl chloride corers. The sediment samples were immediately mixed and 2.0 kg were reserved in a hermetic bag and stored at  $4\text{ }^{\circ}\text{C}$  before arriving at the laboratory.

### 2.2 Analysis of metal(loid)s

In a report by Qin *et al.*,<sup>28</sup> target metal(loid)s except Hg were determined using an inductively-coupled plasma mass spectrometer (ICP-MS, Agilent 7500cx, USA) equipped with an octopole reaction system (ORS). For fish muscle (epaxial muscle on the dorsal surface without skin and backbone) analysis, each 1.0 g of the homogenized sample was digested and mixed using an automatic microwave system with a mixture of  $\text{HNO}_3 : \text{H}_2\text{O}_2 : \text{Milli-Q water} = 5 : 2 : 1$ . After cooling, the sample was diluted with Milli-Q water (Milli-Pore Co., USA; resistivity, 18.2  $\text{M}\Omega\text{ cm}$ ), stored at  $4\text{ }^{\circ}\text{C}$  temperature, and analyzed within 24 h. For sediment samples, each 0.1 g of dried and powdered sediment sample was digested with mixed acids ( $\text{HClO}_4 : \text{HF} : \text{HCl} = 1 : 5 : 1$ ) in an automatic microwave digestion system. After cooling, the sample was distilled to dryness and diluted with Milli-Q water. For analysis of the Hg content, an Automatic Mercury Analyzer Hydra IIC (Teledyne Leeman Labs Hudson, NH, USA) was used.

### 2.3 QA/QC

Nitric acid, peroxide hydrogen and hydrochloric acid utilized in this study were all guaranteed reagents. Glassware used in this study were all immersed in 10% (V/V) of nitric acid overnight, washed three times with Milli-Q water, and then dried in an oven. The certified reference solution ICP Standards of Cr, Ni, Cu, Zn, As, Cd and Pb ( $1000\text{ }\mu\text{g mL}^{-1}$ ) were all purchased from the National Analysis and Testing Center for Nonferrous Metals & Electronic Materials. Yellow croaker (GBW08573), GSS-3 (GBW-07403) and GSS-5 (GBW-07405) were utilized as processing Certified Reference Materials for analysis validation. The results were in good agreement with the certified values and recoveries ranged from 86.5% to 104.3%. Analytical blanks were prepared in parallel in the same way as the samples for evaluating the possibility of pollution and background interference. For fish samples, the limits of detection (LOD,  $\text{mg kg}^{-1}$  wet weight) were the following: Cr 0.005, Ni 0.010, Cu 0.008, Zn 0.015, As 0.003, Cd 0.001, Pb 0.005, and Hg 0.001. For sediment samples, the limits of detection (LOD,  $\text{mg kg}^{-1}$  wet weight) were the following: Cr 0.180, Ni 0.100, Cu 0.100, Zn 0.700, As 0.190, Cd 0.003, Pb 0.040, and Hg 0.001.

### 2.4 Risk assessment

**2.4.1 Estimated daily intake (EDI).** The estimated daily intake (EDI) for each heavy metal element (*i.e.*, Cr, Ni, Cu, Zn, As, Cd, Pb, and Hg) was calculated based on the following equation:

$$\text{EDI} = \frac{\text{FIR} \times C_i}{\text{BW}}$$

where FIR is the food ingestion rate of fish and fish products (g per person per day, wet weight),  $C_i$  is the average concentration of a single heavy metal in a fish muscle ( $\text{mg kg}^{-1}$ ), and BW is the average body weight of an adult.<sup>30</sup> For adults in rural and urban areas of China, the food ingestion rate of fish was 23.7 g per person per day.<sup>31</sup> The average body weight of an adult is 60 kg. If the element concentration was lower than the limit of detection






Fig. 1 Fish sampling locations in the study area.

(LOD), the value would be taken as half of the respective detection limits ( $\frac{1}{2}$  LOD).

**2.4.2 Non-carcinogenic risk.** To express the potential non-carcinogenic risk of fish samples in this study, the target hazard quotient (THQ) was utilized. The THQ is equal to the ratio of an EDI value to an oral reference dose (RfD,  $\mu\text{g}$  per kg per day) value. The RfD was established by the United States Environmental Protection Agency.<sup>32</sup> The THQ for each analyzed metal (Cr, Ni, Cu, Zn, As, Cd, Pb, and Hg) was calculated by using the following formula:

$$\text{THQ} = \frac{\text{EDI}}{\text{RfD}}$$

When the THQ value is less than one, there should be no obvious non-carcinogenic risks for the exposed population. Otherwise, the exposed population would experience adverse health risks. Such concerns would further increase with a higher THQ value.

**2.4.3 Carcinogenic risk.** The carcinogenic risk index (CRI) is used for evaluating the potential carcinogenic risks of fish consumption.<sup>33</sup> The CRIs for elements Cr, As, and Cd are calculated as follows:

$$\text{CRI}_i = \frac{\text{FIR} \times E_F \times E_D}{\text{BW} \times T_A} \times \text{SF}_i \times C_i$$

where  $E_F$  is the exposure frequency (365 d per year),  $E_D$  is the exposure duration (70 year),  $T_A$  is the averaging time ( $365 \times E_D$  d), and  $\text{SF}_i$  is the oral slope factor ( $(\text{mg kg}^{-1} \text{d}^{-1})^{-1}$ ) for a single heavy metal. The oral intake of carcinogenic slope factors for Cr, As, and Cd are 0.50, 0.38 and 1.50, respectively. When the CRI value is less than  $10^{-6}$ , the compound is safe for humans. When the CRI value is between  $10^{-6}$  and  $10^{-4}$ , there are potential carcinogenic risks to the exposed population. When the CRI value is more than  $10^{-4}$ , the exposed people would encounter excess carcinogenic risk.

## 2.5 Statistical analysis

Statistical analysis was performed using the SPSS software (IBM SPSS statistics for Windows, version 20.0; SPSS Inc., Chicago, IL). The Mann-Whitney test was performed to compare the metal(loid)s concentration in fish samples captured from natural waters and that from aquaculture ponds. Metal(loid) concentrations in fish and sediment samples were  $\log(x)$  transformed and their relationships were expressed using the Pearson correlation test. The linear regression analyses were



Table 1 Basic biological information of fish in this study<sup>a</sup>

Scientific name	English name	<i>n</i>	Feeding habits	Foraging habitats	The total length (cm)	Body weights (kg)
<i>Hypomesus olidus</i>	Pond smelt	1	Omnivory	Pelagic fishes	10.20	0.0085
<i>Esox reicherti</i>	Amur pike	6	Sarcophagy	Pelagic fishes	50.60 ± 2.40 (47.00–55.00)	0.64 ± 0.14 (0.52–0.91)
<i>Leuciscus waleckii</i>	Amur ide	3	Omnivory	Pelagic fishes	23.90 ± 3.60 (19.00–29.00)	0.14 ± 0.093 (0.035–0.28)
<i>Ctenopharyngodon idellus</i>	Grass carp	19	Herbivory	Middle-lower layers fishes	53.00 ± 11.20 (38.00–72.00)	1.93 ± 1.14 (0.71–4.23)
<i>Hemibarbus labeo</i>	Barbell steed	2	Omnivory	Bottom fishes	26.00 ± 4.90 (22.50–29.50)	0.16 ± 0.09 (0.095–0.23)
<i>Hemibarbus maculatus</i>	Spotted steed	1	Omnivory	Middle-lower layers fishes	28.00	0.22
<i>Saurogobio dabryi</i>	Chinese lizard gudgeon	1	Omnivory	Middle-lower layers fishes	13.70	0.031
<i>Erythroculter ilishaeformis</i>	Topmouth culter	7	Sarcophagy	Pelagic fishes	54.90 ± 8.50 (42.00–69.00)	1.23 ± 0.46 (0.71–2.06)
<i>Erythroculter mongolicus</i>	Mongolian redfin	1	Sarcophagy	Pelagic fishes	24.50	0.22
<i>Cyprinus carpio</i>	Common carp	39	Omnivory	Bottom fishes	42.80 ± 7.40 (29.00–61.00)	1.29 ± 0.87 (0.32–4.40)
<i>Carassius auratus</i>	Crucian carp	36	Omnivory	Bottom fishes	21.80 ± 5.50 (8.80–29.00)	0.22 ± 0.14 (0.019–0.47)
<i>Hypophthalmichthys molitrix</i>	Silver carp	11	Filter feeder (zooplankton, phytoplankton)	Pelagic fishes	47.30 ± 11.90 (26.00–61.00)	1.22 ± 0.86 (0.15–3.65)
<i>Aristichthys nobilis</i>	Bighead carp	12	Filter feeder (zooplankton, phytoplankton)	Pelagic fishes	51.20 ± 9.40 (33.00–64.00)	2.16 ± 1.13 (0.45–4.19)
<i>Pseudobagrus fulvidraco</i>	Yellow catfish	2	Omnivory	Bottom fishes	15.10 ± 3.40 (12.70–17.50)	32.0 ± 2.83 (30.0–34.0)
<i>Parasilurus asotus</i>	Sheatfish	13	Sarcophagy	Bottom fishes	37.00 ± 9.40 (25.00–50.00)	0.41 ± 0.36 (0.10–1.38)
<i>Protosalanx hyalocranius</i>	Clearhead icefish	1	Sarcophagy	Pelagic fishes	14.50	0.025

<sup>a</sup> *n* means the number of samples.

performed to compare the relationships between metal(loid)s concentration in fish and sediment samples. One-way ANOVA followed by the Dunn's test was performed to determine whether differences in metal(loid) concentrations in different fish species were significant. The statistical significance level was acceptable at  $p < 0.05$ . All sample values were expressed as  $\text{mg kg}^{-1}$  wet weight.

### 3 Results and discussion

#### 3.1 Levels of eight metal(loid)s in fish muscles

The levels of eight metal(loid)s (wet weight) in fish muscles are shown in Table 2. The essential elements (Cu and Zn) are found in all the samples, while the elements that are essential but toxic in excess amounts (Cr and Ni) and the detection rates of toxic elements (As, Cd, Pb and Hg) are 33.55%, 19.35%, 97.42%, 48.39%, 70.97% and 74.19%, respectively. The metal concentration of Zn is the highest ( $7.97 \pm 6.05 \text{ mg kg}^{-1}$ ), followed by Cu ( $0.27 \pm 0.13 \text{ mg kg}^{-1}$ ), Hg ( $0.079 \pm 0.14 \text{ mg kg}^{-1}$ ), As ( $0.052 \pm 0.054 \text{ mg kg}^{-1}$ ) and Pb ( $0.034 \pm 0.074 \text{ mg kg}^{-1}$ ). In contrast, other heavy metals (Cr, Ni and Cd) are of minor importance because of low concentrations ( $<0.02 \text{ mg kg}^{-1}$ ). None of the fish samples have Cr, Cu and Cd levels above the guideline levels and legal limits shown in Table 2.<sup>30,34–37</sup> The concentrations of As, Pb and Hg are in the range of ND–0.39  $\text{mg kg}^{-1}$ , ND–0.70  $\text{mg kg}^{-1}$  and ND–0.82  $\text{mg kg}^{-1}$ , respectively. Total As levels in some of the fish muscle samples exceed the inorganic arsenic

criterion quality standard in China<sup>34</sup> of 0.10  $\text{mg kg}^{-1}$  and overall, 10.97% (17 fish muscle samples) were above the criterion. According to the hypothesis, inorganic arsenic levels could be estimated using a value of 10% of total As. The estimated concentration of inorganic arsenic for the 17 fish muscle samples are all below the criterion of 0.10  $\text{mg kg}^{-1}$ . Nonetheless, the Pb level of one fish muscle sample exceeds the maximum levels set by the quality standard in China and DHO action levels (of 0.50  $\text{mg kg}^{-1}$  wet weight); and 1.94% (3 samples) and 2.58% (4 samples) of the samples are above the Commission Regulation (EC) (0.30  $\text{mg kg}^{-1}$ ) FAO guidelines (0.20  $\text{mg kg}^{-1}$ ), respectively.<sup>30,36</sup> Total Hg levels in 2.58% (4 samples) of the samples were greater than 0.50  $\text{mg kg}^{-1}$ , but only one sample was above the methylmercury criterion set by the quality standard in China because of the maximum level for predatory fish (1.0  $\text{mg kg}^{-1}$ ). Based on the assumption that 75% of total mercury in fish was methylmercury,<sup>38</sup> the estimated methylmercury value of the Hg-excess sample still exceeded the criterion of 0.50  $\text{mg kg}^{-1}$ .

The results presented here are comparable to the studies conducted elsewhere, which also reported Cu and Zn to be the most abundant compounds among detected chemical elements (Table 3).<sup>17,18,39–41</sup> The concentrations of Cr, As, Cd and Pb in most of the samples are either less than previous studies<sup>10,11,22,39,40,42</sup> or undetected. Hg concentrations of fish in this study are higher than those from the Yellow River Estuary,<sup>10</sup>



Table 2 Descriptive statistics of eight metal(loid)s ( $\text{mg kg}^{-1}$  wet weight) in fish muscle samples ( $n = 155$ )<sup>a</sup>

Heavy metals	Min.	Max.	Mean	Standard deviation	Median	Detection rate (%)	Maximum levels (MLs) for heavy metals in fish				
							Quality standard in China	Commission regulation (EC)	FAO	DHO	Over-limit ratio <sup>b</sup> (%)
Cr	ND <sup>c</sup>	0.49	0.018	0.048	0.019	33.55	2.00 <sup>d</sup>				0
Ni	ND	0.78	0.019	0.069	0.039	19.35					
Cu	0.067	0.97	0.27	0.13	0.25	100.00	50 <sup>e</sup>		30		0
Zn	2.49	51.38	7.97	6.05	5.94	100.00			30		
As	ND	0.39	0.052	0.054	0.039	97.42	0.10 <sup>d</sup>			3.00	10.97
Cd	ND	0.009	0.0012	0.0013	0.0010	48.39	0.10 <sup>d</sup>	0.05	0.05	1.00	0
Pb	ND	0.70	0.034	0.074	0.024	70.97	0.50 <sup>d</sup>	0.30	0.20	0.50	0.65
Hg	ND	0.82	0.079	0.14	0.057	74.19	0.50/1.00 <sup>d</sup>	0.50	0.50/1.00 <sup>f</sup>	0.50/1.00 <sup>f</sup>	0.65

<sup>a</sup>  $n$  means the number of muscle samples. <sup>b</sup> According to the quality standard in China. <sup>c</sup> ND: Not detected, means the value is lower than the limit of detection. Concentrations less than the LOD were set to  $\frac{1}{2}$  LOD for statistical analysis. <sup>d</sup> GB 2762-2017 National Food Safety Standard Maximum Levels of Pollutants in Foods (in Chinese).<sup>34</sup> The maximum level for predatory fish is  $1.00 \text{ mg kg}^{-1}$ , but for all other fish it is  $0.50 \text{ mg kg}^{-1}$ . <sup>e</sup> NY 5073-2006 Limited Quantity of Poisonous and Harmful Contents of the National Pollution-Free Aquatic Products (in Chinese).<sup>35</sup> <sup>f</sup> The maximum level for predatory fish including swordfish is  $1.00 \text{ mg kg}^{-1}$  but for all other fish and processed fish, it is  $0.50 \text{ mg kg}^{-1}$ .

South China Sea,<sup>12</sup> East China Sea<sup>43</sup> and Yangtze River,<sup>44</sup> confirming heavy metal pollution in the waters of Northeast China.

### 3.2 Effect of environment on the metal(loid) levels in fish

The target metal(loid) concentrations in wild fish from natural waters and cultured fish from aquaculture ponds are compared in Fig. 2. The Mann-Whitney test of the Ni, Cu and Zn concentrations in fish showed little differences between the means of both separate fish samples ( $\alpha = 0.05$ ). The mean concentrations of Cr in the cultured fish were higher than those in the wild fish and differed significantly among the water types ( $P = 0.003$ ). The accumulation of As was relatively high ( $P = 0.017$ ) in wild fish from rivers and lakes as compared to those from ponds. Significantly higher Cd and Hg levels ( $P < 0.0001$ ) were found in wild fish as compared to those in the cultured fish. The concentration of Pb was much higher ( $P < 0.0001$ ) in cultured fish than that in wild fish. The detection frequencies of these toxic elements (As, Cd and Hg) in wild fish (97.78%, 61.11% and 92.22%, respectively) were higher than those in cultured fish (96.92%, 32.31% and 49.23, respectively). The detection frequency of Pb in wild fish was lower than that in cultured fish (52.22% and 96.92%, respectively). Among the essential elements (Cr, Ni, Cu and Zn), Cu and Zn were detected in all the samples. However, Cr and Ni detection rates (27.78% and 15.56%) in wild fish were lower than in cultured fish (47.69% and 24.62%).

The relationship between metal(loid) concentrations in fish and in their living environment is shown in Fig. 3. The target metal(loid) concentrations in wild fish are not correlated to those in the sediment of their living environment (except for Cd). Cd concentrations in wild fish samples were positively correlated to Cd levels in the sediments of nature water ( $R^2 = 0.081$ ,  $P = 0.012$ ). Positive correlations were observed between Cr, Ni, As and Cd concentrations in the cultured fish and in pond sediment ( $R^2 = 0.099$ ,  $R^2 = 0.070$ ,  $R^2 = 0.038$  and  $R^2 = 0.145$ , respectively,  $P < 0.05$  for all tests). However, correlations between Cu, Zn, Pt and Hg

concentrations in cultured fish and those in pond sediment were relatively weak ( $R^2 = 0.022$ ,  $R^2 = 0.044$ ,  $R^2 = 0.025$  and  $R^2 = 0.004$ , in order,  $P > 0.05$  for all tests).

Metal(loid)s in the fish samples analyzed in this study are closely related to their living environment.<sup>10</sup> In the closed static water environment, cultured fish can absorb the essential elements Cr, Ni, Cu and Zn from aquaculture feed, which results in higher levels and detection rates. This may explain why Cr and Ni levels in cultured fish tend to correlate with those in the pond sediment (mainly composed of feed bait, excrement, plankton particles and other suspended matter). Ponds investigated in this study usually use groundwater or clean spring water as the aquaculture water source. Thus, the significant relationship between the As and Cd concentrations in cultured fish and in pond sediment may indicate that As and Cd in cultured fish originate from the aquaculture feed. The potential routes of metal(loid)s entering a fish include food and non-food particle intake, surface absorption by fish tissues (gill, skin and body mucus) and oral consumption of water.<sup>8,45</sup>

Relative to wild fish, the higher detection rates and levels of Pb in cultured fish are the results of compounded effects from the aquaculture feed and water environment. Metal(loid)s in fish can reflect the water environmental pollution to a certain extent.<sup>10,14</sup> The detection frequency of As in cultured fish was similar to that in fish from natural water, while the As concentration was significantly lower than in wild fish, which indicated As-pollution in natural waters. The Songhua River in Northeastern China is one of the representative Hg-polluted rivers in China.<sup>23</sup> About 149.8 t of Hg from the Jilin chemical plant had been directly discharged into the Songhua River in 1958–1971. Studies on Hg pollution of fish and sediment in the Songhua River and health concerns of local residents have been uninterrupted over the past 40 years.<sup>25,46,47</sup> Recent research showed that the T-Hg pollution level of the Second Songhua River was moderate to severe with high ecological risk, and the pollution levels in the main stream of the Songhua River were



Table 3 Eight metal(loid) concentrations (mg kg<sup>-1</sup> wet weight) in fish in other regions of China and other countries

Location	Number of fish species	Cr	Ni	Cu	Zn	As	Cd	Pb	Hg	References
Northeast of China <sup>a</sup>	16	ND <sup>c</sup> -0.49	ND-0.78	0.067-0.97	2.49-51.38	ND-0.39	ND-0.009	ND-0.70	ND-0.82	Present study
Bangshi River (Bangladesh) <sup>b</sup>		0.47-2.07		8.33-43.18	4.283-418	1.97-6.24	0.09-0.87	1.76-10.27		39
Yangze River (China) <sup>a</sup>	13	0.10-0.24		0.77-1.22	2.8-7.55	NA	0.046-0.12	0.21-0.81		18
Pearl River (China) <sup>a</sup>	7	0.11-4.27		0.15-7.55	8.78-30.3	NA	0.01-0.13	0.09-30.70		40
Nam Co Lake, Yamdro Lake and Lhasa River (China) <sup>a</sup>	1		0.094-0.22	0.33-2.0	2.5-6.9	0.067-0.27	0.013-0.029	0.024-0.079		41
Taihu Lake (China) <sup>a</sup>	11	0.12-0.50		0.39-1.47			0.02-0.11	0.35-1.20	0.01-0.22	17
Yellow River Estuary (China) <sup>a</sup>	11					0.01-0.87				10
Pearl River Delta (China/Hong Kong) <sup>a</sup>	11	0.20-0.65		0.79-2.26	15.20-29.50	0.03-1.53	0.02-0.06	0.03-8.62		22
Hainan coastal area, South China sea (China) <sup>a</sup>	4	1.38-4.36	0.48-0.78	0.07-0.94	1.45-8.07	0.48-6.99	ND-0.012	0.02-0.05		12
Western continental shelf of South China Sea	6	0.86-2.89		0.39-2.65	11.75-15.59		0.027-0.15	0.54-2.17		11
Northeast Mediterranean Sea (Europe) <sup>b</sup>	6	1.24-2.42		2.34-4.41	16.5-37.4		0.37-0.79	2.98-6.12		42

<sup>a</sup> Wet weight. <sup>b</sup> Dry weight. <sup>c</sup> ND means not detected.

mild with low ecological risk.<sup>26</sup> As mentioned above, the Hg levels in wild fish in this study are significantly higher than in the cultured fish ( $P < 0.0001$ ), confirming the high level of Hg pollution and the potential ecological risk of natural waters.

### 3.3 Effects of different species on the metal(loid) levels in fish

The mean concentrations of eight metal(loid)s for each fish species from Northeast China are displayed in Table 4. Because of the different fish species and environmental factors, the data for wild and cultured fishes were analyzed separately. For the wild fish, because the sample sizes of *Ctenopharyngodon idellus* ( $n = 2$ ), *Erythroculter mongolicus* ( $n = 1$ ), *Saurogobio dabryi* ( $n = 1$ ), *Hemibarbus labeo* ( $n = 2$ ), *Leuciscus waleckii* ( $n = 3$ ), *Pseudobagrus fulvidraco* ( $n = 2$ ) and *Hemibarbus maculatus* ( $n = 1$ ) were small, their data were not included in the further analysis for *post hoc* testing. The levels of Pb, Zn, As and Hg significantly differed among the wild species ( $P = 0.0067$ ,  $P < 0.0001$ ,  $P = 0.001$  and  $P = 0.0005$ , in order). Significantly higher Cu levels were found in wild *Carassius auratus* as compared to *Esox reicherti* and *Parasilurus asotus* ( $P < 0.05$  for *post hoc* tests). The level of Zn in wild *Carassius auratus* was significantly higher than in wild *Esox reicherti*, *Hypophthalmichthys molitrix*, *Parasilurus asotus* and *Aristichthys nobilis*, and was high in *Cyprinus carpio* relative to *Hypophthalmichthys molitrix* ( $P < 0.05$ ). The toxic elements As and Hg in fish need to be paid more attention. Significantly higher As levels were found in wild *Carassius auratus*, *Hypophthalmichthys molitrix* and *Cyprinus carpio* compared to those in *Parasilurus asotus* ( $P < 0.05$ ). Mercury was found to be high in *Esox reicherti* as compared to that in *Carassius auratus*, *Aristichthys nobilis* and *Cyprinus carpio* ( $P = 0.007$ ,  $P = 0.016$  and  $P = 0.006$ , respectively). For the cultured fish, since the sample sizes of *Hypophthalmichthys molitrix* ( $n = 1$ ) and *Aristichthys nobilis* ( $n = 3$ ) were small, they were excluded from the *post hoc* test. Only Cu and Zn significantly differed among the cultured species ( $P = 0.0005$  and  $P = 0.030$ , respectively). Much higher Cu levels were found in cultured *Carassius auratus* and *Ctenopharyngodon idellus* as compared to those in *Cyprinus carpio* ( $P = 0.0016$  and  $P = 0.0046$ , respectively). Zinc levels were obviously higher in cultured *Ctenopharyngodon idellus* as compared to that in *Cyprinus carpio* ( $P = 0.039$ ).

Fish species are one of the most important drivers for metal(loid) accumulation, probably because of different feeding habits and foraging habitats of the studied fish. Dietary habits would affect the concentrations of the toxic elements As and Hg in fish from Northeast China. Relatively high As levels are observed in wild filter-feeding (*Hypophthalmichthys molitrix* and *Aristichthys nobilis*) and omnivory (*Carassius auratus* and *Cyprinus carpio*) fish, while the wild sarcophagi fish (*Parasilurus asotus* and *Erythroculter ilishaeformis*) had lower As concentration.




Fig. 2 The target metal(loid) concentrations in wild fish from natural waters and cultured fish from aquaculture ponds.

Differences in As levels among wild fish species could be due to trophic transfer in the food web/sources.<sup>48</sup> The *Hypophthalmichthys molitrix* and *Aristichthys nobilis* were strictly planktivorous and were expected to acquire As from small plankton and zooplankton. *Carassius auratus* and *Cyprinus carpio* do not feed exclusively on the small plankton and zooplankton but also feed on benthic animals. *Parasilurus asotus* and *Erythroculter ilishaeformis* are carnivorous and feed on small fishes. Chen and Folt<sup>20</sup> reported that As could be elevated in lower trophic levels, and lower trophic feeders in the fish feeding strategy experiment would have higher metal burdens (planktivores > omnivores and piscivores). Notably, higher As is found in the muscle of the omnivorous fish, *Carassius auratus*, which could be due to both the foraging habitat and feeding habits. Similar to previous studies, the current study has found that the Hg concentration in wild sarcophagi fish is higher than that in omnivores and filter-feeding. Many studies also reported evidence of the biomagnification and bioaccumulation of mercury with trophic levels.<sup>13,15,49</sup> Particularly, total mercury burdens in muscle tissue are a result of biomagnification through the food web rather than bioaccumulation.<sup>15</sup>

### 3.4 Human health risk assessment of metal(loid)s via fish consumption

Oral reference dose (RfDs) values for Ni, Cu, Zn and Cd are 20, 40, 300 and 1  $\mu\text{g}$  per kg per day, respectively.<sup>32</sup> The RfD of Cr(vi) (3.0  $\mu\text{g}$  per kg per day), inorganic arsenic (0.3  $\mu\text{g}$  per kg per day) and methylmercury (0.1  $\mu\text{g}$  per kg per day) are taken to represent that of Cr, As, and Hg, respectively.<sup>32</sup> Following the European Food Safety Authority,<sup>50</sup> the RfD of Pb is set to 1.5  $\mu\text{g}$  per kg per day. The target hazard quotients (THQs) are calculated using the average, maximum concentrations of eight metal(loid)s (Table 5). Regardless of the exposure levels utilized in the computational formula, all THQs of target elements were below 1.0 through the consumption of either fish alone, with the exception of THQ-Hg at high exposure levels. Hg shows maximum values for individual target hazard quotient (0.10), followed by As (0.069) and Zn (0.0089); whereas, THQs of Pb, Cu, Cr, Cd and Ni were small (below 0.01). The total hazard index (HI) is estimated by summing each THQ value of the target metal(loid)s. The HI at average exposure level was 0.20, indicating a lower health risk associated with the exposure of eight metal(loid)s. However, the HI at high exposure level is greater than 1.0 (1.94), suggesting a high risk from exposure to the maximum concentration of the target metal(loid)s in fish. In a sense, the accumulation of eight metal(loid)s would increase human health risks.

There are significant differences in the contribution rates of the eight metal(loid)s for HI at average exposure levels. Among the heavy metal elements, the contribution rate of Cu is the highest (27.78%), followed by Zn (15.51%), As (13.43%), Cd (13.01%) and Hg (9.62%), while that of Pb, Cr and Ni are quite low (all below 5.00%). While Cu and Zn are essential elements for the human body, their excessive accumulation would be detrimental for human health. The EDIs at the average exposure levels of Cu and Zn are much lower than the RfD values, and the THQs are below 1.0, implying that Cu and Zn are not high-risk elements. Although THQs of As, Cd and Hg are below 1.0, these toxic elements could still cause serious health issues to the general public.




Fig. 3 The relationship between metal(loid) concentrations in fish and in their living environment.



Table 4 Eight meta(loid) concentrations (mg kg<sup>-1</sup> wet weight) in muscle samples of 16 fish species<sup>a</sup>

Scientific name	Cr	Ni	Cu	Zn	As	Cd	Pb	Hg
<i>Carassius auratus</i>	Cultured 0.027 ± 0.043 (ND-0.15)	0.028 ± 0.050 (ND-0.20)	0.34 ± 0.20 (0.15-0.97)	7.93 ± 3.65 (3.83-18.44)	0.035 ± 0.030 (0.003-0.14)	0.0008 ± 0.0006 (ND-0.003)	0.031 ± 0.013 (0.015-0.067)	0.003 ± 0.005 (ND-0.018)
<i>Cyprinus carpio</i>	Wild 0.040 ± 0.12 (ND-0.49)	0.018 ± 0.030 (ND-0.12)	0.37 ± 0.15 (0.12-0.55)	16.59 ± 11.37 (4.94-51.38)	0.075 ± 0.069 (0.009-0.28)	0.002 ± 0.002 (ND-0.008)	0.020 ± 0.021 (ND-0.071)	0.082-0.085 (0.001-0.26)
	Cultured 0.026 ± 0.041 (ND-0.13)	0.008 ± 0.009 (ND-0.035)	0.20 ± 0.067 (0.12-0.35)	6.12 ± 2.71 (3.20-13.74)	0.037 ± 0.034 (ND-0.13)	0.0008 ± 0.0006 (ND-0.003)	0.032 ± 0.02 (0.014-0.085)	0.041 ± 0.18 (ND-0.82)
<i>Ctenopharyngodon idellus</i>	Wild 0.0072 ± 0.0098 (ND-0.031)	0.006 ± 0.002 (ND-0.011)	0.29 ± 0.12 (0.12-0.46)	12.40 ± 8.44 (4.30-29.74)	0.049 ± 0.041 (0.01-0.27)	0.002 ± 0.001 (ND-0.004)	0.044 ± 0.11 (ND-0.70)	0.095-0.053 (ND-0.18)
	Cultured 0.022 ± 0.037 (ND-0.16)	0.061 ± 0.19 (ND-0.78)	0.31 ± 0.095 (0.20-0.50)	8.77 ± 4.53 (4.53-20.01)	0.042 ± 0.030 (0.003-0.14)	0.0006 ± 0.0004 (ND-0.002)	0.082 ± 0.11 (ND-0.43)	0.005 ± 0.006 (ND-0.019)
<i>Esox reicherti</i>	Wild ND	ND	0.25 ± 0.21 (0.11-0.40)	9.26 ± 7.26 (4.13-14.39)	0.074 ± 0.045 (0.042-0.11)	ND	0.036 ± 0.031 (0.014-0.058)	0.010 ± 0.013 (ND-0.019)
	Wild 0.005 ± 0.005 (ND-0.014)	ND	0.18 ± 0.11 (0.074-0.39)	4.50 ± 0.77 (4.61-5.73)	0.085 ± 0.086 (0.027-0.26)	0.0013 ± 0.001 (ND-0.003)	0.004 ± 0.003 (ND-0.01)	0.37 ± 0.099 (0.27-0.56)
<i>Pseudobagrus fulvidraco</i>	Wild 0.013 ± 0.015 (ND-0.023)	ND	0.25 ± 0.12 (0.16-0.33)	7.47 ± 3.43 (5.04-9.89)	0.013 ± 0.006 (0.008-0.017)	0.002 ± 0.001 (0.001-0.003)	0.022 ± 0.002 (0.02-0.023)	0.080 ± 0.004 (0.077-0.082)
	Cultured 0.0025	0.005	0.46	9.78	0.02	0.0005	0.019	ND
<i>Hypophthalmichthys molitrix</i>	Wild 0.005 ± 0.006 (ND-0.019)	0.011 ± 0.013 (ND-0.046)	0.28 ± 0.10 (0.14-0.48)	4.28 ± 2.22 (2.78-10.33)	0.072 ± 0.030 (0.029-0.13)	0.0006 ± 0.0002 (ND-0.001)	0.015 ± 0.028 (ND-0.079)	0.11 ± 0.10 (0.005-0.27)
	Wild 0.016 ± 0.042 (ND-0.15)	0.015 ± 0.027 (ND-0.10)	0.20 ± 0.079 (0.086-0.40)	5.28 ± 1.93 (2.77-10.55)	0.015 ± 0.018 (ND-0.052)	0.001 ± 0.001 (ND-0.004)	0.016 ± 0.012 (ND-0.042)	0.12 ± 0.11 (ND-0.32)
<i>Parasilurus asotus</i>	Wild 0.011 ± 0.014 (ND-0.04)	ND	0.23 ± 0.072 (0.15-0.35)	5.17 ± 1.01 (3.27-6.08)	0.034 ± 0.013 (0.018-0.050)	0.002 ± 0.002 (ND-0.005)	0.030 ± 0.044 (ND-0.12)	0.36 ± 0.25 (0.057-0.63)
<i>Erythroculter tishataformis</i>	Wild ND	ND	0.29 ± 0.059 (0.23-0.34)	5.56 ± 0.73 (4.72-6.05)	0.072 ± 0.016 (0.062-0.09)	0.0008 ± 0.0003 (ND-0.001)	0.004 ± 0.003 (ND-0.008)	0.057 ± 0.030 (0.023-0.075)
<i>Leuciscus waleckii</i>	Wild ND	ND	0.24 ± 0.053 (0.18-0.28)	5.38 ± 2.48 (3.28-7.64)	0.053 ± 0.030 (0.034-0.087)	0.0007 ± 0.0003 (ND-0.001)	0.037 ± 0.18 (0.019-0.054)	ND
<i>Aristichthys nobilis</i>	Cultured 0.023 ± 0.036 (ND-0.065)	0.016 ± 0.019 (ND-0.033)	0.22 ± 0.068 (0.13-0.33)	4.55 ± 1.325 (2.49-6.76)	0.060 ± 0.048 (ND-0.16)	0.001 ± 0.001 (ND-0.004)	0.006 ± 0.007 (ND-0.024)	0.068 ± 0.049 (ND-0.17)
<i>Hemibarbus labeo</i>	Wild ND	ND	0.38 ± 0.26 (0.20-0.56)	5.65 ± 0.96 (4.97-6.32)	0.042 ± 0.016 (0.03-0.053)	0.002 ± 0.0007 (0.001-0.002)	0.011 ± 0.012 (ND-0.02)	0.090 ± 0.053 (0.052-0.13)
	Wild ND	ND	0.26	24.01	0.39	0.007	ND	0.008
<i>Hypomesus olidus</i>	Wild ND	ND	0.067	9.52	0.059	0.004	ND	ND
<i>Protosalanx hyalocranius</i>	Wild ND	ND	0.24	5.51	0.066	0.001	ND	0.12
<i>Hemibarbus maculatus</i>	Wild 0.005	ND	0.18	5.57	0.039	ND	0.01	0.13
<i>Erythroculter mongolicus</i>	Wild ND	ND	0.25	14.56	0.050	0.009	0.007	0.065
<i>Saurogobio dabryi</i>	Wild ND	ND						

<sup>a</sup> ND means not detected.



Table 5 Carcinogenic and non-carcinogenic risk analysis of elements in fish consumption in Northeast China

Elements	RfDs, µg per kg per day	Average exposure level		High exposure level		CRI (10 <sup>-6</sup> )
		EDI, µg per kg per day	THQ	EDI, µg per kg per day	THQ	
Cr	3 <sup>a,b</sup>	0.0071	0.0024	0.19	0.064	3.57 (0.49–95.99)
Ni	20 <sup>a</sup>	0.0076	0.0004	0.31	0.016	
Cu	40 <sup>a</sup>	0.11	0.0027	0.38	0.0096	
Zn	300 <sup>a</sup>	3.15	0.011	20.30	0.068	
As	0.3 <sup>a,c</sup>	0.021	0.069	0.15	0.51	31.04 (0.89–231.08)
Cd	1 <sup>a</sup>	0.0005	0.0005	0.0036	0.0036	0.189 (0.075–1.35)
Pb	1.5 <sup>d</sup>	0.013	0.0089	0.27	0.18	
Hg	0.3 <sup>a,e</sup>	0.031	0.10	0.33	1.08	
HI <sup>f</sup> /CRI <sub>t</sub>			0.20		1.94	34.79 (1.46–232.62)

<sup>a</sup> RfDs obtain from EPA.<sup>32</sup> <sup>b</sup> Standard of hexavalent chromium. <sup>c</sup> Standard of inorganic arsenic. <sup>d</sup> Data from European Food Safety Authority.<sup>50</sup> <sup>e</sup> Standard of mercuric chloride (and other mercury salts). <sup>f</sup> HI (average exposure level) =  $\sum$ THQs; HI (high exposure level) =  $\sum$ THQs; CRI<sub>t</sub> =  $\sum$ CRI<sub>t</sub>.

For the single element, the average carcinogenic risk indexes of Cr, As, and Cd are all lower than 10<sup>-4</sup> (Table 5), indicating that the carcinogenic risk of the residual metal(loid)s in the studied fish are acceptable. The total carcinogenic risk index ranges from a minimum of 1.46 × 10<sup>-6</sup> to a maximum of 232.62 × 10<sup>-6</sup> (an average value of 34.79 × 10<sup>-6</sup>) in fish consumption in Northeast China. The results show the potential carcinogenic risk of fish consumption in Northeast China. The levels of carcinogenic elements (Cr, As, and Cd) in fish in this area should be monitored.

## 4 Conclusions

This study investigated the concentrations of eight metal(loid)s in fish samples from Northeast China and evaluated the health risks from fish consumption. Relative to previous studies conducted in other regions, Cu, Cr, As, Cd and Pb (Ni, Zn and Hg) concentrations are higher (lower) in Northeast China. Apart from individual samples, most of the fish samples have metal(loid) levels below the guideline levels and legal limits. Some sort of relationships exist between the metal(loid)s in fish and their living environment. For instance, Cr, Ni, As and Cd levels in cultured fish appeared to correlate with those in pond sediment. Higher detection frequencies and concentrations of As and Hg in wild fish indicated a certain As-pollution and Hg-pollution in natural waters. The mean concentrations of eight metal(loid)s for each fish species are different, which might be due to different feeding habits and foraging habitats. In particular, Pb is the key risk factor in cultured fish, while As and Hg are the main risk factors in wild fish. Our results also indicate that consuming fish from Northeast China poses carcinogenic risk, and wild fish consumption would probably lead to mercury poisoning.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was financially supported by Open Project of State Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology (No. ES201604), Natural Science Foundation of Heilongjiang Province (No. QC2017033) and the Special Fund for Agro-scientific Research in the Public Interest (No. 201503108).

## References

- 1 M. Hutton and C. Symon, *Sci. Total Environ.*, 1986, **57**, 129–150.
- 2 H. Bing, Y. Wu, J. Zhou, R. Li and J. Wang, *Chemosphere*, 2016, **148**, 211–219.
- 3 J. R. Martín, C. De Arana, J. Ramos-Miras, C. Gil and R. Boluda, *Environ. Pollut.*, 2015, **196**, 156–163.
- 4 S. Cheng, *Environ. Sci. Pollut. Res.*, 2003, **10**, 192–198.
- 5 Y. Guo and S. Yang, *Sci. Total Environ.*, 2016, **543**, 105–115.
- 6 J. Ren, Z. Shang, L. Tao and X. Wang, *Pol. J. Environ. Stud.*, 2015, **24**, 1041–1048.
- 7 W. Wildi, J. Dominik, J. L. Loizeau, R. L. Thomas, P. Y. Favarger, L. Haller, A. Perroud and C. Peytremann, *Lakes Reservoirs: Res. Manage.*, 2004, **9**, 75–87.
- 8 H. Nussey, *Water SA*, 2000, **26**, 269–284.
- 9 A. G. Heath, *Water Pollution and Fish Physiology*, CRC Press, Boca Raton, Florida, U.S.A, 1987.
- 10 Y. Liu, G. Liu, Z. Yuan, H. Liu and P. K. Lam, *Sci. Total Environ.*, 2018, **613**, 462–471.
- 11 Y. G. Gu, Q. Lin, H.-H. Huang, L. g. Wang, J. J. Ning and F. Y. Du, *Mar. Pollut. Bull.*, 2017, **114**, 1125–1129.
- 12 J. L. Liu, X.-R. Xu, Z. H. Ding, J. X. Peng, M. H. Jin, Y. S. Wang, Y. G. Hong and W. Z. Yue, *Ecotoxicology*, 2015, **24**, 1583–1592.
- 13 I. Jæger, H. Hop and G. W. Gabrielsen, *Sci. Total Environ.*, 2009, **407**, 4744–4751.
- 14 N. L. Gale, C. D. Adams, B. G. Wixson, K. A. Loftin and H. Yue-Wern, *Environ. Sci. Technol.*, 2002, **36**, 4262–4268.



- 15 L. Atwell, K. A. Hobson and H. E. Welch, *Can. J. Fish. Aquat. Sci.*, 1998, **55**, 1114–1121.
- 16 V. Sirot, T. Guérin, J. L. Volatier and J.-C. Leblanc, *Sci. Total Environ.*, 2009, **407**, 1875–1885.
- 17 S. Rajeshkumar, Y. Liu, X. Zhang, B. Ravikumar, G. Bai and X. Li, *Chemosphere*, 2018, **191**, 626–638.
- 18 Y.-J. Yi and S.-H. Zhang, *Environ. Sci. Pollut. Res.*, 2012, **19**, 3989–3996.
- 19 A. Mountouris, E. Voutsas and D. Tassios, *Mar. Pollut. Bull.*, 2002, **44**, 1136–1141.
- 20 C. Y. Chen and C. L. Folt, *Environ. Sci. Technol.*, 2000, **34**, 3878–3884.
- 21 A. Goutte, Y. Chereil, C. Churlaud, J.-P. Ponthus, G. Massé and P. Bustamante, *Sci. Total Environ.*, 2015, **538**, 743–749.
- 22 H. Leung, A. Leung, H. Wang, K. Ma, Y. Liang, K. Ho, K. Cheung, F. Tohidi and K. Yung, *Mar. Pollut. Bull.*, 2014, **78**, 235–245.
- 23 G. B. Jiang, J. B. Shi and X. B. Feng, *Environ. Sci. Technol.*, 2006, **40**, 3672–3678.
- 24 L. Bingli, S. Huang, Q. Min, L. Tianyun and W. Zijian, *J. Environ. Sci.*, 2008, **20**, 769–777.
- 25 N. Li, Y. Tian, J. Zhang, W. Zuo, W. Zhan and J. Zhang, *Environ. Sci. Pollut. Res.*, 2017, **24**, 3214–3225.
- 26 D. Qin, H. Jiang, X. Huang, L. Gao, Z. Chen, Z. Mou and L. I. Gang, *Acta Sci. Circumstantiae*, 2016, **36**, 1910–1916.
- 27 K. Pan and W.-X. Wang, *Sci. Total Environ.*, 2012, **421**, 3–16.
- 28 D. Qin, H. Jiang, S. Bai, S. Tang and Z. Mou, *Food Control*, 2015, **50**, 1–8.
- 29 H. Jiang, D. Qin, Z. Chen, S. Tang, S. Bai and Z. Mou, *Bull. Environ. Contam. Toxicol.*, 2016, **97**, 536–542.
- 30 FAO, in *Heavy Metals Regulations Legal Notice No 66/2003*, FAO, Rome, 2003.
- 31 F. Zhai, H. Wang, S. Du, Y. He, Z. Wang, K. Ge and B. M. Popkin, *Nutr. Rev.*, 2009, **67**, S56–S61.
- 32 EPA, in *Regional Screening Level (RSL) Resident Fish Table*, United States Environmental Protection Agency, Washington DC, 2018.
- 33 EPA, *Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories, Risk Assessment and Fish Consumption Limits*, Washington DC, 2000.
- 34 NHFPC, *National standards for food safety-contaminants limit in food*, 2017, GB 2762-2017.
- 35 SAC, *Non-environmental Pollution Food-limit of Poisonous and Harmful Substances in Aquatic Products*, NY, 2006, pp. 5073–2006.
- 36 EC, in *Official Journal of the European Union*, No. 1881/2006, 2015.
- 37 DOH, in *Foodstuffs, cosmetics and disinfectants act, 1972 (Act no. 54 of 1972)*, Government Gazette, 2004.
- 38 R. Nakagawa, Y. Yumita and M. Hiromoto, *Chemosphere*, 1997, **35**, 2909–2913.
- 39 M. S. Rahman, A. H. Molla, N. Saha and A. Rahman, *Food Chem.*, 2012, **134**, 1847–1854.
- 40 C. Ip, X. Li, G. Zhang, C. Wong and W. Zhang, *Environ. Pollut.*, 2005, **138**, 494–504.
- 41 R. Yang, T. Yao, B. Xu, G. Jiang and X. Xin, *Environ. Int.*, 2007, **33**, 151–156.
- 42 M. Canli and G. Atli, *Environ. Pollut.*, 2003, **121**, 129–136.
- 43 K. A. Asante, T. Agusa, H. Mochizuki, K. Ramu, S. Inoue, T. Kubodera, S. Takahashi, A. Subramanian and S. Tanabe, *Environ. Pollut.*, 2008, **156**, 862–873.
- 44 Y. Yujun, W. Zhaoyin, K. Zhang, Y. Guoan and D. Xuehua, *Int. J. Sediment Res.*, 2008, **23**, 338–347.
- 45 R. D. Handy and F. B. Eddy, *J. Fish Biol.*, 1989, **34**, 865–874.
- 46 H. Zhou and M. H. Wong, *Water Res.*, 2000, **34**, 4234–4242.
- 47 H. Zhu, B. Yan, H. Cao and L. Wang, *Environ. Monit. Assess.*, 2012, **184**, 77–88.
- 48 J. Reinfelder, N. Fisher, S. Luoma, J. Nichols and W. X. Wang, *Sci. Total Environ.*, 1998, **219**, 117–135.
- 49 L. M. Campbell, R. J. Norstrom, K. A. Hobson, D. C. Muir, S. Backus and A. T. Fisk, *Sci. Total Environ.*, 2005, **351**, 247–263.
- 50 EFSA, *EFSA J.*, 2010, **8**, 1570.

