

from curd samples were evaluated and optimized for their extracellular digestive enzyme activity.

Materials and Methods

Inoculum preparation

In the present study, the four isolates namely *Bacillus subtilis* GS 1, *Bacillus cereus* GS 3, *Bacillus cereus* GS 199, and *Bacillus subtilis* GS 547 isolated from curd samples were tested for their extracellular hydrolytic enzyme activity. The isolates were grown overnight in nutrient broth at 37°C. After incubation, the cells were harvested by spinning at 7000 rpm/10 min, washed thrice and resuspended in sterile distilled water to obtain OD 1.2 at A₆₀₀ and used as inoculum.

Quantitative assay for amylase, protease, lipase and phytase from *Bacillus* spp.

Spectrophotometric assay of Amylase by DNS method: The amylase was produced by submerged fermentation in the medium containing lactose 40 g l⁻¹, yeast extract 20 g l⁻¹, KH₂PO₄ 0.05 g l⁻¹, MnCl₂ .4H₂O 0.015 g l⁻¹, MgSO₄.7H₂O 0.25 g l⁻¹, CaCl₂.2H₂O 0.05 g l⁻¹, FeSO₄.7H₂O 0.01 g l⁻¹ [25]. 100 mL of autoclaved production medium was inoculated with 2% inoculum and incubated at 37°C/24 h with continuous shaking (200 rpm). After incubation, the broth was centrifuged at 7000 rpm/15 min in a cooling centrifuge. Then the supernatant obtained was used as source for estimation of amylase activity by DNS (3, 5-dinitro salicylic acid) method by monitoring the amount of reducing sugar liberated from starch. To the one milliliter of crude enzyme, 0.5 mL of 1% soluble starch (prepared in 0.1 M phosphate buffer of pH 6) was added and incubated at 37°C for 20 min. Further, 1 mL of DNS reagent was added and boiled for 10 min to stop the reaction. Then, the final volume was made up to 5 mL by adding distilled water and absorbance was measured at 540 nm. One unit of amylase activity was defined as the amount of enzyme per mL of culture supernatant that released 1 µg of maltose per minute. In the assay, heat killed culture supernatant was served as control. The experiment was repeated in triplicates and results were expressed as mean ± standard deviation.

Estimation of *Bacillus* spp. lipase activity using copper soap method: The lipase enzyme was produced by submerged fermentation and quantified using olive oil as a substrate according to Veerapagu. In brief, after 72 h of submerged fermentation, the broth was centrifuged at 1000 rpm/20 min/4°C and supernatant was used as source of enzymes. To the 1 mL of culture supernatant, 2.5 mL of olive oil was added and incubated for 5 min at 37°C. Enzyme activity was arrested by adding 1 mL of 6N HCl and 5 mL benzene. Further, 4 mL of upper layer was carefully collected and to which 1 mL of cupric acetate pyridine was added. In the assay, reaction mixture with heat inactivated supernatant served as control. The lipase activity was determined by measuring the absorbance of free fatty acids dissolved in benzene at 715 nm and compared with oleic acid standard curve. One unit of lipase activity is defined as the amount of enzyme that liberated 1 µmol FFA in 1 min at 37°C.

Spectrophotometric assay of extracellular protease activity of *Bacillus* spp: The submerged fermentation for protease production was carried out according to Ikram-Ul-Haq and Mukthar (2006) and protease activity was assayed by Lowry's method. In brief, after 24 h of incubation at 37°C with continuous shaking at 200 rpm, the fermentation broth was centrifuged at 5000 rpm for 10 min and supernatant was assayed for protease activity. The reaction mixture containing 1 mL of casein (1% solution in 0.1 M phosphate buffer of

pH 6) and 1 mL of culture supernatant was incubated at 37°C/30 min. Control tubes contained supernatants boiled for 10 min. The enzyme activity was arrested by adding 5 mL of 5% trichloro acetic acid and further incubated at 37°C/10 min. Then the reaction mixture was centrifuged at 8000 rpm/10 min to remove insoluble particles and to which 1 mL of 1:1 Folin and Ciocalteu reagent and water was added. After 30 min of incubation, absorbance was measured at 700 nm. The amount of amino acids released was compared with tyrosine standard curve for the determination of protease activity.

Spectrophotometric assay of extracellular phytase activity of *Bacillus* spp: The extra cellular phytase was produced as described by Sreeramulu et al., with slight modification [26]. After fermentation for 72 h, the broth was centrifuged at 6000 rpm/30 min/4°C and supernatant was used for phytase assay. The assay mixture consisted of 1 mL of acetate buffer (pH 5.5) containing 6.82 mM sodium phytate, 0.2 mL of culture supernatant and 0.2 mL 100 mM MgSO₄ and incubated at 37°C for 30 min. Reaction mixture containing heat inactivated culture supernatant served as control. Immediately after stopping the reaction by adding 1 mL of 10% trichloroacetic acid, 1 mL of Taussky shorr color reagent solution prepared as described by Tungala was added and absorbance was measured at 660 nm. Then the absorption values were compared with the standard graph of potassium dihydrogen phosphate (0.1 to 0.5 mg/mL) to determine phytase activity. One unit of phytase activity was defined as the amount of enzyme required to liberate 1 mole of phosphate per min under assay conditions.

Total Protein Estimation

The total protein content was estimated by Lowry's method.

Optimisation of pH, temperature and substrate concentration for amylase, lipase, protease and phytase from *Bacillus* spp.

The optimum pH for all the four enzyme assay was determined by incubating the enzyme-substrate at various pH from 3 to 10 in the following buffers: 50 mM sodium acetate buffer (pH 3-5), 50 mM potassium phosphate buffer (pH 5-7), 50 mM Tris-HCl buffer (pH 7-9), and 50 mM glycine-NaOH buffer (pH 9-10). Enzyme activity in each buffer was measured using the standard assays as described previously. Further, to determine the optimum temperature for the amylase, lipase, protease and phytase, the reaction mixtures were incubated at various temperature ranged from 10 to 60°C and then absorbance were measured according to the standard assays. Furthermore, to determine the effect of substrate concentration on enzyme activities, culture supernatants were incubated with various volumes of specific substrate solutions viz., starch, olive oil, casein and sodium phytate for amylase, lipase, protease and phytase respectively.

Statistical Analyses

All the experiments were conducted in triplicates and results were expressed as mean ± standard deviation. The data were analyzed using two-tailed paired t-test and for all tests, the level of significance was set at p<0.05 (GraphPad Prism version 5.02 for Windows, GraphPad Software, San Diego, California, USA).

Results and Discussion

In the present study the extracellular amylase, lipase, protease and phytase activity of four *Bacillus* spp. were evaluated and optimized. The total protein content and specific activity of all the four extracellular digestive enzymes evaluated (Table 1).

Amylases break down α-linked sugar units in the ingested diet to

Isolates	Amylase		Lipase		Protease		Phytase	
	Protein mg/mL	Specific activity (IU)	Protein mg/mL	Specific activity (IU)	Protein mg/mL	Specific activity (IU)	Protein mg/mL	Specific activity (IU)
<i>B. subtilis</i> GS 1	0.101	1262.04	0.05	653.2	0.098	1602.229	0.046	1337.064
<i>B. cereus</i> GS 3	0.106	1082.77	0.194	142.57	0.093	1512.646	0.038	2138.007
<i>B. cereus</i> GS 199	0.105	994.13	0.045	785.11	0.091	1456.121	0.052	1056.65
<i>B. subtilis</i> GS 547	0.119	759.034	0.193	207.25	0.095	1394.164	0.0403	1966.63

Table 1: Total protein and specific activity.

glucose, maltose, and maltotriose which further undergo fermentation to produce various short chain fatty acids. In case of pancreatic insufficiency, microbial amylases are of best choice in improving the carbohydrate digestion. In the present study, the extracellular amylase activity of the four *Bacillus* spp. were studied and optimized for various physical parameters. As shown in the Figure 1, the amylolytic activity of the tested isolates were ranged between 87 and 128 U. Amongst, *Bacillus subtilis* GS 1 showed highest activity ($127.66 \pm 8.02U$, $p < 0.05$). Further, in optimization studies, the four *Bacillus* spp. tested showed optimum activity at 40°C ranging between 87 and 130 U/mL and subsequently decreased at 50°C with activity ranging from 50 to 73 U/mL. Whereas in another study, *Bacillus* amylase activity as 34 U/mL at 37°C/pH 7 [25], however, Abd-Elhalem et al. reported that the amylase activity of *B. amyloliquefaciens* was 72.5 U/mL with an increased activity at 50°C [27]. During optimization for pH, *B. subtilis* GS 1 and *B. cereus* GS 3 showed maximum activity at pH 6 while *B. subtilis* GS 547 and *B. cereus* GS 199 showed highest activity at pH 7. Further, the V_{max} value for *B. subtilis* GS 1 was 149 ± 4.05 U/mL at a substrate concentration 2 mg/mL/1 mL enzyme solution while for *B. cereus* GS 3 the V_{max} was 122 ± 3.87 U/mL at 1 mg/mL/ 1 mL enzyme solution. However, for *B. subtilis* GS 547 and *B. cereus* GS 199 V_{max} values were 108 ± 5.32 and 95 ± 3.69 U/mL, respectively at a substrate concentration 1.5 mg/mL/ 1 mL enzyme solution. In another study, V_{max} for *B. subtilis* was 100 U/mL which was in agreement with the present study [28] (Figures 1-4).

Lipase hydrolyzes fats into fatty acids and glycerols. In the present study, the crude lipase activity of the four tested *Bacillus* spp. ranged between 27 and 40 U/mL using olive oil as substrate. Amongst, *Bacillus subtilis* GS 547 showed highest activity 40 ± 2.13 U/ml ($p < 0.05$). However, *Bacillus subtilis* showed 6.921 U/mL in soyabean oil, while *Bacillus amyloliquefaciens* exhibited 6.506 U/mL using ground nut oil as substrate at optimum pH 7.5 and temperature 45°C [29]. On storage at temperature above 40°C, pancreatic lipases lose their activity, however, microbial lipases are more resistant to heat inactivation and therefore, thermal stability is one of their desirable characteristics and which was corroborated in the present study [30]. The maximum lipolytic activity was found to be 40°C for all the four isolates tested and indicating the thermal stability at a temperature range between 20 to 50°C and decreased beyond 60°C. Further, optimum pH was 6 for all the isolates except *B. subtilis* GS 2 (pH 7), however, lipase activity was stable at a pH range between 5 and 8 for all the four isolates tested. Further, the maximum rate of reaction V_{max} for *B. cereus* 3 and *B. subtilis* 547 were found to be 31 ± 1.42 and 41 ± 2.36 U/mL respectively. Whereas for *B. subtilis* GS 1 and *B. cereus* GS 199 V_{max} values were 32 ± 3.4 and 34 ± 2.1 U/mL respectively at a substrate concentration 0.5 mL/mL (v/v)/1 mL enzyme solution. However, Hasan et al. (2007) reported the V_{max} value of *B. subtilis* as 0.416 U/mL/min which was quite lower the maximum rate of reaction of the four tested *Bacillus* spp. (Figures 5-7).

The enzyme protease breakdown protein into smaller fragments as peptides and further to amino acids. In the present study, the protease activity of four tested *Bacillus* spp. were ranged between 120 and 137 U/mL. Amongst, *B. subtilis* GS 1 showed the highest enzyme activity




Figure 1: Extracellular digestive enzyme activity of *B. subtilis* GS 1, *B. cereus* GS 3, *B. cereus* GS 199 and *B. subtilis* GS 547.




Figure 2: Temperature optimization for amylase activity.




Figure 3: pH optimization for amylase activity by *Bacillus* spp.

157 ± 3.60 U/ml ($p < 0.05$) at 37°C. Further, the optimum temperature was found to be 40°C and moreover, enzyme activity was stable in the temperature range 20 to 50°C for all the isolates and was in agreement with the report of Kotlar et al. [31]. Further, pH 8 was optimum and the activity was stabilized between pH 6 and 9 for all the tested isolates. Furthermore, the maximum rate of reaction V_{max} were 160 ± 2.34 , 145 ± 1.45 and 130 ± 3.72 U/mL for *B. subtilis* GS 1, *B. cereus* GS 3 and GS 9 respectively at a substrate concentration 2 mg/mL/1 mL enzyme solution. However, Nadeem et al. reported V_{max} value of 61.58 U/ml for *B. licheniformis* [32]. The stability at wide range of temperature and pH suggested that the protease from tested *Bacillus* spp. may function




Figure 4: Effect of substrate concentration on amylase activity.




Figure 8: Temperature optimization for protease activity.




Figure 5: Temperature optimization for lipase activity.




Figure 9: pH optimization for protease activity.




Figure 6: pH optimization for lipase activity.




Figure 10: Effect of substrate concentration on protease activity.




Figure 7: Effect of substrate concentration on lipase activity.




Figure 11: Temperature optimization for phytase activity.

at different pH conditions *in vivo* and also can withstand varying temperature during processing (Figures 8-10).

Phytase catalyzes the hydrolysis of phytic acid into *myo*-inositol and inorganic phosphates. In the present study, all the four tested *Bacillus* spp. showed phytase activity using sodium phytate as

substrate. As shown in the Figure 1, the phytase activity of the four isolates ranged between 54 to 81 U/mL. Out of four isolates evaluated,

B. cereus GS 3 showed highest activity (81.07 ± 2.47 U/mL, $p < 0.05$). In addition, optimum temperature was 40°C , although activity remained fairly stable over temperature range of 30 to 50°C . Interestingly, the activity was found to stabilize at different pH range from 4 to 7 with an optimum pH 5 except for *B. subtilis* GS 547 (pH 6). Further, V_{\max} values for *B. subtilis* GS 1 and *B. cereus* GS 3 were 67.64 and 85.87 U/mL for 4.8 mg/mL/1 mL enzyme solution. Whereas for *B. cereus* GS 199 and *B. subtilis* GS 547, the V_{\max} values were 60.75 and 93.47 U/mL for 6 mg/mL/1 mL enzyme solution. However, in another study, the optimum pH and temperature for phytase from *B. subtilis* were 7 and 60°C respectively and K_m and V_{\max} values for sodium phytate were 0.42 mM and 4.35 $\mu\text{mol}/\text{min}$, respectively. The activity of phytase at optimum pH 4 indicated that the enzyme from the four tested isolates could be used *in vivo* to improve the bioavailability of phosphate from ingested food (Figures 11-13).

Conclusion

In the present study, *B. subtilis* GS 1, *B. cereus* GS 3, *B. cereus* GS 199 and *B. subtilis* GS 547 were evaluated and optimized for their extracellular digestive enzymes viz., amylase, lipase, protease and phytase *in vitro*. The digestive enzymes of all the four isolates were active at wide temperature and pH range. Further, purification and characterization of these digestive enzymes would pave for the development of therapeutic enzyme formulations to alleviate the burden of pancreatic insufficiency and other digestive disorders. In addition, further *in vitro* and *in vivo* evaluation of the safety of tested *Bacillus* spp. and their validation as GRAS (generally regarded as safe) would also helpful in oral administration of these isolates as source of digestive enzymes in improving digestion.

Acknowledgements

The authors sincerely acknowledge the funding support by Department




Figure 12: pH optimization for phytase activity.




Figure 13: Effect of substrate concentration on phytase activity.

of Biotechnology, Government of India, India, R&D project No.BT/PR7089/FNS/20/734/2013.

References

- Fieker A, Philpott J, Armand M (2011) Enzyme replacement therapy for pancreatic insufficiency: present and future. *Clin Exper Gastroenterol* 4: 55-73.
- Roxas M (2008) The role of enzyme supplementation in digestive disorders. *Altern Med Rev* 13: 307-314.
- Bu G, Luo Y, Chen F, Liu K, Zhu T (2013) Milk processing as a tool to reduce cow's milk allergenicity: a mini-review. *Dairy Sci Technol* 93: 211-223.
- Gayathri D and Rashmi BS (2014) Development of Celiac Disease; pathogenesis and Strategies to Control; A Molecular Approach. *J Nutr Food Sci* 4: 310.
- Rashmi BS, Gayathri D (2014) Partial Purification, Characterization of *Lactobacillus* sp G5 Lipase and their Probiotic Potential. *Int Food Res J* 21: 1737-1743.
- Gayathri D, Asha, Devaraja TN (2011) *Lactobacillus* sp. as probiotics for human health with special emphasis on colorectal cancer. *Ind J Sci Technol* 4: 1008-1014.
- Bae HC, Nam MS, Lee JY (2002) Probiotic Characterization of Acid- and Bile-tolerant *Lactobacillus salivarius* sub sp *salivarius* from Korean Faeces. *Asian Australas J Ani Sci* 15: 1798-1807.
- Gayathri D and Rashmi BS (2017) Mechanism of Development of Depression and Probiotics as Adjuvant Therapy for its Prevention and Management. *J Mental Health Prev* (Elsevier)
- Gayathri D (2016a) Short note on Probiotics for Mental health and Anxiety. *J Matern Pediatr Nutr* 2: 114.
- Asha, Gayathri D (2012) Synergistic impact of *Lactobacillus fermentum* and *L. plantarum* in 1, 2- dimethylhydrazine induced colorectal cancer in mice. *Experimental and therapeutics in Medicine* 3: 1049-1054.
- Gayathri D, Bhavyashree G, Rashmi BS (2016) Food Allergy: Immunological Mechanisms and Prevention Strategies. *EC Nutrition* 5: 1058- 1066.
- Gayathri D (2016b) Probiotics for total health: Better today and tomorrow. *J Matern Pediatr Nutr* 2(1).
- Layer P, Keller J (2003) Lipase supplementation therapy: standards, alternatives, and perspectives. *Pancreas* 26: 1-7.
- Griffin SM, Alderson D, Farndon JR (1989) Acid resistant lipase as replacement therapy in chronic pancreatic exocrine insufficiency: a study in dogs. *Gut* 30: 1012-1015.
- Gayathri D, Rashmi BS (2016) Anti-Cancer Properties of Probiotics; A Natural Strategy for Cancer Prevention and Management. *EC Nutrition* 5: 1191-1202.
- De Angelis M, Rizzello CG, Fasano A, Clemente MG, De Simone C, et al. (2006) VSL#3 probiotic preparation has the capacity to hydrolyze gliadin polypeptides responsible for Celiac Sprue. *Biochim Biophys Acta* 1762: 80-93.
- Rosado JL, Solomons NW, Lisker R, Bourges H (1984) Enzyme replacement therapy for primary adult lactase deficiency. Effective reduction of lactose malabsorption and milk intolerance by direct addition of beta-galactosidase to milk at mealtime. *Gastroenterology* 87: 1072-82.
- de Vrese M, Stegelmann A, Richter B, Fenselau S, Laue C, et al. (2001) Probiotics—compensation for lactase insufficiency. *Am J Clin Nutr* 73: 421s-429s.
- Cheryan M (1980) Phytic acid interactions in food systems. *Crit Rev Food Sci* 13: 297-336.
- Sandberg AN, Andlid T (2002) Phytogetic and microbial phytases in human nutrition. *Int J Food Sci Tech* 37: 823-833.
- Ianiro G, Pecere S, Giorgio V, Gasbarrini A, Cammarota G (2016) Digestive Enzyme Supplementation in Gastrointestinal Diseases. *Current Drug Metabolism* 17: 187-193.
- Beckhorn EL, Labbee MD, Underkofler LA (1965) Food Additives, Production and Use of Microbial Enzymes for Food Processing. *J Agric Food Chem* 13: 30-34.
- Schallmeyer M, Singh A, Ward OP (2004) Developments in the use of *Bacillus* species for industrial production. *Can J Microbiol* 50: 1-17.

