

# Modes of Action and Species-Specific Effects of Di-(2-ethylhexyl)Phthalate in the Liver

**Ivan Rusyn**

*Department of Environmental Sciences and Engineering, School of Public Health, University of North Carolina, Chapel Hill, North Carolina, USA*

**Jeffrey M. Peters**

*Department of Veterinary and Biomedical Sciences and the Center for Molecular Toxicology and Carcinogenesis, The Pennsylvania State University, University Park, Pennsylvania, USA*

**Michael L. Cunningham**

*National Toxicology Program, National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina, USA*

The industrial plasticizer di-(2-ethylhexyl)phthalate (DEHP) is used in manufacturing of a wide variety of polyvinyl chloride (PVC)-containing medical and consumer products. DEHP belongs to a class of chemicals known as peroxisome proliferators (PPs). PPs are a structurally diverse group of compounds that share many (but perhaps not all) biological effects and are characterized as non-genotoxic rodent carcinogens. This review focuses on the effect of DEHP in liver, a primary target organ for the pleiotropic effects of DEHP and other PPs. Specifically, liver parenchymal cells, identified herein as hepatocytes, are a major cell type that are responsive to exposure to PPs, including DEHP; however, other cell types in the liver may also play a role. The PP-induced increase in the number and size of peroxisomes in hepatocytes, so called 'peroxisome proliferation' that results in elevation of fatty acid metabolism, is a hallmark response to these compounds in the liver. A link between peroxisome proliferation and tumor formation has been a predominant, albeit questioned, theory to explain the cause of a hepatocarcinogenic effect of PPs. Other molecular events, such as induction of cell proliferation, decreased apoptosis, oxidative DNA damage, and selective clonal expansion of the initiated cells have been also proposed to be critically involved in PP-induced carcinogenesis in liver. Considerable differences in the metabolism and molecular changes induced by DEHP in the liver, most predominantly the activation of the nuclear receptor peroxisome proliferator-activated receptor (PPAR) $\alpha$ , have been identified between species. Both sexes of rats and mice develop adenomas and carcinomas after prolonged feeding with DEHP; however, limited DEHP-specific human data are available, even though exposure to DEHP and other phthalates is common in the general population. This likely constitutes the largest gap in our knowledge on the potential for DEHP to cause liver cancer in humans. Overall, it is believed that the sequence of key events that are relevant to DEHP-induced liver carcinogenesis in rodents involves the following events whereby the combination of the molecular signals and multiple pathways, rather than a single hallmark event (such as induction of PPAR $\alpha$  and peroxisomal genes, or cell proliferation) contribute to the formation of tumors: (i) rapid metabolism of the parental compound to primary and secondary bioactive metabolites that are readily absorbed and distributed throughout the body; (ii) receptor-independent activation of hepatic macrophages and production of oxidants; (iii) activation of PPAR $\alpha$  in hepatocytes and sustained increase in expression of peroxisomal and non-peroxisomal metabolism-related genes; (iv) enlargement of many hepatocellular organelles (peroxisomes, mitochondria, etc.); (v) rapid but transient increase in cell proliferation, and a decrease in apoptosis; (vi) sustained hepatomegaly; (vii) chronic low-level oxidative stress and accumulation of DNA damage; (viii) selective clonal expansion of the initiated cells; (ix) appearance of the pre-neoplastic nodules; (x) development of adenomas and carcinomas.

**Keywords** Cancer, Human, Liver, Mouse, Peroxisome, Proliferators

## ABSORPTION, DISTRIBUTION, METABOLISM, AND EXCRETION OF DEHP AND THEIR RELEVANCE TO EFFECTS ON THE LIVER

Oral exposure of DEHP, primarily through food and water, is the most likely route of exposure in humans [reviewed in (Fay et al., 1999)]. It is the only route of exposure known to lead to liver cancers in rodents (National Toxicology Program, 1982). Inhalation, dermal, and parenteral routes of exposure, as well as hemodialysis, are also possible in humans, but they are not generally considered to be relevant for the potential carcinogenic effect of DEHP in liver or other organs even in susceptible species [reviewed in (Fay et al., 1999)]. However, these routes of exposures may be significant for some industrial workers and individuals undergoing dialysis.

DEHP is a lipophilic compound and can be absorbed through skin and lungs by both humans and rodents. However, the greatest absorption occurs after oral exposure. Once DEHP enters the gastrointestinal tract, it is rapidly metabolized to mono(2-ethylhexyl)phthalate (MEHP) and 2-ethylhexanol via pancreatic lipases (Figure 1). At low concentrations, most of DEHP is absorbed as these two metabolites, but at high doses some unmetabolized DEHP can also be absorbed (Albro, 1986). It was estimated that human absorption is as high as 25% (Schmid and Schlatter, 1985), but rats absorb more than 55% of the oral dose (Rhodes et al., 1986). Furthermore, non-human primates appear to absorb a smaller percentage of the oral dose than rats (Astill, 1989). Once absorbed, DEHP and its metabolites are distributed throughout the body in the blood where unhydrolyzed DEHP can be metabolized via non-specific esterases. Several studies of DEHP distribution after oral administration in rodents and other species, including monkeys, showed that liver contains the greatest amount of DEHP and its metabolites under the condition of repeated exposure (Rhodes et al., 1986).

The primary metabolic conversion of DEHP following oral administration is qualitatively similar in all species studied, including non-human primates and humans, and involves formation of MEHP and 2-ethylhexanol (Figure 1), followed by formation of phthalic acid through the activity of lipases in the gut and many parenchymal organs. Further metabolism can occur via oxidation by cytochrome P450 4A, alcohol and aldehyde dehydrogenases and yields multiple oxidation products of 2-ethylhexanol and the aliphatic side chain of MEHP [Figure 1, reviewed in (Fay et al., 1999)]. The downstream metabolites of DEHP can also be glucuronated before excretion. More than 100-fold differences in activity of DEHP-metabolizing enzymes were reported to exist in liver, intestine, kidney, and lung between rodents and non-human primates (Ito et al., 2005). Notably, significantly less activity is found in non-human primates, suggesting that DEHP is metabolized to its active metabolite (MEHP) to a lesser degree in this species as compared to rodents. This is also consistent with previous studies that indicate that lipase activity and absorption of DEHP is significantly less in humans and/or non-human

primates as compared to rodents [reviewed in (Klaunig et al., 2003)].

Excretion of DEHP and its metabolites occurs through urine and feces (from biliary excretion and unmetabolized/unabsorbed matter). In humans, only urinary excretion was studied and showed that 15–25% of the dose is excreted through this route (Schmid and Schlatter, 1985). Animal studies show that urinary excretion is the predominant route of elimination for DEHP metabolites, while unmetabolized parent compound is predominantly eliminated with feces (Tanaka et al., 1975). More than half of the dose is excreted in the first 24 hrs, and about 80% is excreted within 5–7 days (Astill, 1989; Ikeda et al., 1980; Tanaka et al., 1975). Humans and non-human primates excrete DEHP metabolites predominantly in the form of glucuronidated conjugates, while guinea pigs, hamsters and mice excrete smaller amount of these conjugates, and none of the metabolites in rat urine were found to be conjugated (Albro et al., 1982; Dirven et al., 1993b).

People exposed to DEHP in the workplace routinely test positive for DEHP-related metabolites in the urine (Dirven et al., 1993a). Recent studies have found that humans with no known exposures have detectable concentrations of DEHP metabolites in their urine and serum, and these reference levels have been used to estimate human exposure levels. For example, the concentration of urinary MEHP ranged from undetectable levels up to 66.6 ng/mL, with an average concentration of 3.5 ng/mL (Blount et al., 2000). However, these values reflected both free and glucuronidated forms of MEHP, which could overestimate the actual biologically active form. Others have estimated that while the total concentration of MEHP (free + glucuronidated) in urine ranges from 1.2 to 26.0 ng/mL, with an average concentration of 3.88 ng/mL, the free concentration of MEHP ranged from undetectable to 10.0 ng/mL (Silva et al., 2003). Interestingly, there is good correlation with urinary and serum levels of MEHP (Silva et al., 2003). Based on these biomarker levels of DEHP metabolites, it has been estimated that humans are exposed to DEHP at a concentration ranging from 3–52  $\mu\text{g}/\text{kg}/\text{day}$  (Kavlock et al., 2002; Koch et al., 2003). While these findings demonstrate environmental exposure to DEHP in the general population, the physiological relevance of this exposure is under debate. This is further complicated by the fact that there could be additive effects from other environmental phthalates that may or may not activate PPAR $\alpha$ , and that in some cases (e.g., occupational exposure, dialysis patients, etc.), exposure levels may be significantly greater than those previously reported.

The metabolites of DEHP, primarily MEHP, but not the parental compound per se, are considered to be downstream “biologically active” entities that are responsible for its carcinogenic effect in rodent liver (Deangelo et al., 1986; Ward et al., 1986). In addition, both primary and secondary metabolites of DEHP can activate PPAR $\alpha$  and other isoforms of this nuclear receptor family in both rodents and humans (Hurst and Waxman, 2003; Maloney and Waxman, 1999). Thus, better understanding




FIG. 1. **Metabolic fate of DEHP.** Lypolysis of DEHP by esterases in different tissues results in the formation of MEHP and 2-ethylhexanol. 2-ethylhexanol can be further metabolized with its ultimate fate being acetate and carbon dioxide. MEHP can be further metabolized by esterases to form another molecule of 2-ethylhexanol and phthalic acid. Alternatively, the sidechain of MEHP can be further oxidized to a variety of metabolites. MEHP and any of its downstream metabolites can be glucuronidated and then excreted.

of formation, distribution, and excretion of DEHP metabolites in different species, especially humans, and across various organs is a clear research need that is equally important to studies on species- and tissue-diversity in expression of PPARs (see below).

### HISTOPATHOLOGICAL CHANGES IN LIVER FOLLOWING EXPOSURE TO DEHP

*Rats and mice* are mammalian species that are most susceptible to the characteristic hepatic effects of DEHP and other PPs. Hepatomegaly—marked increase of the liver mass—is the most pronounced early gross pathological change that occurs after continuous oral exposure to DEHP in rodents (Lake et al., 1975; Reddy et al., 1976a). This effect is robust, sustainable over the time course under conditions of continuous exposure, highly reproducible between laboratories, and appears to be a result of both hypertrophy (increase in the size of each cell) and hyperplasia (increase in the number of cells) of liver parenchymal cells. Microscopically, a midzonal to periportal accumulation of fat, a decline in glycogen deposits

in centrilobular areas, structural changes in the bile ducts, and accumulation of lipofuscin granules were reported in rats and mice treated with DEHP for various lengths of time (Mitchell et al., 1985; Ward et al., 1998). Ultrastructurally, hepatocellular hypertrophy is associated with an increase in both size and number of peroxisomes, mitochondria, and lipid-loaded lysosomes (Mitchell et al., 1985; Nair and Kurup, 1987). In parallel, a burst of proliferation is observed in DEHP-treated rats and mice. Elevation in the rate of mitosis of hepatocytes is observed as early as 24 hrs after the initial dose (Smith-Oliver and Butterworth, 1987), and it is gradually increased until about 1 week of exposure but is not sustained over longer time courses in either rats, or mice (Marsman et al., 1988; Smith-Oliver and Butterworth, 1987; Ward et al., 1988). Furthermore, a decrease in apoptosis was also suggested to be an important additional molecular event that may affect the number of cells in rodent liver following exposure to DEHP, although the apoptosis rates were not investigated past 4 days of exposure and thus the time-course of this event is uncertain (Hasmall et al., 2000b; James et al., 1998).

The long-term consequence associated with the above-mentioned events is the development of hepatocellular adenomas and carcinomas in rodents. A two-year rodent bioassay conducted by the National Toxicology Program indicated that there was a statistically significant positive correlation between DEHP exposure and hepatocellular carcinomas or neoplastic nodules of liver in both sexes of mice (B6C3F1) and rats (F344) (National Toxicology Program, 1982). These observations have since been repeated with additional dose- and time-response studies (David et al., 1999) and largely confirmed the original findings. Interestingly, studies have shown that most of the effects of DEHP, such as tumor incidence, liver weight, and peroxisomal proliferation, are reversible after cessation of treatment, which demonstrates a temporal relationship between exposure and liver carcinogenesis (David et al., 1999). Finally, DEHP is known to promote initiated cell in vivo as exemplified by the ability to induce tumors after exposure to a DNA-damaging agent (Oesterle and Deml, 1988; Ward et al., 1983).

Several studies evaluated the pleiotropic responses to prolonged (from 14 days up to 13 weeks) oral administration of relatively high doses of DEHP (500 to 2500 mg/kg) in several species of *non-human primates*. In contrast to results found in rodents (see above), no significant increases in liver weight, induction of peroxisomal enzymes, or proliferation of peroxisomes were reported (Kurata et al., 1998; Pugh, Jr. et al., 2000; Short et al., 1987). Very limited data are available pertaining to hepatic effects in *humans* exposed to DEHP. There are two reports suggesting that dialysis patients exhibit hepatic peroxisome proliferation, which presumably could be due to exposure to DEHP or other phthalates in the plastic tubing (Ganning et al., 1984; Ganning et al., 1987). However, neither quantified data were provided, nor was it indicated whether samples were collected at the time of exposure or at a later time. Thus, the extent of this putative DEHP-dependent effect in human liver is uncertain. Studies that examined patients treated with relatively more potent PPs (e.g., clofibrate, gemfibrozil, or fenofibrate), for prolonged periods of time (i.e., years) are more consistent with the idea that humans do not exhibit peroxisome proliferation in response to exposure to PPs. Similar to findings made in hemodialysis patients, a marginal 50% increase in liver peroxisome number, but not in peroxisome volume is reported in humans treated with clofibrate (Hanefeld et al., 1983). In contrast, the majority of studies examining the effect of PP administration in humans have consistently shown no change in hepatic peroxisome proliferation in liver [reviewed in (Peters et al., 2005)].

There are no known reports of long-term carcinogenesis studies with DEHP in *non-human primates*. One study published in non-peer-reviewed literature reported on a 7-year toxicity study of clofibrate (94 to 263 mg/kg/day) in the marmoset (Tucker and Orton, 1995). While these findings should be interpreted with caution since the number of animals examined was small and very limited data were included in the report, no evidence of

peroxisome proliferation, liver enlargement, or a carcinogenic effect in the liver or any other tissue was observed.

Several studies have evaluated the long-term effects of exposure to DEHP in *humans*. Only one known study evaluated causes of mortality among people exposed to DEHP via inhalation at workplace and found no conclusive evidence of increased death rates from carcinogenesis in this cohort [as reported in (Klaunig et al., 2003); original manuscript not available]. Other known reports investigated workers manufacturing PVC or patients in long-term contact with PVC-containing medical devices. Several large epidemiological studies that examined the relationship between chronic treatment with lipid-lowering PPs gemfibrozil and clofibrate did not find an association with liver cancer [reviewed in (Peters et al., 2005)]. Collectively, while human studies did not show an association between liver cancer and treatment with PPs, they all have important limitations. Thus, it is difficult to draw an unequivocal scientifically-based conclusion for or against a link between long-term DEHP exposure and risk of cancer of liver or other organs (e.g., testes or pancreas) in humans.

#### PLEIOTROPIC RESPONSES TO DEHP IN VITRO IN CULTURED HEPATOCYTES

Extensive studies on the molecular mechanisms of action of PPs, including DEHP metabolites such as MEHP, have been conducted using hepatocytes from rodent, humans, and other species. Such experiments represent the best available direct comparison of responses between rodents and humans. However, it is important to note that the overall magnitude of response to PPs in cultured hepatocytes is typically weaker than that observed in vivo in whole liver, which raises a question of whether other cell types (e.g., Kupffer cells, see below) are necessary to achieve the full spectrum of responses [reviewed in (Rose et al., 1999b)].

With respect to the effects of MEHP in cultured isolated *rat hepatocytes*, several studies reported a significant increase in DNA replication, peroxisomal gene induction, and suppression of apoptosis. While the induction of peroxisomal genes in vitro in rodent hepatocytes is robust and comparable to that in whole liver (Goll et al., 1999; Hasmall et al., 1999), the increase in DNA synthesis, a marker of cell proliferation in vitro, has been reported to be only 120–200%, which is markedly less than that observed in vivo (Goll et al., 1999; Hasmall et al., 1999). Suppression of apoptosis studies in cultured rat hepatocytes were conducted under both “spontaneous” and growth factor (e.g., TGF $\beta$ 1)-induced conditions (Goll et al., 1999; Hasmall et al., 1999; Hasmall et al., 2000b). However, while the qualitative effect of MEHP was similar to that of DEHP in whole rat liver, the magnitude of response was muted even though the rates of apoptosis in culture were much higher than in vivo (Hasmall et al., 2000b).

One study compared the effect of MEHP and metabolites on the ability of rat, rabbit, guinea pig, and *monkey hepatocytes* to induce changes in gene expression indicative

of peroxisome proliferation (Dirven et al., 1993c). It was concluded that hepatocytes from all species included in the study were responsive to MEHP by inducing peroxisomal palmitoyl-CoA oxidase, but rat hepatocytes were about 30-fold more sensitive based on the dose-response studies. The effects of MEHP on *human hepatocytes* have also been studied in vitro by several groups (Goll et al., 1999; Hasmall et al., 1999; Hasmall et al., 2000b). A general conclusion from these experiments is that human hepatocytes are largely non-responsive to MEHP when DNA synthesis, apoptosis, or peroxisomal gene induction are evaluated. While a trend for activation of  $\beta$ -oxidation capacity of human cells (2- to 4-fold) was shown, these changes were not statistically significant (David et al., 1999; Hasmall et al., 2000b). In addition, hepatocytes from one individual exhibited a small but highly significant increase in DNA replication in response to both MEHP and another phthalate, di-isononyl phthalate (DINP) (David et al., 1999). It should be noted that a limited number of humans were sampled, which may have affected the significance levels of these studies due to presumably higher genetic variability in human population. In addition, an unknown condition of the livers used for cell isolations, as well as duration and condition of storage of human cells before the experiments, may also be considered as confounding variables. A larger database from more representative human samples would improve our understanding, and these datasets should also demonstrate that cultured cells are responsive to PPAR $\alpha$ -dependent molecular events known to occur in human liver.

#### PROPOSED MODES OF ACTION FOR DEHP IN HEPATOCYTES

Several major modes of action have been proposed to be responsible for short- and long-term effects of PPs, including DEHP, in the hepatocyte, a target cell for liver carcinogenesis due to these compounds. Below, the molecular events that are thought to be involved in each pathophysiological process are discussed and compared between species.

##### Activation of PPAR $\alpha$

A discovery of a family of the nuclear receptors, peroxisome proliferator-activated receptors (PPARs) (Issemann and Green, 1990), shed light on the mechanism of action of PP compounds, as well as opened new opportunities for therapeutic interventions in many human diseases (Berger et al., 2005). These steroid hormone-like receptors act as transcription factors upon ligand binding. PPARs play a number of important roles in normal physiology [reviewed in (Berger et al., 2005)]. For example, PPAR $\alpha$  serves a fundamental role in mammals by acting as a central modulator of signaling molecules that mediate changes in gene expression to maintain lipid homeostasis (Duval et al., 2004). PPARs generally form a heterodimer with the retinoid-X receptor (RXR) before translocating to the nucleus and binding to the peroxisome proliferator response elements (PPRE) on

DNA, and subsequently activating transcription of specific target genes. PPARs are expressed in all tissues, though the distribution of different isoforms varies between organs and individual cell types. Of the three isoforms that have been cloned to date,  $\alpha$ ,  $\beta/\delta$ , and  $\gamma$ , the former is the primary form responsible for mediating PP-induced effects in the liver, and it is ubiquitously expressed in hepatocytes in all species (Corton et al., 2000). Still, significant cross-talk between different forms of PPARs and other transcription factors exists (Shipley and Waxman, 2004), and PPs, including MEHP, are able to activate other PPAR isoforms, particularly PPAR $\gamma$  (Bility et al., 2004; Hurst and Waxman, 2003).

All synthetic PPs, including some of the downstream metabolites of DEHP, function as PPAR $\alpha$  ligands. For example, MEHP and 2-ethylhexanoic acid can both transactivate PPAR $\alpha$  reporter constructs in transiently transfected cells (Maloney and Waxman, 1999), but there is a large species difference as rodent PPAR $\alpha$  is transactivated by MEHP with lower concentrations and with greater efficacy as compared to human PPAR $\alpha$  (Bility et al., 2004; Hurst and Waxman, 2003; Maloney and Waxman, 1999). Treatment of transfected cells with DEHP caused no change in reporter activity, demonstrating that DEHP must be metabolized in order to transactivate PPAR $\alpha$  (Maloney and Waxman, 1999). The ability of DEHP metabolites to transactivate PPAR $\alpha$  also correlates well with the ability to increase PPAR $\alpha$ -dependent target gene expression. Increased activity of PPAR $\alpha$ -regulated genes by MEHP, but not unmetabolized parent compound DEHP, was first reported to occur in COS1 monkey kidney epithelial cells (Issemann and Green, 1990). A number of other studies have also shown good correlation between transactivation of PPAR $\alpha$  in vitro with increased expression of PPAR $\alpha$  target gene expression. Additionally, these changes also confirm the reported species differences as changes in transactivation and target gene expression occur with lower concentrations and result in changes in target gene expression with greater efficacy with rodent PPAR $\alpha$  as compared to human PPAR $\alpha$  (Bility et al., 2004; Hasmall et al., 2000b; Hurst and Waxman, 2003; Lapinskas et al., 2005; Maloney and Waxman, 1999; Yu et al., 2001). Interestingly, MEHP, but not 2-ethylhexanoic acid, can also transactivate PPAR $\gamma$ , and this also correlates well with the ability to induce functional changes in adipocyte differentiation indicative of PPAR $\gamma$  activity (Bility et al., 2004; Hasmall et al., 2000b; Hurst and Waxman, 2003; Lapinskas et al., 2005; Maloney and Waxman, 1999; Yu et al., 2001). Lastly, MEHP is also reported to transactivate PPAR $\beta/\delta$  (Bility et al., 2004). However, the induction of many target genes induced by DEHP is mediated by PPAR $\alpha$  as these changes in gene expression do not occur in PPAR $\alpha$ -null mice (Ward et al., 1998). Thus, while there is good reason to examine the PPAR $\alpha$ -mediated effects induced by DEHP and DEHP metabolites, it remains possible that DEHP and DEHP metabolites could alter cellular functions by interacting with other PPARs and/or through other "non-PPAR"-mediated events (discussed below).

MEHP is a relatively weak PPAR $\alpha$  agonist as compared to fibrates or more highly specific ligands such as GW7647 [reviewed in (Klaunig et al., 2003)]; still, both rodent and human PPAR $\alpha$  can be transactivated by MEHP in vitro. Mouse PPAR $\alpha$  is more sensitive to MEHP ( $EC_{50} = 0.6 \mu\text{M}$ ) than human PPAR $\alpha$  ( $EC_{50} = 3.2 \mu\text{M}$ ) (Hurst and Waxman, 2003). Since it was reported that the average concentration of MEHP found in human serum is about  $\sim 18 \text{ nM}$ ; (Silva et al., 2003), it appears that the amount of MEHP found in human tissues is about 100-times lower than that required to activate the hPPAR $\alpha$ . While comparing the concentration of MEHP in human liver would be more informative, these data are not available.

It has been hypothesized that there are at least three molecular levels of possible *species-specific differences* in pleiotropic responses to PPs that concern PPARs, including the relative expression levels of the receptor, differences in the ability of PPAR $\alpha$  to activate target gene expression, and differences in the ability of the receptor to interact with other transcriptional proteins. While it appears that human hepatocytes contain fully functional PPAR $\alpha$  (Mukherjee et al., 1994; Sher et al., 1993), the relative expression levels of this receptor in human liver is about 10-fold lower than that in the rat or mouse (Palmer et al., 1998; Sher et al., 1993). However, another report suggests that human liver may express comparable levels of PPAR $\alpha$  as compared to rodents (Walgren et al., 2000) which suggests that significant variability in PPAR $\alpha$  expression between individuals in human population is likely. In addition, truncated and polymorphic forms of the protein have been found in human population (Flavell et al., 2000; Gervois et al., 1999; Palmer et al., 1998; Roberts et al., 2000; Sapone et al., 2000; Vohl et al., 2000). In one case, the truncated form of PPAR $\alpha$  was shown to act as a dominant negative (Gervois et al., 1999). Still, MEHP can bind to the human PPARs, but there are clear differences in the ability of MEHP to transactivate PPAR $\alpha$  between rodents and humans (Bility et al., 2004; Hassmall et al., 2000b; Hurst and Waxman, 2003; Maloney and Waxman, 1999; Yu et al., 2001). Thus, while there is evidence suggesting that humans may express less PPAR $\alpha$  as compared to rodents, in addition to polymorphic alleles, the functional significance of these variants is uncertain [reviewed in (Peters et al., 2005)].

The ability of the activated receptor to bind to PPREs on the genes responsible for the pleiotropic response of PPs has also been postulated to underlie the reported species difference in PPAR $\alpha$  activity. When activated, both rodent and human PPARs are capable to transactivate reporter genes driven by rodent-specific PP-responsive promoters (Hurst and Waxman, 2003; Kohli et al., 1999; Maloney and Waxman, 1999). Conversely, human PPAR $\alpha$  rescues the PP-induced peroxisomal proliferation response in liver of PPAR $\alpha$ -null mice when it is over-expressed transiently (with adenovirus injection) or permanently (in transgenic mice) (Cheung et al., 2004; Yu et al., 2001). However, several human-specific PPREs in peroxisomal genes studied to date were found to be largely non-responsive to transactivation by either mouse or rodent

PPAR $\alpha$  (Hassall et al., 2000b; Woodyatt et al., 1999). Thus, there is some uncertainty regarding the role of polymorphisms in functional PPREs in the reported species differences observed in response to PPs. Importantly, genes responsible for lipid-lowering effects of several PP compounds contain PPREs that are highly responsive to activation in humans, but not rats (Vu-Dac et al., 1998; Vu-Dac et al., 2003). Clearly, this discordance underscores the importance of this level of regulation of PPAR $\alpha$ -mediated events in different species. This is also consistent with working showing that expression of acyl-CoA oxidase mRNA is not increased in human liver following exposure to fibrate therapy (Roglans et al., 2002), while it is well documented that this mRNA increases significantly in fibrate-treated rodents.

It has also been hypothesized that other proteins that interact with PPAR $\alpha$  could alter the specificity of gene activation, and this may differ between cell types and species (Hertz and Bar-Tana, 1998). It was shown that some proteins, such as heat shock protein (Hsp)90 and the hepatitis virus B X-associated protein 2 (XAP2), complex with PPAR $\alpha$  in the cytoplasm and may serve to suppress ligand activation (Sumanasekera et al., 2003b; Sumanasekera et al., 2003a); however, no data for potential species differences in this mechanism are available at this time.

On a more functional level, PPAR $\alpha$  activation is intertwined with other modes of action for pharmacological and unwanted (e.g., liver carcinogenesis) effects of DEHP and other PPs. The development of the PPAR $\alpha$ -null mouse opened new research opportunities that helped to define the causal links that exist in rodent liver between PPs, induction of lipid and xenobiotic metabolism genes, proliferation of peroxisomes and other organelles in hepatocytes, increases in cell proliferation, and decreases in apoptosis. Most notably, the PPAR $\alpha$ -null mouse was instrumental for demonstrating that the receptor is required for liver tumorigenesis induced by a model PP and potent rodent hepatocarcinogen WY-14,643. Specifically, a long-term study in PPAR $\alpha$  homozygous and nullizygous mice showed that feeding WY-14,643 for up to 11 months resulted in all wild-type mice developing hepatocellular adenomas or carcinomas (Peters et al., 1997). PPAR $\alpha$ -null mice administered the same treatment did not develop tumors, nor were any of the other pleiotropic responses associated with PP-treatment observed. Similar negative results were obtained recently for another PP, bezafibrate (Hays et al., 2005). While no long-term carcinogenesis study was performed with DEHP in PPAR $\alpha$ -null mice, it was shown that for up to 24 weeks of feeding a carcinogenic dose of DEHP (12,000 ppm) failed to induce proliferation of peroxisomes or liver enlargement in PPAR $\alpha$ -null mice (Ward et al., 1998).

Most recently, the examination of the mechanisms determining species differences in peroxisome proliferator response between mice and humans became possible with a PPAR $\alpha$ -humanized mouse line (Cheung et al., 2004). Human PPAR $\alpha$  was expressed in liver of PPAR $\alpha$ -null mouse strain, thus

providing an *in vivo* model to examine the reported species difference. In early studies where these mice were fed WY-14,643 or another PP, fenofibrate, the *PPAR* $\alpha$ -humanized and wild-type mice responded to treatment with *PPAR* $\alpha$  ligands as revealed by induction of genes encoding peroxisomal and mitochondrial fatty acid metabolizing enzymes and resultant decrease of serum triglycerides. However, only the wild-type mice and not the *PPAR* $\alpha$ -humanized mice exhibited hepatocellular proliferation and hepatomegaly (Cheung et al., 2004). More recently, long-term feeding studies demonstrate that the humanized *PPAR* $\alpha$  mice are also resistant to Wy-14,643-induced liver cancer (Morimura et al., 2006). This demonstrates that it is highly likely that there are large species differences in the receptor function, and establishes that following ligand activation, the *PPAR* $\alpha$ -mediated pathways controlling lipid metabolism are independent from those controlling the cell proliferation pathways that lead to hepatocellular carcinogenesis (see below).

#### **Proliferation of Peroxisomes and Induction of Peroxisomal Proteins**

Peroxisome proliferators, as the name implies, cause an increase in the number and size of peroxisomes. These ubiquitous organelles are present in essentially all cell types and are largely responsible for fatty acid  $\beta$ -oxidation of very long chain fatty acids, oxidation of cholesterol derivatives, D- and L- amino acids oxidation, and purine catabolism. In livers of rodents, the peroxisomes house an array of  $H_2O_2$ -generating oxidases, including acyl-coA oxidase, along with  $H_2O_2$ -degrading enzyme catalase (Schrader and Fahimi, 2004). *PPAR* $\alpha$  acts as a transcription factor involved in regulating expression of several target genes in peroxisomes, most of which contain a PPRE in their promoter region (Desvergne and Wahli, 1999; Reddy and Chu, 1996). Many of the genes identified as *PPAR* $\alpha$  target genes are involved in fatty acid metabolism (Lee et al., 1995). For example, mitochondrial and peroxisomal fatty acid enzyme expression is affected by *PPAR* $\alpha$ .

Activity of enzymes involved in lipid  $\beta$ -oxidation (i.e., acyl-coA oxidases) is dramatically increased in livers of rats and mice following even a single-dose treatment with PP compounds (Reddy et al., 1980). By far, this hallmark feature associated with administration of *PPAR* $\alpha$  ligands in rats and mice is the best and well-accepted phenotype that serves as a biomarker that determines the classification of chemical compounds into the class of peroxisome proliferators. A number of investigators over the past 30 years demonstrated that DEHP exhibits similar effects on peroxisomal oxidases and other enzymes, as well as density and number of peroxisomes in hepatocytes, in both *rats and mice* (Barber et al., 1987; David et al., 1999; Mitchell et al., 1985; Reddy et al., 1976a). Multiple strains of rats and mice and both sexes in these species are responsive, and both time- and dose-response have been reported.

*Non-human primates* and *humans* are considered to be far less responsive species when proliferation of peroxisomes is

considered as an endpoint. While some reports of modest, often non-significant, induction of peroxisomal oxidases in cultured human hepatocytes and *in vivo* in patients on chronic hemodialysis have been brought forward (see above), it is well accepted that proliferation of peroxisomes is a minor and not reliable phenotype caused by DEHP or other PPs in people. It should be noted, however, that ciprofibrate, when given to cynomolgus monkeys for 15 days in doses much higher than therapeutic (up to 400 mg/kg/day), caused a treatment-related increase in liver weight, periportal hepatocellular hypertrophy, increase in the number of peroxisomes (~3 times) and mitochondria (~2.5 times), and increase in activity of peroxisomal enzymes (Hoivik et al., 2004). A 3-year long study in marmosets with lower (2–20 mg/kg/day) doses of ciprofibrate found an up to 2.5-fold induction of activity of peroxisomal proteins, but no increase in either liver weight or peroxisomal number and size (Graham et al., 1994). Collectively, no direct inter-species comparison of plasma concentrations of ciprofibrate has been performed. However, based on the values reported by Hoivick et al. (2004) for monkeys and studies of Easton et al. (1989) in the rat, we estimate that the highest dose studied in non-human primates is comparable to the dose that is carcinogenic in the rat.

While peroxisome proliferation and induction of fatty acid metabolism are hallmark responses to PPs, a causal link between this response and liver tumor formation, even in rodents, is debated. One of the original hypotheses that linked proliferation of peroxisomes with liver carcinogenesis was the fact that  $H_2O_2$ -generating oxidases are induced disproportionately compared to  $H_2O_2$ -degrading enzyme catalase, thus causing a state of chronic oxidative stress in the liver that contributes to carcinogenesis via oxidative DNA damage [see below and (Reddy et al., 1980)]. Although it is clear that peroxisome proliferation occurs in response to treatment with PPs, including DEHP, the role of this organelle in the hepatocarcinogenic response does not appear to be a sole causative event. This is based on the finding that the degree of peroxisome proliferation observed in rats fed peroxisome proliferators does not correlate well with tumorigenicity (Marsman et al., 1988). Thus, peroxisomal induction continues to be regarded as an important associative endpoint (Klaunig et al., 2003).

#### **Induction of Non-Peroxisomal Metabolism Proteins**

In addition to peroxisomal fatty acid catabolism, PPs are known to also affect fatty acid  $\beta$ -oxidation in mitochondria, and  $\omega$ -hydroxylation in microsomes. These effects, together with the recently established role of *PPAR* $\alpha$  in fatty acid uptake and transport, ketogenesis, and lipogenesis in liver [reviewed in (Mandard et al., 2004)], clearly point to the overall importance of *PPAR* $\alpha$ -ligands for physiological regulation of lipid and energy metabolism. Specifically, it was shown that *PPAR* $\alpha$  regulates mitochondrial fatty acid oxidation by controlling expression of a key oxidative enzyme, medium-chain acyl-CoA dehydrogenase (Gulick et al., 1994). The  $\omega$ -hydroxylation is governed by a

subfamily of cytochrome P450 4A enzymes that are known to be regulated by PPAR $\alpha$  (Aldridge et al., 1995; Kroetz et al., 1998). A number of electron transport chain proteins in the mitochondria, as well as CYP4A isoforms, are known to be “leaky” enzymes that can generate secondary reactive oxygen species and thus potentially contribute to oxidative DNA damage and carcinogenesis due to PPs (see below); thus, these two non-peroxisomal pathways are important to consider in the evaluation of the mechanisms of DEHP and related compounds.

Indeed, DEHP and its metabolites MEHP and 2-ethylhexanol were shown to cause a significant dose-related induction of CYP4A1 and lauric acid  $\omega$ -hydroxylase activity in *rat* liver in vivo even at about 1/4 of a dose used for the 2-year NTP carcinogenesis study (Dirven et al., 1990; Dirven et al., 1992). Similar effects were observed in cultured rat hepatocytes (Dirven et al., 1993c), and in the mouse (Currie et al., 2005; Ward et al., 1998). Induction of CYP4A genes by DEHP is dependent on the presence of functional PPAR $\alpha$  (Ward et al., 1998). Interestingly, MEHP fails to induce  $\omega$ -hydroxylase activity in cultured hepatocytes from *other species*, such as guinea pig, rabbit, and monkey (Dirven et al., 1993c). No experiments have been performed on human hepatocytes to address the effects of DEHP or its metabolites on these endpoints.

The effects of DEHP on the mitochondria appear to be more complex than just regulation of expression of  $\beta$ -oxidation proteins. Despite the fact that expression of some genes encoding mitochondrial fatty acid  $\beta$ -oxidation enzyme is regulated by the PPAR $\alpha$  (Aoyama et al., 1998; Gulick et al., 1994), numerous studies showed that phthalates and other PPs inhibit mitochondrial  $\beta$ -oxidation in *rodent hepatocytes* as a result of diminished mitochondrial respiration at the level of cytochrome c reductase (Badr et al., 1990; Keller et al., 1992; Winberg and Badr, 1995). While this effect on the mitochondria may explain the accumulation of lipids in the rodent liver observed following administration of phthalates (Hinton et al., 1986), it is not known what effect this may have on the carcinogenesis. Furthermore, it is not presently known whether similar effects are also observed in human cells. It remains possible that inhibition of mitochondria function by PPs, including DEHP, could occur through PPAR $\alpha$ -independent mechanism, which will be discussed later.

It is likely that other genes that modulate metabolism in hepatocytes are also inducible by DEHP in a PPAR $\alpha$ -dependent manner. For example, a recent report showed that CYP1A1, a known Ah receptor target gene, is highly inducible by PPAR $\alpha$  ligands, including MEHP, in *human cell lines* CaCo-2 (derived from colon adenocarcinoma), HepG2 (hepatoma), A549 (lung adenocarcinoma), as well as primary human keratinocytes (Seree et al., 2004). This induction specifically involved PPAR $\alpha$  and required 2 PPRE sites that were located within the CYP 1A1 promoter. Whether this also occurs in normal human hepatocytes is yet to be established. CYP 1A1 is a major P450 isoform that is responsible for bioactivation of a number of environmental pro-carcinogens, and it was reported that DEHP

and other PPs enhance DNA adduct formation in rat liver when co-administered with benzo[*a*]pyrene (Voskoboinik et al., 1997). Thus, the fact that MEHP and other PPs are capable of regulating CYP 1A1 in PPAR $\alpha$ -mediated manner in human cells may have significant implications for the assessment of potential carcinogenic risk by environmental compounds that are PPAR $\alpha$  ligands. However, others have shown that clofibrate inhibits expression of CYP1A1 and CYP1A2 in liver cells (Shaban et al., 2004). This would suggest that treatment with PPs such as DEHP would reduce bioactivation of chemical carcinogens. Thus, it is currently unclear how PPs regulate other important CYPs in the liver, and whether these changes result in physiological significant changes.

### Induction of Cell Proliferation

Several classes of rodent non-genotoxic liver carcinogens, such as PPs and phenobarbital-like compounds, exhibit a characteristic pattern of changes in hepatocyte proliferation rates. This process is characterized by a rapid, within hours of a single dose of a compound, and remarkable surge in the number of proliferating cells. The increase reaches a peak of several- to more than ten-fold over control levels within a week after initiation of continuous exposure, yet it subsides gradually to nearly control levels despite the prolongation of the treatment (Marsman et al., 1988). The proliferative response appears to be mitogenic in nature, as there is no evidence of cell necrosis in the first weeks of treatment with PPs and other compounds. It is well accepted that induction of cell proliferation is an important step in the tumorigenesis by genotoxic and non-genotoxic carcinogens. Increased cell replication facilitates the fixation of DNA damage (either from direct interaction with DNA, or indirect oxidative damage) and silencing of tumor suppressor genes. These conditions promote clonal expansion of mutated cells and development of focal lesions in the liver that progress to adenomas and carcinomas.

Induction of cell proliferation is a phenotype characteristic of exposure to PPs in *rat and mouse liver* (Marsman et al., 1988; Reddy et al., 1976b; Reddy et al., 1979). Extensive time- and dose-response studies of proliferative effects of DEHP are available in the *rat* (Marsman et al., 1988; Mitchell et al., 1985; Smith-Oliver and Butterworth, 1987). A rapid burst of cell proliferation (peak at 3 days) occurs in a dose-dependent manner and subsides by 7–14 days of treatment (Mitchell et al., 1985). Sustained increase in cell proliferation is not observed; however, there are some reports of an increased proliferation at 52 weeks of treatment (Marsman et al., 1988). In the *mouse*, several reports provide data showing that DEHP causes an increase in cell proliferation early after initiation of treatment (David et al., 1999; James et al., 1998), and that such an increase may also be present at 40 weeks of exposure (Ward et al., 1988). However, no conclusive dose- and/or time-response study was conducted with DEHP in mice to elucidate the shape of the dose-/time-response curve of DEHP on cell proliferation in the liver.

There is no *in vivo* evidence for cell proliferation or hyperplasia (i.e., liver enlargement) in response to DEHP in **human, or non-human primate liver**. The only available studies that evaluated cell proliferation (number of proliferating cell, or liver size), and/or markers of increased cell cycle in the liver were performed with other PPs. One report showed that clinical hepatomegaly did not occur in 12 human patients treated with fenofibrate for 4–86 months (Gariot et al., 1987). In studies on non-human primates, administration of clofibrate, ciprofibrate, or fenofibrate for either 2 or 13 weeks had no significant effect on relative liver weight with doses at therapeutic levels (Hoivik et al., 2004). Dose of these drugs that exceeded therapeutic by 4- to 9-fold did cause a significant increase in liver weight. This was, however, attributed to hypertrophy of hepatocytes due to induction of peroxisomes and mitochondria, but not increased proliferation since no evidence of increased mitotic figures, Ki-67-positive cells, or mRNA markers of enhanced cell proliferation (e.g., proliferating cell nuclear antigen, cyclin D, c-myc, etc.) was detected in liver (Hoivik et al., 2004). In another study in which the effects of di-isononyl phthalate (DINP) and DEHP were evaluated in young adult male cynomolgus monkeys after 14 days of treatment, there were no statistically significant changes in total hepatic DNA synthesis or liver weights (Pugh, Jr. et al., 2000). It should be noted, however, that the time points that were considered by these studies may not have been the most appropriate since in rodents the rapid burst of proliferation in liver subsides by 7–14 days of treatment (Mitchell et al., 1985).

In **cultured rat hepatocytes**, MEHP shows a modest (up to 2-fold) yet significant effect on replicative DNA synthesis (Goll et al., 1999; Hasmall et al., 1999; Roberts, 1999). It should be noted, however, that the magnitude of response is much lower than that seen in rat liver *in vivo*, and it has been hypothesized that other cells in liver (e.g., Kupffer cells) may play an important role by potentiating the proliferative response of the hepatocytes by producing mitogenic cytokines (see below). Studies with **cultured human hepatocytes** failed to produce any evidence of increased proliferation even though they were generally conducted in parallel with rat hepatocyte assays and at comparable doses (Goll et al., 1999; Hasmall et al., 1999; Roberts, 1999). It should be noted that a limited number of humans were sampled, which may have affected the significance levels of these studies due to presumably higher genetic variability in human population. In addition, an unknown condition of the livers used for cell isolations, as well as duration and condition of storage of human cells before the experiments, may also be considered as confounding variables.

Induction of cell proliferation by PPs, including DEHP, is absolutely dependent on the presence of the functional PPAR $\alpha$  in hepatocytes. Feeding of the carcinogenic doses of DEHP or WY-14,643 failed to induce cell proliferation or liver enlargement in PPAR $\alpha$ -null mice (Peters et al., 1997; Ward et al., 1998). A similar loss of proliferative response is seen in cultured hepatocytes from PPAR $\alpha$ -null mice (Hasmall

et al., 2000a). Interestingly, a recent study that used hepatocyte transplantation to generate chimeric livers composed of PPAR $\alpha$ -null and -positive hepatocytes in PPAR $\alpha$ -null or -positive mice examined a relationship between PPAR $\alpha$  status and hepatocytes' ability to proliferate in response to PPs *in vivo* (Weglarz and Sandgren, 2004). When treated with WY-14,643 for 7 days, both PPAR $\alpha$ -null and -positive hepatocytes in chimeric livers displayed elevated DNA synthesis regardless of host receptor status, as long as at least some hepatocytes contained the receptor. These findings suggest that the mitogenic response to peroxisome proliferators does not require the presence of active PPAR $\alpha$  in all hepatocytes. Most recently, it was shown that PPAR $\alpha$ -humanized mice exhibited neither hepatocellular proliferation nor hepatomegaly when fed WY-14,643 or another PP, fenofibrate (Cheung et al., 2004).

Collectively, cell proliferation appears to be the PP-induced phenotype that best correlates with the carcinogenic potential of PPs in rats and mice (Marsman et al., 1988). Induction of cell cycle appears to be linked, at least partially, to PPAR $\alpha$ -mediated signaling in the hepatocyte; however, the exact mechanism of such interaction is not presently known. Several hypotheses suggesting a link via mitogen-activated kinases, such as p38 (Roberts et al., 2001), and/or Ras-mediated signaling (Wheeler et al., 2003), have been recently brought forward. These leads, while intriguing, require further validation, and it is not clear whether any of these mechanisms are relevant for the mode of action of DEHP. Furthermore, it is plausible that other, non-PPAR $\alpha$ -mediated events in hepatocytes and/or other cell types in liver may be important for induction of cell proliferation in rodent liver (see below).

### Suppression of apoptosis

Apoptosis is a physiological process that is highly conserved due to its importance in maintaining correct cell number in various tissues and removal of damaged or unwanted cells. In adult animals, apoptosis acts as a safeguard to prevent cells with damaged DNA from progressing to a tumor. Several studies have shown that agonists of PPAR $\alpha$ , including DEHP, suppress rates of apoptosis in **rat and mouse liver *in vivo*** (Bursch et al., 1984; Gerbracht et al., 1990; Hasmall et al., 2000b). Thus, it was hypothesized that the cells that would normally be removed by apoptosis may then persist for further mitogenic stimulation by non-genotoxic carcinogens, giving rise to tumors. Interestingly, one study that quantified the age-dependent effect of PPAR $\alpha$  agonists on liver apoptosis revealed that only livers of middle age and senescent rats were sensitive, while no effect was observed in young animals (Youssef et al., 2003). There are no known reports on the anti-apoptotic effects of DEHP or other PPs in **livers of humans or non-human primates**.

In **cultured rodent hepatocytes**, MEHP and other PPs suppress both spontaneous hepatocyte apoptosis and that induced by the physiological negative regulator of liver growth, transforming growth factor (TGF) $\beta$ 1, DNA-damaging drugs etoposide or hydroxyurea, or anti-Fas antibody (Gill et al.,

1998; Goll et al., 1999; Hasmall et al., 1999). Furthermore, viability assays from isolated cultured hepatocyte have shown that cells from PP-treated rats are viable in standard culture for at least 4 weeks, as compared to ~8 days for cells from control animals, and signs of apoptosis (i.e., condensed or fragmented nuclei) were less frequently observed in PP-treated cultures (Roberts et al., 1995). It appears that PPAR $\alpha$  is required to mediate inhibition of apoptosis since hepatocytes from PPAR $\alpha$ -null mice are refractory to the effects of PPs (nafenopin) on spontaneous, or TGF $\beta$ 1-induced apoptosis (Hasmall et al., 2001; Hasmall et al., 2000a). The inhibition of apoptosis by PPAR $\alpha$ -mediated signaling can be eliminated by introducing a negative effector regulator of this nuclear receptor (Roberts et al., 1998). Conversely, the anti-apoptotic effects of PPs are also dependent on non-parenchymal cells (NPCs, most likely Kupffer cells, see below). Highly purified hepatocytes show no response, but the phenotype is rescued by co-culturing with NPCs (Hasmall et al., 2001; Hasmall et al., 2000c). Interestingly, the PPAR $\alpha$  status of NPCs is irrelevant for this effect since PPAR $\alpha$ -wild type and -null NPCs produced a similar result (Hasmall et al., 2001), and Kupffer cells do not express PPAR $\alpha$  (Peters et al., 2000). Collectively, whereas it is clear that DEHP and other PPAR $\alpha$  agonists lead to inhibition of apoptosis, the specific target genes mediating these events remain unidentified.

Several published reports provide conclusive evidence that *cultured human hepatocytes* are non-responsive to anti-apoptotic effects of MEHP (Hasmall et al., 1999) and other PPs (Goll et al., 1999). The lack of an effect is true for both spontaneous and cytokine-induced (TGF $\beta$ 1, or TNF $\alpha$ /amanitine) apoptosis. It should be noted, however, that human hepatocyte cultures used in these experiments may have contained up to 5% of non-parenchymal cells (Goll et al., 1999).

### Production of Reactive Oxygen Species

Reactive oxygen species are thought to be intimately associated with the mechanism of tumorigenesis by PPs, including DEHP. This assumption is based to a large degree on a fact that various proteins that are induced by these chemicals in liver parenchymal cells (peroxisomes, mitochondria, and microsomes, see above) are prone to formation of hydrogen peroxide and other oxidants. Indeed, it was hypothesized that such overproduction of oxidants might cause DNA damage and lead to mutations and cancer (Reddy and Rao, 1989; Yeldandi et al., 2000). In addition, recent discoveries show that reactive oxygen species play an important signaling role in a rapid increase of parenchymal cell proliferation caused by peroxisome proliferators (Rusyn et al., 2000b). Collectively, it appears that oxidant-related molecular events could interact with other pathways activated by peroxisome proliferators in vivo in rodent liver, and thus it is critically important to understand their precise mechanism of action.

In *rat and mouse liver in vivo* and in *isolated cultured hepatocytes*, peroxisome proliferators induce disproportionate increases in the activity of enzymes that generate and degrade

hydrogen peroxide, oxidize DNA, lipids, and other molecules. Reddy et al. originally proposed that fatty acyl-CoA oxidase in the peroxisome is the enzyme responsible for oxidative stress by peroxisome proliferators (Nemali et al., 1988; Reddy et al., 1986); however, mice lacking this protein, instead of being protected from chemically induced liver cancer, develop liver tumors spontaneously, possibly as a result of a hyperactivation of PPAR $\alpha$  by unmetabolized lipids (Fan et al., 1998). A number of indirect confirmations for DEHP-initiated increases in oxidants have been collected over the past two decades (Conway et al., 1989; Eagon et al., 1999; Elliott et al., 1986; Lake et al., 1987; Takagi et al., 1990; Tomaszewski et al., 1986). It should be noted, however, that causative relevance of some of such evidence to the carcinogenic effect of PP in general, and DEHP in particular, has been questioned, and contrasting views have been presented (Cattley and Glover, 1993; Handler et al., 1992; Hegi et al., 1990; Sausen et al., 1995; Soliman et al., 1997). It has been shown that PPAR $\alpha$  is required for generation of oxidants in mouse liver after prolonged [weeks, (Rusyn et al., 2004)] but not acute [hours, (Rusyn et al., 2001)] exposure to PPs. In addition, Kupffer cells have been suggested to be a potential source of oxidants in rat and mouse liver after treatment with DEHP or other PPs [see below, (Rose et al., 2000; Rusyn et al., 2001)]. Collectively, despite the fact that few question a role of oxidative stress in the mechanism of action of these compounds, precise molecular source(s) of reactive oxygen species that are activated by peroxisome proliferators are still a subject of intense debate.

There are no known reports on the ability of DEHP or other PPs to induce production of reactive oxygen species in *livers of humans or non-human primates*, or *in cultured liver cells* from these species.

### Oxidative DNA Damage

Oxidative stress to DNA has been hypothesized to be a common pathway for many non-genotoxic chemical carcinogens (Klaunig et al., 1998). An overwhelming number of studies draw a connection between chemical exposure, DNA damage, and cancer based on detection of 8-hydroxy-2'-deoxyguanosine (8-OHdG), a highly mutagenic lesion, in DNA isolated from organs of in vivo treated animals [reviewed in (Kasai, 1997)]. Kasai et al. (Kasai et al., 1989) showed that treatment with ciprofibrate, a potent peroxisome proliferator, leads to a two-fold increase in the number of 8-OHdG adducts *in vivo in rat liver* genomic DNA. Similar observations were made for DEHP (Qu et al., 2001; Takagi et al., 1990). In addition to 8-OHdG, increases in other DNA alterations that were detectable by <sup>32</sup>P-postlabeling assay have been reported after chronic administration of PPs clofibrate and WY-14,643 (Randerath et al., 1991). Despite the ever-growing number of published reports that describe high levels of oxidative stress-induced DNA adducts in livers of rats and mice treated with peroxisome proliferators, a concern exists as to whether increases in 8-OHdG represent damage to genomic DNA, a confounding contamination with mitochondrial DNA, or an experimental

artifact (Elliott and Elcombe, 1987; Rusyn et al., 2004; Sausen et al., 1995; Soliman et al., 1997). Recent reports indicate that earlier estimates of ubiquitous persistence of 8-OHdG in genomic DNA were greatly (up to 1000-fold) exaggerated, most likely due to significant artifactual oxidation of guanine bases during isolation of DNA ((ESCODD) European Standards Committee on Oxidative DNA Damage, 2002). These recent developments cause a serious concern as to whether previous studies on oxidative DNA adducts, especially 8-OHdG, should be reconsidered.

One possible explanation to an apparent lack of convincing increases in oxidant-modified DNA bases despite the fact that conclusive evidence of increased oxidant production after treatment with PPs has been obtained in a variety of studies including direct *in vivo* measurements (Rusyn et al., 2001), is that oxidative stress to DNA in a living animal is a chronic event that is much weaker in its magnitude than treatment of cells *in vitro* with a concentrated solution of hydrogen peroxide. Indeed, it was shown that chronic treatment with PPs, including DEHP, elicits an adaptive response such as increased DNA repair in both rat and mouse liver (Rusyn et al., 2000a). The degree of induction of DNA repair gene expression correlated with a dose and a carcinogenic potency of the PP compounds. Furthermore, it was shown that oxidative DNA damage is a PPAR $\alpha$ -dependent event (Rusyn et al., 2004).

In view of possible induction of DNA damage by PPs, compensatory induction of DNA repair may be an important factor to consider in the mechanism of action of these compounds. First, increased repair may mask actual levels of oxidative DNA damage by removing the bulk of the lesions that are introduced by a chemical treatment. Second, in the process of DNA repair by a base-excision mechanism, additional mutagenic lesions, such as abasic sites and single strand breaks, are introduced and if the induction of enzymes that participate at different steps of the repair is uneven, there is potential for accumulation of mutagenic intermediates (Posnick and Samson, 1999). Third, if DNA polymerase  $\beta$ , a repair-specific enzyme that has low fidelity for DNA replication, is present in significant amounts in the nucleus of a proliferating cell, mutations can be introduced by erroneous base pairing by this error prone protein (Canitrot et al., 1998). Collectively, oxidative stress to DNA is most likely involved in the mechanism of carcinogenesis by PPs. It appears that gene expression for oxidative stress-related DNA repair proteins, but not levels of DNA adducts, is a reliable biomarker that can be used as proof of a carcinogenic potential of PPs.

There are no published reports on the effects of PPs on oxidative stress to DNA in *cultured rodent, human, or non-human primate hepatocytes*.

### **Inhibition of Gap Junctional Intercellular Communication**

Gap junctional intercellular communication can be modulated during the cancer process, and it decreases progressively during multi-stage rat hepatocarcinogenesis [reviewed

(Klaunig and Kamendulis, 2004)]. Blockage of cell-to-cell communication has particularly been associated with the tumor promotion stage of chemical carcinogenesis and has been suggested as a mechanism of action for the tumor promotion process (Klaunig and Ruch, 1990). Additionally, the inhibition of gap junctional intercellular communication observed following treatment with tumor promoting compounds correlates with species and strain sensitivity of the chemical (Klaunig and Ruch, 1990).

All of the available data on this mode of action of PPs were generated by J.E. Klaunig and co-workers. DEHP, DINP, and other PPs were shown to inhibit gap junctional intercellular communication in the *liver of rats or mice* (Smith et al., 2000). This inhibition persists while DEHP treatment continues, but reverses when treatment is stopped (Isenberg et al., 2001). It also correlates with the dose- and species (rat, mouse and hamster)-specific tumorigenicity of DEHP (Isenberg et al., 2000). MEHP has also been shown to inhibit gap junctional intercellular communication in *cultured rat and mouse hepatocytes* in a concentration-dependent manner (Kamendulis et al., 2002; Klaunig et al., 1988). However, it is not known whether this effect is dependent on PPAR $\alpha$ . Inhibition of gap junctional intercellular communication in rodent cells was substantially reversed within 24 hrs of chemical removal (Kamendulis et al., 2002). In contrast, cell-to-cell communication is not inhibited by MEHP in *cultured hamster, cynomolgus, or human hepatocytes* (Kamendulis et al., 2002), or *in vivo in non-human primate* studies (Pugh, Jr. et al., 2000).

### **Clonal Expansion of Previously Initiated Cells**

Induction of cell proliferation and reduction of apoptosis that are caused by PPs in rodent liver (see above) likely play a central role in further genetic alterations in hepatocytes that are initiated either by PP-induced oxidative stress, or due to other endogenous factors (e.g., replication errors, normal oxidative metabolism, etc). DEHP and other PPs are commonly classified as tumor promoters based on their ability to promote clonal expansion of the initiated cells both *in vivo* and *in vitro* in rodent liver. PPs selectively stimulate growth of cells within PP-induced basophilic foci and adenomas in *rat liver* (Grasl-Kraupp et al., 1993; Oesterle and Deml, 1988). DEHP and other PPs promote hepatocellular neoplastic growth induced by genotoxic compounds in mouse and rat liver (Cattley and Popp, 1989; Ward et al., 1983; Ward et al., 1986). Furthermore, progression from initiated cell to hepatic carcinomas is dependent on the continued presence of the PPs (Grasl-Kraupp et al., 1997). Since initiation is considered to occur spontaneously and irreversibly, old animals may have a greater accumulation of spontaneously initiated hepatocytes than young ones. Indeed, when the carcinogenicity of the PP compound WY-14,643 was compared in young (starting age 2 months) and old (starting age 15 months) rats, old rats had a 5- to 7-fold higher yield of grossly visible hepatic tumors following 22 weeks of dietary WY-14,643 when compared to young rats (Cattley et al., 1991).

Potent PPs, such as nafenopin and WY-14,643, can promote the clonal expansion of anchorage-independent *rat and mouse hepatocytes* (James and Roberts, 1994; James and Roberts, 1995) and other cells (Ward et al., 1986) *in vitro*, which suggests that these compounds perturb the balance of mitosis and apoptosis, leading to the net outgrowth of initiated clones. In contrast, hamster, guinea pig, and *human hepatocytes* do not respond to mitogenic effects of nafenopin (James and Roberts, 1995); however, no studies have been performed with MEHP or other metabolites of DEHP *in vitro* in animal or human hepatocytes. Furthermore, potential confounding variables (e.g., condition of human livers used for the experiments, etc.) in the published report that used human cells have not been adequately considered. It is not clear whether this mode of action is PPAR $\alpha$ - and/or Kupffer cell- (see below) dependent as there are no published reports that examined the relationship between receptor status and promotional activity of PPAR $\alpha$  agonists.

#### PROPOSED MODES OF ACTION FOR DEHP IN KUPFFER CELLS

DEHP and other PPs increase proliferation of rodent liver parenchymal cells both *in vivo* and *in vitro*; however, the effect on isolated hepatocytes from rats and mice is much less robust and persistent (i.e., 8- to 10-fold increases *in vivo* versus only up to 2-fold increases *in vitro*), regardless of the dose of the compound used (Goll et al., 1999; Hasmall et al., 1999; Marsman et al., 1988). Most interestingly, in highly purified rodent hepatocytes, PPs fail to further increase DNA synthesis (Hasmall et al., 2001; Hasmall et al., 2000c; Parzefall et al., 2001). The possible explanation of this difference is the involvement of nonparenchymal cell(s) in whole liver that is lost in isolated cultured hepatocytes. It was hypothesized that such cell-cell interaction occurs via PP-induced activation of resident hepatic macrophages, Kupffer cells that are the predominant source of mitogens and co-mitogens in liver [reviewed in (Rose et al., 1999b)]. Several laboratories in the past decade have demonstrated that Kupffer cells are involved in a number of acute PP-mediated pleiotropic responses in rodent liver. It was shown that Kupffer cell activation by PPs (i) is independent of PPAR $\alpha$ , (ii) involves generation of reactive oxygen species, and (iii) leads to production of mitogenic cytokines [reviewed in (Rusyn et al., 2000b)]. It is not presently known whether Kupffer cell-specific events play a role in long-term effects of DEHP and other PPs, and if these mechanisms are operational in species other than rats and mice. Finally, it should be noted that apart from one *in vitro* study that used MEHP (Rose et al., 1999a), and two *in vivo* studies that used DEHP (Currie et al., 2005; Rusyn et al., 2001), all other reports cited below in support of this mode of action have used other PPs.

#### Activation of Kupffer Cells is Independent of PPAR $\alpha$

Tumor necrosis factor (TNF) $\alpha$  is a cytokine that is one of the master regulators of liver growth and regeneration, and it is produced by Kupffer cells in liver (Diehl, 2000). The first

evidence of activation of Kupffer cells by PPs was shown in isolated perfused rat liver. It was demonstrated that nafenopin and WY-14,643 doubled uptake of particulate colloidal carbon, reflecting activation of Kupffer cells (Bojes and Thurman, 1996). Then, it was reported that neutralizing antibody to TNF $\alpha$  blocked WY-14,643-induced cell proliferation in the *rat liver in vivo*, and that TNF $\alpha$  was detected immunohistochemically exclusively in Kupffer cells following the treatment (Bojes et al., 1997). It was also shown that TNF $\alpha$  stimulates proliferation and decreases apoptosis in cultured rat hepatocytes, mimicking the effects of PPs with the exception of induction of peroxisomes (Rolfe et al., 1997). Indeed, studies in the rat demonstrated increases in whole-liver TNF $\alpha$  mRNA (Rose et al., 1997b) and serum TNF $\alpha$  protein (Adinehzadeh and Reo, 1998) levels following treatment with PPs. It should be noted, however, that one study that reports an extensive time-course study with WY-14,643 in mice (from 2 to 72 hrs) found no induction of TNF $\alpha$ , or other cytokine mRNAs in mouse liver (Anderson et al., 2001).

Inactivation of Kupffer cells in rat liver *in vivo* by methyl palmitate or glycine prevented the increases in both TNF $\alpha$  mRNA and protein, and cell proliferation due to the peroxisome proliferator WY-14,643 (Rose et al., 1997a; Rose et al., 1997b). It appears that activation of Kupffer cells in whole liver occurs very rapidly. In rat liver, activity of the transcription factor NF- $\kappa$ B, a major regulator of TNF $\alpha$  transcription, is transiently increased in 2–8 hrs followed by a steady decline to near control levels at 36 hrs following treatment with WY-14,643 (Rusyn et al., 1998). As early as 2 hrs after WY-14,643 treatment, the active form of NF- $\kappa$ B was localized predominantly in Kupffer cells with values 20 to 25 times greater than in hepatocytes where a small increase in NF- $\kappa$ B binding was observed, but only 8 hrs later. A study that evaluated the ability of DEHP to activate production of reactive oxygen species in rat and mouse liver also concluded that DEHP activates Kupffer cells within hours of treatment and before induction of peroxisomal enzymes (Rusyn et al., 2001). Finally, recent gene expression profiling study examined transcriptional changes induced by DEHP in mouse liver (Currie et al., 2005). In addition to many genes that have been traditionally associated with hepatocyte-specific responses to PPs, a number of known components of the TNF/IL-1 signaling pathways, including *Irak2*, *Myd88*, *Ikkkg*, and others were induced very early (2 hrs) and declined at later times (24 hrs) after acute treatment with DEHP, consistent with other studies showing a time-course of Kupffer cell activation.

Studies with cultured rodent hepatocytes, Kupffer cells, or mixed cell populations strengthened the evidence for the role of Kupffer cells. Both WY-14,643 and MEHP directly activate *rat Kupffer cells in vitro* to produce oxidants (Rose et al., 1999a). Furthermore, it was shown that LY-171883, a peroxisome proliferator, was taken up by Kupffer cells via phagocytosis, leading to their activation [as reported in (Rose et al., 1999b)]. Addition of non-parenchymal cells, or

conditioned medium from non-parenchymal cell cultures, increased DNA synthesis 2–3 fold and suppressed TGF $\beta$ 1-induced apoptosis by 50–70% in highly purified rat hepatocytes (Hasmall et al., 2000c). Furthermore, in both rats and mice, removal of non-parenchymal cells from normal hepatocyte cultures prevented both the nafenopin- and TNF $\alpha$ -induced increases in DNA synthesis and suppression of hepatocyte apoptosis; this response was restored by returning non-parenchymal cells to the purified hepatocytes (Hasmall et al., 2001; Hasmall et al., 2000c).

It appears that Kupffer cell activation by PPs in PPAR $\alpha$ -independent. In rat Kupffer cells, from naïve or WY-14,643-treated rats, neither PPAR $\alpha$  mRNA, nor protein could be detected (Peters et al., 2000). Moreover, when Kupffer cells from wild-type or PPAR $\alpha$ -null mice were treated with WY-14,643 in vitro, superoxide production was similar (Peters et al., 2000). In addition, non-parenchymal cells isolated from PPAR $\alpha$ -null mice, like those isolated from the wild-type mice, restored the proliferative hepatocyte response to nafenopin that is lost in highly purified liver parenchymal cells (Hasmall et al., 2001). However, as expected, PPAR $\alpha$ -null hepatocytes remained non-responsive to PPs, irrespective of the genotype of the added non-parenchymal cells. Finally, one study examined the potentiation of the effect of TNF $\alpha$  and a peroxisome proliferator WY-14,643 on cell proliferation in purified cultured rat hepatocytes (Parzefall et al., 2001). It was concluded that an increase in mitogenic cytokine production by Kupffer cells is necessary for stimulation of DNA synthesis in purified rat parenchymal cells since the presence of both the PP and TNF $\alpha$  is required for a maximal proliferative response similar to that in whole liver.

### Kupffer Cell-Mediated Production of Reactive Oxygen Species

Stimulation of Kupffer cells results in activation of NADPH oxidase and production of superoxide anion (Decker, 1990). Indeed, both WY-14,643 and MEHP are able to increase superoxide production by isolated rat Kupffer cells in a dose-dependent manner, indicating that they can affect Kupffer cells directly (Rose et al., 1999a). This effect of WY-14,643 was from activation of NADPH oxidase, since Kupffer cells isolated from *p47<sup>phox</sup>*-null mice, that lack the ability to activate this oxidant-generating protein, show no increase in superoxide production (Rusyn et al., 2000c). In addition, direct evidence for increased oxidant production in vivo after treatment with DEHP was obtained using a spin-trapping technique and electron spin resonance (ESR) spectroscopy (Rusyn et al., 2001). Specifically, when rats were given DEHP acutely for 2 hrs, an ESR-detectable radical adduct signal was detected. No increase in the radical signal due to DEHP was observed when Kupffer cells were inactivated in vivo with glycine pre-treatment, or in *p47<sup>phox</sup>*-null mice. The rapid DEHP-induced free radical production in vivo occurred long before H<sub>2</sub>O<sub>2</sub>-generating enzymes in peroxisomes were induced, and it was not dependent on PPAR $\alpha$  status.


It was suggested that Kupffer cell-derived oxidants may play a role in signaling increases in cell proliferation caused by PPs via a mechanism involving TNF $\alpha$  and NF- $\kappa$ B (Rose et al., 2000). Indeed, both activation of NF- $\kappa$ B and increases in cell proliferation due to a single dose of WY-14,643 were prevented completely when rats were pretreated with diphenyleneiodonium, an inhibitor of NADPH oxidase (Rusyn et al., 2000c). Furthermore, WY-14,643-induced activation of NF- $\kappa$ B, increase in TNF $\alpha$  mRNA, and acute increases in liver weight and cell proliferation did not occur in *p47<sup>phox</sup>*-null mice. Combined, these results provide strong evidence for a role of NADPH oxidase in Kupffer cells as a source of oxidants increased by treatment with PPs.

Since PPAR $\alpha$  is not involved in activation of Kupffer cells by PPs (see above), it is not clear how exactly such activation occurs upstream of the NADPH oxidase. No increase in hepatic BrDU labeling index was detected in PPAR $\alpha$ -null mice fed the WY-14,643 diet for 1 or 5 weeks, which suggests that the effects of peroxisome proliferators on replicative DNA synthesis in rodent liver are mediated by PPAR $\alpha$  but not Kupffer cells (Peters et al., 1997). Furthermore, several published reports questioned the role for TNF $\alpha$  in PP-induced cell proliferation in rodent liver, and the possibility of involvement of the Kupffer cell-specific events in PP-induced carcinogenesis is uncertain (see below).

### Lack of Data on a Sustained Role of the Kupffer Cell-Derived Cytokines in PP-Induced Rodent Liver Carcinogenesis

There remains uncertainty in establishing a causal link between Kupffer cell-mediated production of cytokines and long-term effects of PPs as there is no published report that examined such a possible connection. The only published report that suggests that Kupffer cell activation may persist for longer than a few days compared the effects of WY-14,643 and DEHP that both elevate hepatocyte replication during the first few days of treatment; however, only WY-14,643 sustains rates of proliferation with long-term treatment [as reported in (Marsman et al., 1988)]. It was found that Kupffer cells isolated from rats fed WY-14,643 generated superoxide at rates significantly greater than cells from controls for up to 3 weeks of treatment; however, superoxide production was not stimulated by feeding DEHP for the same period of time (Rose et al., 1999a).

Two separate studies attempted to define the role of TNF $\alpha$  signaling in hepatocellular growth induced by PPs by administering the potent peroxisome proliferator, WY-14 643 (for up to 4 days), to mice nullizygous for TNF-receptor 1 (TNFR1), TNFR2, both receptors, or TNF $\alpha$  protein (Anderson et al., 2001; Lawrence et al., 2001). Neither study found any evidence of abrogated PP-induced proliferative response in mouse liver as a result of a null genotype. It should be noted, however, that since cytokine-induced signaling is highly redundant, the knockout models used in these studies may have compensated for the lack of a particular signaling molecule. When cytokine expression



**FIG. 2. Postulated mechanisms of DEHP-mediated toxicity in liver.** (A) DEHP causes liver cancer in rodent models via activation of the nuclear receptor, PPAR $\alpha$ , which leads to changes in gene expression that cause increased cell proliferation/inhibition of apoptosis and/or oxidative stress. DNA mutations that may or may not be due to the increased oxidative stress are “fixed” in cells due to the increased cell proliferation leading to liver tumors. DEHP and other PPAR $\alpha$  ligands can also activate Kupffer cells and influence cell proliferation, independent of PPAR $\alpha$ . (B) In addition to this well-documented mechanism of action, DEHP could also cause PPAR $\alpha$ -independent changes including activation of intracellular kinases, enzyme inhibition, or interactions with other molecular targets that could potentially influence DEHP-dependent toxicity. However, the role, if any, of these changes is still uncertain.

was examined in non-tumorous liver tissue or adenomas in wild type mice fed with a carcinogenic dose of WY-14,643 for 52 weeks, no difference in expression of TNF $\alpha$ , IL-6, or TNFR1&2 was found (Anderson et al., 2001). However, IL-1 $\beta$  mRNA was significantly elevated, and it was postulated that cytokines other than TNF $\alpha$  may be important liver co-mitogens in PP-treated rodents.

The potential for the Kupffer cell-derived oxidants to contribute to oxidative DNA damage following exposure to PPs has also been questioned. The analysis of expression of base excision DNA repair genes was used to assess whether this sensitive *in vivo* biomarker of oxidative stress to DNA can be used to determine the source of DNA-damaging oxidants following treatment with PPs. Using PPAR $\alpha$ - and *p47<sup>phox</sup>*-null mice treated with WY-14,643 for 4 weeks, the report concluded that DNA-damaging oxidants are generated by enzymes that are induced after activation of PPAR $\alpha$ , such as those involved in lipid metabolism in peroxisomes, and are not the result of activation of NADPH oxidase in Kupffer cells (Rusyn et al., 2004).

#### PPAR $\alpha$ -INDEPENDENT MECHANISMS

It is clear from the previous discussion that many of the biological effects resulting from exposure to DEHP and its metabolites are due to PPAR $\alpha$ -mediated events. However, it is essential to point out that there are other possible mechanisms of action that may also contribute to DEHP associated toxicity. Since prolonged exposure to DEHP can cause testicular toxicity in the absence of PPAR $\alpha$  expression (Ward et al., 1983), this demonstrates that this compound can cause toxicity independent of receptor activation. Indeed, PPAR ligands can cause mitochondrial dysfunction, activate intracellular kinases signaling pathway, inhibit enzymes, and possibly interact with other receptors or unidentified intracellular targets, and there is good evidence that these effects are mediated independent of receptor activation (Peraza et al., 2005). For example, DEHP and other PPAR $\alpha$  ligands cause direct depolarization of mitochondrial membrane potential leading to inhibition of ATP synthesis (Zhou and Wallace, 1999), and MEHP is a more effective uncoupler of mitochondrial oxidative phosphorylation than is DEHP (Melnick and Schiller, 1985). While the potential

for DEHP to activate kinase signaling, inhibit enzymes, or interact with other unidentified molecular targets has not been examined to date, since other PPAR $\alpha$  ligands have been shown to have similar effects (Peraza et al., 2005), it remains possible that these pathways contribute to DEHP-dependent toxicity. However, it is also worth noting that the biological significance of many of these changes with respect to modulating toxicity is largely unknown.

## SUMMARY

DEHP is clearly carcinogenic in liver in both sexes of rats and mice. It is evident that a number of molecular signals and multiple pathways in several cell types in the liver, rather than a single molecular event such as activation of PPAR $\alpha$  and peroxisomal genes, or cell proliferation, contribute to the formation of liver tumors in susceptible rodent species. While exposure to DEHP and other phthalates is common in general population in humans, only limited DEHP-specific human data are available. The majority of experimental human data that are available in the peer-reviewed literature comes from in vitro studies in cultured human liver cells, and these studies suggest that key pathways that are responsible for liver carcinogenesis in rats and mice either do not occur in human cells, or the response is considerably weaker. The lack of adequate epidemiological studies on long-term effects of DEHP and other PPs constitutes an important gap in our knowledge on the potential for these agents to cause human liver tumors. Activation of PPAR $\alpha$  and the subsequent downstream events mediated by this transcription factor may be one central mechanism of action (Figure 2), but the average concentration of DEHP and DEHP metabolites found in humans is considerably lower than that required to transactivate PPAR $\alpha$  in vitro. Coupled with the observations that humans appears to be relatively refractory to many of the effects induced by exposure to DEHP, due to a variety of species differences, suggests that the average human population may not be at risk for PPAR $\alpha$ -mediated liver cancer. However, certain exposures (e.g., occupational or dialysis exposures) could still represent significant concern. More importantly, given the potential for DEHP and other PPAR $\alpha$  ligands to induce alterations independent of PPAR $\alpha$  activation, it remains possible that these pathways contribute significantly to toxicity (Figure 2). This is an area of research that deserves evaluation.

## ACKNOWLEDGMENTS

This work was supported, in part, by the grants from NIH (ES12686, ES11660 and ES11391) and the Intramural Research Program of the NIH, and NIEHS.

## REFERENCES

(ESCODD) European Standards Committee on Oxidative DNA Damage. (2002). Comparative analysis of baseline 8-oxo-7,8-dihydroguanine in mammalian cell DNA, by different

- methods in different laboratories: an approach to consensus. *Carcinogenesis* **23**(12):2129–2133.
- Adinehzadeh, M., and Reo, N.V. (1998). Effects of peroxisome proliferators on rat liver phospholipids: sphingomyelin degradation may be involved in hepatotoxic mechanism of perfluorodecanoic acid. *Chem. Res. Toxicol.* **11**(5):428–440.
- Albro, P.W. (1986). Absorption, metabolism, and excretion of di(2-ethylhexyl) phthalate by rats and mice. *Environ Health Perspect.* **65**:293–298.
- Albro, P.W., Corbett, J.T., Schroeder, J.L., Jordan, S., and Matthews, H.B. (1982). Pharmacokinetics, interactions with macromolecules and species differences in metabolism of DEHP. *Environ. Health Perspect.* **45**:19–25.
- Aldridge, T.C., Tugwood, J.D., and Green, S. (1995). Identification and characterization of DNA elements implicated in the regulation of CYP4A1 transcription. *Biochem. J.* **306**(Pt 2):473–479.
- Anderson, S.P., Dunn, C.S., Cattley, R.C., and Corton, J.C. (2001). Hepatocellular proliferation in response to a peroxisome proliferator does not require TNF $\alpha$  signaling. *Carcinogenesis* **22**(11):1843–1851.
- Aoyama, T., Peters, J.M., Iritani, N., Nakajima, T., Furihata, K., Hashimoto, T., and Gonzalez, F.J. (1998). Altered constitutive expression of fatty acid-metabolizing enzymes in mice lacking the peroxisome proliferator-activated receptor alpha (PPAR $\alpha$ ). *J. Biol. Chem.* **273**(10):5678–5684.
- Astill, B.D. (1989). Metabolism of DEHP: effects of prefeeding and dose variation, and comparative studies in rodents and the cynomolgus monkey (CMA studies). *Drug. Metab. Rev.* **21**(1):35–53.
- Badr, M.Z., Handler, J.A., Whittaker, M., Kauffman, F.C., and Thurman, R.G. (1990). Interactions between plasticizers and fatty acid metabolism in the perfused rat liver and in vivo. Inhibition of ketogenesis by 2-ethylhexanol. *Biochem Pharmacol.* **39**(4):715–721.
- Barber, E.D., Astill, B.D., Moran, E.J., Schneider, B.F., Gray, T.J., Lake, B.G., and Evans, J.G. (1987). Peroxisome induction studies on seven phthalate esters. *Toxicol. Ind. Health.* **3**(2):7–24.
- Berger, J.P., Akiyama, T.E., and Meinke, P.T. (2005). PPARs: therapeutic targets for metabolic disease. *Trends Pharmacol. Sci.* **26**(5):244–251.
- Bility, M.T., Thompson, J.T., McKee, R.H., David, R.M., Butala, J.H., Vanden Heuvel, J.P., and Peters, J.M. (2004). Activation of mouse and human peroxisome proliferator-activated receptors (PPARs) by phthalate monoesters. *Toxicol. Sci.* **82**(1):170–182.
- Blount, B.C., Silva, M.J., Caudill, S.P., Needham, L.L., Pirkle, J.L., Sampson, E.J., Lucier, G.W., Jackson, R.J., and Brock, J.W. (2000). Levels of seven urinary phthalate metabolites in a human reference population. *Environ Health Perspect.* **108**(10):979–982.
- Bojes, H.K., Germolec, D.R., Simeonova, P., Bruccoleri, A., Schoonhoven, R., Luster, M.I., and Thurman, R.G. (1997). Antibodies to tumor necrosis factor alpha prevent increases in cell replication in liver due to the potent peroxisome proliferator, WY-14,643. *Carcinogenesis* **18**(4):669–674.
- Bojes, H.K., and Thurman, R.G. (1996). Peroxisome proliferators activate Kupffer cells in vivo. *Cancer Res.* **56**(1):1–4.
- Bursch, W., Lauer, B., Timmermann-Trosiener, I., Barthel, G., Schuppler, J., and Schulte-Hermann, R. (1984). Controlled death (apoptosis) of normal and putative preneoplastic cells in rat liver

- following withdrawal of tumor promoters. *Carcinogenesis* **5**(4):453–458.
- Canitrot, Y., Cazaux, C., Frechet, M., Bouayadi, K., Lesca, C., Salles, B., and Hoffmann, J.S. (1998). Overexpression of DNA polymerase beta in cell results in a mutator phenotype and a decreased sensitivity to anticancer drugs. *Proc. Natl. Acad. Sci. USA*. **95**(21):12586–12590.
- Cattley, R.C., and Glover, S.E. (1993). Elevated 8-hydroxydeoxyguanosine in hepatic DNA of rats following exposure to peroxisome proliferators: Relationship to carcinogenesis and nuclear localization. *Carcinogenesis* **14**:2495–2499.
- Cattley, R.C., Marsman, D.S., and Popp, J.A. (1991). Age-related susceptibility to the carcinogenic effect of the peroxisome proliferator WY-14,643 in rat liver. *Carcinogenesis* **12**(3):469–473.
- Cattley, R.C., and Popp, J.A. (1989). Differences between the promoting activities of the peroxisome proliferator WY-14,643 and phenobarbital in rat liver. *Cancer Res.* **49**:3246–3251.
- Cheung, C., Akiyama, T.E., Ward, J.M., Nicol, C.J., Feigenbaum, L., Vinson, C., and Gonzalez, F.J. (2004). Diminished hepatocellular proliferation in mice humanized for the nuclear receptor peroxisome proliferator-activated receptor alpha. *Cancer Res.* **64**(11):3849–3854.
- Conway, J.G., Tomaszewski, K.E., Olson, M.J., Cattley, R.C., Marsman, D.S., and Popp, J.A. (1989). Relationship of oxidative damage to the hepatocarcinogenicity of the peroxisome proliferators di(2-ethylhexyl)phthalate and Wy-14,643. *Carcinogenesis* **10**(3):513–519.
- Corton, J.C., Lapinskas, P.J., and Gonzalez, F.J. (2000). Central role of PPARalpha in the mechanism of action of hepatocarcinogenic peroxisome proliferators. *Mutat. Res.* **448**(2):139–151.
- Currie, R.A., Bombail, V., Oliver, J.D., Moore, D.J., Lim, F.L., Gwilliam, V., Kimber, I., Chipman, K., Moggs, J.G., and Orphanides, G. (2005). Gene ontology mapping as an unbiased method for identifying molecular pathways and processes affected by toxicant exposure: application to acute effects caused by the rodent non-genotoxic carcinogen diethylhexylphthalate. *Toxicol. Sci.* **86**(2):453–469.
- David, R.M., Moore, M.R., Cifone, M.A., Finney, D.C., and Guest, D. (1999). Chronic peroxisome proliferation and hepatomegaly associated with the hepatocellular tumorigenesis of di(2-ethylhexyl)phthalate and the effects of recovery. *Toxicol. Sci.* **50**(2):195–205.
- Deangelo, A.B., Garrett, C.T., Manolukas, L.A., and Yario, T. (1986). Di-n-octyl phthalate (DOP), a relatively ineffective peroxisome inducing straight chain isomer of the environmental contaminant di(2-ethylhexyl)phthalate (DEHP), enhances the development of putative preneoplastic lesions in rat liver. *Toxicology* **41**(3):279–288.
- Decker, K. (1990). Biologically active products of stimulated liver macrophages (Kupffer cells). *Eur. J. Biochem.* **192**(2):245–261.
- Desvergne, B., and Wahli, W. (1999). Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr. Rev.* **20**(5):649–688.
- Diehl, A.M. (2000). Cytokine regulation of liver injury and repair. *Immunol. Rev.* **174**:160–171.
- Dirven, H.A., van den Broek, P.H., Arends, A.M., Nordkamp, H.H., de Lepper, A.J., Henderson, P.T., and Jongeneelen, F.J. (1993a). Metabolites of the plasticizer di(2-ethylhexyl)phthalate in urine samples of workers in polyvinylchloride processing industries. *Int. Arch. Occup. Environ. Health.* **64**(8):549–554.
- Dirven, H.A., van den Broek, P.H., and Jongeneelen, F.J. (1990). Effect of di(2-ethylhexyl)phthalate on enzyme activity levels in liver and serum of rats. *Toxicology.* **65**(1–2):199–207.
- Dirven, H.A., van den Broek, P.H., and Jongeneelen, F.J. (1993b). Determination of four metabolites of the plasticizer di(2-ethylhexyl)phthalate in human urine samples. *Int. Arch. Occup. Environ. Health.* **64**(8):555–560.
- Dirven, H.A., van den Broek, P.H., Peeters, M.C., Peters, J.G., Mennes, W.C., Blaauboer, B.J., Noordhoek, J., and Jongeneelen, F.J. (1993c). Effects of the peroxisome proliferator mono(2-ethylhexyl)phthalate in primary hepatocyte cultures derived from rat, guinea pig, rabbit, and monkey. *Biochem. Pharmacol.* **45**(12):2425–2434.
- Dirven, H.A., van den Broek, P.H., Peters, J.G., Noordhoek, J., and Jongeneelen, F.J. (1992). Microsomal lauric acid hydroxylase activities after treatment of rats with three classical cytochrome P450 inducers and peroxisome proliferating compounds. *Biochem. Pharmacol.* **43**(12):2621–2629.
- Duval, C., Fruchart, J.C., and Staels, B. (2004). PPAR alpha, fibrates, lipid metabolism and inflammation. *Arch. Mal. Coeur. Vais.* **97**(6):665–672.
- Eagon, P.K., Teepe, A.G., Elm, M.S., Tadic, S.D., Epley, M.J., Beiler, B.E., Shinozuka, H., and Rao, K.N. (1999). Hepatic hyperplasia and cancer in rats: alterations in copper metabolism. *Carcinogenesis* **20**(6):1091–1096.
- Eason, C.T., Powles, P., Henry, G., Spencer, A.J., Pattison, A., and Bonner, F.W. (1989). The comparative pharmacokinetics and gastric toxicity of bezafibrate and ciprofibrate in the rat. *Xenobiotica* **19**(8):913–925.
- Elliott, B.M., Dodd, N.J., and Elcombe, C.R. (1986). Increased hydroxyl radical production in liver peroxisomal fractions from rats treated with peroxisome proliferators. *Carcinogenesis* **7**(5):795–799.
- Elliott, B.M., and Elcombe, C.R. (1987). Lack of DNA damage or lipid peroxidation measured in vivo in the rat liver following treatment with peroxisomal proliferators. *Carcinogenesis* **8**(9):1213–1218.
- Fan, C.Y., Pan, J., Usuda, N., Yeldandi, A.V., Rao, M.S., and Reddy, J.K. (1998). Steatohepatitis, spontaneous peroxisome proliferation and liver tumors in mice lacking peroxisomal fatty acyl-CoA oxidase. Implications for peroxisome proliferator-activated receptor alpha natural ligand metabolism. *J. Biol. Chem.* **273**(25):15639–15645.
- Fay, M., Donohue, J.M., and De Rosa, C. (1999). ATSDR evaluation of health effects of chemicals. VI. Di(2-ethylhexyl)phthalate. Agency for Toxic Substances and Disease Registry. *Toxicol. Ind. Health.* **15**(8):651–746.
- Flavell, D.M., Pineda, T.I., Jamshidi, Y., Evans, D., Diamond, J.R., Elkeles, R.S., Bujac, S.R., Miller, G., Talmud, P.J., Staels, B., and Humphries, S.E. (2000). Variation in the PPARalpha gene is associated with altered function in vitro and plasma lipid concentrations in Type II diabetic subjects. *Diabetologia* **43**(5):673–680.
- Ganning, A.E., Brunk, U., and Dallner, G. (1984). Phthalate esters and their effect on the liver. *Hepatology* **4**(3):541–547.
- Ganning, A.E., Brunk, U., Edlund, C., Elhammer, A., and Dallner, G. (1987). Effects of prolonged administration of phthalate ester on the liver. *Environ. Health Perspect.* **73**:251–258.
- Gariot, P., Barrat, E., Drouin, P., Genton, P., Pointel, J.P., Foliguet, B., Kolopp, M., and Debry, G. (1987). Morphometric study of human hepatic cell modifications induced by fenofibrate. *Metabolism* **36**(3):203–210.

- Gerbracht, U., Bursch, W., Kraus, P., Putz, B., Reinacher, M., Timmermann-Trosiener, I., and Schulte-Hermann, R. (1990). Effects of hypolipidemic drugs nafenopin and clofibrate on phenotypic expression and cell death (apoptosis) in altered foci of rat liver. *Carcinogenesis* **11**(4):617–624.
- Gervois, P., Torra, I.P., Chinetti, G., Grotzinger, T., Dubois, G., Fruchart, J.C., Fruchart-Najib, J., Leitersdorf, E., and Staels, B. (1999). A truncated human peroxisome proliferator-activated receptor alpha splice variant with dominant negative activity. *Mol. Endocrinol.* **13**(9):1535–1549.
- Gill, J.H., James, N.H., Roberts, R.A., and Dive, C. (1998). The non-genotoxic hepatocarcinogen nafenopin suppresses rodent hepatocyte apoptosis induced by TGFbeta1, DNA damage and Fas. *Carcinogenesis* **19**(2):299–304.
- Goll, V., Alexandre, E., Viollon-Abadie, C., Nicod, L., Jaeck, D., and Richert, L. (1999). Comparison of the effects of various peroxisome proliferators on peroxisomal enzyme activities, DNA synthesis, and apoptosis in rat and human hepatocyte cultures. *Toxicol. Appl. Pharmacol.* **160**(1):21–32.
- Graham, M.J., Wilson, S.A., Winham, M.A., Spencer, A.J., Rees, J.A., Old, S.L., and Bonner, F.W. (1994). Lack of peroxisome proliferation in marmoset liver following treatment with ciprofibrate for 3 years. *Fund. Appl. Toxicol.* **22**:58–64.
- Grasl-Kraupp, B., Huber, W., Just, W., Gibson, G., and Schulte-Hermann, R. (1993). Enhancement of peroxisomal enzymes, cytochrome P-450 and DNA synthesis in putative preneoplastic foci of rat liver treated with the peroxisome proliferator nafenopin. *Carcinogenesis* **14**(5):1007–1012.
- Grasl-Kraupp, B., Ruttikay-Nedecky, B., Mullauer, L., Taper, H., Huber, W., Bursch, W., and Schulte-Hermann, R. (1997). Inherent increase of apoptosis in liver tumors: implications for carcinogenesis and tumor regression. *Hepatology* **25**(4):906–912.
- Gulick, T., Cresci, S., Caira, T., Moore, D.D., and Kelly, D.P. (1994). The peroxisome proliferator-activated receptor regulates mitochondrial fatty acid oxidative enzyme gene expression. *Proc. Natl. Acad. Sci. U S A.* **91**(23):11012–11016.
- Handler, J.A., Seed, C.B., Bradford, B.U., and Thurman, R.G. (1992). Induction of peroxisomes by treatment with perfluorooctanoate does not increase rates of H<sub>2</sub>O<sub>2</sub> production in intact liver. *Toxicol. Lett.* **60**(1):61–68.
- Hanefeld, M., Kemmer, C., and Kadner, E. (1983). Relationship between morphological changes and lipid-lowering action of p-chlorophenoxyisobutyric acid (CPIB) on hepatic mitochondria and peroxisomes in man. *Atherosclerosis* **46**(2):239–246.
- Hasmall, S., James, N., Hedley, K., Olsen, K., and Roberts, R. (2001). Mouse hepatocyte response to peroxisome proliferators: dependency on hepatic nonparenchymal cells and peroxisome proliferator activated receptor alpha (PPARalpha). *Arch. Toxicol.* **75**(6):357–361.
- Hasmall, S.C., James, N.H., Macdonald, N., Gonzalez, F.J., Peters, J.M., and Roberts, R.A. (2000a). Suppression of mouse hepatocyte apoptosis by peroxisome proliferators: role of PPARalpha and TNFalpha. *Mutat. Res.* **448**(2):193–200.
- Hasmall, S.C., James, N.H., Macdonald, N., Soames, A.R., and Roberts, R.A. (2000b). Species differences in response to diethylhexylphthalate: suppression of apoptosis, induction of DNA synthesis and peroxisome proliferator activated receptor alpha-mediated gene expression. *Arch. Toxicol.* **74**(2):85–91.
- Hasmall, S.C., James, N.H., Macdonald, N., West, D., Chevalier, S., Cosulich, S.C., and Roberts, R.A. (1999). Suppression of apoptosis and induction of DNA synthesis in vitro by the phthalate plasticizers monoethylhexylphthalate (MEHP) and diisononylphthalate (DINP): a comparison of rat and human hepatocytes in vitro. *Arch. Toxicol.* **73**(8–9):451–456.
- Hasmall, S.C., West, D.A., Olsen, K., and Roberts, R.A. (2000c). Role of hepatic non-parenchymal cells in the response of rat hepatocytes to the peroxisome proliferator nafenopin in vitro. *Carcinogenesis* **21**(12):2159–2165.
- Hays, T., Rusyn, I., Burns, A.M., Kennett, M.J., Ward, J.M., Gonzalez, F.J., and Peters, J.M. (2005). Role of peroxisome proliferator-activated receptor-alpha (PPARalpha) in bezafibrate-induced hepatocarcinogenesis and cholestasis. *Carcinogenesis* **26**(1):219–227.
- Hegi, M.E., Ulrich, D., Sagelsdorff, P., Richter, C., and Lutz, W.K. (1990). No measurable increase in thymidine glycol or 8-hydroxydeoxyguanosine in liver DNA of rats treated with nafenopin or choline-devoid low-methionine diet. *Mutat. Res.* **238**(3):325–329.
- Hertz, R., and Bar-Tana, J. (1998). Peroxisome proliferator-activated receptor (PPAR)  $\alpha$  activation and its consequences in humans. *Toxicol. Lett.* **102–103**:85–90.
- Hinton, R.H., Mitchell, F.E., Mann, A., Chescoe, D., Price, S.C., Nunn, A., Grasso, P., and Bridges, J.W. (1986). Effects of phthalic acid esters on the liver and thyroid. *Environ Health Perspect.* **70**:195–210.
- Hoivik, D.J., Qualls, Jr., C.W., Cariello, N.F., Kimbrough, C.L., Colton, H.M., Anderson, S.P., Santostefano, M.J., Morgan, R.J., Dahl, R.R., Brown, A.R., Zhao, Z., Mudd, Jr., P.N., Oliver, Jr., W.B., Brown, H.R., and Miller, R.T. (2004). Fibrates induce hepatic peroxisome and mitochondrial proliferation without overt evidence of cellular proliferation and oxidative stress in cynomolgus monkeys. *Carcinogenesis* **25**(9):1757–1769.
- Hurst, C.H., and Waxman, D.J. (2003). Activation of PPARalpha and PPARgamma by environmental phthalate monoesters. *Toxicol. Sci.* **74**(2):297–308.
- Ikeda, G.J., Sapienza, P.P., Couvillion, J.L., Farber, T.M., and van Loon, E.J. (1980). Comparative distribution, excretion and metabolism of di-(2-ethylhexyl) phthalate in rats, dogs and miniature pigs. *Food Cosmet. Toxicol.* **18**(6):637–642.
- Isenberg, J.S., Kamendulis, L.M., Ackley, D.C., Smith, J.H., Pugh, Jr., G.W., McKee, R.H., and Klaunig, J.E. (2001). Reversibility and persistence of di-2-ethylhexyl phthalate (DEHP)- and phenobarbital-induced hepatocellular changes in rodents. *Toxicol. Sci.* **64**(2):192–199.
- Isenberg, J.S., Kamendulis, L.M., Smith, J.H., Ackley, D.C., Pugh, Jr., G., Lington, A.W., and Klaunig, J.E. (2000). Effects of Di-2-ethylhexyl phthalate (DEHP) on gap-junctional intercellular communication (GJIC), DNA synthesis, and peroxisomal beta oxidation (PBOX) in rat, mouse, and hamster liver. *Toxicol. Sci.* **56**(1):73–85.
- Issemann, I., and Green, S. (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* **347**(6294):645–650.
- Ito, Y., Yokota, H., Wang, R., Yamanoshita, O., Ichihara, G., Wang, H., Kurata, Y., Takagi, K., and Nakajima, T. (2005). Species differences in the metabolism of di(2-ethylhexyl) phthalate (DEHP) in several organs of mice, rats, and marmosets. *Arch. Toxicol.* **79**(3):147–154.

- James, N.H., and Roberts, R.A. (1994). The peroxisome proliferator class of non-genotoxic hepatocarcinogens synergize with epidermal growth factor to promote clonal expansion of initiated rat hepatocytes. *Carcinogenesis* **15**(12):2687–2694.
- James, N.H., and Roberts, R.A. (1995). Species differences in the clonal expansion of hepatocytes in response to the coaction of epidermal growth factor and nafenopin, a rodent hepatocarcinogenic peroxisome proliferator. *Fundam. Appl. Toxicol.* **26**(1):143–149.
- James, N.H., Soames, A.R., and Roberts, R.A. (1998). Suppression of hepatocyte apoptosis and induction of DNA synthesis by the rat and mouse hepatocarcinogen diethylhexylphthalate (DEHP) and the mouse hepatocarcinogen 1,4-dichlorobenzene (DCB). *Arch. Toxicol.* **72**(12):784–790.
- Kamendulis, L.M., Isenberg, J.S., Smith, J.H., Pugh, Jr., G., Lington, A.W., and Klaunig, J.E. (2002). Comparative effects of phthalate monoesters on gap junctional intercellular communication and peroxisome proliferation in rodent and primate hepatocytes. *J. Toxicol. Environ. Health A.* **65**(8):569–588.
- Kasai, H. (1997). Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutat. Res.* **387**(3):147–163.
- Kasai, H., Okada, Y., Nishimura, S., Rao, M.S., and Reddy, J.K. (1989). Formation of 8-hydroxydeoxyguanosine in liver DNA of rats following long-term exposure to a peroxisome proliferator. *Cancer Res.* **49**(10):2603–2605.
- Kavlock, R., Boekelheide, K., Chapin, R., Cunningham, M., Faustman, E., Foster, P., Golub, M., Henderson, R., Hinberg, I., Little, R., Seed, J., Shea, K., Tabacova, S., Tyl, R., Williams, P., and Zacharewski, T. (2002). NTP Center for the Evaluation of Risks to Human Reproduction: phthalates expert panel report on the reproductive and developmental toxicity of di(2-ethylhexyl) phthalate. *Reprod. Toxicol.* **16**(5):529–653.
- Keller, B.J., Yamanaka, H., and Thurman, R.G. (1992). Inhibition of mitochondrial respiration and oxygen-dependent hepatotoxicity by six structurally dissimilar peroxisomal proliferating agents. *Toxicology* **71**(1–2):49–61.
- Klaunig, J.E., Babich, M.A., Baetcke, K.P., Cook, J.C., Corton, J.C., David, R.M., Deluca, J.G., Lai, D.Y., McKee, R.H., Peters, J.M., Roberts, R.A., and Fenner-Crisp, P.A. (2003). PPARalpha agonist-induced rodent tumors: modes of action and human relevance. *Crit. Rev. Toxicol.* **33**(6):655–780.
- Klaunig, J.E., and Kamendulis, L.M. (2004). The role of oxidative stress in carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.* **44**:239–267.
- Klaunig, J.E., and Ruch, R.J. (1990). Role of inhibition of intercellular communication in carcinogenesis. *Lab. Invest.* **62**(2):135–146.
- Klaunig, J.E., Ruch, R.J., Deangelo, A.B., and Kaylor, W.H. (1988). Inhibition of mouse hepatocyte intercellular communication by phthalate monoesters. *Cancer Lett.* **43**(1–2):65–71.
- Klaunig, J.E., Xu, Y., Isenberg, J.S., Bachowski, S., Kolaja, K.L., Jiang, J., Stevenson, D.E., and Walborg, Jr., E.F. (1998). The role of oxidative stress in chemical carcinogenesis. *Environ. Health Perspect.* **106**(Suppl 1):289–295.
- Koch, H.M., Drexler, H., and Angerer, J. (2003). An estimation of the daily intake of di(2-ethylhexyl)phthalate (DEHP) and other phthalates in the general population. *Int. J. Hyg. Environ. Health.* **206**(2):77–83.
- Kohli, V., Selzner, M., Madden, J.F., Bentley, R.C., and Clavien, P.A. (1999). Endothelial cell and hepatocyte deaths occur by apoptosis after ischemia-reperfusion injury in the rat liver. *Transplantation* **67**(8):1099–1105.
- Kroetz, D.L., Yook, P., Costet, P., Bianchi, P., and Pineau, T. (1998). Peroxisome proliferator-activated receptor alpha controls the hepatic CYP4A induction adaptive response to starvation and diabetes. *J. Biol. Chem.* **273**(47):31581–31589.
- Kurata, Y., Kidachi, F., Yokoyama, M., Toyota, N., Tsuchitani, M., and Katoh, M. (1998). Subchronic toxicity of Di(2-ethylhexyl)phthalate in common marmosets: lack of hepatic peroxisome proliferation, testicular atrophy, or pancreatic acinar cell hyperplasia. *Toxicol. Sci.* **42**(1):49–56.
- Lake, B.G., Gangolli, S.D., Grasso, P., and Lloyd, A.G. (1975). Studies on the hepatic effects of orally administered di-(2-ethylhexyl) phthalate in the rat. *Toxicol. Appl. Pharmacol.* **32**(2):355–367.
- Lake, B.G., Kozlen, S.L., Evans, J.G., Gray, T.J., Young, P.J., and Gangolli, S.D. (1987). Effect of prolonged administration of clofibrilic acid and di-(2-ethylhexyl)phthalate on hepatic enzyme activities and lipid peroxidation in the rat. *Toxicology* **44**(2):213–228.
- Lapinskas, P.J., Brown, S., Leesnitzer, L.M., Blanchard, S., Swanson, C., Cattley, R.C., and Corton, J.C. (2005). Role of PPARalpha in mediating the effects of phthalates and metabolites in the liver. *Toxicology* **207**(1):149–163.
- Lawrence, J.W., Wollenberg, G.K., and Deluca, J.G. (2001). Tumor necrosis factor alpha is not required for WY14,643-induced cell proliferation. *Carcinogenesis* **22**(3):381–386.
- Lee, S.S., Pineau, T., Drago, J., Lee, E.J., Owens, J.W., Kroetz, D.L., Fernandez-Salguero, P.M., Westphal, H., and Gonzalez, F.J. (1995). Targeted disruption of the alpha isoform of the peroxisome proliferator-activated receptor gene in mice results in abolishment of the pleiotropic effects of peroxisome proliferators. *Mol. Cell. Biol.* **15**(6):3012–3022.
- Maloney, E.K., and Waxman, D.J. (1999). Trans-activation of PPARalpha and PPARgamma by structurally diverse environmental chemicals. *Toxicol. Appl. Pharmacol.* **161**(2):209–218.
- Mandard, S., Muller, M., and Kersten, S. (2004). Peroxisome proliferator-activated receptor alpha target genes. *Cell. Mol. Life Sci.* **61**(4):393–416.
- Marsman, D.S., Cattley, R.C., Conway, J.G., and Popp, J.A. (1988). Relationship of hepatic peroxisome proliferation and replicative DNA synthesis to the hepatocarcinogenicity of the peroxisome proliferators di(2-ethylhexyl)phthalate and [4-chloro-6-(2,3-xylylidino)-2-pyrimidinylthio]acetic acid (Wy-14,643) in rats. *Cancer Res.* **48**(23):6739–6744.
- Melnick, R.L., and Schiller, C.M. (1985). Effect of phthalate esters on energy coupling and succinate oxidation in rat liver mitochondria. *Toxicology* **34**(1):13–27.
- Mitchell, F.E., Price, S.C., Hinton, R.H., Grasso, P., and Bridges, J.W. (1985). Time and dose-response study of the effects on rats of the plasticizer di(2-ethylhexyl) phthalate. *Toxicol. Appl. Pharmacol.* **81**(3 Pt 1):371–392.
- Morimura, K., Cheung, C., Ward, J.M., Reddy, J.K., and Gonzalez, F.J. (2006). Differential susceptibility of mice humanized for peroxisome proliferator-activated receptor {alpha} to Wy-14,643-induced liver tumorigenesis. *Carcinogenesis*. (in Press)
- Mukherjee, R., Jow, L., Noonan, D., and McDonnell, D.P. (1994). Human and rat peroxisome proliferator activated receptors (PPARs) demonstrate similar tissue distribution but different responsiveness

- to PPAR activators. *J. Steroid. Biochem. Mol. Biol.* **51**(3-4):157-166.
- Nair, N., and Kurup, C.K. (1987). Effect of administration of diethylhexyl phthalate on the function and turnover of rat hepatic mitochondria. *Biochim. Biophys. Acta.* **925**(3):332-340.
- National Toxicology Program. (1982). Carcinogenesis Bioassay of Di(2-ethylhexyl)phthalate (CAS No. 117-81-7) in F344 Rats and B6C3F1 Mice (Feed Studies). *Natl. Toxicol. Program. Tech. Rep. Ser.* **217**:1-127.
- Nemali, M.R., Usuda, N., Reddy, M.K., Oyasu, K., Hashimoto, T., Osumi, T., Rao, M.S., and Reddy, J.K. (1988). Comparison of constitutive and inducible levels of expression of peroxisomal beta-oxidation and catalase genes in liver and extrahepatic tissues of rat. *Cancer Res.* **48**(18):5316-5324.
- Oesterle, D., and Deml, E. (1988). Promoting activity of di(2-ethylhexyl)phthalate in rat liver foci bioassay. *J. Cancer Res. Clin. Oncol.* **114**(2):133-136.
- Palmer, C.N., Hsu, M.H., Griffin, K.J., Raucy, J.L., and Johnson, E.F. (1998). Peroxisome proliferator activated receptor-alpha expression in human liver. *Mol. Pharmacol.* **53**(1):14-22.
- Parzefall, W., Berger, W., Kainzbauer, E., Teufelhofer, O., Schulte-Hermann, R., and Thurman, R.G. (2001). Peroxisome proliferators do not increase DNA synthesis in purified rat hepatocytes. *Carcinogenesis* **22**(3):519-523.
- Peraza, M.A., Burdick, A.D., Marin, H.E., Gonzalez, F.J., and Peters, J.M. (2006). The Toxicology of Ligands for Peroxisome Proliferator-Activated Receptors (PPAR). *Toxicol. Sci.* **90**(2):269-295.
- Peters, J.M., Cattley, R.C., and Gonzalez, F.J. (1997). Role of PPAR alpha in the mechanism of action of the nongenotoxic carcinogen and peroxisome proliferator Wy-14,643. *Carcinogenesis* **18**(11):2029-2033.
- Peters, J.M., Cheung, C., and Gonzalez, F.J. (2005). Peroxisome proliferator-activated receptor-alpha and liver cancer: where do we stand?. *J. Mol. Med.* **83**(10):774-785.
- Peters, J.M., Rusyn, I., Rose, M.L., Gonzalez, F.J., and Thurman, R.G. (2000). Peroxisome proliferator-activated receptor alpha is restricted to hepatic parenchymal cells, not Kupffer cells: implications for the mechanism of action of peroxisome proliferators in hepatocarcinogenesis. *Carcinogenesis* **21**(4):823-826.
- Posnick, L.M., and Samson, L.D. (1999). Imbalanced base excision repair increases spontaneous mutation and alkylation sensitivity in *Escherichia coli*. *J. Bacteriol.* **181**(21):6763-6771.
- Pugh, Jr., G., Isenberg, J.S., Kamendulis, L.M., Ackley, D.C., Clare, L.J., Brown, R., Lington, A.W., Smith, J.H., and Klaunig, J.E. (2000). Effects of di-isonyl phthalate, di-2-ethylhexyl phthalate, and clofibrate in cynomolgus monkeys. *Toxicol. Sci.* **56**(1):181-188.
- Qu, B., Li, Q., Wong, K.P., Tan, T.M., and Halliwell, B. (2001). Mechanism of clofibrate hepatotoxicity: Mitochondrial damage and oxidative stress in hepatocytes. *Free Radic. Biol. Med.* **31**(5):659-669.
- Randerath, E., Randerath, K., Reddy, R., Danna, T.F., Rao, M.S., and Reddy, J.K. (1991). Induction of rat liver DNA alterations by chronic administration of peroxisome proliferators as detected by <sup>32</sup>P-postlabeling. *Mutat. Res.* **247**:65-76.
- Reddy, J.K., Azarnoff, D.L., and Hignite, C.E. (1980). Hypolipidaemic hepatic peroxisome proliferators form a novel class of chemical carcinogens. *Nature* **283**(5745):397-398.
- Reddy, J.K., and Chu, R. (1996). Peroxisome proliferator-induced pleiotropic responses: pursuit of a phenomenon. *Ann. N. Y. Acad. Sci.* **804**:176-201.
- Reddy, J.K., Goel, S.K., Nemali, M.R., Carrino, J.J., Laffler, T.G., Reddy, M.K., Sperbeck, S.J., Osumi, T., Hashimoto, T., and Lalwani, N.D. (1986). Transcription regulation of peroxisomal fatty acyl-CoA oxidase and enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase in rat liver by peroxisome proliferators. *Proc. Natl. Acad. Sci. U S A.* **83**(6):1747-1751.
- Reddy, J.K., Moody, D.E., Azarnoff, D.L., and Rao, M.S. (1976a). Di-(2-ethylhexyl)phthalate: an industrial plasticizer induces hypolipidemia and enhances hepatic catalase and carnitine acetyltransferase activities in rat and mice. *Life Sci.* **18**(9):941-945.
- Reddy, J.K., and Rao, M.S. (1989). Oxidative DNA damage caused by persistent peroxisome proliferation: its role in hepatocarcinogenesis. *Mutat. Res.* **214**(1):63-68.
- Reddy, J.K., Rao, M.S., Azarnoff, D.L., and Sell, S. (1979). Mitogenic and carcinogenic effects of a hypolipidemic peroxisome proliferator, [4-chloro-6-(2,3-xylidino)-2-pyrimidinylthio]acetic acid (Wy-14, 643), in rat and mouse liver. *Cancer Res.* **39**(1):152-161.
- Reddy, J.K., Rao, S., and Moody, D.E. (1976b). Hepatocellular carcinomas in acatalasemic mice treated with nafenopin, a hypolipidemic peroxisome proliferator. *Cancer Res.* **36**(4):1211-1217.
- Rhodes, C., Orton, T.C., Pratt, I.S., Batten, P.L., Bratt, H., Jackson, S.J., and Elcombe, C.R. (1986). Comparative pharmacokinetics and subacute toxicity of di(2-ethylhexyl) phthalate (DEHP) in rats and marmosets: extrapolation of effects in rodents to man. *Environ. Health Perspect.* **65**:299-307.
- Roberts, R.A. (1999). Peroxisome proliferators: Mechanisms of adverse effects in rodents and molecular basis for species differences. *Arch. Toxicol.* **73**(8-9):413-418.
- Roberts, R.A., James, N.H., Cosulich, S., Hasmall, S.C., and Orphanides, G. (2001). Role of cytokines in non-genotoxic hepatocarcinogenesis: cause or effect?. *Toxicol. Lett.* **120**(1-3):301-306.
- Roberts, R.A., James, N.H., Hasmall, S.C., Holden, P.R., Lambe, K., Macdonald, N., West, D., Woodyatt, N.J., and Whitcome, D. (2000). Apoptosis and proliferation in nongenotoxic carcinogenesis: species differences and role of PPARalpha. *Toxicol. Lett.* **112-113**:49-57.
- Roberts, R.A., James, N.H., Woodyatt, N.J., Macdonald, N., and Tugwood, J.D. (1998). Evidence for the suppression of apoptosis by the peroxisome proliferator activated receptor alpha. *Carcinogenesis* **19**(1):43-48.
- Roberts, R.A., Soames, A.R., Gill, J.H., James, N.H., and Wheeldon, E.B. (1995). Non-genotoxic hepatocarcinogens stimulate DNA synthesis and their withdrawal induces apoptosis, but in different hepatocyte populations. *Carcinogenesis* **16**(8):1693-1698.
- Roglans, N., Bellido, A., Rodriguez, C., Cabrero, A., Novell, F., Ros, E., Zambon, D., and Laguna, J.C. (2002). Fibrate treatment does not modify the expression of acyl coenzyme A oxidase in human liver. *Clin. Pharmacol. Ther.* **72**(6):692-701.
- Rolfé, M., James, N.H., and Roberts, R.A. (1997). Tumour necrosis factor alpha (TNF alpha) suppresses apoptosis and induces DNA synthesis in rodent hepatocytes: a mediator of the hepatocarcinogenicity of peroxisome proliferators?. *Carcinogenesis* **18**(11):2277-2280.
- Rose, M.L., Germolec, D., Arteel, G.E., Schoonhoven, R., and Thurman, R.G. (1997a). Dietary glycine prevents increases in hepatocyte proliferation caused by the peroxisome proliferator WY-14,643. *Chem. Res. Toxicol.* **10**(10):1198-1204.
- Rose, M.L., Germolec, D.R., Schoonhoven, R., and Thurman, R.G. (1997b). Kupffer cells are causally responsible for the mitogenic effect of peroxisome proliferators. *Carcinogenesis* **18**(8):1453-1456.

- Rose, M.L., Rivera, C.A., Bradford, B.U., Graves, L.M., Cattley, R.C., Schoonhoven, R., Swenberg, J.A., and Thurman, R.G. (1999a). Kupffer cell oxidant production is central to the mechanism of peroxisome proliferators. *Carcinogenesis* **20**(1):27–33.
- Rose, M.L., Rusyn, I., Bojes, H.K., Belyea, J., Cattley, R.C., and Thurman, R.G. (2000). Role of Kupffer cells and oxidants in signaling peroxisome proliferator-induced hepatocyte proliferation. *Mutat. Res.* **448**(2):179–192.
- Rose, M.L., Rusyn, I., Bojes, H.K., Germolec, D.R., Luster, M., and Thurman, R.G. (1999b). Role of Kupffer cells in peroxisome proliferator-induced hepatocyte proliferation. *Drug. Metab. Rev.* **31**(1):87–116.
- Rusyn, I., Asakura, S., Pachkowski, B., Bradford, B.U., Denissenko, M.F., Peters, J.M., Holland, S.M., Reddy, J.K., Cunningham, M.L., and Swenberg, J.A. (2004). Expression of base excision DNA repair genes is a sensitive biomarker for in vivo detection of chemical-induced chronic oxidative stress: Identification of the molecular source of radicals responsible for DNA damage by peroxisome proliferators. *Cancer Res.* **64**(3):1050–1057.
- Rusyn, I., Denissenko, M.F., Wong, V.A., Butterworth, B.E., Cunningham, M.L., Upton, P.B., Thurman, R.G., and Swenberg, J.A. (2000a). Expression of base excision repair enzymes in rat and mouse liver is induced by peroxisome proliferators and is dependent upon carcinogenic potency. *Carcinogenesis* **21**(12):2141–2145.
- Rusyn, I., Kadiiska, M.B., Dikalova, A., Kono, H., Yin, M., Tsuchiya, K., Mason, R.P., Peters, J.M., Gonzalez, F.J., Segal, B.H., Holland, S.M., and Thurman, R.G. (2001). Phthalates rapidly increase production of reactive oxygen species in vivo: role of Kupffer cells. *Mol. Pharmacol.* **59**(4):744–750.
- Rusyn, I., Rose, M.L., Bojes, H.K., and Thurman, R.G. (2000b). Novel role of oxidants in the molecular mechanism of action of peroxisome proliferators. *Antiox. Redox. Signal.* **2**:607–621.
- Rusyn, I., Tsukamoto, H., and Thurman, R.G. (1998). WY-14,643 rapidly activates nuclear factor kappaB in Kupffer cells before hepatocytes. *Carcinogenesis* **19**(7):1217–1222.
- Rusyn, I., Yamashina, S., Segal, B.H., Schoonhoven, R., Holland, S.M., Cattley, R.C., Swenberg, J.A., and Thurman, R.G. (2000c). Oxidants from nicotinamide adenine dinucleotide phosphate oxidase are involved in triggering cell proliferation in the liver due to peroxisome proliferators. *Cancer Res.* **60**(17):4798–4803.
- Sapone, A., Peters, J.M., Sakai, S., Tomita, S., Papiha, S.S., Dai, R., Friedman, F.K., and Gonzalez, F.J. (2000). The human peroxisome proliferator-activated receptor alpha gene: identification and functional characterization of two natural allelic variants. *Pharmacogenetics* **10**(4):321–333.
- Sausen, P.J., Lee, D.C., Rose, M.L., and Cattley, R.C. (1995). Elevated 8-hydroxydeoxyguanosine in hepatic DNA of rats following exposure to peroxisome proliferators: relationship to mitochondrial alterations. *Carcinogenesis* **16**(8):1795–1801.
- Schmid, P., and Schlatter, C. (1985). Excretion and metabolism of di(2-ethylhexyl)phthalate in man. *Xenobiotica* **15**(3):251–256.
- Schrader, M., and Fahimi, H.D. (2004). Mammalian peroxisomes and reactive oxygen species. *Histochem. Cell. Biol.* **122**(4):383–393.
- Seree, E., Villard, P.H., Pascucci, J.M., Pineau, T., Maurel, P., Nguyen, Q.B., Fallone, F., Martin, P.M., Champion, S., Lacarelle, B., Savouret, J.F., and Barra, Y. (2004). Evidence for a new human CYP1A1 regulation pathway involving PPAR-alpha and 2 PPRE sites. *Gastroenterology* **127**(5):1436–1445.
- Shaban, Z., El Shazly, S., Ishizuka, M., Kimura, K., Kazusaka, A., and Fujita, S. (2004). PPARalpha-dependent modulation of hepatic CYP1A by clofibrac acid in rats. *Arch. Toxicol.* **78**(9):496–507.
- Sher, T., Yi, H.F., McBride, O.W., and Gonzalez, F.J. (1993). cDNA cloning, chromosomal mapping, and functional characterization of the human peroxisome proliferator activated receptor. *Biochemistry* **32**(21):5598–5604.
- Shipley, J.M., and Waxman, D.J. (2004). Simultaneous, bidirectional inhibitory crosstalk between PPAR and STAT5b. *Toxicol. Appl. Pharmacol.* **199**(3):275–284.
- Short, R.D., Robinson, E.C., Lington, A.W., and Chin, A.E. (1987). Metabolic and peroxisome proliferation studies with di(2-ethylhexyl)phthalate in rats and monkeys. *Toxicol. Ind. Health.* **3**(2):185–195.
- Silva, M.J., Barr, D.B., Reidy, J.A., Kato, K., Malek, N.A., Hodge, C.C., Hurtz, D.III., Needham, L.L., and Brock, J.W. (2003). Glucuronidation patterns of common urinary and serum monoester phthalate metabolites. *Arch. Toxicol.* **77**(10):561–567.
- Smith, J.H., Isenberg, J.S., Pugh, Jr., G.M., Ackley, D., Lington, A.W., and Klaunig, J.E. (2000). Comparative in vivo hepatic effects of Di-isononyl phthalate (DINP) and related C7-C11 dialkyl phthalates on gap junctional intercellular communication (GJIC), peroxisomal beta-oxidation (PBOX), and DNA synthesis in rat and mouse liver. *Toxicol. Sci.* **54**(2):312–321.
- Smith-Oliver, T., and Butterworth, B.E. (1987). Correlation of the carcinogenic potential of di(2-ethylhexyl)phthalate (DEHP) with induced hyperplasia rather than with genotoxic activity. *Mutat. Res.* **188**(1):21–28.
- Soliman, M.S., Cunningham, M.L., Morrow, J.D., Roberts, L.J., and Badr, M.Z. (1997). Evidence against peroxisome proliferation-induced hepatic oxidative damage. *Biochem. Pharmacol.* **53**(9):1369–1374.
- Sumanasekera, W.K., Tien, E.S., Davis, J.W., Turpey, R., Perdew, G.H., and Vanden Heuvel, J.P. (2003a). Heat shock protein-90 (Hsp90) acts as a repressor of peroxisome proliferator-activated receptor-alpha (PPARalpha) and PPARbeta activity. *Biochemistry* **42**(36):10726–10735.
- Sumanasekera, W.K., Tien, E.S., Turpey, R., Vanden Heuvel, J.P., and Perdew, G.H. (2003b). Evidence that peroxisome proliferator-activated receptor alpha is complexed with the 90-kDa heat shock protein and the hepatitis virus B X-associated protein 2. *J. Biol. Chem.* **278**(7):4467–4473.
- Takagi, A., Sai, K., Umemura, T., Hasegawa, R., and Kurokawa, Y. (1990). Significant increase of 8-hydroxydeoxyguanosine in liver DNA of rats following short-term exposure to the peroxisome proliferators di(2-ethylhexyl)phthalate and di(2-ethylhexyl)adipate. *Jpn. J. Cancer. Res.* **81**(3):213–215.
- Tanaka, A., Adachi, T., Takahashi, T., and Yamaha, T. (1975). Biochemical studies on phthalic esters I. Elimination, distribution and metabolism of di-(2-ethylhexyl)phthalate in rats. *Toxicology* **4**(2):253–264.
- Tomaszewski, K.E., Agarwal, D.K., and Melnick, R.L. (1986). In vitro steady-state levels of hydrogen peroxide after exposure of male F344 rats and female B6C3F1 mice to hepatic peroxisome proliferators. *Carcinogenesis* **7**(11):1871–1876.
- (1995). Comparative toxicology of hypolipidaemic fibrates, Taylor & Francis, London.

- Vohl, M.C., Lepage, P., Gaudet, D., Brewer, C.G., Betard, C., Perron, P., Houde, G., Cellier, C., Faith, J.M., Despres, J.P., Morgan, K., and Hudson, T.J. (2000). Molecular scanning of the human PPAR $\alpha$  gene: association of the L162v mutation with hyperapobetalipoproteinemia. *J. Lipid. Res.* **41**(6):945–952.
- Voskoboinik, I., Ooi, S.G., Drew, R., and Ahokas, J.T. (1997). Peroxisome proliferators increase the formation of BPDE-DNA adducts in isolated rat hepatocytes. *Toxicology* **122**(1–2):81–91.
- Vu-Dac, N., Chopin-Delannoy, S., Gervois, P., Bonnelye, E., Martin, G., Fruchart, J.C., Laudet, V., and Staels, B. (1998). The nuclear receptors peroxisome proliferator-activated receptor alpha and Rev-erbalpha mediate the species-specific regulation of apolipoprotein A-I expression by fibrates. *J. Biol. Chem.* **273**(40):25713–25720.
- Vu-Dac, N., Gervois, P., Jakel, H., Nowak, M., Bauge, E., Dehondt, H., Staels, B., Pennacchio, L.A., Rubin, E.M., Fruchart-Najib, J., and Fruchart, J.C. (2003). Apolipoprotein A5, a crucial determinant of plasma triglyceride levels, is highly responsive to peroxisome proliferator-activated receptor alpha activators. *J. Biol. Chem.* **278**(20):17982–17985.
- Walgren, J.E., Kurtz, D.T., and McMillan, J.M. (2000). Expression of PPAR(alpha) in human hepatocytes and activation by trichloroacetate and dichloroacetate. *Res. Commun. Mol. Pathol. Pharmacol.* **108**(1–2):116–132.
- Ward, J.M., Diwan, B.A., Ohshima, M., Hu, H., Schuller, H.M., and Rice, J.M. (1986). Tumor-initiating and promoting activities of di(2-ethylhexyl) phthalate in vivo and in vitro. *Environ. Health Perspect.* **65**:279–291.
- Ward, J.M., Hagiwara, A., Anderson, L.M., Lindsey, K., and Diwan, B.A. (1988). The chronic hepatic or renal toxicity of di(2-ethylhexyl) phthalate, acetaminophen, sodium barbital, and phenobarbital in male B6C3F1 mice: autoradiographic, immunohistochemical, and biochemical evidence for levels of DNA synthesis not associated with carcinogenesis or tumor promotion. *Toxicol. Appl. Pharmacol.* **96**(3):494–506.
- Ward, J.M., Peters, J.M., Perella, C.M., and Gonzalez, F.J. (1998). Receptor and nonreceptor-mediated organ-specific toxicity of di(2-ethylhexyl)phthalate (DEHP) in peroxisome proliferator-activated receptor alpha-null mice. *Toxicol. Pathol.* **26**(2):240–246.
- Ward, J.M., Rice, J.M., Creasia, D., Lynch, P., and Riggs, C. (1983). Dissimilar patterns of promotion by di(2-ethylhexyl)phthalate and phenobarbital of hepatocellular neoplasia initiated by diethylnitrosamine in B6C3F1 mice. *Carcinogenesis* **4**(8):1021–1029.
- Weglarz, T.C., and Sandgren, E.P. (2004). Cell cross-talk mediates PPARalpha null hepatocyte proliferation after peroxisome proliferator exposure. *Carcinogenesis* **25**(1):107–112.
- Wheeler, M.D., Smutney, O.M., Check, J.F., Rusyn, I., Schulte-Hermann, R., and Thurman, R.G. (2003). Impaired Ras membrane association and activation in PPARalpha knockout mice after partial hepatectomy. *Am. J. Physiol. Gastrointest. Liver Physiol.* **284**(2):G302–G312.
- Winberg, L.D., and Badr, M.Z. (1995). Mechanism of phthalate-induced inhibition of hepatic mitochondrial beta-oxidation. *Toxicol. Lett.* **76**(1):63–69.
- Woodyatt, N.J., Lambe, K.G., Myers, K.A., Tugwood, J.D., and Roberts, R.A. (1999). The peroxisome proliferator (PP) response element upstream of the human acyl CoA oxidase gene is inactive among a sample human population: significance for species differences in response to PPs. *Carcinogenesis* **20**(3):369–372.
- Yeldandi, A.V., Rao, M.S., and Reddy, J.K. (2000). Hydrogen peroxide generation in peroxisome proliferator-induced oncogenesis. *Mutat. Res.* **448**(2):159–177.
- Youssef, J.A., Bouziane, M., and Badr, M.Z. (2003). Age-dependent effects of nongenotoxic hepatocarcinogens on liver apoptosis in vivo. *Mech. Aging Dev.* **124**(3):333–340.
- Yu, S., Cao, W.Q., Kashireddy, P., Meyer, K., Jia, Y., Hughes, D.E., Tan, Y., Feng, J., Yeldandi, A.V., Rao, M.S., Costa, R.H., Gonzalez, F.J., and Reddy, J.K. (2001). Human peroxisome proliferator-activated receptor alpha (PPARalpha) supports the induction of peroxisome proliferation in PPARalpha-deficient mouse liver. *J. Biol. Chem.* **276**(45):42485–42491.
- Zhou, S., and Wallace, K.B. (1999). The effect of peroxisome proliferators on mitochondrial bioenergetics. *Toxicol. Sci.* **48**(1):82–89.