

Research Article

Basis Set Effects in the Description of the Cl-O Bond in ClO and XClO/ClOX Isomers (X = H, O, and Cl) Using DFT and CCSD(T) Methods

Kenneth Irving, Martina Kieninger, and Oscar N. Ventura 

CCBG, DETEMA, Facultad de Química, Isidoro de María 1616, 11800 Montevideo, Uruguay

Correspondence should be addressed to Oscar N. Ventura; oscar.n.ventura@gmail.com

Received 1 October 2018; Accepted 16 January 2019; Published 26 February 2019

Academic Editor: Davut Avci

Copyright © 2019 Kenneth Irving et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The performance of a group of density functional methods of progressive complexity for the description of the ClO bond in a series of chlorine oxides was investigated. The simplest ClO radical species and the two isomeric structures XClO/ClOX for each X = H, Cl, and O were studied using the PW91, TPSS, B3LYP, PBE0, M06, M06-2X, BMK, and B2PLYP functionals. Geometry optimizations and reaction enthalpies and enthalpies of formation for each species were calculated using Pople basis sets and the (aug)-cc-pVnZ Dunning sets, with $n = D, T, Q, 5,$ and 6 . For the calculation of enthalpies of formation, atomization and isodesmic reactions were employed. Both the precision of the methods with respect to the increase of the basis sets, as well as their accuracy, were gauged by comparing the results with the more accurate CCSD(T) calculations, performed using the same basis sets as for the DFT methods. The results obtained employing composite chemical methods (G4, CBS-QB3, and W1BD) were also used for the comparisons, as well as the experimental results when they are available. The results obtained show that error compensation is the key for successful description of molecular properties (geometries and energies) by carefully selecting the method and basis sets. In general, expansion of the one-electron basis set to the limit of completeness does not improve results at the DFT level, but just the opposite. The enthalpies of formation calculated at the CCSD(T)/aug-cc-pV6Z for the species considered are generally in agreement with experimental determinations and the most accurate theoretical values. Different sources of error in the calculations are discussed in detail.

1. Introduction

Ozone loss over the Antarctic was reported by Farman et al. in 1985 [1]. Several hypotheses were proposed to explain the Antarctic ozone hole, of which the ones obtaining the largest success were those related to halogen-catalyzed ozone loss [2–4]. Ground-based measurements of ClO, HCl, ClONO₂, and OClO obtained in Antarctica during 1986 indicated that ozone depletion is associated with elevated abundances of ClO. A plethora of research on the behavior of chlorine and bromine compounds in the atmosphere was then started.

It is known today that stratospheric ozone is largely destroyed by radical reactions involving XO radicals (X = H, Cl, or Br). Self-reaction of these radicals, with the same or different X, gives rise to XYO₂ species (with and also H, Cl, or Br). Two of the most interesting species are HClO₂ and

Cl₂O₂ [5]. These species exhibit a similar behavior, with several possible stable isomers. Many theoretical studies have been performed, but it is not absolutely clear whether the most stable isomer computed is actually the observed one. For instance, it is well known that chlorous acid has the structure HOClO, at least in solution. However, the most stable isomer in the gas phase seems to be the HOCl peroxide. In the case of Cl₂O₂ instead, the most stable isomer seems to be ClClO₂ instead of the peroxide structure.

Theoretical studies are aimed fundamentally at determining thermochemical and kinetic data, which are essential for building models of the catalytic cycles. Calculating accurate data, which rival the experimental determinations when these latter are known, is not a trivial matter. Geometries, spectroscopic properties, and energetic information obtained depend heavily on the theoretical

methods employed and the one-electron basis sets used. Recent papers by Somers and Simmie [6] and by Rogers et al. [7] discuss some of the complexities associated with the study of enthalpies of formation, central to the formulation of atmospheric kinetic models.

In the case of composite methods, good accuracy is obtained through the addition of energies calculated at different theoretical levels using different basis sets [7]. In the case of direct calculations, high-level molecular orbital methods (CCSD(T) usually) or density functional methods (DFT) are combined with isogyric or isodesmic reactions [8, 9].

It is well known that molecular orbital methods depend heavily on the basis set chosen. It is less commonly recognized that this is also true in certain cases for DFT methods. In this work, we have performed a careful investigation of the effects of methods and basis sets on the description of the ClO bond in ClO and XClO ($X = \text{H, O, and Cl}$) species, using molecular orbital methods (MP2 and CCSD(T)), chemical models (the composite methods CBS-QB3, G4, and W1BD), and DFT methods, representing different rungs of Jacob's ladder (PW91PW, TPSS, B3LYP, PBE0, M06, M06-2X, BMK, and B2PLYP) [10, 11]. Our purpose is to compare the performance of DFT methods with that of molecular orbital methods, as well as their rate of convergence with increasingly extended basis sets. To this end, we have used Pople's basis sets and Dunning's correlation consistent basis sets from cc-pVDZ to aug-cc-pV6Z.

Results are reported for geometries, IR spectra, and energies of ClO, XClO, and XOCl, where $X = \text{H, O, or Cl}$. Whenever possible, the theoretical results have been compared with the experimental ones. In cases when these results are not available, a rational guess of the estimated values with their expected error bars is provided.

2. Methods

Geometry optimization and calculation of the energies were performed at the molecular orbital and DFT levels. MP2, CCSD(T) [12, 13], and composite models CBS-QB3 [14], G4 [15], and W1BD [16] were employed on the molecular orbital side. On the DFT side, methods belonging to different rungs in Jacob's ladder [17] were used. The methods chosen were the GGA PW91 [18] (for both exchange and correlation); the meta-GGA TPSS [19]; the hybrid GGAs PBE0 [20] and B3LYP [21]; the hybrid meta-GGAs M06 [22], M06-2X [22], and BMK [23]; and the double hybrid B2PLYP [24].

Two Pople basis sets, 6-31+G(d,p) and 6-311++G(2df,2pd), were chosen as representative of those commonly employed in theoretical chemistry calculations. Convergence of different properties was investigated using Dunning's cc-pVnZ and aug-cc-pVnZ, with $n = D, T, Q, 5$, and 6 (which includes up to i functions on the chlorine atom) (see [25, 26] and references therein).

Geometry optimization of all the species was fully performed for each and all models (method/basis set combination). Forces were reduced until a precision of at least 10^{-4} Å was obtained in the Cartesian coordinates.


Thermochemical properties were obtained within the harmonic oscillator/rigid rotor model. Analytical second derivatives were used whenever possible.

Calculations were performed using the Gaussian 09 set of computer programs [27] for the DFT and model chemistry calculations and the Molpro code [28, 29] for the post-Hartree-Fock calculations. A mixed cluster of Xeon and Opteron machines was employed for the calculations, with a maximum of 32 cores per job and 4 GB per core.


3. Results and Discussion

ClO is the simplest chlorine oxide possible. The experimental bond length was determined in 1968 by Amano et al. [30] as 1.5696 Å and more recently by Drouin et al. [31] as 1.569539 Å. The heat of formation at 298.15 K was measured experimentally as 24.29 ± 0.03 kcal/mol [32], while it is listed as 24.192 kcal/mol in the JANAF data tables [33]. Grant et al. [34] calculated this enthalpy of formation as 24.9 kcal/mol using the R/UCCSD(T) method [12, 13] with the augmented correlation consistent basis sets aug-cc-pVnZ ($n = D, T, Q$, and 5). The best theoretical value was reported by Karton et al. [35], who obtained 24.19 ± 0.19 kcal/mol at 0 K and 24.18 ± 0.03 at 298 K using the W4 method (see [36] and references therein).

In this paper, we calculated the enthalpy of formation employing both the atomization reaction



and the reaction of formation from the diatomics



The necessary experimental enthalpies of formation of the atoms in reaction (1) are $\Delta_f H^0(\text{Cl}, g) = 28.992 \pm 0.002$ kcal/mol and $\Delta_f H^0(\text{O}, g) = 59.555 \pm 0.024$ kcal/mol according to Cox et al. [37]. The standard enthalpies of formation of the diatomics in reaction (2) are zero by definition.

Different degrees of error compensation are expected in the determination of the enthalpy of formation using reactions (1) and (2). Errors associated with the representation of the triplet state oxygen atom or molecule within each method/basis set model will not be compensated neither in reaction (1) nor in (2) since no other triplet species is present. However, reaction (1) does present a doublet species in the right-hand side (*rhs*) and, therefore, would be expected to compensate for the errors associated to the description of the nonpaired single electron in ClO. Neither of these reactions are homodesmic or isogyric so that error compensation is expected to be low. Moreover, to obtain a reasonable estimate of the bonding energy of ClO, which allows accurate calculation of the enthalpy of formation from reaction (1), one should include in the CCSD method full calculation of triples and perturbation estimation of quadruples (CCSDT(Q)), correct for core correlation, introduce relativistic corrections, and account for anharmonicity. This is clearly not the goal of this work, a task already performed by the application of W4 methods, but just the

comparison between DFT and CCSD(T) methods to describe the ClO bond.

The calculated optimum bond length of ClO and the enthalpies of reaction for reactions (1) and (2) are deployed in Table 1. Experimental results for the enthalpies of reaction, obtained from the experimental enthalpies of formation of the species involved, are listed in the last entry of the table. The calculated enthalpies of formation at 298.15 K of ClO using the different models were obtained from the calculated enthalpies of reaction and the experimental enthalpies of formation of the species involved. These are also listed in the table, as well as the signed error with respect to the experimental result.

As we expected, the CCSD(T) values obtained with the large aug-cc-pV6Z are extremely accurate. The CCSD(T)/aug-cc-pV6Z model works well for both reactions, but the enthalpy of reaction derived from reaction (1), 25.00 kcal/mol, is slightly worse than that derived from reaction (2), 24.91 kcal/mol. Some of the errors incurred are similar in both reactions (core correlation or lack of quadruple excitations for instance), while others are clearly different (spin orbit corrections for the atoms, for instance). All of these errors are either too small or compensate so that the value obtained at the CCSD(T)/aug-cc-pV6Z level for the enthalpy of formation is only 0.7 kcal/mol above the experimental result.

Actually, it can be shown that the error compensation of post-CCSD(T) contributions is larger. With the addition of the complete triple, quadruple, and quintuple excitations contributions to the TAE (total atomization energy) plus the core correlation, relativistic and spin orbit contributions amount only to -0.01 kcal/mol (see Table 2 in [35]). The TAE calculated at 0 K in [35] using the W4 method is 64.51 kcal/mol, while the one we obtain at the CCSD(T)/aug-cc-pV6Z level is 63.27 kcal/mol. The difference between these values is coming from two sources. On the one side, Karton et al. extrapolate to the CBS limit, while we are using only the values obtained at the aug-cc-pV6Z level. However, as can be seen in their Table 3, the different energies are already converged below 0.10 kcal/mol when the 5Z basis set is used. This fact is also supported by the difference we observed in the TAE between the results using the cc-pV6Z and aug-cc-pV6Z basis sets, only 0.23 kcal/mol. There is a higher probability that the difference is coming from the slightly erroneous geometry at which the calculations to obtain the W4 results were performed. The geometry optimization done by Karton and coworkers was at the CCSD(T)/cc-pV(Q+d)Z level only, and the ClO bond length 1.575 Å is larger than the 1.5695 Å experimental value. Our own result, at the CCSD(T)/aug-cc-pV6Z level, is 1.5694 Å. This is an error unaccounted for in the W4 calculations.

Moreover, it must be noticed that the results obtained with CCSD(T) using Pople's basis sets are far off the mark and very unreliable.

The three chemical models used, CBS-QB3, G4, and W1BD, are much more accurate than the MP2/6-311++G(2df,2pd) or CCSD(T)/6-311++G(2df,2pd) models for the calculation of the enthalpies of formation. The W1BD calculation is in fact even more accurate than the

CCSD(T)/aug-cc-pV6Z result, at a fraction of the cost. In this particular case, W1BD outperforms G4, contrary to the more general situation reported by Somers [6]. The optimum geometries in each case are not so good, but the ad hoc corrections to the energies actually improve very much the accuracy of the derived enthalpy of formation.

The DFT methods perform somehow erratically. The B2PLYP/6-311++G(2df,2pd) model, for instance, performs worse than the M06/6-311++G(2df,2pd) model, notwithstanding that the former procedure should be better and is considerably more costly. This observation notwithstanding all DFT results obtained using reaction (2) are extremely good, even using Pople's basis sets. The largest error was observed for M06-2X and was only 2.4 kcal/mol. For the TAE, the results are similar, except that, in this case, neither the PW91 or TPSS methods with any basis set nor the other DFT methods with the smaller basis set give good results. It can be observed that the DFT method may or may not improve with the increase of the basis set, an aspect we will discuss further in the following.

To shed some more light on the behavior of the molecular orbital and DFT calculations, we performed an extended study in which the series of basis sets cc-pVnZ and aug-cc-pVnZ, with n from 2 to 6, were used to compute the optimum geometry of the ClO, Cl₂, and O₂ molecules, their enthalpies, and those of the O and Cl atoms. The results are best displayed in graphic form, showing the difference between the calculated and experimental values. They are displayed in Figures 1-3.

Figure 1 shows the signed error for the calculated bond lengths. The CCSD(T) values show a very nice convergence toward the experimental value. A fairly large basis set, namely, aug-cc-pV5Z, is needed to decrease the error to about 2×10^{-3} Å.

The PW91, TPSS, B3LYP, and BMK methods also converge to a limit value with the increase of the basis set. Only the latter, however, is able to produce errors similar to the CCSD(T) in the limit of the large basis set.

Contrary to the previous methods, the PBE0, M06, M06-2X, and B2PLYP methods do not converge toward the experimental value, but to a shorter bond length. In these cases, a combination of this potential with the aug-cc-pVTZ basis set is the best choice to get fairly accurate results at minimum cost. Purely from the point of view of obtaining the best bond length, the M06-2X/cc-pVTZ or M06/aug-cc-pVTZ models are the best options. If the CCSD(T) method was used for the optimizations with these same basis sets, the error would be ten times larger. Thus, it is clear that error compensation makes sufficiently complex DFT methods to have the advantage over the more theoretically rigorous CCSD(T) procedure when performing geometry optimizations. Let us see what happens in terms of energies.

When the enthalpy of formation of ClO is obtained from reaction (1) the results are those depicted in Figure 2. CCSD(T) converges monotonically toward the experimental value, while all the DFT methods converge toward a smaller enthalpy of formation. The results obtained with the PW91 and TPSS rule out these methods with any basis set. In fact, the results worsen systematically with the increase of the

TABLE 1: Enthalpy of reaction for reactions (1) and (2) and enthalpy of formation of ClO derived from these reactions using different models (in kcal/mol) and ClO bond length (in Å).

Method	Basis set	Cl + O \longrightarrow ClO			Cl ₂ + O ₂ \longrightarrow 2ClO			Bond length	
		$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)	δ^a	$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)	δ^a	r	δ^a
MP2	6-31G(d,p)	-46.42	42.13	17.84	65.22	32.61	8.3	1.6088	0.0392
	6-311++G(2df,2pd)	-56.99	30.59	6.44	68.76	34.38	10.2	1.5614	-0.0082
CCSD(T)	6-31G(d,p)	-48.13	39.45	15.30	47.24	23.62	-0.5	1.6287	0.0591
	6-311++G(2df,2pd)	-56.95	30.63	6.48	52.17	26.08	1.9	1.5885	0.0189
	aug-cc-pV6Z	-63.54	24.04	-0.11	49.82	24.91	0.8	1.5694	-0.0002
CBS-QB3		-63.47	24.11	-0.04	52.52	26.26	2.1	1.6102	0.0406
G4		-62.70	24.88	0.73	50.60	25.30	1.2	1.5925	0.0229
W1BD		-64.03	23.55	-0.60	49.61	24.80	0.7	1.5810	0.0114
PW91	6-31G(d,p)	-73.61	13.97	-10.18	49.76	24.88	0.7	1.5850	0.0154
	6-311++G(2df,2pd)	-79.68	7.90	-16.25	47.43	23.72	-0.4	1.5848	0.0152
	aug-cc-pV6Z	-81.37	6.21	-17.94	45.54	22.77	-1.4	1.5778	0.0082
TPSS	6-31G(d,p)	-64.48	23.10	-1.05	47.54	23.77	-0.4	1.5780	0.0084
	6-311++G(2df,2pd)	-70.30	17.28	-6.87	45.69	22.84	-1.3	1.5778	0.0082
	aug-cc-pV6Z	-71.71	15.87	-8.28	43.87	21.94	-2.2	1.5848	0.0152
B3LYP	6-31G(d,p)	-58.76	28.82	4.67	49.53	24.76	0.6	1.6195	0.0499
	6-311++G(2df,2pd)	-64.48	23.10	-1.05	47.87	23.94	-0.2	1.5844	0.0148
	aug-cc-pV6Z	-66.26	21.32	-2.83	45.98	22.99	-1.2	1.5736	0.0040
PBE0	6-31G(d,p)	-84.29	3.29	-20.86	51.77	25.88	1.7	1.5999	0.0303
	6-311++G(2df,2pd)	-65.58	22.00	-2.15	50.57	25.29	1.1	1.5665	-0.0031
	aug-cc-pV6Z	-67.38	20.20	-3.95	48.60	24.30	0.2	1.5563	-0.0133
M06	6-31G(d,p)	-58.66	28.92	4.77	51.26	25.63	1.5	1.5970	0.0274
	6-311++G(2df,2pd)	-64.26	23.32	-0.83	50.34	25.17	1.0	1.5661	-0.0035
	aug-cc-pV6Z	-65.54	22.04	-2.11	50.05	25.02	0.9	1.5563	-0.0133
M06-2X	6-31G(d,p)	-56.56	31.02	6.87	54.49	27.25	3.1	1.5967	0.0271
	6-311++G(2df,2pd)	-62.91	24.67	0.52	53.35	26.67	2.5	1.5633	-0.0063
	aug-cc-pV6Z	-64.38	23.20	-0.95	51.35	25.67	1.5	1.5545	-0.0151
BMK	6-31G(d,p)	-57.76	29.82	5.67	54.24	27.12	3.0	1.6199	0.0503
	6-311++G(2df,2pd)	-63.79	23.79	-0.36	53.25	26.62	2.5	1.5842	0.0146
	aug-cc-pV6Z	-65.36	22.22	-1.93	51.37	25.68	1.5	1.5693	-0.0003
B2PLYP	6-31G(d,p)	-54.20	33.38	9.23	52.43	26.21	2.1	1.6093	0.0397
	6-311++G(2df,2pd)	-61.61	25.97	1.82	52.52	26.26	2.1	1.5758	0.0062
	aug-cc-pV6Z							1.5631	-0.0065
		-63.43	24.29 ^b		48.30	24.29 ^b		1.5695 ^c	

^aDifference between the calculated value and the experimental value; ^b[32]; ^c[31].

basis set. The best results in this case are obtained with the M06/cc-pVQZ model. Similar results are obtained with the B3LYP/cc-pVTZ, PBE0/cc-pVTZ, and BMK/cc-pVQZ models although in these cases it is clear that the convergence of the models with the increase of the basis set is toward a much smaller enthalpy of formation. Both the M06-2X and the B2-PLYP method afford a good value with a relatively small basis set and converge to a value around 1 kcal/mol lower than the experimental one. B2PLYP calculations are, of course, much more demanding than M06-2X (although not so much as the CCSD(T) ones). Although larger, the cc-pVQZ basis set is comparable to the 6-311++G(2df,2pd) Pople set. The results obtained with the M06 methods are then comparable also (24.83 and 24.29 kcal/mol) and close to the experimental one.

If the enthalpy of formation is obtained from the enthalpy of reaction of the diatomics, the results are those depicted in Figure 3. The largest error of each model is now much lower than when the enthalpy of formation was obtained from the atomization energy, a fact probably due to the less than perfect description of the atoms provided by the

smaller basis sets. When the basis sets are large enough, the results are definitely better. For instance, the largest difference between theoretical calculations and the experimental value is -2.4 kcal/mol for the TPSS/aug-cc-pV6Z model, which is not too much worse than the 0.8 kcal/mol obtained at the much more demanding CCSD(T)/aug-cc-p6Z level and much better than the -7.4 kcal/mol error obtained when the enthalpy of formation was obtained at the same level from the TAE (reaction (1)).

In the case of reaction (2), there is a residual error in the calculation of the enthalpy of formation, due to the presence of the doublet radicals in the left-hand side (*lhs*) of the equation, while a triplet is present in the *rhs*. PW91, TPSS, B3LYP, and PBE0 exhibit some error compensation, and then models like PW91/aug-cc-pVTZ, TPSS/cc-pVTZ, B3LYP/cc-pVTZ, and PBE0/cc-pV5Z or larger give results very similar to the experimental ones. For the rest of the methods, only M06 and B2PLYP with basis sets cc-pV5Z (or larger) are able to give results with similar accuracy than the CCSD(T) calculations at the same basis set level. However, M06 using the cc-pVQZ basis set is able to give also an error

TABLE 2: Relative energies (in kcal/mol) of the isomers of ClO₂ and enthalpies of formation at 298K determined from reactions (16)–(18).

Species	Method	Basis set	$\Delta H(0)$	$\Delta_f H^0$, reac. (16a)	$\Delta_f H^0$, reac. (16b)	$\Delta_f H^0$, reac. (17a)	$\Delta_f H^0$, reac. (17b)	$\Delta_f H^0$, reac. (18)	Average
ClOO	G4		2.99	28.1	28.1	26.2	26.1	27.6	27.2 ± 1.8
	W1BD		4.50	28.3	29.1	27.2	27.7	27.8	28.1 ± 1.1
	B3LYP	Aug6Z	-1.65	24.4	24.2	22.6	24.6	20.4	23.2 ± 3.1
	PBE0	Aug6Z	2.41	24.6	26.4	23.2	25.5	19.9	23.9 ± 4.5
	M06	Aug6Z	3.03	23.3	25.9	23.8	24.3	22.4	23.9 ± 2.4
	M06-2X	Aug6Z	1.85	30.0	31.9	28.9	29.6	28.6	29.8 ± 2.3
	BMK	Aug6Z	-0.26	27.5	30.2	27.0	27.9	25.7	27.7 ± 2.9
	B2PLYP	Aug6Z	4.32	27.6	28.1	24.0	26.1	23.3	25.8 ± 3.8
	CCSD(T)	6Z	2.46	25.5	26.8	25.7	25.2	26.5	25.9 ± 1.2
	W4	CBS ¹	0.55						25.1
	Exper ²								23.4 ± 1.0
OCIO	G4			25.3	24.6	22.7	22.7	24.1	23.9 ± 2.1
	W1BD			23.9	24.1	22.7	22.7	22.8	23.2 ± 1.3
	B3LYP	Aug6Z		25.5	25.3	23.7	25.7	21.6	24.4 ± 3.1
	PBE0	Aug6Z		21.7	23.5	20.3	22.6	17.0	21.0 ± 4.5
	M06	Aug6Z		19.8	22.4	20.3	20.7	18.8	20.4 ± 2.4
	M06-2X	Aug6Z		27.9	29.8	26.8	27.5	26.5	27.7 ± 2.3
	BMK	Aug6Z		27.6	30.2	27.0	27.9	25.8	27.7 ± 2.9
	B2PLYP	Aug6Z		22.4	22.8	18.8	20.9	18.1	20.6 ± 3.8
	CCSD(T)	Aug6Z		22.4	23.7	22.6	22.2	23.1	22.8 ± 1.1
	W4	CBS ¹							23.9
	Exper. ³								22.6 ± 0.3
	Exper. ⁴								23.53 ± 0.24

¹[35]; ²[38]; ³[39]; ⁴[40].

TABLE 3: Standard enthalpies of formation of HClO (in kcal/mol) obtained from the reaction of atomization and reactions (3), (4), (7), and (8) are reported in the first five columns of the table. The energy difference between HClO and HOCl (without zero-point energies) is shown in the sixth column. The last column shows the average of the enthalpies of formation calculated using reactions of atomization, (4), (7), and (8). Values obtained with reaction (3) were not included in this average. The last file shows the averages, for each reaction, of the values of the enthalpy of reaction obtained with each method, excluding the PW91 and TPSS values.

	Reaction					ΔE	
	TAE	(4)	(8)	(7)	(3)	HClO-HOCl	Average
PW91	16.3	29.7	37.4	33.7	12.0	48.1	29.3
TPSS	26.6	33.3	34.6	34.4	3.5	48.6	32.2
B3LYP	33.1	34.2	32.1	33.8	6.5	51.7	33.3
PBE0	32.3	32.8	34.4	33.8	32.4	50.6	33.3
M06	31.9	32.5	35.8	33.3	19.5	51.8	33.4
M06-2X	33.7	34.2	30.1	33.1	20.4	54.3	32.8
BMK	34.9	36.3	31.6	35.8	23.1	55.0	34.6
B2PLYP	32.9	33.9	37.2	32.5	21.3	54.8	34.2
CBS-QB3	32.3	34.0	35.8	32.5	34.7	51.6	33.7
G4	33.9	33.6	34.7	31.8	34.8	51.6	33.5
W1BD		32.7	34.7	33.1	34.4	51.6	33.5
CCSD(T)	33.5	33.1	34.5	33.0	40.6	53.5	33.5
Average	33.2	33.7	34.1	33.3	26.8		

below 1.5 kcal/mol which is acceptable in terms of cost/benefit analysis.

Taking the three sets of results, the bond length and the enthalpy of formation were obtained from both reactions (1) and (2), and we may conclude that the M06/cc-pVQZ calculations give a reasonable description of the ClO

bond. The respective errors are -0.003 \AA , 0.5 kcal/mol, and 1.5 kcal/mol. Using the same basis set, the errors of the CCSD(T) calculations are 0.012 \AA , 4.4 kcal/mol, and 1.7 kcal/mol, with a much larger computing effort.

3.1. HClO/HOCl Isomers. The reaction of ClO with a hydrogen atom is the simplest possible route for the formation of a closed shell species. Two isomers can be formed depending on the oxidation state of chlorine: H-O-Cl and H-Cl=O. These isomers have been extensively studied, both theoretically and, at least for the first isomer, also experimentally.

Ashby in 1967 studied experimentally the IR spectrum of HOCl at high resolution [41]. They determined the bent geometry of this molecule as $r(\text{OH}) = 0.974 \pm 0.02 \text{ \AA}$, $r(\text{OCl}) = 1.689 \pm 0.006 \text{ \AA}$, and $\angle \text{HOCl} = 104.78$ degrees. Mirri et al. [42] a little later refined this structure studying the rotational spectra of several isotopic variants of HOCl. The geometric parameters found were $r(\text{OH}) = 0.975 \pm 0.003 \text{ \AA}$, $r(\text{OCl}) = 1.6895 \pm 0.0035 \text{ \AA}$, and $\angle \text{HOCl} = 102.48 \pm 0.45$ degrees. We chose to use the parameters selected by the authors themselves because a relatively large error in the $r(\text{OH})$ distance was noticed depending on which inertia defects were chosen for the derivation. A value as small as 0.960 \AA was reported for the $r(\text{OH})$ bond length. More recently, a detailed analysis of the vibration-rotation spectra of deuterated HOCl was performed by Deely [43]. The r_e structure determined yield the values $r(\text{OH}) = 0.9643 \pm 0.0005 \text{ \AA}$, $r(\text{OCl}) = 1.6891 \pm 0.0002 \text{ \AA}$, and $\angle \text{HOCl} = 102.96 \pm 0.08$ degrees. We will use these values to compare against the theoretically calculated ones.




FIGURE 1: Difference between the optimum bond length calculated with each theoretical model and the experimental value (in Å).




FIGURE 2: Difference between the enthalpy of formation of ClO at 298.15 K obtained from the reaction of atomization (1) and the experimental value (in kcal/mol).






FIGURE 3: Difference between the enthalpy of formation of ClO at 298.15 K obtained from the reaction of diatomic chlorine and oxygen, reaction (2), and the experimental value (in kcal/mol).

Experimental determination of the enthalpy of formation of HOCl has been performed several times. The values determined at 298 K, -17.8 kcal/mol [44, 45], -18.0 kcal/mol [46], -17.9 kcal/mol [47], and -18.0 ± 0.9 kcal/mol [48] are quite consistent. These determinations were obtained based on the equilibrium constant of the reaction:




The most recent determination at 0 K is that of Joens using thermochemical cycles [49], who obtained a value of 17.68 ± 0.03 kcal/mol. The net reaction used by Joens was



Several theoretical studies have been performed on this molecule. Hirsch et al. [50] performed MRDCI calculations with small basis sets on this species in 1977. They assumed a fixed geometry using the microwave-determined values of Mirri et al. [42] and calculated the enthalpy of formation of HClO among other properties. Depending on the reaction scheme used to calculate the enthalpy of formation, they obtained values ranging from -24.1 to -14.8 kcal/mol. Francisco and Sander [51] performed QCISD(T) calculations using the Pople 6-311++G(3df,3pd) basis set on top of a geometry optimized at a simpler level. This geometry reproduced reasonably well the H-O bond (0.963 \AA) but not the Cl-O bond which was about 0.05 \AA longer than the

experimental one. The enthalpy of formation was obtained using the following isodesmic reaction:



The best result obtained was -19.9 ± 2 kcal/mol, in reasonable agreement with the experimental determinations. Lee in 1994 [52] performed CCSD(T) geometry optimizations with a TZ2P basis set and CCSD(T) with ANO basis sets on top of those optimized geometries to obtain different properties of HClO and other related molecules. Again, the $r(\text{OH})$ distance was well represented, while the $r(\text{OCl})$ bond was too large. Denis in 2006 [53] estimated the enthalpy of formation of HOCl from the atomization reaction, employing CCSD(T) with correlation consistent basis sets and corrections for core-valence, relativistic, and anharmonic effects. The correlation consistent cc-pVQZ basis set was used for geometry optimizations, and extrapolation to the complete basis set was used for the energies. The enthalpy of formation of HOCl was calculated as -18.1 ± 0.3 kcal/mol which may be considered to be the most accurate value calculated up to date. This value is in very good agreement with the value calculated later by Karton et al. [35], -18.20 ± 0.14 kcal/mol, using essentially the same methodology. Finally, another work worth citing was published by Meyer and Kass in 2010 [54] who performed a comparison between the results obtained with several density functional methods and those obtained at the CCSD(T) level using correlation consistent basis sets up to

the cc-pV(Q+d)Z level. The best result obtained at the W4 level was -18.2 ± 0.1 kcal/mol (same, of course, as Karston et al. did at the same level), to be compared with M06-2X and M06 results of -17.7 and -16.5 kcal/mol using the same basis set.

There is not so detailed information about the less stable HClO isomer. No experimental values are available, neither for the geometry of this species nor for the enthalpy of formation. Theoretical studies do exist, starting with the aforementioned study of Lee [52], another one performed by Hernandez et al. in 1996 [55], and the most recent one by Peterson and coworkers [56] at the MRCI + Q/CBS level. The ClO bond is calculated as 0.13 \AA shorter in HClO than in HOCl, and both isomers are separated by a 73.5 kcal/mol barrier. At this level of calculation, the HClO isomer is 53.7 kcal/mol less stable than HOCl. A similar result was reported by Jalbout [57] who performed B3LYP and CCSD(T) calculations using Pople's 6-311++G(3df,3pd) basis set. He found activation barriers of 76.1 kcal/mol and 70.1 kcal/mol at each level, while the HClO isomer was found to lie between 54.5 kcal/mol and 52.5 kcal/mol, respectively, above the HOCl isomer.


The geometry optimization at the CCSD(T)/aug-cc-pV6Z performed in this paper for HOCl is the more extensive we are aware of. Even if the molecule is pretty small, the calculation of integrals with up to i Gaussian functions for oxygen and chlorine as well as h functions for hydrogen represents a sizeable challenge. Similar to the case of ClO, for which the calculated ClO distance was only 0.0001 \AA smaller than the experimental one, in this case, the difference is also very small. The geometric parameters obtained at the CCSD(T)/aug-cc-pV6Z level, $r(\text{OH}) = 0.9644 \text{ \AA}$, $r(\text{OCl}) = 1.6899 \text{ \AA}$, and $\angle \text{HOCl} = 102.95 \text{ deg}$ are in extraordinarily good agreement with Deely's experimental values $r(\text{OH}) = 0.9643 \pm 0.0005 \text{ \AA}$, $r(\text{OCl}) = 1.6891 \pm 0.0002 \text{ \AA}$, and $\angle \text{HOCl} = 102.96 \pm 0.08 \text{ deg}$ [43]. A graphical comparison of the errors in the estimation of the $r(\text{OCl})$ bond length with different DFT methods and the CCSD(T) method using the Dunning basis sets is depicted in Figure 4.

The optimum geometry calculated using the cc-pVQZ basis set at the CCSD(T) level, as done previously, affords an absolute error which is similar to that obtained at the B2PLYP or PBE0 levels and larger than that obtained using the BMK method. From the aspect of the graphs, it is clear that this is due to error compensation. As in the case of ClO, CCSD(T) calculation of geometries using a moderately large basis set like cc-pVQZ is not better than those obtained with the much cheaper DFT calculations. However, in this case, neither M06 nor M06-2X is able to give as good results as those obtained for the ClO radical.

As we explained before, there is no experimental value for the structure of HClO. Our best result at the CCSD(T)/aug-cc-pV6Z level is $r(\text{HCl}) = 1.3094 \text{ \AA}$, $r(\text{ClO}) = 1.5568 \text{ \AA}$, and $\angle \text{HOCl} = 107.75 \text{ degree}$. The parameters calculated by Peterson and coworkers at the MRCI + Q level with an aug-cc-pVQZ basis set are $r(\text{HCl}) = 1.3066 \text{ \AA}$, $r(\text{ClO}) = 1.5625 \text{ \AA}$, and $\angle \text{HOCl} = 107.81 \text{ degree}$. The most interesting result to be observed from these data concerns the bond order of the ClO bond in the ClO radical.



In Table 4, we show the optimum ClO bond lengths obtained at different levels of theory for the three species HClO, ClO, and HOCl. The ClO bond in HOCl is a single bond, while in HClO, we have a double bond. What is then the bond order in ClO? Two models were chosen. One was the simple linear correlation between bond order and bond length, while the second was Pauling's exponential correlation between the bond order and the difference of the observed bond length and that of a single bond. In the first case, the bond order derived from the CCSD(T)/aug-cc-pV6Z model (same results are obtained from DFT methods) is 1.89 , while in the second case is 1.87 . These congruent results imply that the bond order in ClO is much nearer to a double bond than to a single bond and the structure is more similar to $\bullet\text{Cl}=\text{O}$ than to $\text{Cl}-\text{O}\bullet$. It means that the solitary electron density distribution is located mainly on the chlorine atom and not the oxygen atom.

Several reactions can be used to calculate the enthalpy of formation of HClO and HOCl. Obviously the reaction of atomization, as well as reaction (4), can be used to obtain the enthalpy of formation of both isomers. Reaction (3) is isodesmic between closed shell molecules and can be used to calculate the enthalpy of formation of HOCl. Reaction (5) in turn is also isodesmic and isogyric (a doublet radical species is present in both the lhs and rhs of the equation). Combining reactions (3) and (5), one can obtain another reaction, which can be written as



Comparison of the enthalpies of formation obtained from reactions (5) and (6) with those obtained from reaction (3) will provide us with some indications about errors involved in the calculations.

For the case of HClO, we can also formulate a pair of other reactions which can be used to obtain the enthalpy of formation. These reactions are



Both for reactions (6) and (8), one needs to calculate two of the XOCl isomers ($X=\text{O}$ and Cl), which we will describe in detail later. For the moment, we will use their enthalpies to calculate the derived enthalpies of reaction we need in this section. As in the case of the other molecules, full geometry optimization at each level of calculation (method + basis set) was performed for H_2 , HO , H_2O , HCl , OCIO , and ClOCl , the additional species one needs to obtain the enthalpies of reactions (3)–(8).

The full table with the enthalpies of reaction and derived enthalpies of formation for HOCl and HClO is given in the Additional Material. In the following, we will analyze those numbers in two ways. On the one side, we are interested in knowing the enthalpies converge with the basis set, given a certain method, for all reactions. On the other side, we are




FIGURE 4: Signed error of the calculated $r(\text{OCl})$ bond distance with respect to the experimental one determined by Deely (see text (in Å)).

interested in knowing how do the results converge for all methods and basis sets in the case of each of the reactions.

A general overview of the results obtained for the enthalpy of formation of HOCl is shown in Figures 5 and 6. In Figure 5, we show the convergence with the increase of the basis set of the enthalpy of formation of HOCl extracted from each one of the reactions at the CCSD(T) level. Full lines show the results obtained from the reactions in which mostly closed shell species participate (the only exception is the triplet oxygen molecule). The results obtained from reaction (3) are slightly too negative, the best value being -18.80 kcal/mol which is within the error limits determined experimentally by Ennis and Birks, -18.0 ± 0.9 kcal/mol [48], using the equilibrium constant of the same reaction. As we explained before, this isodesmic reaction can be considered as the composing of reactions (5) and (6); the results are plotted in Figure 5 using dotted lines. It is clear that the error is larger in both cases, but to opposite directions, so that they compensate in the calculation derived from reaction (3).

The limit result obtained from reaction (4) is -18.45 kcal/mol. The errors with each basis set are similar to those of the determination from reaction (3). The less negative result for the enthalpy of formation is in agreement with the experimental determination of Joens [49] (the difference in the enthalpy of formation at 0 and 298 K is around 0.05 kcal/mol).

Finally, the dashed line represents the enthalpy of formation obtained from the TAE (without any of the corrections used by Denis [53], Karton et al. [35], or Meyer

and Kass [54] using the W4 method). Our best result is -18.15 kcal/mol, within the error limits of the values reported by the former authors, -18.1 ± 0.3 kcal/mol [53], -18.20 ± 0.14 kcal/mol [35], and -18.2 ± 0.1 kcal/mol [54]. It may be concluded then that, as was discussed in the case of ClO, no significant improvement is obtained in the determination of the enthalpy of reaction of HOCl using methods beyond CCSD(T) with an extended basis set both for geometries and energies.

Since all the reactions employed for obtaining the enthalpy of formation seem to exhibit some degree of error and error cancellation, one can aim at obtaining the enthalpy of formation as an average of the values derived from the different reactions. In Figure 6, there is a graphical representation of the convergence of the average, as well as maximum and minimum values, for the CCSD(T) calculations. On the one side, the values obtained from the five reactions considered were averaged. This is shown in blue. On the other side, only the values obtained from reactions (3) and (4) (red dotted lines) were averaged (red continuous line). The conclusion is that the average obtained from all the reactions, -18.05 kcal/mol, is smaller and more in agreement with the experimental results than that obtained from only reactions (3) and (4), -18.64 kcal/mol, but in all cases in reasonable agreement with previous accurate theoretical determinations. Figure S1 in the Supplementary Information shows an overview of the results obtained for all the reactions using different methods and basis sets.

A final analysis can be performed comparing the results obtained using the same methods and basis sets for HOCl

TABLE 4: Optimum ClO bond length (in Å) calculated at different theoretical levels for the three species HClO, ClO, and HOCl.

Species	Method	6-31G(d)	6-311++G(2df,2pd)	aug-cc-pV6Z
HClO	MP2	1.5961	1.5492	
	CCSD(T)	1.6493	1.5806	1.5568
	PW91PW91	1.6095	1.5720	1.5566
	TPSSTPSS	1.6200	1.5785	1.5626
	B3LYP	1.6262	1.5775	1.5594
	PBE0	1.5955	1.5539	1.5381
	M06	1.5944	1.5496	1.5310
	M06-2X	1.6117	1.5643	1.5501
	BMK	1.6350	1.5869	1.5646
	B2PLYP	1.6216	1.5715	1.5540
ClO	MP2	1.6088	1.5614	
	CCSD(T)	1.6287	1.5885	1.5694
	PW91PW91			1.5778
	TPSSTPSS			1.5848
	B3LYP	1.6195	1.5844	1.5736
	PBE0	1.5999	1.5665	1.5563
	M06	1.5970	1.5661	1.5563
	M06-2X	1.5967	1.5633	1.5545
	BMK	1.6199	1.5842	1.5693
	B2PLYP	1.6093	1.5758	1.5631
HOCl	MP2	1.7183	1.6946	
	CCSD(T)	1.7355	1.7072	1.6899
	PW91PW91	1.7384	1.7158	1.7088
	TPSSTPSS	1.7419	1.7197	1.7128
	B3LYP	1.7272	1.7053	1.6980
	PBE0	1.7010	1.6807	1.6739
	M06	1.6987	1.6785	1.6717
	M06-2X	1.6940	1.6765	1.6706
	BMK	1.7125	1.6917	1.6830
	B2PLYP	1.7235	1.7017	1.6924




FIGURE 5: Enthalpies of formation of HOCl (in kcal/mol) obtained at the CCSD(T) level using different basis sets and reactions (3)–(6).

and HOH. In Table 5 are shown the results obtained for the enthalpy of formation of HOH and HOCl using the CCSD(T) method with the series of Dunning basis sets,

obtained from the enthalpy of reaction of the atomization reaction, and the formation from the diatomics in each case.




FIGURE 6: Average values and maxima and minima of the enthalpy of formation of HOCl obtained from the five reactions considered (in red) or only from reactions (3) and (4) (in blue).

TABLE 5: Enthalpies of reaction and derived standard enthalpies of formation (in kcal/mol) for water and hypochlorous acid using different models.

Basis set	H ₂ O				HOCl			
	HOH \rightarrow 2H + O		HOH \rightarrow H ₂ + 1/2O ₂		HOCl \rightarrow H + Cl + O		2HOCl \rightarrow H ₂ + Cl ₂ + O ₂	
	$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)	$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)	$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)	$\Delta_r H^0$ (298K)	$\Delta_f H^0$ (298K)
cc-pVDZ	197.41	6.05	47.85	-8.15	132.90	7.75	23.09	-11.55
aug-cc-pVDZ	208.10	-4.64	56.56	-16.86	142.63	-1.98	36.66	-18.33
cc-pVTZ	213.61	-10.15	54.52	-14.82	150.00	-9.35	32.72	-16.36
aug-cc-pVTZ	216.91	-13.45	57.01	-17.31	153.20	-12.55	36.20	-18.10
cc-pVQZ	218.44	-14.98	56.84	-17.14	155.41	-14.76	35.46	-17.73
aug-cc-pVQZ	219.79	-16.33	57.71	-18.01	156.93	-16.28	36.83	-18.41
cc-pV5Z	220.09	-16.63	57.65	-17.95	157.81	-17.16	36.59	-18.30
aug-cc-pV5Z	220.49	-17.03	57.80	-18.11	158.43	-17.78	36.94	-18.47
cc-pV6Z	220.62	-17.16	57.78	-18.09	158.66	-18.01	36.81	-18.40
aug-cc-pV6Z	220.77	-17.31	57.81	-18.11		-18.15		-18.45

The results are depicted graphically in Figure 7, where the values for the enthalpy of formation of water has been shifted upward by the difference of the experimental enthalpies of formation of both species, so as to compare the curves more easily.

Several conclusions can be obtained from this comparison. On the one hand, the convergence of the values for the enthalpy of formation of HOCl with both reactions is more precise than that for H₂O. On the other hand, although the series converge to very similar values in each case, the values obtained with the participation of the diatomics is converged much faster than when the enthalpy of formation is obtained from the TAE. Using the cc-pVQZ basis set, the values obtained from reaction (4) are well converged, while those obtained from the TAE are still several kcal/mol off the mark. Finally, the agreement between the values obtained from both reactions show that, at least for closed shell molecules, it is unnecessary to include post-(T) effects, since triple, quadruple, and quintuple excitation contributions tend to cancel with the contributions from core correlation, relativistic, and spin-orbit corrections.

In the case of HClO, there are no experimental values to compare with. We will derive the enthalpy of formation using the TAE, the decomposition to diatomics (similar to equation (4) with HClO instead of HOCl), and the other two isodesmic reactions (7) and (8). Additionally, we will also use reaction (3) which is not isodesmic in the case of HClO. It will provide us with a measure of the suppression of errors produced by the isodesmicity.

We show in Table 3 the enthalpies of formation obtained from each of the reactions and the energy difference between the isomers, employing different methods and the largest basis set in each case.

The energy difference between the isomers, calculated at the CCSD(T)/cc-pV6Z level, 53.5 kcal/mol, is in complete agreement with that found by Peterson et al. [56]. Using the smaller basis set employed by Jalbout [57], the result is only 1 kcal/mol lower so that there is not a large influence of the basis set in this property. Except for the less precise PW91 and TPSS models, all the DFT and composite ab initio calculations give results in the 50–55 kcal/mol range. Thus, the barrier is not heavily dependent on the method used either.




FIGURE 7: Convergence of the CCSD(T) standard enthalpies of formation of H_2O and HOCl (in kcal/mol) obtained from the atomization reaction and the decomposition toward the diatomics. The values for water have been shifted upwards by the difference between the experimental enthalpies of formation 39.7 kcal/mol.

The result obtained averaging the values for the enthalpy of formation across methods and reactions is 33.6 ± 2.8 kcal/mol (error calculated as 2σ). Because of their obvious inaccuracy, we have left out of this average both the results obtained with the PW91 and TPSS methods and the results obtained from reaction (3). The lack of isodesmicity of this reaction for HClO implies that no error cancellation is occurring and the results are thus much worse than when the same reaction was used for calculating the enthalpy of formation of the HOCl isomer, for which (3) is truly isodesmic.

The behavior of different methods of calculation of the enthalpy of formation of HClO can be seen in Figure 8.


The most important observation is the very small dispersion of the values of the enthalpy of formation calculated with different chemical models from reaction (7). This is another indication that the unpaired electron in the ClO radical is located mainly on the chlorine atom, since the errors associated with the radical in the rhs of the reaction are clearly compensated by those associated with the description of the chlorine atom in the lhs.

3.2. Cl_2O Isomers. Hypochlorous acid, HOCl , is in equilibrium with its anhydride, Cl_2O , on the surface of ice crystals in the atmosphere [58], according to the already presented reaction (3). It is known to have C_{2v} symmetry, and the experimental geometry was determined by microwave spectroscopy by Tanaka et al. [59] as $r(\text{OCl}) = 1.69587(7)$ Å and $\angle\text{ClOCl} = 110.886(6)$ degrees. A second isomer, with the structure $\text{Cl}-\text{Cl}=\text{O}$, was observed spectroscopically already in 1967 by Rochkind and Pimentel [60] who determined two of the vibrational fundamentals ($962\text{--}952\text{ cm}^{-1}$ and $368\text{--}377\text{ cm}^{-1}$). The third one was determined by Chi and Andrews [61] in 1973 (241 cm^{-1}). The UV and IR spectra of this species were studied further by Johnsson et al. [62] in 1995, who proposed that the

formation occurred by geminal recombination of ClO and Cl , produced previously in the photolysis of the more stable isomer ClOCl . Later, a resonant Raman study of Cl_2O photochemistry in CCl_4 solution was performed by Esposito et al. in 1999 [63]. Their results confirmed that photo dissociation of Cl_2O produced ClO and Cl , followed by partial geminate recombination of those fragments, resulting in ClClO formation.

The potential energy surface and the interconversion of the isomers has been studied several times, for instance, by Collaveri et al. [64], Grant et al. [34], Xu and Lin [65], and An et al. [66]. The standard enthalpies of formation of both Cl_2O and ClClO at 298 K are available experimentally. Thorn et al. [67] determined a value of 18.44 ± 0.4 kcal/mol for ClOCl . Hassanzadeh and Irikura in 1997 revised the value of the enthalpy of formation of HOCl and obtained a slightly smaller value than the one used previously in this work, namely, 17.9 ± 0.3 kcal/mol. Using equation (3), they derived an experimental value of 19.4 ± 0.4 kcal/mol [68] and 1 kcal/mol above the value of Thorn et al. On the contrary, Chase reported the enthalpy of formation of ClClO as 22 ± 7 kcal/mol [33]. Using the usual composite methods in which CCSD(T) energies at lower levels are extrapolated to the complete basis set level and several corrections for missing contributions are added, Matus et al. [69] determined values of 18.9 kcal/mol and 31.9 kcal/mol for Cl_2O and ClClO , respectively.

For the calculation of the enthalpy of formation in this work, we will use the TAEs and the diatomic molecules as mentioned before



and the decompositions








FIGURE 8: Convergence of the values of the enthalpy of formation of HClO (in kcal/mol) for different reactions and models. The CCSD(T) and M06 results are shown in the upper two panels for each of the four reactions considered. The lower four panels show the convergence of all methods for each one of the four reactions.



For the ClOCl species, we will use also the isodesmic hydration reaction (3), employing now the experimental enthalpy of formation of HOCl. Finally, we will also use the addition reactions:



In the case of ClClO, reaction (3) is not isodesmic and may lead to large errors. Instead of that, we used another reaction:



Reaction (15) is isodesmic, but we do not have an experimental value for the enthalpy of formation of HClO. We will use our best CCSD(T) value instead, 33.6 kcal/mol.

As we did previously with the other species, the error in the Cl-O bond length of the calculations at different levels with respect to the experimental value is shown in Figure 9. The best value obtained at the CCSD(T)/aug-cc-pV6Z level differs in only 6×10^{-4} Å from the experimental one, similar to the error obtained in HClO (8×10^{-4} Å). With respect to the performance of the different methods, there is no noticeable difference between the results obtained for HOCl and ClOCl. Thus, the same considerations done previously apply in this case.

Similarly to what we did for HOCl, we show in Figure 10 the CCSD(T) calculations of the enthalpy of formation of Cl₂O as obtained from the different reactions using the series of correlation consistent basis sets. Full lines show the convergence of the values obtained from reactions (3) and (10) which involve only closed shell species. The value obtained from reaction (10) at the CCSD(T)/aug-cc-pV6Z level is 18.45 kcal/mol, in excellent agreement with the experimental value of Thorn et al. [67]. On the contrary, the value obtained from the isodesmic reaction (3) at the same level is 19.17 kcal/mol, nearer to the value of Hassanzadeh and Irikura [68].

The dashed lines in Figure 10 represent the convergence of the values obtained from reactions (9) and (11)–(14) which involve open shell species. The value obtained from the atomization reaction is 18.57 kcal/mol, also in agreement with Thorn et al. [67], and lower than the theoretical value of Matus et al. [69]. All the other reactions involving radicals give values near the experimental value of Thorn et al. Averaging the results of all the reactions (the black dotted line in Figure 10), we find a value for $\Delta_f H_{298}^0(\text{Cl}_2\text{O})$ of 18.7 ± 0.5 kcal/mol within the error limits of both experimental determinations and in agreement to the theoretically derived value of Matus et al. [69].

The results obtained with the largest basis set used for all methods and reactions are collected in Table 6. It is clear that no DFT method can provide the same accuracy as CCSD(T)

when employing a very large basis set. Of all the methods listed, M06-2X is the one that gives more similar results, in agreement with what we discussed previously. In the same line of thought as before, a combination of the M06-2X method with a not so large basis set (cc-pVQZ) gives similar or better results than a CCSD(T) calculation with the same basis set. This can be appreciated, in the case of Cl₂O in the error distribution presented in the panels of Figure S2 in the Supplementary Information.

In the case of ClClO, we have no experimental data, except the rough estimation made by Chase $\Delta_f H_{298}^0(\text{ClClO}) = 22 \pm 7$ kcal/mol [33]. Therefore, we will assume that the CCSD(T) values have a similar accuracy than that shown in the case of ClOCl. The convergence of the enthalpy of formation obtained from reactions (9)–(15) using the different correlation consistent basis sets is shown in Figure 11. Reasonable convergence of all values is obtained for basis sets larger than cc-pV6Z, and an average value of 32.1 ± 0.4 kcal/mol is obtained as a result, in agreement with the recommendation of Matus et al. [69].

The enthalpies of formation obtained with the largest basis set are shown in Table 7. Similarly to the case of ClOCl, none of the DFT methods exhibits such a small dispersion as the CCSD(T) results. Moreover, the M06-2X method in this case, although exhibiting also the smallest dispersion of the DFT results, is not able to give a reasonably accurate result neither with the aug-cc-pV6Z nor at the simpler cc-pVQZ level.

3.3. ClO₂ Isomers. The last species studied in this paper are the radical isomers chlorine dioxide, OClO, and chlorine superoxide, ClOO. The most stable structure and the most important one from a practical point of view is the C_{2v} isomer OClO. This species is extensively used in pulp bleaching and in decontamination of drinkable water. The C_s chlorine superoxide, on the other side, presents some intriguing properties that make it a challenge for theoretical studies.

Very precise studies of the geometrical structure of these species exist. Chlorine dioxide has been experimentally known since the nineteenth century and well studied. Tanoura et al. [70] studied the ground state rotational transitions of ³⁵ClO₂ and ³⁷ClO₂, while Miyazaki et al. [71] studied the microwave spectra of ³⁵ClO₂ and ³⁷ClO₂ in the excited vibrational states, ν_1 , ν_2 , ν_3 , and $2\nu_2$, and determined the equilibrium structure of the OClO radical as $r_e = 1.46984 \pm 0.00001$ Å and $\angle\text{OClO} = 117.403 \pm 0.003$ degrees. The infrared and rotational spectra of this species—as well as that of the isomer—were studied for instance by Müller and Wilmer in 1993 [72] and Müller et al. [73]. Theoretically, this species was studied several times, by Luke [74], Zhu and Lin [75], Karton et al. [36], Peterson and Werner [76, 77], and Grant et al. [34], among others. References to the latest experimental and theoretical work can be found in the recent work by Grein [78].

On the contrary, chlorine superoxide has been a more elusive species. The existence of this radical was suggested as






FIGURE 9: Signed error of the calculated $r(\text{ClO})$ bond length with respect to the experimental value (in Å) for each level of calculation.

a reactive intermediate in the gas-phase photolysis reactions of Cl_2 with O_2 by Porter and Wright in the fifties [79]. Arkell and Schwager [40] in 1967 were the first to isolate the species in an Ar solid matrix and to determine its IR spectrum. In this paper, there is already present some speculation about an intermediate ClOO^* structure which would be responsible of some lines in the O-O stretching region. A theoretical discussion of the isomerization $\text{ClOO} \leftrightarrow \text{OCLO}$, in solid matrices, implying the presence of an excited state $^2A'$ —the ground state is $^2A''$ —was presented by Gole in 1980 on the basis of Hartree-Fock calculations [80]. Müller and Wilner [72] compared the spectra of both isomers in Ne and Ar matrices, obtaining the superoxide by irradiation of the dioxide with visible light. Experimental geometric parameters for ClOO were determined as $r_e(\text{ClO}) = 2.139 \text{ \AA}$, $r_e(\text{OO}) = 1.201 \text{ \AA}$, and $\angle \text{ClOO} = 115.7$ degrees. Later, experimental work was performed by Johnsson et al. [81], who also observed ClOO^* as a different species than ClOO , and by Thomsen et al. [82]. The most recent experimental work on this radical was performed by Fourier transform infrared spectroscopy and FT microwave-millimeter wave double resonance spectroscopy by Suma et al. [83], who found the parameters $r_e(\text{ClO}) = 2.084 \pm 0.001 \text{ \AA}$, $r_e(\text{OO}) = 1.206 \pm 0.002 \text{ \AA}$, and $\angle \text{ClOO} = 115.4 \pm 0.1$ degrees.

From the thermochemical point of view, the most interesting topic is the relative stability of both isomers. This can be studied comparing directly their absolute energies or determining their respective enthalpies of formation using these reactions:



In fact, while the asymmetrical chlorine superoxide seems to be thermodynamically more stable than chlorine dioxide, it is kinetically unstable, and the study of this radical by high resolution spectroscopy in the gas phase has been currently unsuccessful, due probably to the decomposition as in reaction (16). Baer et al. [84] studied the thermodynamic and kinetic properties of reaction (16) and determined that the bond dissociation energy was only $4.83 \pm 0.05 \text{ kcal/mol}$. In contrast, the bond energy of the Cl-O bond in chlorine dioxide was found to be much higher, $55.2 \pm 2.0 \text{ kcal/mol}$ [85].

Other thermochemical property which we can obtain from reactions (16)–(18) are the enthalpies of formation of OCLO and ClOO . Experimentally, the enthalpy of formation of OCLO has been reported as $22.6 \pm 0.3 \text{ kcal/mol}$ by Sander et al. [39] and as 23.53 ± 0.24 by Arkell and Schwager [40]. For ClOO , Wahner et al. [38] determined a value of $23.4 \pm 1.0 \text{ kcal/mol}$. The most recent and precise calculation of these values, done by Karton et al. [35] at the W4 level [36], yielded values of $\Delta H_f^0(298 \text{ K}) = 23.9 \text{ kcal/mol}$ for OCLO and 25.1 kcal/mol for ClOO .




FIGURE 10: CCSD(T) calculation of the enthalpy of formation of Cl_2O (in kcal/mol) using different basis sets and reactions. Full lines refer to reactions between closed-shell molecules. Dashed lines are the results for reactions involving radicals. The dotted black line is the average for all the reactions for each basis set.

TABLE 6: Enthalpies of formation of ClOCl (in kcal/mol) obtained from reactions (3) and (9)–(14) using the indicated method and the aug-cc-pV6Z basis set. The average value of those obtained for all the reactions and the standard deviation are shown in the last two columns of the table.

Method	(3)	(10)	(9)	(11)	(12)	(13)	(14)	Average	Standard deviation
PW91	17.92	17.10	-2.00	15.79	5.21	33.58	23.00	15.80	11.57
TPSS	18.65	17.99	11.65	19.77	14.35	27.89	22.48	18.97	5.30
B3LYP	19.33	20.55	21.45	24.12	18.46	26.80	21.14	21.69	2.88
PBE0	19.95	21.87	18.24	22.04	19.44	25.84	23.24	21.54	2.57
M06	19.18	20.81	17.55	19.51	20.23	21.47	22.19	20.13	1.55
M06-2X	20.04	19.42	17.41	18.21	18.60	19.01	19.41	18.87	0.88
BMK	20.09	22.29	18.50	20.27	21.29	22.05	23.06	21.08	1.56
B2PLYP	18.90	20.10	19.51	20.70	17.86	21.90	19.06	19.72	1.32
CCSD(T)	19.17	18.45	18.57	18.53	18.73	18.49	18.69	18.66	0.24




FIGURE 11: CCSD(T) calculation of the enthalpy of formation of ClClO (in kcal/mol) using different basis sets and reactions. Full lines refer to reactions between closed-shell molecules. Dashed lines are the results for reactions involving radicals. The dotted black line is the average for all the reactions for each basis set.

TABLE 7: Enthalpies of formation of ClClO (in kcal/mol) obtained from reactions (3) and (9)–(14) using the indicated method and the aug-cc-pV6Z basis set. The average value of those obtained for all the reactions and the standard deviation are shown in the table.

Method	(9)	(10)	(11)	(12)	(13)	(14)	(15)	Average	Standard deviation
PW91	3.81	22.91	21.60	11.03	39.39	28.82	28.26	22.26	11.83
TPSS	19.58	25.92	27.70	22.28	35.83	30.41	29.15	27.27	5.36
B3LYP	32.71	31.81	35.38	29.72	38.06	32.40	31.80	33.12	2.74
PBE0	28.23	31.86	32.03	29.43	35.83	33.23	32.60	31.89	2.49
M06	27.00	30.26	28.96	29.68	30.92	31.64	31.50	29.99	1.64
M06-2X/a6	34.26	36.27	35.06	35.46	35.87	36.26	36.45	35.66	0.79
M06-2X/Q	36.72	38.38	36.71	37.15	36.70	37.14	35.88	36.95	0.76
BMK	32.46	36.25	34.23	35.25	36.01	37.02	34.53	35.11	1.52
B2PLYP	31.23	31.82	32.42	29.58	33.62	30.78	31.72	31.60	1.27
CCSD(T)	32.01	31.90	3.97	32.17	31.94	32.13	32.46	32.08	0.19




FIGURE 12: Total energies (in Hartree) for the isomers CIOO and OCIO calculated at the CCSD(T) level employing cc and aug-cc basis sets. The inset shows an enlargement of the region of results with the larger basis sets.

It is known that nondynamical correlation is an important factor in the determination of the structure of these species [35]. Therefore, in this work, we will discuss only the best results obtained for the bond energies and enthalpies of formation. A deeper analysis will be published elsewhere.

We show in Figure 12 the convergence of the total energies (in Hartree) of the optimum geometries of CIOO and OCIO at the CCSD(T) level with the correlation consistent and augmented correlation consistent basis sets. Two effects are immediately obvious. One is that both cc and aug-cc sets do converge slowly toward the CBS (complete basis set) limit. The second is that, at the CCSD(T) level, OCIO is less stable than CIOO, but the situation is reversed for large basis sets. This conclusion is reached also when the DFT methods are used (Figure 13). The convergence of the free energy difference with all methods shows a similar pattern, but two of them (B3LYP and BMK) predict CIOO to be more stable than OCIO even with the larger basis sets. All the other methods, as well as CCSD(T), predict the inverse result, with a free energy difference between 1 and 5 kcal/mol (1.8 kcal/mol at the CCSD(T)/CBS level using Feller's extrapolation method [86]).

We have collected in Table 2 the enthalpy differences between the isomers as well as the enthalpy of formation at room temperature obtained with all the methods employed in this paper. Only the results obtained with the larger basis sets are reported.

Several observations can be done on these data. On the one side, the agreement between the CCSD(T) results and the more recent experimental determinations of the enthalpy of formation is better for OCIO than for CIOO. This is related to the difficulty in describing the Cl-O bond using monoconfigurational MO approaches. On the other side, one sees that the values obtained at the CCSD(T) level with

the larger basis sets are very much in agreement with the W4 method which includes many more corrections and is a much more demanding procedure. It is also observed that neither for OCIO nor for CIOO are the DFT methods able to give a result as accurate as the CCSD(T) method with the largest basis sets. We have estimated the precision of the calculations as $2 * \sigma$, where σ is the standard deviation from the arithmetic mean of the results obtained for the enthalpy of formation from the five reactions (16)–(18). Both in terms of precision and accuracy, the CCSD(T) results are the only ones comparable to experiment. Notice that since the W4 uses atomization energies, the difference of enthalpies of formation between the isomers (1.2 kcal/mol) is about half the experimental one (2.5 kcal/mol). On the contrary, the CCSD(T)/cc-pV6Z results obtained as done in this work (i.e., geometry optimization performed at the same level than the calculations) are much nearer to the experimental result (3.1 kcal/mol).

As we have observed in the other cases, DFT methods with smaller basis sets provide better results than when a large basis is used. In Figure 14, we present the results for the series of basis sets and different functionals, as compared to the results obtained at the CCSD(T) level using the same basis sets.

We observe again that the error in some of the DFT calculations change sign with the increase of the basis set. Thus, there is an error compensation between the potential and the basis set although this is not the case for all methods. One important point, however, is that when a small basis set is used (anything smaller than aug-cc-pV5Z) almost all DFT calculations give a smaller error than the CCSD(T) calculations. In all cases, the error is unacceptably large (more than 5 kcal/mol), but it is larger with CCSD(T) methods.




FIGURE 13: Free energy difference (in kcal/mol) between the ClOO and OClO isomers at their optimum geometries, calculated with several methods using cc and aug-cc basis sets.




FIGURE 14: Unsigned error (with respect to the experimental value) in the enthalpy of formation of OClO at 298.15 K (in kcal/mol) as compared to the CCSD(T) calculations with the cc and aug-cc basis sets.

With a carefully selected choice of method and basis sets (M06 or B2PLYP with the cc-pV5Z basis set, for instance), the error is less than 1 kcal/mol, an error obtainable at the

CCSD(T) level only if basis cc-pV6Z or aug-cc-pV6Z are used. Needless to say, the latter calculations are much slower and more demanding than the DFT ones.

4. Conclusions

MP2, CCSD(T), density functional, and composite chemical methods were employed in this paper to study the ClO radical species as well as the two isomeric structures XClO/ClOX for each $X = \text{H}, \text{Cl}, \text{and O}$.

The main purpose of the study was to compare the performance of several DFT methods (PW91, TPSS, B3LYP, PBE0, M06, M06-2X, BMK, and B2PLYP) to describe the Cl-O bond in radicals of importance in atmospheric chemistry. Geometry optimizations as well as reaction enthalpies and enthalpies of formation for each species were calculated using Pople basis sets and the (aug)-cc-pVnZ Dunning sets, with $n = 2-6$. For the calculation of enthalpies of formation, atomization and isodesmic reactions were employed.

Both the precision of the methods with respect to the increase of the basis sets, as well as their accuracy, were estimated by comparing the results with the more accurate CCSD(T) calculations performed using the same basis sets as for the DFT methods. The results obtained employing composite chemical methods (G4, CBS-QB3, and W1BD) were also used for the comparisons, as well as experimental data when they are available.

The results obtained show that error compensation is the key for successful description of molecular properties (geometries and energies) by carefully selecting the method and the basis set. In general, expansion of the one-electron basis set to the limit of completeness does not improve results at the DFT level—as it occurs with precise post-Hartree-Fock MO methods—but just the opposite. The enthalpies of formation calculated at the CCSD(T)/aug-cc-pV6Z for the species considered are generally in agreement with experimental determinations and the most accurate calculated values. In the case of the DFT methods, we found that a reasonable description of the Cl-O bond across the species studies can be obtained at M06/aug-cc-pVQZ or M06/cc-pV5Z. Other methods can occasionally produce better results but are not generally as accurate in all cases.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Disclosure

Part of the data contained in this study were obtained from the Thesis of Kenneth Irving. Some of the calculations reported in this paper were performed in ClusterUY, a newly installed platform for high-performance scientific computing at the National Supercomputing Center, Uruguay.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We thank the permanent financial support of ANII, CSIC, and Pedeciba for the completion of this work.

Supplementary Materials

The Supplementary Information contains two figures, showing comparative charts of the calculated enthalpies of formation of HOCl and Cl₂O using different reactions, methods, and basis sets to obtain them. (*Supplementary Materials*)





References

- [1] J. C. Farman, B. G. Gardiner, and J. D. Shanklin, "Large losses of total ozone in Antarctica reveal seasonal ClO_x/NO_x interaction," *Nature*, vol. 315, no. 6016, pp. 207–210, 1985.
- [2] S. Solomon, R. R. Garcia, F. S. Rowland, and D. J. Wuebbles, "On the depletion of Antarctic ozone," *Nature*, vol. 321, no. 6072, pp. 755–758, 1986.
- [3] M. B. McElroy, R. J. Salawitch, S. C. Wofsy, and J. A. Logan, "Reductions of Antarctic ozone due to synergistic interactions of chlorine and bromine," *Nature*, vol. 321, no. 6072, pp. 759–762, 1986.
- [4] L. T. Molina and M. J. Molina, "Production of chlorine oxide (Cl₂O₂) from the self-reaction of the chlorine oxide (ClO) radical," *Journal of Physical Chemistry*, vol. 91, no. 2, pp. 433–436, 1987.
- [5] M. von Hobe, R. J. Salawitch, T. Canty et al., "Understanding the kinetics of the ClO dimer cycle," *Atmospheric Chemistry and Physics*, vol. 7, no. 12, pp. 3055–3069, 2007.
- [6] K. P. Somers and J. M. Simmie, "Benchmarking Compound Methods (CBS-QB3, CBS-APNO, G3, G4, W1BD) against the active thermochemical tables: formation enthalpies of radicals," *Journal of Physical Chemistry A*, vol. 119, no. 33, pp. 8922–8933, 2015.
- [7] D. W. Rogers, A. A. Zavitsas, and N. Matsunaga, "Determination of enthalpies ("Heats") of formation," *Wiley Interdisciplinary Reviews: Computational Molecular Science*, vol. 3, no. 1, pp. 21–36, 2013.
- [8] W. J. Hehre, R. Ditchfield, L. Radom, and J. A. Pople, "Molecular orbital theory of the electronic structure of organic compounds. V. Molecular theory of bond separation," *Journal of the American Chemical Society*, vol. 92, no. 16, pp. 4796–4801, 1970.
- [9] S. E. Wheeler, K. N. Houk, P. V. R. Schleyer, and W. D. Allen, "A hierarchy of homodesmotic reactions for thermochemistry," *Journal of the American Chemical Society*, vol. 131, no. 7, pp. 2547–2560, 2009.
- [10] J. P. Perdew, A. Ruzsinszky, L. A. Constantin, J. Sun, and G. I. Csonka, "Some fundamental issues in ground-state density functional theory: a guide for the perplexed," *Journal of Chemical Theory and Computation*, vol. 5, no. 4, pp. 902–908, 2009.
- [11] R. O. Jones, "Density functional theory: its origins, rise to prominence, and future," *Reviews of Modern Physics*, vol. 87, no. 3, pp. 897–923, 2015.
- [12] P. J. Knowles, C. Hampel, and H. J. Werner, "Coupled cluster theory for high spin, open shell reference wave functions," *Journal of Chemical Physics*, vol. 99, no. 7, pp. 5219–5227, 1993.
- [13] M. J. O. Deegan and P. J. Knowles, "Perturbative corrections to account for triple excitations in closed and open shell coupled cluster theories," *Chemical Physics Letters*, vol. 227, no. 3, pp. 321–326, 1994.
- [14] J. A. Montgomery, M. J. Frisch, J. W. Ochterski, and G. A. Petersson, "A complete basis set model chemistry. VII. Use of the minimum population localization method,"

- Journal of Chemical Physics*, vol. 112, no. 15, pp. 6532–6542, 2000.
- [15] L. A. Curtiss, P. C. Redfern, and K. Raghavachari, “Gaussian-4 theory,” *Journal of Chemical Physics*, vol. 126, no. 8, article 084108, 2007.
- [16] E. C. Barnes, G. A. Petersson, J. A. Montgomery, M. J. Frisch, and J. M. L. Martin, “Unrestricted coupled cluster And brueckner doubles variations of W1 theory,” *Journal of Chemical Theory and Computation*, vol. 5, no. 10, pp. 2687–2693, 2009.
- [17] J. P. Perdew and K. Schmidt, “Jacob’s ladder of density functional approximations for the exchange-correlation energy,” *AIP Proceedings*, vol. 577, no. 1, p. 1, 2001.
- [18] J. P. Perdew, *de Electronic Structure of Solids ’91*, P. Z. A. H. Eschrig, Ed., Akademie Verlag, Berlin, Germany, 1991.
- [19] J. M. Tao, J. P. Perdew, V. N. Staroverov, and G. E. Scuseria, “Climbing the density functional ladder: nonempirical meta-generalized gradient approximation designed for molecules and solids,” *Physical Review Letters*, vol. 91, no. 14, article 146401, 2003.
- [20] C. Adamo and V. Barone, “Toward reliable density functional methods without adjustable parameters: the PBE0 model,” *Journal of Chemical Physics*, vol. 110, no. 13, pp. 6158–6170, 1999.
- [21] A. D. Becke, “Density-functional thermochemistry. III. The role of exact exchange,” *Journal of Chemical Physics*, vol. 98, no. 7, pp. 5648–5652, 1993.
- [22] Y. Zhao and D. G. Truhlar, “The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals,” *Theoretical Chemistry Accounts*, vol. 120, no. 1–3, pp. 215–241, 2008.
- [23] A. D. Boese and J. M. L. Martin, “Development of density functionals for thermochemical kinetics,” *Journal of Chemical Physics*, vol. 121, no. 8, pp. 3405–3416, 2004.
- [24] S. Grimme, “Semiempirical hybrid density functional with perturbative second-order correlation,” *Journal of Chemical Physics*, vol. 124, no. 3, article 034108, 2006.
- [25] K. A. Peterson, D. E. Woon, and J. T. H. Dunning, “Benchmark calculations with correlated molecular wave functions. IV. The classical barrier height of the $H+H_2 \rightarrow H_2+H$ reaction,” *Journal of Chemical Physics*, vol. 100, no. 10, pp. 7410–7415, 2004.
- [26] A. K. Wilson, T. van Mourik, and T. H. Dunning Jr, “Gaussian Basis Sets for use in Correlated Molecular Calculations. VI. Sextuple zeta correlation consistent basis sets for boron through neon,” *Journal of Molecular Structure: Theochem*, vol. 388, no. 1–3, pp. 339–349, 1996.
- [27] M. J. Frisch, G. W. Trucks, and H. B. Schlegel, *Gaussian 09, Revision D.01*, Gaussian, Inc., Wallingford, CT, USA, 2009.
- [28] H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, and M. Schütz, “Molpro: a general-purpose quantum chemistry program package,” *Wiley Interdisciplinary Reviews: Computational Molecular Science*, vol. 2, no. 2, pp. 242–253, 2012.
- [29] H.-J. Werner, P. J. Knowles, G. Knizia, and F. R. Manby, MOLPRO, version 2012.1, A Package of ab Initio Programs, <http://www.molpro.net>, 2012.
- [30] T. Amano, E. Hirota, and Y. Morino, “Microwave spectrum of the ClO radical,” *Journal of Molecular Spectroscopy*, vol. 27, no. 1–4, pp. 257–265, 1968.
- [31] B. J. Drouin, C. E. Miller, E. A. Cohen, G. Wagner, and M. Birk, “Further investigations of the ClO rotational spectrum,” *Journal of Molecular Spectroscopy*, vol. 207, no. 1, pp. 4–9, 2001.
- [32] S. Abramowitz and M. W. Chase, “Thermodynamic properties of gas phase species of importance to ozone depletion,” *Pure and Applied Chemistry*, vol. 63, no. 10, pp. 1449–1454, 1991.
- [33] M. W. Chase Jr, “NIST-JANAF thermochemical tables,” in *Journal of Physical and Chemical Reference Data, Monograph*, Vol. 9, pp. 1–1951, National Institute of Standards and Technology, Gaithersburg, MA, USA, 4th edition, 1998.
- [34] D. J. Grant, E. B. Garner III, M. H. Matus et al., “Thermodynamic properties of the XO_2 , X_2O , XYO , X_2O_2 , and XYO_2 ($X, Y = Cl, Br, \text{ and } I$) isomers,” *Journal of Physical Chemistry A*, vol. 114, no. 12, pp. 4254–4265, 2010.
- [35] A. Karton, S. Parthiban, and J. M. L. Martin, “Post-CCSD(T) ab initio thermochemistry of halogen oxides and related hydrides XOX , $XOOX$, HOX , XOn , and $HXOn$ ($X = F, Cl$), and evaluation of DFT methods for these systems†,” *Journal of Physical Chemistry A*, vol. 113, no. 16, pp. 4802–4816, 2009.
- [36] A. Karton, E. Rabinovich, J. M. L. Martin, and B. Ruscic, “W4 theory for computational thermochemistry: in pursuit of confident sub-kJ/mol predictions,” *Journal of Chemical Physics*, vol. 125, no. 14, article 144108, 2006.
- [37] J. Cox, D. Wagman, and V. Medvedev, *CODATA Key Values for Thermodynamics*, Hemisphere Publishing Corporation, New York, NY, USA, 1984.
- [38] A. Wahner, G. S. Tyndall, and A. R. Ravishankara, “Absorption cross sections for symmetric chlorine dioxide as a function of temperature in the wavelength range 240–480 nm,” *Journal of Physical Chemistry*, vol. 91, no. 11, pp. 2734–2738, 1987.
- [39] S. P. Sander, R. R. Friedl, A. R. Ravishankara et al., *Chemical Kinetics and Photochemical Data for use in Atmospheric Studies: Evaluation Number 14*, California: National Aeronautics and Space Administration, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA, 2003.
- [40] A. Arkell and I. Schwager, “Matrix infrared study of the chlorine superoxide radical,” *Journal of the American Chemical Society*, vol. 89, no. 24, pp. 5999–6006, 1967.
- [41] R. A. Ashby, “The infrared spectrum of HOCl,” *Journal of Molecular Spectroscopy*, vol. 23, no. 4, pp. 439–447, 1967.
- [42] A. M. Mirri, F. Scappini, and G. Cazzoli, “Microwave spectrum, structure and electric quadrupole coupling constants of HOCl,” *Journal of Molecular Spectroscopy*, vol. 38, no. 2, pp. 218–227, 1971.
- [43] C. M. Deely, “Vibration-rotation spectra of deuterated hypochlorous acid and the determination of the equilibrium structure,” *Journal of Molecular Spectroscopy*, vol. 122, pp. 481–489, 1987.
- [44] J. M. W. Chase, C. A. D. J. J. R. Davies, F. D. J. R. McDonald, and A. N. Syverud, “NIST-JANAF thermochemical tables, suppl. 1,” *Journal of Physical Chemistry Reference Data*, vol. 14, pp. 1–1856, 1985.
- [45] H. D. Knauth, H. Alberti, and H. Clausen, “Equilibrium constant of the gas reaction chlorine oxide (Cl_2O) + water = hypochlorous acid ($2HOCl$) and the ultraviolet spectrum of hypochlorous acid,” *Journal of Physical Chemistry*, vol. 83, no. 12, pp. 1604–1612, 1979.
- [46] L. T. Molina and M. J. Molina, “Ultraviolet spectrum of hypochlorous acid,” *Journal of Physical Chemistry*, vol. 82, no. 22, pp. 2410–2414, 1978.
- [47] H. Niki, P. D. Maker, C. M. Savage, and L. P. Breitenbach, “Fourier transform IR spectroscopic determination of the equilibrium constant for $H_2O + Cl_2O = 2HOCl$,” *Chemical Physics Letters*, vol. 66, no. 2, pp. 325–328, 1979.

- [48] C. A. Ennis and J. W. Birks, "Applications of a new laboratory source of gaseous hypochlorous acid (HOCl): product distribution in the atomic chlorine + hypochlorous acid (HOCl) reaction and equilibrium constant for the reaction chlorine oxide (Cl_2O) + water = 2 hypochlorous acid (2HOCl)," *Journal of Physical Chemistry*, vol. 89, no. 1, pp. 186–191, 1985.
- [49] J. A. Joens, "The dissociation energy of $\text{OH}(\text{X}^2\Pi^{3/2})$ and the enthalpy of formation of $\text{OH}(\text{X}^2\Pi^{3/2})$, ClOH , and BrOH from thermochemical cycles," *Journal of Physical Chemistry A*, vol. 105, no. 49, pp. 11041–11044, 2001.
- [50] G. Hirsch, P. J. Bruna, S. D. Peyerimhoff, and R. J. Buenker, "Ab initio CI study of the stability and electronic spectrum of the HOCl molecule," *Chemical Physics Letters*, vol. 52, no. 3, pp. 442–448, 1977.
- [51] J. S. Francisco and S. P. Sander, "Structure and thermochemistry of hydrochlorous acid, HOCl," *The Journal of Chemical Physics*, vol. 99, no. 8, pp. 6219–6220, 1993.
- [52] T. J. Lee, "A Comparative Coupled-Cluster Study of the XOCl and XClO ($\text{X} = \text{H}, \text{F}, \text{Cl}$) isomers: an investigation of hypervalent chlorine compounds," *Journal of Physical Chemistry*, vol. 98, no. 14, pp. 3697–3700, 1994.
- [53] P. A. Denis, "Thermochemistry of the hypobromous and hypochlorous acids, HOBr and HOCl," *The Journal of Physical Chemistry A*, vol. 110, no. 17, pp. 5887–5892, 2006.
- [54] M. M. Meyer and S. R. Kass, "Experimental and theoretical gas-phase acidities, bond dissociation energies, and heats of formation of HClO_x , $x = 1-4$," *Journal of Physical Chemistry A*, vol. 114, no. 12, pp. 4086–4092, 2010.
- [55] M. L. Hernandez, C. Redondo, A. Laganà, G. Ochoa de Aspuru, M. Rosi, and A. Sgamellotti, "Anabinitiostudy of the $\text{O}(\text{1D})+\text{HCl}$ reaction," *Journal of Chemical Physics*, vol. 105, no. 7, pp. 2710–2718, 1996.
- [56] K. A. Peterson, S. Skokov, and J. M. Bowman, "A theoretical study of the vibrational energy spectrum of the HOCl/HClO system on an accurate ab initio potential energy surface," *Journal of Chemical Physics*, vol. 111, no. 16, pp. 7446–7456, 1999.
- [57] A. F. Jalbout, "A comparative coupled cluster and density functional theory study of the $\text{HOCl} \rightarrow \text{HClO}$ transition state," *Molecular Physics*, vol. 100, no. 24, pp. 3785–3788, 2002.
- [58] J. B. Burkholder, "Ultraviolet absorption spectrum of HOCl," *Journal of Geophysical Research: Atmospheres*, vol. 98, no. D2, pp. 2963–2974, 1993.
- [59] M. Nakata, M. Sugie, H. Takeo, C. Matsumura, T. Fukuyama, and K. Kuchitsu, "Structure of dichlorine monoxide as studied by microwave spectroscopy. determination of equilibrium structure by a modified mass dependence method," *Journal of Molecular Spectroscopy*, vol. 86, no. 1, pp. 241–249, 1981.
- [60] M. M. Rochkind and G. C. Pimentel, "Photolysis of matrix-isolated dichlorine monoxide: infrared spectra of ClClO and $(\text{ClO})_2$," *Journal of Chemical Physics*, vol. 46, no. 11, pp. 4481–4490, 1967.
- [61] F. K. Chi and L. Andrews, "Argon matrix Raman spectra of Cl_2O and its photolysis products ClO and ClClO . Infrared matrix spectra of ClO and $(\text{ClO})_2$," *Journal of Physical Chemistry*, vol. 77, no. 26, pp. 3062–3067, 1973.
- [62] K. Johnsson, A. Engdahl, and B. Nelander, "The UV and IR spectra of the ClClO molecule," *Journal of Physical Chemistry*, vol. 99, no. 12, pp. 3965–3968, 1995.
- [63] A. P. Esposito, P. J. Reid, and K. W. Rousslang, "A resonance Raman study of Cl_2O photochemistry in solution: evidence for ClClO formation," *Journal of Photochemistry and Photobiology A: Chemistry*, vol. 129, no. 1-2, pp. 9–15, 1999.
- [64] C. Collaveri, G. Granucci, M. Persico, and A. Toniolo, "Theoretical study of the photochemistry of Cl_2O ," *Journal of Chemical Physics*, vol. 115, no. 3, pp. 1251–1263, 2001.
- [65] Z. F. Xu and M. C. Lin, "Ab initio chemical kinetic study on $\text{Cl}+\text{ClO}$ and related reverse processes," *Journal of Physical Chemistry A*, vol. 114, no. 43, pp. 11477–11482, 2010.
- [66] H.-S. An, K.-H. Lee, and K.-K. Baeck, "Ab initio study of the transition state of direct isomerization from ClOCl to ClClO on the S_0 and S_1 State," *Bulletin of the Korean Chemical Society*, vol. 32, no. 2, pp. 701–704, 2011.
- [67] R. P. Thorn Jr., L. J. Stief, S.-C. Kuo, and R. B. Klemm, "Ionization energy of Cl_2O and ClO , appearance energy of ClO^+ (Cl_2O), and heat of formation of Cl_2O ," *Journal of Physical Chemistry*, vol. 100, no. 33, pp. 14178–14183, 1996.
- [68] P. Hassanzadeh and K. K. Irikura, "Nearly ab initio thermochemistry: the use of reaction schemes. application to IO and HOI ," *Journal of Physical Chemistry A*, vol. 101, no. 8, pp. 1580–1587, 1997.
- [69] M. H. Matus, M. T. Nguyen, D. A. Dixon, K. A. Peterson, and J. S. Francisco, " ClClO_2 Is the most stable isomer of Cl_2O_2 . Accurate coupled cluster energetics and electronic spectra of Cl_2O_2 isomers," *Journal of Physical Chemistry A*, vol. 112, no. 40, pp. 9623–9627, 2008.
- [70] M. Tanoura, K. Chiba, K. Tanaka, and T. Tanaka, "Microwave spectroscopy of chlorine dioxide," *Journal of Molecular Spectroscopy*, vol. 95, no. 1, pp. 157–181, 1982.
- [71] K. Miyazaki, M. Tanoura, K. Tanaka, and T. Tanaka, "Microwave spectrum of chlorine dioxide in excited vibrational states," *Journal of Molecular Spectroscopy*, vol. 116, no. 2, pp. 435–449, 1986.
- [72] H. Müller and H. Willner, "Vibrational and electronic spectra of chlorine dioxide, OClO , and chlorine superoxide ClOO , isolated in cryogenic matrices," *Journal of Physical Chemistry*, vol. 97, no. 41, pp. 10589–10598, 1993.
- [73] H. S. P. Müller, G. O. Sørensen, and M. Birk, "The rotational spectrum and anharmonic force field of chlorine dioxide, OClO ," *Journal of Molecular Spectroscopy*, vol. 186, no. 1, pp. 177–188, 1997.
- [74] B. T. Friedl, "A theoretical investigation of chlorine-oxygen compounds. Part 1. G_2 and $\text{G}_2(\text{MP}2)$ energies of ClO , Cl_2O , ClO_2 and Cl_2O_2 ," *Journal of Molecular Structure: THEOCHEM*, vol. 332, no. 3, pp. 283–289, 1995.
- [75] R. S. Zhu and M. C. Lin, "Ab initio studies of ClO_x reactions. VIII. Isomerization and decomposition of ClO_2 radicals and related bimolecular processes," *Journal of Chemical Physics*, vol. 119, no. 4, pp. 2075–2082, 2003.
- [76] K. A. Peterson and H. J. Werner, "The photodissociation of ClO_2 : potential energy surfaces of $\text{OClO} \rightarrow \text{Cl} + \text{O}_2$," *Journal of Chemical Physics*, vol. 105, no. 22, pp. 9823–9832, 1996.
- [77] K. A. Peterson and H. J. Werner, "Multireference configuration interaction calculations of the low-lying electronic states of ClO_2 ," *Journal of Chemical Physics*, vol. 96, no. 12, pp. 8948–8961, 1992.
- [78] F. Grein, "Multireference configuration interaction studies on higher valence and Rydberg states of OClO , ionization potentials, and electron detachment energies," *Journal of Chemical Physics*, vol. 135, no. 4, p. 044304, 2011.
- [79] G. Porter and F. J. Wright, "Studies of free radical reactivity by the methods of flash photolysis. The photochemical reaction between chlorine and oxygen," *Discussions of the Faraday Society*, vol. 14, pp. 23–34, 1953.
- [80] J. L. Gole, "Photochemical isomerization of chlorine dioxide and the low-lying electronic state of the asymmetric isomer. Possible implications for matrix isolation spectroscopy,"

- Journal of Physical Chemistry*, vol. 84, no. 11, pp. 1333–1340, 1980.
- [81] K. Johnsson, A. Engdahl, and B. Nelander, “A matrix isolation study of the chlorine superoxide radical,” *Journal of Physical Chemistry*, vol. 97, no. 38, pp. 9603–9606, 1993.
- [82] C. L. Thomsen, P. J. Reid, and S. R. Keiding, “Quantum yield for ClOO formation following photolysis of aqueous OClO,” *Journal of the American Chemical Society*, vol. 122, no. 51, pp. 12795–12801, 2000.
- [83] K. Suma, Y. Sumiyoshi, and Y. Endo, “Fourier transform microwave spectroscopy and Fourier transform microwave-millimeter wave double resonance spectroscopy of the ClOO radical,” *Journal of Chemical Physics*, vol. 121, no. 17, pp. 8351–8359, 2004.
- [84] S. Baer, H. Hippler, R. Rahn, M. Siefke, N. Seitzinger, and J. Troe, “Thermodynamic and kinetic properties of the reaction $\text{Cl} + \text{O}_2 + \text{M} = \text{ClOO} + \text{M}$ in the range 160–300 K and 1–1000 bar,” *Journal of Chemical Physics*, vol. 95, no. 9, pp. 6463–6470, 1991.
- [85] I. P. Fischer, “Mass spectrometry study of intermediates in thermal decomposition of perchloric acid and chlorine dioxide,” *Transactions of the Faraday Society*, vol. 63, pp. 684–692, 1967.
- [86] D. Feller, “Benchmarks of improved complete basis set extrapolation schemes designed for standard CCSD(T) atomization energies,” *Journal of Chemical Physics*, vol. 138, no. 7, article 074103, 2013.



Hindawi

Submit your manuscripts at
www.hindawi.com

